

---

Teleinformatics Engineering Department, Federal University of Ceará

# Introduction to Gaussian Processes

Filipe P. de Farias, IC  
filipepfarias@fisica.ufc.br

October 24, 2018

The **Gaussian Processes** are the widely used stochastic processes for modeling dependent data observed over time, space or even time and space. Here, we'll initiate our study with a **Probability and Random Process Theory Review** taking some point to base our journey, going through **Linear Regression** and finally the **Gaussian Processes**.

The material here presented isn't sufficient to guide you over basic probability, so it's recommended to have some knowledge, once we'll just take a simple review.

## 1 Probability and Random Process Theory Review

- 1.1 Basic Concepts of Probability Theory
- 1.2 The Gaussian distribution
- 1.3 Independence of two random variables

## 2 Linear Regression

- 2.1 Basic Concepts of Curve Fitting
- 2.2 Bayesian Curve Fitting

# Probability and Random Process Theory Review

A key concept in the field of pattern recognition is that of **uncertainty**, that arises from both through noise on measurements, as well as through the finite size of data sets. Let's begin from a simple example.




Figure: Considering in the table  $X = x_i$  and  $Y = y_j$

Let's choose a cell in the 6. We define the probability of choose a cell in a given column is  $p(X = x_i) = c_i/N$ , being  $N$  the total number of cells and  $c_i = \sum_j n_{ij}$ .




Figure: Considering in the table  $X = x_i$  and  $Y = y_j$

Let's choose a cell in the 6. We define the probability of choose a cell in a given column is  $p(X = x_i) = c_i/N$ , being  $N$  the total number of cells and  $c_i = \sum_j n_{ij}$ .

We could say that the probability of choose a cell in a given a row is defined as  $p(Y = y_j|X = x_i) = n_{ij}/c_i$ .




Figure: Considering in the table  $X = x_i$  and  $Y = y_j$



Let's choose a cell in the 6. We define the probability of choose a cell in a given column is  $p(X = x_i) = c_i/N$ , being  $N$  the total number of cells and  $c_i = \sum_j n_{ij}$ .

We could say that the probability of choose a cell in a given a row is defined as  $p(Y = y_j|X = x_i) = n_{ij}/c_i$ .

And so, the probability of choose a cell is defined as  $p(X = x_i, Y = y_j) = n_{ij}/N$ .




Figure: Considering in the table  $X = x_i$  and  $Y = y_j$

Here, we could see some properties that we call **The Rules of Probability**

## The Rules of Probability

- Sum Rule:

$$p(X = x_i) = \sum_j n_{ij}/N \quad (1)$$

- Product Rule:

$$p(Y = y_j, X = x_i) = n_{ij}/N \quad (2)$$

Here, we could see some properties that we call **The Rules of Probability**

## The Rules of Probability

- Sum Rule:

$$p(X = x_i) = \sum_j n_{ij}/N \Rightarrow p(X) = \sum_Y p(X, Y) \quad (1)$$

- Product Rule:

$$p(Y = y_j, X = x_i) = n_{ij}/N \quad (2)$$

Here, we could see some properties that we call **The Rules of Probability**

## The Rules of Probability

- Sum Rule:

$$p(X = x_i) = \sum_j n_{ij}/N \Rightarrow p(X) = \sum_Y p(X, Y) \quad (1)$$

- Product Rule:

$$p(Y = y_j, X = x_i) = n_{ij}/N = n_{ij}/c_i \cdot c_i/N \quad (2)$$

Here, we could see some properties that we call **The Rules of Probability**

## The Rules of Probability

- Sum Rule:

$$p(X = x_i) = \sum_j n_{ij}/N \Rightarrow p(X) = \sum_Y p(X, Y) \quad (1)$$

- Product Rule:

$$p(Y = y_j, X = x_i) = n_{ij}/N = n_{ij}/c_i \cdot c_i/N \Rightarrow p(X, Y) = p(Y|X)p(X) \quad (2)$$

And by the **Product Rule** we prove that

## Bayes' Rule

$$p(X, Y) = p(Y|X)p(X) = p(X|Y)p(Y) \Rightarrow p(Y|X) = \frac{p(X|Y)p(Y)}{p(X)} \quad (3)$$

The **Gaussian distribution** is defined as

$$\mathcal{N}(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left\{ -\frac{1}{2\sigma^2} (x - \mu)^2 \right\} \quad (4)$$

## Sentence

**X and Y are independent random variables** if *any* event  $A_1$  defined in terms of X is independent of *any* event  $A_2$  defined in terms of Y



## Sentence

**X and Y are independent random variables** if *any* event  $A_1$  defined in terms of X is independent of *any* event  $A_2$  defined in terms of Y

The sentence above is equivalent to say mathematically that

## Sentence

**X and Y are independent random variables** if *any* event  $A_1$  defined in terms of  $X$  is independent of *any* event  $A_2$  defined in terms of  $Y$

The sentence above is equivalent to say mathematically that

$$P[X \text{ in } A_1, Y \text{ in } A_2] = P[X \text{ in } A_1]P[Y \text{ in } A_2] \quad (5)$$

## Sentence

**X and Y are independent random variables** if *any* event  $A_1$  defined in terms of X is independent of *any* event  $A_2$  defined in terms of Y

The sentence above is equivalent to say mathematically that

$$P[X \text{ in } A_1, Y \text{ in } A_2] = P[X \text{ in } A_1]P[Y \text{ in } A_2] \quad (5)$$

that means in other words that *if X and Y are independent discrete random variables, then the **joint probability mass function (pmf)** is equal to the product of the marginal pmf's.*

# Linear Regression

So, we'll start to look the regression with a statistical approach. To encourage you, let's take the sentence.

So, we'll start to look the regression with a statistical approach. To encourage you, let's take the sentence.

## Sentence

*If we could update the **regression weights** as we acquire some new values of the experiment?*

Let's take a look again at the Bayes Theorem

Let's take a look again at the Bayes Theorem

## Bayes Theorem

$$p(\mathbf{w}|\mathcal{D}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w})}^{\text{the weights probability}} \overbrace{p(\mathbf{w})}^{\text{the data probability}}}{\underbrace{p(\mathcal{D})}_{\text{the data probability}}} \quad (6)$$



Let's take a look again at the Bayes Theorem

## Bayes Theorem

$$p(\mathbf{w}|\mathcal{D}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w})}^{\text{the weights probability}} \overbrace{p(\mathbf{w})}^{\text{the data probability}}}{\underbrace{p(\mathcal{D})}_{\text{the data probability}}} \quad (6)$$

So, if **we have the probability** of the data, we'll could estimate the **future weights**.

Let's take a look again at the Bayes Theorem

## Bayes Theorem

$$p(\mathbf{w}|\mathcal{D}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w})}^{\text{the weights probability}} \overbrace{p(\mathbf{w})}^{\text{the weights prior}}}{\underbrace{p(\mathcal{D})}_{\text{the data probability}}} \quad (6)$$

So, if we have the probability of the data, we'll could estimate the future weights.  
**But, how?**

Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.

Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.

Now we'll try to view the same problem with a *probabilistic perspective*. We're trying to make predictions for the target value  $\mathbf{t}$  given some new values of  $x$ .

Taking some steps back, let's re-visit the **Curve Fitting**. There, the strategy was minimize the error function.

Now we'll try to view the same problem with a *probabilistic perspective*. We're trying to make predictions for the target value  $\mathbf{t}$  given some new values of  $x$ .

A good idea is to express our target values  $\mathbf{t}$  in terms of **gaussians distributions** with the mean equals to  $y(x, \mathbf{w})$ .




Figure: Schematic of the polynomial function  $y(x, \mathbf{w})$  and the gaussian distribution  $p$ .

By the 15, we assume the relation

(8)

(9)

By the 15, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1}) \quad (7)$$

(8)

(9)



By the 15, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1}) \quad (7)$$

and then, assume that the training data  $\{\mathbf{x}, \mathbf{t}\}$  is independent and identically distributed (i.i.d.) and put on **product form**, i.e. the joint probability is

$$p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = \mathcal{N}(t_0|y(x_1, \mathbf{w}), \beta^{-1}) \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1}) \dots \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1}) \quad (8)$$

(9)

By the 15, we assume the relation

$$p(t|x, \mathbf{w}, \beta) = \mathcal{N}(t|y(x, \mathbf{w}), \beta^{-1}) \quad (7)$$

and then, assume that the training data  $\{\mathbf{x}, \mathbf{t}\}$  is independent and identically distributed (i.i.d.) and put on **product form**, i.e. the joint probability is

$$p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = \mathcal{N}(t_0|y(x_1, \mathbf{w}), \beta^{-1}) \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1}) \dots \cap \mathcal{N}(t_n|y(x_0, \mathbf{w}), \beta^{-1}) \quad (8)$$

$$= \prod_{n=1}^N \mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1}) \quad (9)$$

regarding that  $\beta^{-1} = \sigma^2$ .

And we'll have a function to maximize if we apply the logarithm function to  $p$ , so

And we'll have a function to maximize if we apply the logarithm function to  $p$ , so

$$\ln(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = \sum_{n=1}^N \ln\left(\mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1})\right) \quad (10)$$

And we'll have a function to maximize if we apply the logarithm function to  $p$ , so

$$\ln(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = \sum_{n=1}^N \ln\left(\mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1})\right) \quad (10)$$

Applying the **Gaussian distribution** (see 4) will result

$$\ln(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = -\frac{\beta}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{N}{2} \ln(\beta) - \frac{N}{2} \ln(2\pi) \quad (11)$$

And taking the derivatives with respect to  $\beta$  to minimize the error

(12)

(14)

And taking the derivatives with respect to  $\beta$  to minimize the error

$$\frac{\partial}{\partial \beta} \ln (p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = 0 \quad (12)$$

(14)

And taking the derivatives with respect to  $\beta$  to minimize the error

$$\frac{\partial}{\partial \beta} \ln(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = 0 \quad (12)$$

$$-\frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w} - t_n)\}^2 + \frac{N}{2} \frac{1}{\beta} = 0 \quad (13)$$

$$(14)$$



And taking the derivatives with respect to  $\beta$  to minimize the error

$$\frac{\partial}{\partial \beta} \ln(p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)) = 0 \quad (12)$$

$$-\frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w} - t_n)\}^2 + \frac{N}{2} \frac{1}{\beta} = 0 \quad (13)$$

$$\frac{1}{N} \sum_{n=1}^N \{y(x_n, \mathbf{w} - t_n)\}^2 = \frac{1}{\beta_{ML}} \quad (14)$$

Where  $\beta_{ML}$  is the maximum likelihood.