

FYS4411 - COMPUTATIONAL QUANTUM MECHANICS

SPRING 2016

Project 1; Variational Monte Carlo Studies of Bosonic systems

TEMPORARY REPORT

Sean Bruce Sangolt Miller

s.b.s.miller@fys.uio.no

Filip Henrik Lasren

filiphenriklarsen@gmail.com

Date: February 23, 2016

Abstract

Some text that is abstact

Contents

1	Introduction	1
2	Theory and methods	1
2.1	Preliminary derivations	1
2.1.1	Simplified problem	1
2.1.2	Full problem	2
3	Results	4
4	Conclusions	4
5	Appendix	4

1 Introduction

2 Theory and methods

2.1 Preliminary derivations

2.1.1 Simplified problem

The local energy is defined as:

$$E_L(\mathbf{R}) = \frac{1}{\Psi_T(\mathbf{R})} H \Psi_T(\mathbf{R}), \quad (1)$$

As a first approximation, it is assumed there is no interaction term in the Hamiltonian, which means the hard sphere bosons have no physical size (the hard-core diameter is zero). It is also assumed that no magnetic field is applied to the bosonic gas, leaving a perfectly spherically symmetrical harmonic trap. Inserting this new Hamiltonian into the local energy gives:

$$E_L(\mathbf{R}) = \frac{1}{\Psi_T(\mathbf{R})} \sum_i^N \left(\frac{-\hbar^2}{2m} \nabla_i^2 + \frac{1}{2} m \omega_{ho}^2 r_i^2 \right) \Psi_T(\mathbf{R}) \quad (2)$$

The potential term is trivial since this is a scalar, i.e. the denominator will cancel the wavefunction. A more challenging problem is to find an expression for $\nabla_i^2 \Psi_T(\mathbf{R})$. The trial wavefunction shown in equation (...), with the aforementioned approximations, is:

$$\Psi_T(\mathbf{R}) = \prod_i e^{-\alpha r_i^2} \quad (3)$$

where α is the variational parameter for VCM. The first derivative is:

$$\nabla_j \prod_i e^{-\alpha r_i^2} = -2\alpha \mathbf{r}_j e^{-\alpha r_j^2} \prod_{i \neq j} e^{-\alpha r_i^2} \quad (4)$$

$$= -2\alpha \mathbf{r}_j \prod_i e^{-\alpha r_i^2}. \quad (5)$$

The second derivative then follows:

$$\nabla_j^2 \prod_i e^{-\alpha r_i^2} = \nabla_j \left(-2\alpha \mathbf{r}_j \prod_i e^{-\alpha r_i^2} \right) \quad (6)$$

$$= (4\alpha^2 r_j^2 - 2d\alpha) \prod_i e^{-\alpha r_i^2}. \quad (7)$$

where d is the number of dimensions. Inserting this into back into the local energy (equation (2)), the final expression can be derived:

$$\begin{aligned} E_L(\mathbf{R}) &= \frac{1}{\Psi_T(\mathbf{R})} \sum_i^N \left(\frac{-\hbar^2}{2m} \nabla_i^2 + \frac{1}{2} m \omega_{ho}^2 r_i^2 \right) \Psi_T(\mathbf{R}) \\ &= \sum_{i=1}^N \left[\frac{-\hbar^2}{2m} (4\alpha^2 r_i^2 - 2d\alpha) + \frac{1}{2} m \omega_{ho}^2 r_i^2 \right] \end{aligned}$$

The drift force (quantum force), still with the approximations above, is defined by:

$$F = \frac{2\nabla \Psi_T}{\Psi_T} \quad (8)$$

The gradient here is defined as

$$\nabla \equiv (\nabla_1, \nabla_2, \dots, \nabla_N)$$

i.e. a vector of dimension Nd . The gradient with respect to a single particle's position is already given in equation 5, so it's not too hard to realise:

$$\begin{aligned}
F &= \frac{-4\alpha}{\Psi_T} (\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) \Psi_T \\
&= -4\alpha (\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)
\end{aligned}$$

2.1.2 Full problem

The full local energy¹ is a bit more tedious to derive. The first step is to rewrite the trial wavefunction to the following form:

$$\Psi_T(\mathbf{R}) = \prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \quad (9)$$

where, in order for this to fit with the previous wavefunction, $u(r_{ij}) \equiv \ln[f(r_{ij})]$ and $\phi(\mathbf{r}_i) \equiv g(\alpha, \beta, \mathbf{r}_i)$. The gradient with respect to the k -th coordinate set is:

$$\nabla_k \Psi_T = \nabla_k \prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \quad (10)$$

$$= \nabla_k \phi_k \left[\prod_{i \neq k} \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \right] + \left[\prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \nabla_k \left(\sum_{i'' < j''} u_{i''j''} \right) \right] \quad (11)$$

$$= \nabla_k \phi_k \left[\prod_{i \neq k} \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \right] + \left[\prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \left(\sum_{i'' < j''} \nabla_k u_{i''j''} \right) \right] \quad (12)$$

The function u_{ij} is symmetric under permutation $i \leftrightarrow j$, as one can see from the definitions of u_{ij} and $f(r_{ij})$. Therefore, the last sum above can have a different indexing: All terms without an index k , will give zero when taking the derivative ∇_k , so only $i = k$ or $j = k$ remains (remember $i \neq j$). Due to the symmetry of u_{ij} , one can simply always say $i = k$ and let j be the summation index:

$$\nabla_k \Psi_T = \nabla_k \phi_k \left[\prod_{i \neq k} \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \right] + \left[\prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \quad (13)$$

The second derivative now becomes (where ∇_k only acts on the first parenthesis to its right):

$$\begin{aligned}
\nabla_k^2 \Psi_T &= (\nabla_k^2 \phi_k) \left[\prod_{i \neq k} \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \right] + (\nabla_k \phi_k) \left[\prod_{i \neq k} \phi(\mathbf{r}_i) \nabla_k e^{\sum_{i' < j'} u(r_{i'j'})} \right] \\
&+ \left[\nabla_k \left(\prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \right) \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \\
&+ \left[\prod_i \phi(\mathbf{r}_i) e^{\sum_{i' < j'} u(r_{i'j'})} \left(\sum_{j'' \neq k} \nabla_k^2 u_{kj''} \right) \right]
\end{aligned}$$

While a bit of a nuisance to read, the expression above simply the product rule for $\nabla_k(\nabla_k \Psi_T)$. Written in terms of Ψ_T , the above can be a bit simplified²:

$$\begin{aligned}
\nabla_k^2 \Psi_T &= (\nabla_k^2 \phi_k) \frac{\Psi_T}{\phi(\mathbf{r}_k)} + (\nabla_k \phi_k) \left[\prod_{i \neq k} \phi(\mathbf{r}_i) \nabla_k e^{\sum_{i' < j'} u(r_{i'j'})} \right] \\
&+ \left[(\nabla_k \Psi_T) \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \\
&+ \left[\Psi_T \left(\sum_{j'' \neq k} \nabla_k^2 u_{kj''} \right) \right]
\end{aligned}$$

¹The "full local energy" means not making any assumptions on the particle interactions or the potential.

²For the more mathematically concerned nitpicker, the product symbol only runs over the functions $\phi(\mathbf{r}_i)$, not the following exponential. This is easy to realise by recalling how Ψ_T was defined, and is important to know when inserting Ψ_T as done now.

The second term above is equal to the second term in equation 13, divided by $\phi(\mathbf{r}_k)$. Furthermore, the gradient $\nabla_k \Psi_T$ is already calculated above, and in terms of Ψ_T is:

$$\nabla_k \Psi_T = \frac{\Psi_T}{\phi(\mathbf{r}_k)} \nabla_k \phi_k + \Psi_T \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \quad (14)$$

Inserting all this back into the second derivative yields:

$$\begin{aligned} \nabla_k^2 \Psi_T &= (\nabla_k^2 \phi_k) \frac{\Psi_T}{\phi(\mathbf{r}_k)} + (\nabla_k \phi_k) \left[\frac{\Psi_T}{\phi(\mathbf{r}_k)} \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \\ &+ \left[\frac{\Psi_T}{\phi(\mathbf{r}_k)} \nabla_k \phi_k + \Psi_T \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \\ &+ \left[\Psi_T \left(\sum_{j'' \neq k} \nabla_k^2 u_{kj''} \right) \right] \end{aligned}$$

Giving:

$$\begin{aligned} \frac{1}{\Psi_T} \nabla_k^2 \Psi_T &= (\nabla_k^2 \phi_k) \frac{1}{\phi(\mathbf{r}_k)} + (\nabla_k \phi_k) \left[\frac{1}{\phi(\mathbf{r}_k)} \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \\ &+ \left[\frac{1}{\phi(\mathbf{r}_k)} \nabla_k \phi_k + \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \right] \left(\sum_{j'' \neq k} \nabla_k u_{kj''} \right) \\ &+ \sum_{j'' \neq k} \nabla_k^2 u_{kj''} \end{aligned}$$

Which can be rewritten to:

$$\frac{1}{\Psi_T} \nabla_k^2 \Psi_T = \frac{\nabla_k^2 \phi_k}{\phi(\mathbf{r}_k)} + \frac{2 \nabla_k \phi_k}{\phi(\mathbf{r}_k)} \left(\sum_{i \neq k} \nabla_k u_{ki} \right) + \left(\sum_{j \neq k} \nabla_k u_{kj} \right)^2 + \sum_{l \neq k} \nabla_k^2 u_{kl} \quad (15)$$

The gradients $\nabla_k u_{ki}$ can be rewritten using partial differentiation (where $r_{k,i}$ is coordinate i of \mathbf{r}_k):

$$\begin{aligned} \nabla_k u_{ki} &= \left(\frac{\partial}{\partial r_{k,1}}, \frac{\partial}{\partial r_{k,2}}, \dots \right) u_{ki} \\ &= \frac{\partial u_{ki}}{\partial r_{ki}} \left(\frac{\partial r_{ki}}{\partial r_{k,1}}, \frac{\partial r_{ki}}{\partial r_{k,2}}, \dots \right) \\ &= \frac{\partial u_{ki}}{\partial r_{ki}} \left(\frac{\partial}{\partial r_{k,1}} \left(\sqrt{[(r_{k,1} - r_{i,1})\hat{e}_1 + \dots]^2} \right), \dots \right) \\ &= \frac{\partial u_{ki}}{\partial r_{ki}} \left(\frac{r_{k,1} - r_{i,1}}{r_{ki}}, \dots \right) \\ &= \frac{\partial u_{ki}}{\partial r_{ki}} \frac{\mathbf{r}_k - \mathbf{r}_i}{r_{ki}} \\ &= \frac{\mathbf{r}_k - \mathbf{r}_i}{r_{ki}} u'_{ki} \quad , \quad u'_{ki} \equiv \frac{\partial u_{ki}}{\partial r_{ki}} \end{aligned}$$

While the second derivative of u_{ki} is:

$$\begin{aligned} \nabla_k^2 u_{ki} &= \nabla_k (\nabla_k u_{ki}) \\ &= u'_{ki} \nabla_k \left(\frac{\mathbf{r}_k - \mathbf{r}_i}{r_{ki}} \right) + \frac{\mathbf{r}_k - \mathbf{r}_i}{r_{ki}} \nabla_k u'_{ki} \\ &= u'_{ki} \left(\frac{d}{dr_{ki}} - \right) \end{aligned}$$

Which finally gives:

$$\frac{1}{\Psi_T} \nabla_k^2 \Psi_T = \frac{\nabla_k^2 \phi_k}{\phi(\mathbf{r}_k)} + \frac{2 \nabla_k \phi_k}{\phi(\mathbf{r}_k)} \left(\sum_{i \neq k} \frac{\mathbf{r}_k - \mathbf{r}_i}{r_{ki}} u'_{ki} \right) + \sum_{i, j \neq k} \frac{(\mathbf{r}_k - \mathbf{r}_i)(\mathbf{r}_k - \mathbf{r}_j)}{r_{ki} r_{kj}} u'_{ki} u'_{kj} + \sum_{l \neq k} \nabla_k^2 u_{kl} \quad (16)$$

3 Results

N	D	Analytical	Numerical	Variance	Time
1	1	0			
1	2	0			
1	3	0.88			
10	1				
10	2	10	10	7.4e-13	1.35
10	3	15	15	3.7e-13	1.41
100	1				
100	2				
100	3	150	150	8.5e-9	119.38
500	1				
500	2				
500	3	250	250	4.1e-8	316.19

Table 1: some caption

4 Conclusions

5 Appendix