


Fast and accurate parameter estimation of high-redshift sources with the Einstein Telescope

In its triangular configuration, the Einstein Telescope will provide more precise distance measurements than sky localization.

Background: The Einstein Telescope will be a key instrument for detecting GWs in the coming decades [Punturo et al. 2010, Maggiore et al. 2020, Branchesi et al. 2023, Abac et al. 2025]. However, analyzing the data and estimating source parameters will be challenging, especially given the large number of expected detections – of order 10^5 per year – which makes current methods based on stochastic sampling impractical [Couvares et al. 2021]

Main result

Posterior samples from Dingo-IS [Dax et al. 2021] and Bilby [Romero-Shaw et al. 2020] indicate that the two distributions are effectively identical. The corner plot shows also multimodalities that arise due to the geometry of a triangular detector [Singh and Bulik 2021]. These modes are absent by construction when using the Fisher information matrix approximation [GWFish+Priors, Dupletsa et al. 2025].




Methods

In Santoliquido et al. 2025, we use Dingo-IS to perform Neural Posterior Estimation (NPE) of high-redshift events observable with the Einstein Telescope. NPE is a likelihood-free inference technique that leverages normalizing flows to approximate posterior distributions [Dax et al. 2023]. Once trained, this approach enables fast and accurate inference—typically requiring only a few minutes per source. To correct for any residual bias, we employ importance sampling,

$$w_i = \frac{\mathcal{L}(d|\theta)\pi(\theta)}{q(\theta|d)},$$

where $\mathcal{L}(d|\theta)$ is the likelihood, $\pi(\theta)$ the prior and $q(\theta|d)$ is the neural approximation of the posterior.

Sky modes



Filippo Santoliquido, Jacopo Tissino, Ulyana Dupletsa, Marica Branchesi, Jan Harms, Manuel Arca Sedda, Maximilian Dax, Annalena Kofler, Stephen R. Green, Nihar Gupte, Isobel M. Romero-Shaw, Emanuele Berti

