

# Introduction to **GWAS**

## Data Pre-Processing: Initial & Exploratory Data Analysis

Christian Werner

*(Quantitative geneticist and biostatistician)* **EiB, CIMMYT**, Texcoco (Mexico)

Filippo Biscarini

*(Biostatistician, bioinformatician and quantitative geneticist)* **CNR-IBBA**, Milan (Italy)



HerrFalloppio

Oscar González-Recio

*(Computational biologist and quantitative geneticist)* **INIA-UPM**, Madrid (Spain)



OscarGenomics



**Initial data analysis** mainly focuses on data cleaning, a first screening, and transformation (if necessary) to ensure data quality and confirm that our data set meets the relevant distributional and model assumptions. Data cleaning may include steps such as elimination of duplicate records, handling of missing values, identification of systematic errors, or correction of coding inconsistencies.

**Exploratory data analysis** is used to examine data sets and summarize their main characteristics. EDA helps to discover patterns in the data, spot anomalies and outliers, test a hypothesis and check our assumptions. EDA tells us what data can reveal beyond the formal modeling or hypothesis testing and provides a better understanding of data set variables and their interactions. It can also help to determine if the statistical techniques you are considering for data analysis are appropriate.



# Some basic data handling – **plink** (run in the shell)

Basic **plink** command structure:

***./plink --function specification***

Call program from path

Prefix for input files

dogs.ped and dogs.map are the basic input files

***./plink --dog --file dogs --recode vcf --out dogs***

Specify a nonhuman  
chromosome set:  
**--dog** = --chr-set 38

Recode .ped  
and .map file  
to .vcf file

Prefix for  
output files

# Some basic data handling – **plink** (run in the shell)

Basic **plink** command structure: ***./plink --function specification***

Problem with this command: `./plink --dog --file dogs --recode vcf --out dogs`

**POP1** **SAMPLE1** 0 0 0 -9 C C

**POP1** **SAMPLE2** 0 0 0 -9 C C

Vs.

#CHROM POS ID REF ALT QUAL FILTER INFO FORMAT NAME1 NAME2 NAME3....

`./plink --dog --file dogs --recode vcf-fid --out dogs`

`./plink --dog --file dogs --recode vcf-iid --out dogs`

**'vcf-fid'** and **'vcf-iid'** cause family IDs and within-family IDs respectively to be used for the sample IDs  
**'vcf'** merges both IDs and puts an underscore between them

# Some basic data handling – **plink** (run in the shell)

Basic **plink** command structure: ***./plink --function specification***

***When ped and map have different names:***

```
plink --dog --ped dogs.ped --map dogs.map --recode vcf --out dogs
```

***plink reads vcf too!***

```
./plink --vcf dogs.vcf --recode --out dogs
```

# Some basic data handling – **vcftools** (run in the shell)

Basic command structure: ***./vcftools --function specification***

***./vcftools --vcf*** <path to vcf file> ***--plink*** --out <path to out file>

***vcftools --vcf*** dogs.vcf ***--plink*** --out dogs\_plink

(only biallelic markers will be in the output)

# Initial and Exploratory Data Analysis

## ("crap in, crap out")

# Initial and Exploratory Data Analysis – IDA & EDA

Before we conduct a GWAS, there are **two types of data** to explore

- **Genotypic data**
- **Phenotypic data**



# Initial and Exploratory Data Analysis – IDA & EDA

## Genotypic data

# IDA & EDA - **genotypic data**

Various metrics – statistics are performed either across SNP or across samples

Key concept: detect SNP and samples that should be removed prior to GWAS

Use of some metrics and thresholds is species- and population-specific

Still some level of subjectivity in the thresholds

# IDA & EDA - **genotypic data**

**Also referred to as Quality Control (QC)**

Some parameters to look at...

- Genotype calling and signal intensities (not covered here...)
- Marker allele frequencies
- Missing rate per marker and per individual
- Hardy-Weinberg equilibrium
- Heterozygosity

# IDA & EDA - **genotypic data**

## **Marker allele frequencies**

- Allele counts & genotype counts
- Minor allele frequency (MAF)
  - Some SNPs will be monomorphic
  - One of the alleles may be at very low frequency
  - Might be due to genotyping errors
  - Power to detect the association is very low

Common MAF thresholds are between 1 – 5%

In samples with known group structure, MAF should be checked within groups

# IDA & EDA - **genotypic data**

## **Missing rate per marker & per individual**

**SNPs might be of poor quality if their genotyping failed in many individuals**

- Should be investigated separately for all study groups (if known)
- Common thresholds are 2 – 5% (based on sample size & SNP number)

**Sample DNA might be of poor quality if there are many missing SNPs in an individual**

- Too many missing SNPs per individual can be an indication of poor DNA quality
- (or true deletions...)
- Common call rate thresholds are between 2 – 5% (based on sample size & SNP number)
- Includes monomorphic SNP !!

# Exploratory data – genotypes

Marker	Individuals												
	1	2	3	4	5	6	7	8	9	10	11	12	13
2	2	2	2	2	2	2	2	2	2	2	2	2	2
2	2	2	2	2	2	-1	2	2	2	2	2	2	2
2	-1	2	2	2	2	2	1	0	2	1	2	2	2
2	2	2	2	2	2	2	2	2	2	2	2	2	2
0	1	0	0	0	0	-1	0	0	0	0	0	0	0
0	0	0	0	0	0	-1	0	0	0	0	0	0	0
2	1	1	0	0	0	-1	0	0	1	0	0	0	1
0	2	1	1	1	2	2	1	1	2	0	1	1	2
1	0	1	0	0	0	0	0	0	0	0	0	0	0
1	2	1	1	1	2	2	1	2	2	1	2	1	1

# Exploratory data – genotypes

Monomorphic marker

Marker	Individuals												
	1	2	3	4	5	6	7	8	9	10	11	12	13
1	2	2	2	2	2	2	2	2	2	2	2	2	2
2	2	2	2	2	2	-1	2	2	2	2	2	2	2
3	2	-1	2	2	2	2	1	0	2	1	2	2	2
4	2	2	2	2	2	2	2	2	2	2	2	2	2
5	0	1	0	0	0	-1	0	0	0	0	0	0	0
6	0	0	0	0	0	-1	0	0	0	0	0	0	0
7	2	1	1	0	0	-1	0	0	1	0	0	0	1
8	0	2	1	1	2	2	1	1	2	0	1	1	2
9	1	0	1	0	0	0	0	0	0	0	0	0	0
10	1	2	1	1	2	2	1	2	2	1	2	1	1

# Exploratory data – genotypes

Marker	Individuals												
	1	2	3	4	5	6	7	8	9	10	11	12	13
	2	2	2	2	2	2	2	2	2	2	2	2	2
	2	2	2	2	2	-1	2	2	2	2	2	2	2
	2	-1	2	2	2	2	1	0	2	1	2	2	2
	2	2	2	2	2	2	2	2	2	2	2	2	2
	0	1	0	0	0	-1	0	0	0	0	0	0	0
	0	0	0	0	0	-1	0	0	0	0	0	0	0
	2	1	1	0	0	-1	0	0	1	0	0	0	1
	0	2	1	1	2	2	1	1	2	0	1	1	2
	1	0	1	0	0	0	0	0	0	0	0	0	0
	1	2	1	1	2	2	1	2	2	1	2	1	1

Monomorphic marker

Low MAF



# Exploratory data – genotypes

Monomorphic marker

Individuals

Marker	2	2	2	2	2	2	2	2	2	2	2	2	2
	2	2	2	2	2	-1	2	2	2	2	2	2	2
	2	-1	2	2	2	2	1	0	2	1	2	2	2
	2	2	2	2	2	2	2	2	2	2	2	2	2
	0	1	0	0	0	-1	0	0	0	0	0	0	0
	0	0	0	0	0	-1	0	0	0	0	0	0	0
	2	1	1	0	0	-1	0	0	1	0	0	0	1
	0	2	1	1	2	2	1	1	2	0	1	1	2
	1	0	1	0	0	0	0	0	0	0	0	0	0
	1	2	1	1	2	2	1	2	2	1	2	1	1

Low MAF


Many missing markers

# IDA & EDA - genotypic data

**Hardy-Weinberg equilibrium:** relationship between allele and genotype frequencies

$$(p + q)^2 = p^2 + 2pq + q^2$$

- diploid genomes
- autosomal loci
- large population
- random mating
- equal frequencies in both sexes
- no selection
- no migration
- no mutations



# IDA & EDA - **genotypic data**

## Deviations from HW equilibrium

- systematic genotyping errors
- violation of assumptions

## Test for HW deviation

- chi-squared ( $\chi^2$ ) test
- Fisher's (exact) test
- many, many more ...

**But also selection, assortative mating, population structure and inbreeding cause deviations from HWE!**

**HWE is, in most cases, NOT a reasonable assumption...**

# IDA & EDA - genotypic data

## Heterozygosity

- Proportion of heterozygotes
- Heterozygosity can be checked per locus & per marker

### **Very high sample heterozygosity can be an indication of DNA contamination**

- But also could be that a small proportion of samples are truly very different from the rest...
- Removal of samples that depart  $\pm 3$  SD from the mean

### **Very high heterozygosity per marker could also indicate poor DNA quality, but also be due to...**

- the breeding scheme (e.g. hybrid breeding in plants, or very low heterozygosity in lines)
- Genome duplications

IDA & EDA - **genotypic data**

**Phenotypic data**

# IDA & EDA - phenotypic data

## Data type

- Continuous (e.g. height)
- Binary (e.g. case/control)
- Categorical (e.g. scores (ordered), eye colour (ordered))

**Measure of centrality:** mean, mode, median

**Measures of dispersion:** range, variance, standard deviation

## Distribution of the data

- Distribution of values as expected? Outliers?
- representative sample of the population?
- Other explanatory covariables

# IDA & EDA - phenotypic data

## Covariables

Are there any variables which may have a relationship with the phenotype?

- E.g. sex, breed, age, treatments, year effects, ... (population structure)

**The data needs to be corrected for these effects.** Otherwise they can be confounded with allelic variants with an effect on the phenotype which we try to identify.

Covariables with significant effects on the phenotype can be identified using ANOVA (requires balanced datasets - ANOVA is outdated) or **linear mixed models**. However, a comprehensive preparation of phenotypic data including model comparison is not covered here...

# IDA & EDA - **phenotypic data**

## **Assumptions for continuous variables** (different for binary traits...)

- Normally distributed **residuals** (prerequisite of GWAS model assumptions)
- Homogeneity of variance (differences in variance might indicate a factor that has not been included in the phenotype processing)



# IDA & EDA - phenotypic data

## Outliers

- Apparently rare phenotypes are often a result of errors or poor models rather than true outliers
- However, the values might be real - outliers should be investigated thoroughly rather than relying on statistical tests


## Data transformation

- **Positively skewed distributions of the residual** with the long tail into the positive direction can be corrected with a logarithmic or square root transformation.
- **Negatively skewed distributions of the residual** that have a long tail in the negative direction can be corrected with cubing or squaring
- Transformations only if really necessary...

# IDA & EDA - phenotypic data


**Histograms,**

Histogram of arrivals




**Scatter plots,**

Scatterplot for quality characteristic XXX



**Box plots,**



# IDA & EDA - phenotypic data

## Normality of residuals

- Look at the data (Histogram, QQ-Plot)
- Don't rely on Shapiro-Wilk test


## Variance homogeneity

- Look at the data (Scatterplot)
- Don't rely on Leven's test

Tests are conservative and might indicate a violation of the assumptions of normality and variance homogeneity in a suitable “real-world” dataset.


# IDA & EDA - phenotypic data

Almost perfectly distributed residuals




# IDA & EDA - phenotypic data

Still good...



# IDA & EDA - phenotypic data

There might be something wrong here....



# Summary of pre-processing for genotype data

# Pre-processing genotypic data – **Standard procedures**

## **missing rate**

- per-sample (e.g. max 10% missing SNP genotypes per sample)
- per-site (e.g. max 5% missing genotype per variant/SNP)
- stricter/looser thresholds depending on data/experiment (e.g. SNP array, GBS, quality of reference sequence, ...)

MAF

Hardy Weinberg equilibrium

others



# Pre-processing genotypic data – Standard procedures

missing rate

## MAF

- remove monomorphic variants → non-informative
- remove variants at low frequency (“rare”) → spurious associations
- threshold may depend on sample size
- usually (re)done after imputation

Hardy Weinberg equilibrium

others

# Pre-processing genotypic data – Standard procedures

missing rate

MAF

## Hardy Weinberg equilibrium

- set **low threshold** for p-value (e.g.  $\exp(-10)$ )
- **questionable**: some of the forces driving HW disequilibrium are characteristic for breeding (selection, migration, mutation, adaptation etc.)

others

# Pre-processing genotypic data – Standard procedures

missing rate

MAF

Hardy Weinberg equilibrium

**others**

- sex chromosomes (might need to be removed / analyzed separately)
- Mendelian errors
- quality scores (vcf files)
- relatedness (between samples - check for duplicates)