RESEARCH

Methods to compute the composite log-likelihood (CLL) of allelic frequencies for the detection of signatures of selection in diploid genomes

Filippo Biscarini^{1*}, Nelson Nazzicari¹ and Alessandra Stella¹

*Correspondence:

filippo.biscarini@tecnoparco.org

¹Department of Bioinformatics,
PTP, Via Einstein - Loc. Cascina
Codazza, Lodi, Italy
Full list of author information is
available at the end of the article

† Equal contributor

Abstract

First part title: Text for this section. **Second part title:** Text for this section.

Keywords: composite log-likelihood; signatures of selection; diploid genomes

Background

Text and results for this section, as per the individual journal's instructions for authors.

Section title

Text for this section ...

Sub-heading for section

Text for this sub-heading ...

Sub-sub heading for section

Text for this sub-sub-heading \dots

Sub-sub-sub heading for section Text for this sub-sub-heading... In this section we examine the growth rate of the mean of Z_0 , Z_1 and Z_2 . In addition, we examine a common modeling assumption and note the importance of considering the tails of the extinction time T_x in studies of escape dynamics. We will first consider the expected resistant population at vT_x for some v > 0, (and temporarily assume $\alpha = 0$)

$$E[Z_1(vT_x)] = E\left[\mu T_x \int_0^{v \wedge 1} Z_0(uT_x) \exp(\lambda_1 T_x(v-u)) du\right].$$

If we assume that sensitive cells follow a deterministic decay $Z_0(t) = xe^{\lambda_0 t}$ and approximate their extinction time as $T_x \approx -\frac{1}{\lambda_0} \log x$, then we can heuristically estimate the expected value as

$$E\left[Z_1(vT_x)\right] = \frac{\mu}{r}\log x \int_0^{v\wedge 1} x^{1-u} x^{(\lambda_1/r)(v-u)} du$$

Biscarini et al. Page 2 of 2

$$= \frac{\mu}{r} x^{1-\lambda_1/\lambda_0 v} \log x \int_0^{v \wedge 1} x^{-u(1+\lambda_1/r)} du$$

$$= \frac{\mu}{\lambda_1 - \lambda_0} x^{1+\lambda_1/r v} \left(1 - \exp\left[-(v \wedge 1) \left(1 + \frac{\lambda_1}{r} \right) \log x \right] \right). \quad (1)$$

Thus we observe that this expected value is finite for all v > 0 (also see [1, 2, 3, 4, 5]).

Competing interests

The authors declare that they have no competing interests.

Author's contributions

Text for this section ...

Acknowledgements

Text for this section ...

Author details

¹Department of Bioinformatics, PTP, Via Einstein - Loc. Cascina Codazza, Lodi, Italy. ²Marine Ecology Department, Institute of Marine Sciences Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany.

References

- 1. Koonin, E.V., Altschul, S.F., Bork, P.: Brca1 protein products: functional motifs. Nat Genet 13, 266-267 (1996)
- 2. Kharitonov, S.A., Barnes, P.J.: Clinical Aspects of Exhaled Nitric Oxide. in press
- Zvaifler, N.J., Burger, J.A., Marinova-Mutafchieva, L., Taylor, P., Maini, R.N.: Mesenchymal cells, stromal derived factor-1 and rheumatoid arthritis [abstract]. Arthritis Rheum 42, 250 (1999)
- 4. Jones, X.: Zeolites and synthetic mechanisms. In: Smith, Y. (ed.) Proceedings of the First National Conference on Porous Sieves: 27-30 June 1996; Baltimore, pp. 16–27 (1996). Stoneham: Butterworth-Heinemann
- 5. Margulis, L.: Origin of Eukaryotic Cells. Yale University Press, New Haven (1970)

Figures

 $\textbf{Figure 1 Sample figure title.} \ \textbf{A} \ \textbf{short description of the figure content should go here}.$

Figure 2 Sample figure title. Figure legend text.

Tables

Table 1 Sample table title. This is where the description of the table should go.

| | В1 | B2 | B3 |
|----|-----|-----|-----|
| A1 | 0.1 | 0.2 | 0.3 |
| A2 | | | |
| A3 | | | |

Additional Files

Additional file 1 — Sample additional file title

Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.

Additional file 2 — Sample additional file title Additional file descriptions text.