

Developing a parsimonious predictor for binary traits in sugar beet (*Beta vulgaris*)

Filson Nazzarini · Simone Marini ·
Piergiorgio Stevanato · Nelppo Biscicari

Received: 05 August 2014 / Accepted:

Abstract Insert your abstract here. Include keywords, PACS and mathematical subject classification numbers as needed.

Keywords binary traits · genomic predictions · parsimonious predictor · sugar beet

1 Introduction

The primary goal of breeding schemes in farm animals and crops is generally to increase the agricultural output. Production traits are typically quantitative continuous variables (e.g. milk yield in dairy cattle, or per hectare yield in maize and rice). Many traits of importance in plant and animal breeding follow nonetheless a discrete categorical distribution, both ordered (e.g. calving ease in cattle, grain texture in rice) and unordered (e.g. grain pigmentation in rice, coat colour in cattle). A special case is that of binomial traits, which can take up only two different values, like disease resistance/susceptibility or presence/absence of a morphological characteristic. Annual bolting (flowering) behaviour and root vigor are examples of binomial traits of agronomic importance in sugar beet (*Beta vulgaris*). [move this?]

Filippo Biscarini and Nelson Nazzicari contributed equally to the work.

F. Biscarini, N. Nazzicari
Fondazione Parco Tecnologico Padano
E-mail: filippo.biscarini@tecnoparco.org

S. Marini
Dipartimento di Ingegneria Industriale e dell'Informazione
Università di Pavia

P. Stevanato
DAFNE, Università di Padova
24105 Padova, Italy

Advances in biotechnology and genomics, and the advent of high-density molecular markers (especially single-nucleotide polymorphisms, SNP) genotyping have led to the emergence of molecular breeding [14]. One exciting application of molecular breeding is genomic selection: the possibility of predicting the genetic value and future performance of selection candidates solely from their genotypes ([13]). The predictive equations are trained on reference individuals with both genotypic and phenotypic data and then applied to selection candidates with genotypes only. Genomic selection may bring about multiple benefits in breeding programmes such as shorter breeding cycles or more efficient (e.g. traits difficult or expensive to phenotype) and more accurate (e.g. traits with low heritability) estimation of breeding values/selection ([6,9]). Key to the application of genomic selection to breeding programmes are reliable genomic predictions. The recent publication of the reference genome for *Beta vulgaris* genome [4] is facilitating the application of molecular breeding also in this crop species. Pioneering studies on genomic predictions for both continuous ([10,19]) and binary ([2]) traits in sugar beet have already been published.

Genomic predictions are being based on increasing number of molecular markers (e.g. 777K SNP-chip in cattle, 56K SNP-chip in maize, whole-genome sequence data). When a huge number of potential predictors is available, it may be useful to select a subset to limit laboratory and bioinformatics costs, and the time of analysis, while at the same time improving interpretability of results. There is therefore interest in finding the minimum necessary set of information for a specific problem. The principle of parsimony states that a model needs to be simpler than the data it explains (think for instance of K-nearest neighbors -KNN- classifier with $k=1$), and according to Occam's razor, given two models that explain the data equally well, the simplest has to be chosen ([3]).

The objective of this paper is to present the development of a parsimonious predictor for the binary trait root vigor in a population of sugar beet accessions. SNPs in the panel were ranked based on their relevance and used to classify observations: one SNP at a time was removed, progressively reducing the number of SNPs in the predictive model. We found that it was possible to strongly reduce the dimension of the predictor and still achieve high accuracy.

2 Material and methods

2.1 Plant material and SNP genotypes

The available population comprised 123 individual sugar beet (*B. vulgaris*) plants, 100 with high- and 24 with low-root vigor. Root vigor is related to nutrient uptake from the soil and plant productivity ([18]) and is recorded as a binary trait (either high or low) based on the root elongation rate of eleven-days-old seedlings. No predetermined root elongation rate threshold was used to classify sugar beets into high or low root vigour. The classification

was subjectively made upon phenotypic inspection and has nevertheless been shown to be robust over time ([18]). The plant material was provided by Lion Seeds Ltd. (UK).

All plants were genotyped for 192 SNP markers with the high-throughput marker array QuantStudio 12K Flex system coupled with Taqman OpenArray technology. Additional details on the genotyping procedure are described in Stevanato et al., 2013 ([17]). After imputation and editing (call-rate $\geq 95\%$, MAF $\leq 2.5\%$) 175 SNPs were left for the analysis. A more detailed description of SNP genotypes and editing procedure can be found in Biscarini et al. ([2]).

2.2 Predictor development procedure

A two-step approach was adopted for the construction of a parsimonious predictor for root vigor in sugar beet. First, the 175 SNP available for the analysis after data editing were ranked based on their relevance for predicting the trait under study. In the second step the set of predictors was progressively reduced by removing the least useful predictors one at a time. At each iteration logistic regression was used to classify observations with the given set of SNPs. As many classification results as the number of SNPs (i.e. 175) were therefore obtained.

2.2.1 Rank of predictors

When many predictors are available -especially if $p > n$ - it may be of interest to reduce the dimensionality of the problem by choosing the optimal subset of predictors that best describe the relationship between dependent and independent variables [or that best model the response variable, or are most informative with respect to the outcome to be predicted, etc ...]. A Bayesian model selection method, the binary outcome stochastic search (BOSS) algorithm [16], was applied to identify the best set of predictive SNPs by repeated sampling in a Markov Chain Monte Carlo (MCMC) approach. SNPs were ranked based on their probability of inclusion in the best predictive model.

In BOSS the relationship between binary observations and predictors is described by a Gaussian latent variable model with a probit link function:

$$P(Y = [0/1]|X) = \phi(X\beta) \quad (1)$$

where $P(Y = [0/1]|X)$ is the probability of having low or high root vigor given the SNP genotypes X , β is a vector of regression coefficients, and ϕ is the normal cumulative distribution function.

The n independent latent variables were normally distributed as $Z_i \sim N(X_i^T \beta, 1)$ (with $i = 1 \dots n$), and used to model the relationships between SNP genotypes and root vigor:

$$Y_i = \begin{cases} 0 & Z_i \leq 0 \\ 1 & Z_i > 0 \end{cases} \quad \mathbf{Z} = \mathbf{X}\beta + \epsilon \quad (2)$$

BOSS extensively explores the model space to identify relevant predictors, similar to what is done in Best Subset Selection ([8]).

The selection of predictors was performed by introducing a vector of indicator variables $\gamma = (\gamma_1, \dots, \gamma_p)$ such that $\gamma_i = 0$ if $\beta_i = 0$ and $\gamma_i = 1$ if $\beta_i \neq 0$. A predictor was included in the model if its regression coefficient was not null and the associated indicator variable $\gamma = 1$. The model space to be searched was therefore given by the 2^p possible combinations of SNPs (either included or excluded). This yielded a vector β_γ of size p_γ with only the coefficients for which $\gamma_i = 1$.

The prior for vector γ was chosen from the following beta-binomial (B) distribution:

$$f(\gamma) = \frac{B(p_\gamma + a, p - p_\gamma + b)}{B(a, b)} \quad (3)$$

with parameters a and b related to model size p_γ .

Knowing that the posterior conditional density for γ is proportional to its prior and the marginal likelihood of \mathbf{Z} , the following holds:

$$f(\gamma|\mathbf{Z}) \propto f(\gamma) \cdot f(\mathbf{Z}|\gamma) \quad (4)$$

from which vector γ can be sampled.

The BOSS algorithm is summarised below:

1. initialize the latent variables \mathbf{Z} ;
2. sample γ through a Metropolis-Hastings sampler (check the Theory that would not die) from its conditional posterior distribution in equation 4;
3. given \mathbf{Z} and γ , sample β_γ through standard Bayesian linear regression ($\mathbf{Z} = \mathbf{X}_\gamma \beta_\gamma + \epsilon$);
4. given β_γ and \mathbf{Y} , sample the vector of latent variables \mathbf{Z} from equation 2;
5. restart from step 2 until $m = 1\,000\,000$ iterations are completed

From the $m = 1\,000\,000$ MCMC iterations the inclusion probabilities for the 175 SNPs were computed and used to rank the predictors. A more detailed description of the BOSS algorithm can be found in Russu et al. ([16]).

2.2.2 Selection of predictors and classification method

SNPs were selected based on the BOSS rank (see section 2.2.1) by progressively removing one SNP at a time. The full set of 175 SNPs was used at first in the prediction model. Subsequently, the m^{th} least relevant SNP (from the BOSS rank) was removed each time, and the resulting $175 - m$ SNPs model was fitted. As a result of this, 175 different predictive models (from 175 to 1 predictor) were fitted.

With each set of SNP, a logistic regression model for binary outcomes was used to classify observations into low and high root vigor based on the SNP genotypes. The probability having high root vigor ($P(Y = 1|X) = p(x)$) was

modeled as a linear combination of the predictors (SNPs) through a *logit* link-function in a generalised linear model:

$$\text{logit}(p(x_i)) = \log\left(\frac{p(x_i)}{1-p(x_i)}\right) = \mu + \sum_{j=1}^m z_{ij} \text{SNP}_j \quad (5)$$

where $p(x_i)$ is the $P(Y = 1|X)$ for individual i with vector of predictors x_i ; SNP_j is the effect of the j_{th} marker; z_{ij} is the genotype of individual i at locus j (0, 1 or 2 for AA, AB and BB genotypes). Equation 5 returns the odds of $p(x)$ which are backtransformed to $P(Y = 1|X)$ through the cumulative distribution function of the logistic distribution (i.e. the logistic function). Individuals with $p(x) > / < 0.5$ were classified as high/low root vigor individual plants.

Since there were more predictors than observations in model 5 ($p > n$), a ridge logistic regression fitting method [12] was adopted, which consisted in maximizing the following penalized log-likelihood:

$$L(\mu, \text{SNP}) = \sum_{i=1}^n [y_i \log(p_i) + (1 - y_i) \log(1 - p_i)] - \frac{1}{2} \lambda \sum_{j=1}^p \text{SNP}_j^2 \quad (6)$$

where μ is the intercept, SNP is the vector of SNP effects (i.e. the regression coefficients of SNP genotypes) and λ is a tuning parameter that was specified through cross-validation and controls the amount of penalization.

2.2.3 Predictive ability

For each set of SNPs (175 to 1), the predictive ability of the ridge logistic regression model (equation 5) was assessed through a two-fold cross-validation. The 123 samples were randomly split into a training and testing set of approximately equal size. The training set was used to fit the model and estimate SNP effects which were then used to predict root vigor in the testing set. This process was repeated 100 times, each time sampling different training and testing sets. The test error rate in each replicate was computed as:

$$\text{ER}_{(n)} = \frac{1}{n} \sum_{i=1}^n \text{Err}_i \quad (7)$$

where n is the number of observations in the test set and $\text{Err}_i = I(y_i \neq \hat{y}_i)$, with $I(\cdot)$ an indicator function which returns a value of 1 if the predicted and observed phenotypes are different, 0 otherwise. The cross-validation error rate was then estimated averaging the test error rate over all $k = 100$ replicates:

$$\text{CV}_k = \frac{1}{k} \sum_{i=1}^k \text{ER}_i \quad (8)$$

The prediction accuracy was then defined as $1 - \text{CV}_k$. Explanation of error rate and other parameters (ROC?)




Fig. 1 Accuracy (1 - error rate) of prediction as a function of the number of SNPs included in the classifier: BOSS (blue line) vs logistic regression (red line)

2.3 Comparison with another method to rank predictors

Another ranker: why use one, and its description. P value and SNP effect (as it is done in GWAS)

SNP variance [5]

2.4 Software

R [15], weka [7], perl.

3 Results

Supplementary table with BOSS rank of SNPs (Table 2: use SNP name instead of order, add scaffold and chromosome). A few comments on the most important SNPs (which ones, on which chromosomes, how many per chromosome ...).

Figure 1: Accuracy as a function of the number of predictors, BOSS vs logistic [improve plot: no need to go down to 0.0 in the y-axis; legend names and position; color of the lines? The “bump” at around 20-30 SNPs is not visible]

Table 1: TER, FPR, FNR for the first 30/35 SNPs + average for the rest of the SNP (error close to 0). BOSS + GWAS (6 columns)

Probability of assignment as a function of predictors: Figure 2. Better a table? Maybe in discussion?

From ROC curves only the AUC. No plot, use AUC as result in the text (e.g. comparison between ranker: overall average AUC, average AUC per # SNPs + std). Table?




Fig. 2 Distribution of $P(Y = 1|x)$ as a function of the number of SNPs in the classifier

4 Discussion

General overview why error rates are not evenly distributed? Reminder: it works very well because of LD and H2

Unstable below 30/40 SNPs; little “bump” around 20 SNPs: more marked with BOSS, but also visible with GWAS. Why there? SNPs with large effect on the trait, but low significance? SNPs with large effect but low LD (with the QTL)? In the latter case, the marker might sometimes be in the opposite phase. Look also at marker frequency.

Based on results, a panel of 30-35-40 SNPs is recommended for accurate prediction of root vigor (move to breeding applications? Together with development of a custom-chip?)

4.1 Relative performance of rankers

why using Pvalues and not other standard rankers (e.g. backward stepwise selection)? Because of the specific nature of the problem. AUC?

Comparison of rankers: spearman correlation + plot (ranker1 vs ranker2).

Figure 4




Fig. 3 Average linkage disequilibrium (LD) for increasing number of SNPs in the predictive model




Fig. 4 Comparison of BOSS and logistic regression in terms of relative rank position of relevant SNPs

4.2 SNP effects [alternative: LD and probabilities]

Manhattan plot with BOSS weights and weights from the other articles, somehow compared (same chart? two charts? only ten best?).

Let's not use effects from GWAS (the paper is on BOSS!!): something on the top ranked SNPs (position ...)?

Do the peaks make sense from the biological perspective? (something on sugar beet genes?)

Variance of SNPs vs genetic variance: → missing heritability? (cite Brachi 2011, Manolio 2009?).

BOSS probability: 1 big peak + smaller peaks. Compare against SNP density? Maybe the big peak corresponds to a physically isolated SNP, whereas smaller peaks correspond to a cluster of SNPs in LD which individually account for a smaller part of the variability, but together play an important predictive role.

4.3 Genotyping strategies and applications to breeding

Several technology choices are commonly available when genotyping strategies must be decided. Assuming knowledge of SNP flanking sequences, we examine four options: SNP chips, genotyping by sequencing (GBS), targeted sequencing (TS), and a commercial solution, Illumina BeadXpress.

Genotyping by sequencing is

genotyping strategies: Costs, possible technologies (gbs, snp chip, macroarrays), implications

applications to breeding: why is it important root vigor early detection. Other binomial traits (e.g. disease resistance) May be applied to bolting (another trait which exhibits binomial distribution), which has been shown to be controlled by multiple genes and influenced by environmental factors ([1]).

sugar beet: 30% of world's sugar production (cite Dohm? FAO?). Root vigor linked to yield.

Sugar beet: sugar + energy (citation?)

Other binomial traits: resistance to viral and fungal diseases, bolting (cite Dohm? Someone else?)

Breeding has shaped the genome of sugar beet (comparison with *Beta maritima*, [4]).

Extensions to multinomial traits? Examples?

potential and challenges of genomic selection in plant breeding ([11])

5 Conclusions

Concluding remarks.

Acknowledgements This research was financially supported by the Marie Curie European Reintegration Grant “NEUTRADAPT”.

Table 2: SNP rank and inclusion probability from BOSS

SNP	probability
94	0.59
161	0.24
108	0.19
35	0.17

Table 1 Total error rate (TER), false positive (FPR) and false negative (FNR) rates as a function of the number of SNPs ranked according to BOSS or logistic regression

# SNPs	TER	FPR	FNR
1	0.114	0.065	0.049
2	0.085	0.037	0.047
3	0.008		
4	0.005		
5	0.003		
6	0.005		
7	0.003		
8	0.002		
9	0.003		
10	0.002		
11	0.002		
12	0.008		
13	0.009		
14	0.016		
15	0.012		
16	0.007		
17	0.005		
18	0.004		
19	0.002		
20	0.001		
21–30	0.002		
31–40	0.001		
41–50	0.004		
51–60	0.002		
61–70	0.003		
71–80	0.004		
81–90	0.002		
91–100	0.001		
101–175	0.001		

References

1. Abou-Elwafa, S., Büttner, B., Kopisch-Obuch, F., Jung, C., Müller, A.: Genetic identification of a novel bolting locus in *Beta vulgaris* which promotes annuality independently of the bolting gene *B*. *Molecular Breeding* **29**, 989–998 (2012)
2. Biscarini, F., Stevanato, P., Broccanello, C., Stella, A., Saccomani, M.: Genome-enabled predictions for binomial traits in sugar beet populations. *BMC Genetics* **18**(5), 1–9 (2014)
3. Chaitin, G.: The limits of reason. *Scientific American* **294**(3), 74–81 (2006)
4. Dohm, J.C., Minoche, A.E., Holtgräwe, D., Capella-Gutiérrez, S., Zakrzewski, F., Tafer, H., Rupp, O., Sörensen, T.R., Stracke, R., Reinhardt, R., et al.: The genome of the recently domesticated crop plant sugar beet (*beta vulgaris*). *Nature* (2013)
5. Gianola, D., de Los Campos, G., Hill, W.G., Manfredi, E., Fernando, R.: Additive genetic variability and the bayesian alphabet. *Genetics* **183**(1), 347–363 (2009)
6. Goddard, M., Hayes, B.: Genomic selection. *Journal of Animal Breeding and Genetics* **124**(6), 323–330 (2007)
7. Hall, M., Frank, E., Holmes, G., Pfahringer, B., Reutemann, P., Witten, I.H.: The weka data mining software: an update. *ACM SIGKDD explorations newsletter* **11**(1), 10–18 (2009)
8. Hastie, T., Tibshirani, R., Friedman, J.: Model assessment and selection. In: *The elements of statistical learning*, pp. 219–260. Springer, New York, NY, USA (2009)

9. Heffner, E.L., Lorenz, A.J., Jannink, J.L., Sorrells, M.E.: Plant breeding with genomic selection: gain per unit time and cost. *Crop science* **50**(5), 1681–1690 (2010)
10. Hofheinz, N., Borchardt, D., Weissleder, K., Frisch, M.: Genome-based prediction of test cross performance in two subsequent breeding cycles. *Theoretical and Applied Genetics* **125**(8), 1639–1645 (2012)
11. Jonas, E., de Koning, D.J.: Does genomic selection have a future in plant breeding? *Trends in biotechnology* **31**(9), 497–504 (2013)
12. Liu, Z., Shen, Y., Ott, J.: Multilocus association mapping using generalized ridge logistic regression. *BMC bioinformatics* **12**(1), 384 (2011)
13. Meuwissen, T., Hayes, B., Goddard, M.: Prediction of total genetic value using genome-wide dense marker maps. *Genetics* **157**(4), 1819–1829 (2001)
14. Moose, S.P., Mumm, R.H.: Molecular plant breeding as the foundation for 21st century crop improvement. *Plant physiology* **147**(3), 969–977 (2008)
15. R Development Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria (2008). URL <http://www.R-project.org>. ISBN 3-900051-07-0
16. Russu, A., Malovini, A., Puca, A.A., Bellazzi, R.: Stochastic model search with binary outcomes for genome-wide association studies. *Journal of the American Medical Informatics Association* **19**(e1), e13–e20 (2012)
17. Stevanato, P., Broccanello, C., Biscarini, F., Del Corvo, M., Sablok, G., Panella, L., Stella, A., Concheri, G.: High-throughput rad-snp genotyping for characterization of sugar beet genotypes. *Plant Molecular Biology Reporter* pp. 1–6 (2013)
18. Stevanato, P., Trebbi, D., Saccomani, M.: Root traits and yield in sugar beet: identification of aflp markers associated with root elongation rate. *Euphytica* **173**(3), 289–298 (2010)
19. Würschum, T., Reif, J.C., Kraft, T., Janssen, G., Zhao, Y.: Genomic selection in sugar beet breeding populations. *BMC genetics* **14**(1), 85 (2013)