


CONSTANT GAUSSIAN CURVATURE SURFACES IN HYPERBOLIC 3-MANIFOLDS

Filippo Mazzoli (University of Virginia)
November 3rd, 2020
Pangolin Seminar

SKETCH


M quasi-Fuchsian



SKETCH

M quasi-Fuchsian

$\partial C(M)$ & $\partial_\infty M$




SKETCH

M quasi-Fuchsian

$\partial C(M) \& \partial_\infty M$

$M \setminus C(M) = \bigcup_k \Sigma_k$




SKETCH

M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$

$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$$



SKETCH


M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$


$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$$

immersion data of $\Sigma_k \in T^*\mathcal{T}$



SKETCH



M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$


$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$$

immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

SKETCH



M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$

$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$$

immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

symplectic structure of $T^*\mathcal{T}$


PRELIMINARIES

THE HYPERBOLIC SPACE

$\mathbb{H}^n \cong (H^n, h)$, with

$$H^n := \{(x, t) \in \mathbb{R}^{n-1} \times \mathbb{R} \mid t > 0\},$$

$$h = \frac{|dx|^2 + dt^2}{t^2}.$$




THE HYPERBOLIC SPACE

$\mathbb{H}^n \cong (H^n, h)$, with

$$H^n := \{(x, t) \in \mathbb{R}^{n-1} \times \mathbb{R} \mid t > 0\},$$

$$h = \frac{|dx|^2 + dt^2}{t^2}.$$




$$\partial_\infty \mathbb{H}^2 \cong \mathbb{R} \cup \infty = \mathbb{RP}^1, \quad \partial_\infty \mathbb{H}^3 \cong \mathbb{C} \cup \infty = \mathbb{CP}^1$$

THE HYPERBOLIC SPACE

$\mathbb{H}^n \cong (H^n, h)$, with

$$H^n := \{(x, t) \in \mathbb{R}^{n-1} \times \mathbb{R} \mid t > 0\},$$

$$h = \frac{|dx|^2 + dt^2}{t^2}.$$



$$\partial_\infty \mathbb{H}^2 \cong \mathbb{R} \cup \infty = \mathbb{RP}^1, \quad \partial_\infty \mathbb{H}^3 \cong \mathbb{C} \cup \infty = \mathbb{CP}^1$$

$$\text{Iso}^+(\mathbb{H}^2) \cong \text{Aut}(\mathbb{RP}^1) = \mathbb{P}\text{SL}_2 \mathbb{R},$$

$$\text{Iso}^+(\mathbb{H}^3) \cong \text{Aut}(\mathbb{CP}^1) = \mathbb{P}\text{SL}_2 \mathbb{C},$$

where

$$\begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} \cdot z := \frac{\alpha z + \beta}{\gamma z + \delta}.$$

HYPERBOLIC MANIFOLDS

A hyperbolic manifold is a $M = \mathbb{H}^n / \Gamma$, with $\Gamma \overset{\text{d. t.}}{<} \text{Iso}^+(\mathbb{H}^n)$.

A hyperbolic manifold is a $M = \mathbb{H}^n / \Gamma$, with $\Gamma \overset{\text{d. t.}}{<} \text{Iso}^+(\mathbb{H}^n)$.

EXAMPLES FOR $n = 2$

Let Σ be a oriented surface of genus ≥ 2 . $\forall c$ conformal str

$$(\widetilde{\Sigma}, \tilde{c}) \cong H^2, \quad (\text{Unif. thm})$$

and $\exists \rho : \pi_1 \Sigma \rightarrow \Gamma \overset{\text{d. t.}}{<} \text{Aut}(H^2) = \mathbb{P} \text{SL}_2 \mathbb{R}$.

HYPERBOLIC MANIFOLDS

A hyperbolic manifold is a $M = \mathbb{H}^n / \Gamma$, with $\Gamma \overset{\text{d. t.}}{<} \text{Iso}^+(\mathbb{H}^n)$.

EXAMPLES FOR $n = 2$

Let Σ be a oriented surface of genus ≥ 2 . $\forall c$ conformal str

$$(\widetilde{\Sigma}, \tilde{c}) \cong H^2, \quad (\text{Unif. thm})$$

and $\exists \rho : \pi_1 \Sigma \rightarrow \Gamma \overset{\text{d. t.}}{<} \text{Aut}(H^2) = \mathbb{P} \text{SL}_2 \mathbb{R}$. Therefore

$$(\Sigma, c) \text{ Riemann surface} \underset{\text{Unif. thm}}{\leftrightarrow} (\Sigma, h) \text{ hyperbolic surface}$$

Two structures s, s' on Σ are equivalent if $\exists (\Sigma, s) \rightarrow (\Sigma, s')$ isomorphism isotopic to id_{Σ} .

Two structures s, s' on Σ are equivalent if $\exists(\Sigma, s) \rightarrow (\Sigma, s')$ isomorphism isotopic to id_{Σ} .

We define the **Teichmüller space** to be

$$\begin{aligned}\mathcal{T}^c(\Sigma) &:= \{\text{equiv. classes of conf. str's}\} \\ &\cong \{\text{equiv. classes of hyp. str's}\} \quad (\text{Unif. thm}) \\ &=: \mathcal{T}^h(\Sigma).\end{aligned}$$

Two structures s, s' on Σ are equivalent if $\exists(\Sigma, s) \rightarrow (\Sigma, s')$ isomorphism isotopic to id_{Σ} .

We define the **Teichmüller space** to be

$$\begin{aligned}\mathcal{T}^c(\Sigma) &:= \{\text{equiv. classes of conf. str's}\} \\ &\cong \{\text{equiv. classes of hyp. str's}\} \quad (\text{Unif. thm}) \\ &=: \mathcal{T}^h(\Sigma).\end{aligned}$$

$\mathcal{T} \cong_{\text{homeo}} \mathbb{R}^{6(g-1)}$ is Kähler, distances $d_{WP}, d_{\mathcal{T}}, d_{Th}$...

$$\mathcal{M} := \mathcal{T} / \mathcal{MCG}. \quad (\text{Moduli space})$$

QUASI-FUCHSIAN MANIFOLDS


Let $\rho_0 := \rho : \pi_1 \Sigma \rightarrow \mathbb{P} \mathrm{SL}_2 \mathbb{R} < \mathbb{P} \mathrm{SL}_2 \mathbb{C}$ be a hyperbolic structure.

QUASI-FUCHSIAN MANIFOLDS

Let $\rho_0 := \rho : \pi_1 \Sigma \rightarrow \mathbb{P} \mathrm{SL}_2 \mathbb{R} < \mathbb{P} \mathrm{SL}_2 \mathbb{C}$ be a hyperbolic structure.

$\rho_0(\pi_1 \Sigma)$ acts on \mathbb{H}^3 preserving a $\mathbb{H}^2 \subset \mathbb{H}^3$.

$$M_0 := \mathbb{H}^3 / \rho_0(\pi_1 \Sigma). \quad (\text{Fuchsian mfd})$$




QUASI-FUCHSIAN MANIFOLDS

Let $\rho_0 := \rho : \pi_1 \Sigma \rightarrow \mathbb{P} \mathrm{SL}_2 \mathbb{R} < \mathbb{P} \mathrm{SL}_2 \mathbb{C}$ be a hyperbolic structure.

$\rho_0(\pi_1 \Sigma)$ acts on \mathbb{H}^3 preserving a $\mathbb{H}^2 \subset \mathbb{H}^3$.

$$M_0 := \mathbb{H}^3 / \rho_0(\pi_1 \Sigma). \quad (\text{Fuchsian mfd})$$

Now $\rho_0 \rightsquigarrow (\rho_t)_t$, with $\rho_t(\pi_1 \Sigma)$ discrete and ρ_t injective.



QUASI-FUCHSIAN MANIFOLDS

Let $\rho_0 := \rho : \pi_1\Sigma \rightarrow \mathbb{P}SL_2\mathbb{R} < \mathbb{P}SL_2\mathbb{C}$ be a hyperbolic structure.


$\rho_0(\pi_1\Sigma)$ acts on \mathbb{H}^3 preserving a $\mathbb{H}^2 \subset \mathbb{H}^3$.


$$M_0 := \mathbb{H}^3 / \rho_0(\pi_1\Sigma). \quad (\text{Fuchsian mfd})$$

Now $\rho_0 \rightsquigarrow (\rho_t)_t$, with $\rho_t(\pi_1\Sigma)$ discrete and ρ_t injective. Then

$$M_t := \mathbb{H}^3 / \rho_t(\pi_1\Sigma)$$

is a **quasi-Fuchsian mfd** (for t small).





M quasi-Fuchsian

$\partial C(M) \& \partial_\infty M$

$M \setminus C(M) = \bigcup_k \Sigma_k$

$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$


immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

symplectic structure of $T^*\mathcal{T}$

CONVEX CORE

$$\Lambda(\Gamma) := \overline{\Gamma \cdot x_0}^{\mathbb{H}^n} \cap \partial_{\infty} \mathbb{H}^n \quad (\text{Limit set})$$




M is quasi-Fuchsian iff $\Lambda(\Gamma)$ is a Jordan curve.

CONVEX CORE

$$\Lambda(\Gamma) := \overline{\Gamma \cdot x_0}^{\mathbb{H}^n} \cap \partial_\infty \mathbb{H}^n \quad (\text{Limit set})$$

$$C(\Gamma) := \text{CH}_{\overline{\mathbb{H}^n}}(\Lambda(\Gamma)) \cap \mathbb{H}^n \quad (\text{Convex hull})$$




M is quasi-Fuchsian iff $\Lambda(\Gamma)$ is a Jordan curve.

CONVEX CORE

$$\Lambda(\Gamma) := \overline{\Gamma \cdot x_0}^{\mathbb{H}^n} \cap \partial_\infty \mathbb{H}^n \quad (\text{Limit set})$$

$$C(\Gamma) := \text{CH}_{\overline{\mathbb{H}^n}}(\Lambda(\Gamma)) \cap \mathbb{H}^n \quad (\text{Convex hull})$$

$$C(M) := C(\Gamma)/\Gamma \quad (\text{Convex core})$$




M is quasi-Fuchsian iff $\Lambda(\Gamma)$ is a Jordan curve.

CONVEX CORE

$$\Lambda(\Gamma) := \overline{\Gamma \cdot x_0}^{\mathbb{H}^n} \cap \partial_\infty \mathbb{H}^n \quad (\text{Limit set})$$

$$C(\Gamma) := \text{CH}_{\overline{\mathbb{H}^n}}(\Lambda(\Gamma)) \cap \mathbb{H}^n \quad (\text{Convex hull})$$

$$C(M) := C(\Gamma)/\Gamma \quad (\text{Convex core})$$




M is quasi-Fuchsian iff $\Lambda(\Gamma)$ is a Jordan curve.

CONVEX CORE

$$\Lambda(\Gamma) := \overline{\Gamma \cdot x_0}^{\mathbb{H}^n} \cap \partial_\infty \mathbb{H}^n \quad (\text{Limit set})$$

$$C(\Gamma) := \text{CH}_{\mathbb{H}^n}(\Lambda(\Gamma)) \cap \mathbb{H}^n \quad (\text{Convex hull})$$

$$C(M) := C(\Gamma)/\Gamma \quad (\text{Convex core})$$




M is quasi-Fuchsian iff $\Lambda(\Gamma)$ is a Jordan curve.

$C(M)$ is either a hyperbolic surface (if M Fuchsian), or is homeo $\Sigma \times [-1, 1]$.

BOUNDARY OF THE CONVEX CORE

hyp metric $m^\pm \in \mathcal{T}$




BOUNDARY OF THE CONVEX CORE

hyp metric $m^\pm \in \mathcal{T}$

bending measure

$$\mu^\pm := \sum_i \vartheta_i \cdot \gamma_i \in \mathcal{ML}$$



BOUNDARY OF THE CONVEX CORE


hyp metric $m^\pm \in \mathcal{T}$

bending measure

$$\mu^\pm := \sum_i \vartheta_i \cdot \gamma_i \in \mathcal{ML}$$

hyperbolic length

$$L_\mu(m) := \sum_i \vartheta_i \cdot L_{\gamma_i}(m)$$



BOUNDARY AT ∞

$$\Omega(\Gamma) := \partial_\infty \mathbb{H}^3 \setminus \Lambda(\Gamma).$$

$$\Omega(\Gamma) := \partial_\infty \mathbb{H}^3 \setminus \Lambda(\Gamma).$$

The ideal boundary $\partial_\infty M := \Omega(\Gamma)/\Gamma$ is a $C\mathcal{P}$ -surface. It is encoded by

$$(c_\infty^\pm, q_\infty^\pm) \in T^*\mathcal{T},$$

$$\Omega(\Gamma) := \partial_\infty \mathbb{H}^3 \setminus \Lambda(\Gamma).$$

The ideal boundary $\partial_\infty M := \Omega(\Gamma)/\Gamma$ is a $C\mathcal{P}$ -surface. It is encoded by

$$(c_\infty^\pm, q_\infty^\pm) \in T^*\mathcal{T},$$

where:

- c_∞^\pm conformal structures on $\partial_\infty M$;

$$\Omega(\Gamma) := \partial_\infty \mathbb{H}^3 \setminus \Lambda(\Gamma).$$

The ideal boundary $\partial_\infty M := \Omega(\Gamma)/\Gamma$ is a $C\mathcal{P}$ -surface. It is encoded by

$$(c_\infty^\pm, q_\infty^\pm) \in T^*\mathcal{T},$$

where:

- c_∞^\pm conformal structures on $\partial_\infty M$;
- q_∞^\pm Schwarzian at infinity:

$$\Omega(\Gamma) := \partial_\infty \mathbb{H}^3 \setminus \Lambda(\Gamma).$$

The ideal boundary $\partial_\infty M := \Omega(\Gamma)/\Gamma$ is a $C\mathcal{P}$ -surface. It is encoded by

$$(c_\infty^\pm, q_\infty^\pm) \in T^*\mathcal{T},$$

where:

- c_∞^\pm conformal structures on $\partial_\infty M$;
- q_∞^\pm Schwarzian at infinity: given $f_\pm : \Omega(\Gamma)^\pm \rightarrow \Delta$ uniformization map, define

$$\tilde{q}_\infty^\pm := \left(\left(\frac{f''_\pm}{f'_\pm} \right)' - \frac{1}{2} \left(\frac{f''_\pm}{f'_\pm} \right)^2 \right) dz^2.$$

∂CM vs $\partial_\infty M$

On ∂CM	On $\partial_\infty M$
$m^\pm \in \mathcal{T}$	$c_\infty^\pm \in \mathcal{T}$
$\mu^\pm \in \mathcal{ML}$	$\mathcal{F}_\infty^\pm = \text{Hor}(q_\infty^\pm) \in \mathcal{MF}$
$L_\mu(m)$	$\text{ext}_{\mathcal{F}_\infty}(c_\infty)$

∂CM vs $\partial_\infty M$

On ∂CM	On $\partial_\infty M$
$m^\pm \in \mathcal{T}$	$c_\infty^\pm \in \mathcal{T}$
$\mu^\pm \in \mathcal{ML}$	$\mathcal{F}_\infty^\pm = \text{Hor}(q_\infty^\pm) \in \mathcal{MF}$
$L_\mu(m)$	$\text{ext}_{\mathcal{F}_\infty}(c_\infty)$

Theorem (Bers)

$Q\mathcal{F} \ni M \mapsto (c_\infty^+, c_\infty^-) \in \mathcal{T} \times \overline{\mathcal{T}}$ is a homeo!

∂CM vs $\partial_\infty M$

On ∂CM	On $\partial_\infty M$
$m^\pm \in \mathcal{T}$	$c_\infty^\pm \in \mathcal{T}$
$\mu^\pm \in \mathcal{ML}$	$\mathcal{F}_\infty^\pm = \text{Hor}(q_\infty^\pm) \in \mathcal{MF}$
$L_\mu(m)$	$\text{ext}_{\mathcal{F}_\infty}(c_\infty)$

Theorem (Bers)

$Q\mathcal{F} \ni M \mapsto (c_\infty^+, c_\infty^-) \in \mathcal{T} \times \overline{\mathcal{T}}$ is a homeo!

Conjecture (Thurston)

Is $Q\mathcal{F}(\Sigma) \ni M \mapsto (m^+, m^-) \in \mathcal{T}^\flat \times \mathcal{T}^\flat$ a homeo ?

M quasi-Fuchsian

$\partial C(M)$ & $\partial_\infty M$


$M \setminus C(M) = \bigcup_k \Sigma_k$

$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$

immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

symplectic structure of $T^*\mathcal{T}$



k-SURFACES

k -SURFACES

A k -surface is an embedded surface $\iota: \Sigma \rightarrow M$ such that the Gauss curvature of ι^*g_M is $\equiv k$.


k -SURFACES

A k -surface is an embedded surface $\iota: \Sigma \rightarrow M$ such that the Gauss curvature of ι^*g_M is $\equiv k$.

Theorem (Labourie '91)

Let M be q-Fuchsian. Then $M \setminus CM$ is foliated by k -surfaces

$$\Sigma_k = \Sigma_k^+ \sqcup \Sigma_k^-, \text{ with } k \in (-1, 0).$$



IMMERSION DATA

Let $\iota: \Sigma_k \rightarrow M$ be a k -surface, n_k normal vector field.

Let $\iota: \Sigma_k \rightarrow M$ be a k -surface, n_k normal vector field.

$$I_k := \iota^* g_M,$$

$$B_k := -\nabla_{\bullet} n_k > 0,$$

$$II_k := I_k(B_k \cdot, \cdot),$$

$$III_k := I_k(B_k \cdot, B_k \cdot).$$

Let $\iota: \Sigma_k \rightarrow M$ be a k -surface, n_k normal vector field.

$$I_k := \iota^* g_M,$$

$$B_k := -\nabla_{\bullet} n_k > 0,$$

$$\mathcal{II}_k := I_k(B_k \cdot, \cdot),$$

$$\mathcal{III}_k := I_k(B_k \cdot, B_k \cdot).$$

Fact : \mathcal{III}_k has constant curvature too!

Let $\iota: \Sigma_k \rightarrow M$ be a k -surface, n_k normal vector field.

$$\begin{aligned} I_k &:= \iota^* g_M, & B_k &:= -\nabla_{\bullet} n_k > 0, \\ II_k &:= I_k(B_k \cdot, \cdot), & III_k &:= I_k(B_k \cdot, B_k \cdot). \end{aligned}$$

Fact : III_k has constant curvature too! Moreover

$$(\Sigma_k, I_k) \xleftarrow{id} (\Sigma_k, II_k) \xrightarrow{id} (\Sigma_k, III_k)$$

are harmonic and

$$q_k := +\text{Hopf}(id: II_k \rightarrow I_k) \dot{-} -\text{Hopf}(id: II_k \rightarrow III_k) \in QD(\Sigma_k, [II_k]).$$

Let $\iota: \Sigma_k \rightarrow M$ be a k -surface, n_k normal vector field.

$$I_k := \iota^* g_M,$$

$$B_k := -\nabla_{\bullet} n_k > 0,$$

$$\mathbb{I}_k := I_k(B_k \cdot, \cdot),$$

$$\mathbb{III}_k := I_k(B_k \cdot, B_k \cdot).$$

Fact : \mathbb{III}_k has constant curvature too! Moreover

$$(\Sigma_k, I_k) \xleftarrow{id} (\Sigma_k, \mathbb{I}_k) \xrightarrow{id} (\Sigma_k, \mathbb{III}_k)$$

are harmonic and


$$q_k := +\text{Hopf}(id: \mathbb{I}_k \rightarrow I_k) \dot{-} -\text{Hopf}(id: \mathbb{III}_k \rightarrow \mathbb{I}_k) \in Q\mathcal{D}(\Sigma_k, [\mathbb{I}_k]).$$

Rmk: All statements are ok up to multiply by the correct function of k ...

FOLIATION DATA IN \mathcal{T}

Fix a component of $M \setminus CM$.

$$l_k \xrightarrow{k \rightarrow -1} m$$




FOLIATION DATA IN \mathcal{T}

Fix a component of $M \setminus CM$.

$$l_k \xrightarrow{k \rightarrow -1} m$$

$$m_k \xrightarrow{k \rightarrow -1} [\mu] \quad (\text{Belraouti '17})$$




FOLIATION DATA IN \mathcal{T}

Fix a component of $M \setminus CM$.

$$l_k \xrightarrow{k \rightarrow -1} m$$

$$III_k \xrightarrow{k \rightarrow -1} [\mu] \quad (\text{Belraouti '17})$$

$$l_k, II_k, III_k \xrightarrow{k \rightarrow 0} c_\infty$$



FOLIATION DATA IN \mathcal{T}


Fix a component of $M \setminus CM$.

$$l_k \xrightarrow{k \rightarrow -1} m$$

$$|||_k \xrightarrow{k \rightarrow -1} [\mu] \quad (\text{Belraouti '17})$$

$$l_k, ||_k, |||_k \xrightarrow{k \rightarrow 0} c_\infty$$


$$\hat{q}_k \xrightarrow{k \rightarrow 0} q_\infty \quad (\text{Quinn '20})$$



INTERPOLATION BETWEEN ∂CM AND $\partial_\infty M$

On ∂CM	$-1 \leftarrow k$	On ∂M_k	$k \rightarrow 0$	On $\partial_\infty M$
$m \in \mathcal{T}$		$I_k \& c_k$		$c_\infty \in \mathcal{T}$
$\mu \in \mathcal{ML}$		$III_k \& \mathcal{F}_k = \text{Hor}(q_k)$		$\mathcal{F}_\infty \in \mathcal{MF}$
$L_\mu(m)$		$\int H_k \& \text{ext}_{\mathcal{F}_k}(c_k)$		$\text{ext}_{\mathcal{F}_\infty}(c_\infty)$

SKETCH



M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$

$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \hookleftarrow \Sigma_k \rightsquigarrow \partial_\infty M$$

immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

symplectic structure of $T^*\mathcal{T}$

VOLUMES

VOLUMES

Let $M \in Q\mathcal{F}(\Sigma)$. Notion of V^* related to the duality $\mathbb{H}^3 \leftrightarrow d\mathbb{S}^3$.

Let $M \in Q\mathcal{F}(\Sigma)$. Notion of V^* related to the duality $\mathbb{H}^3 \leftrightarrow d\mathbb{S}^3$.

Given $N \subset M$ convex with regular ∂ ,

$$V^*(N) := V(N) - \frac{1}{2} \int_{\partial N} H \, da,$$

$$W(N) := \frac{(V + V^*)(N)}{2} = V(N) - \frac{1}{4} \int_{\partial N} H \, da.$$

Rmk: $H = \text{tr } B$.

Let $M \in QF(\Sigma)$. Notion of V^* related to the duality $\mathbb{H}^3 \leftrightarrow d\mathbb{S}^3$.

Given $N \subset M$ convex with regular ∂ ,

$$V^*(N) := V(N) - \frac{1}{2} \int_{\partial N} H \, da,$$

$$W(N) := \frac{(V + V^*)(N)}{2} = V(N) - \frac{1}{4} \int_{\partial N} H \, da.$$

Rmk: $H = \text{tr } B$.

$t \mapsto W(N_t)$ grows linearly in t (not exponential as V or $-V^*$).

VOLUMES

Renormalized volume: motivated by AdS/CFT correspondence (Witten, Graham). Krasnov and Schlenker studied it for convex co-compact hyp 3-manifolds, highlighted its geometric properties.

VOLUMES

Renormalized volume: motivated by AdS/CFT correspondence (Witten, Graham). Krasnov and Schlenker studied it for convex co-compact hyp 3-manifolds, highlighted its geometric properties.

- for any $g \in c_\infty$, \exists equidistant foliation $(S_t)_t$ near $\partial_\infty M$

$$W(M, e^t g) = W(M, S_t) - t\pi |\chi(\partial M)|.$$

Renormalized volume: motivated by AdS/CFT correspondence (Witten, Graham). Krasnov and Schlenker studied it for convex co-compact hyp 3-manifolds, highlighted its geometric properties.

- for any $g \in c_\infty$, \exists equidistant foliation $(S_t)_t$ near $\partial_\infty M$

$$W(M, e^t g) = W(M, S_t) - t\pi |\chi(\partial M)|.$$

- we maximize over c_∞ :

$$V_R(M) := \sup_{g \in c_\infty} W(M, g) = W(M, g_{\text{hyp}}).$$

VOLUMES

Renormalized volume: motivated by AdS/CFT correspondence (Witten, Graham). Krasnov and Schlenker studied it for convex co-compact hyp 3-manifolds, highlighted its geometric properties.

- for any $g \in c_\infty$, \exists equidistant foliation $(S_t)_t$ near $\partial_\infty M$

$$W(M, e^t g) = W(M, S_t) - t\pi |\chi(\partial M)|.$$

- we maximize over c_∞ :

$$V_R(M) := \sup_{g \in c_\infty} W(M, g) = W(M, g_{\text{hyp}}).$$

Dual volume of the convex core: $V_C^*(M) := V(M) - \frac{1}{2}L_\mu(m)$.

SCHLÄFLI FORMULAE

$$dV_C^* = -\frac{1}{2} dL_\mu (\dot{m}) \quad (\text{Krasnov-Schlenker '09, M '18})$$

(Gardiner's formula)

SCHLÄFLI FORMULAE

$$dV_C^* = -\frac{1}{2} dL_\mu(\dot{m}) \quad (\text{Krasnov-Schlenker '09, M '18})$$

$$dV_R = -\operatorname{Re}\langle q_\infty, \dot{c}_\infty \rangle \quad (\text{Krasnov-Schlenker '08})$$

$$= -\frac{1}{2} d \operatorname{ext}_{\mathcal{F}_\infty}(\dot{c}_\infty) \quad (\text{Gardiner's formula})$$

$$dV_C^* = -\frac{1}{2} dL_\mu(\dot{m}) \quad (\text{Krasnov-Schlenker '09, M '18})$$

$$dV_R = -\operatorname{Re}\langle q_\infty, \dot{c}_\infty \rangle \quad (\text{Krasnov-Schlenker '08})$$

$$= -\frac{1}{2} d \operatorname{ext}_{\mathcal{F}_\infty}(\dot{c}_\infty) \quad (\text{Gardiner's formula})$$

Proposition (M)

$$dW_k(\dot{M}) = -\operatorname{Re}\langle q_k, \dot{c}_k \rangle = -\frac{1}{2} d \operatorname{ext}_{\mathcal{F}_k}(\dot{c}_k),$$

$$dV_k^*(\dot{M}) = -\frac{1}{2} dL_{\mathcal{M}_k}(i_k).$$

CONSEQUENCES: THE RENORMALIZED VOLUME

The (renormalized) quadr diff \hat{q}_k converges to q_∞ , the Schwartzian at infinity (Quinn '20).

CONSEQUENCES: THE RENORMALIZED VOLUME

The (renormalized) quadr diff \hat{q}_k converges to q_∞ , the Schwartzian at infinity (Quinn '20).

Theorem (M)

$$V_R(M) = \lim_{t \rightarrow +\infty} W(M_{k(t)}) - t\pi|\chi(\partial M)|,$$

where $k(t) = -1/\cosh^2 t$.

CONSEQUENCES: THE RENORMALIZED VOLUME

The (renormalized) quadr diff \hat{q}_k converges to q_∞ , the Schwartzian at infinity (Quinn '20).

Theorem (M)

$$V_R(M) = \lim_{t \rightarrow +\infty} W(M_{k(t)}) - t\pi|\chi(\partial M)|,$$


where $k(t) = -1/\cosh^2 t$.

No need to study equidistant foliations!

Proof : $dW_k = d\text{ext}_{\mathcal{F}_k}(\dot{c}_k) \rightarrow d\text{ext}_{\mathcal{F}_\infty}(\dot{c}_\infty) = dV_R$.

$W_k(M) = 0 = V_k(M)$ for every M Fuchsian.

SKETCH



M quasi-Fuchsian

$$\partial C(M) \& \partial_\infty M$$

$$M \setminus C(M) = \bigcup_k \Sigma_k$$

$$\partial C(M) \leftrightsquigarrow \Sigma_k \rightsquigarrow \partial_\infty M$$

immersion data of $\Sigma_k \in T^*\mathcal{T}$

volumes V^* and W and their
Schläfli formulae

symplectic structure of $T^*\mathcal{T}$

A component E of $M \setminus CM$ is a hyperbolic end.

HAMILTONIAN FLOW

A component E of $M \setminus CM$ is a hyperbolic end.

$(\Sigma_k)_k$ foliation of $E \leftrightarrow \gamma_E(k) := (c_k, q_k)$ path in $T^*\mathcal{T}$

A component E of $M \setminus CM$ is a hyperbolic end.

$(\Sigma_k)_k$ foliation of $E \leftrightarrow \gamma_E(k) := (c_k, q_k)$ path in $T^*\mathcal{T}$

Theorem F (M)

There exists a time-dep Hamiltonian vector field $(X_k)_k$ on $(T^*\mathcal{T}, \omega_{\text{cot}})$ whose integral curves are precisely the γ_E 's.

A component E of $M \setminus CM$ is a hyperbolic end.

$(\Sigma_k)_k$ foliation of $E \leftrightarrow \gamma_E(k) := (c_k, q_k)$ path in $T^*\mathcal{T}$

Theorem F (M)

There exists a time-dep Hamiltonian vector field $(X_k)_k$ on $(T^*\mathcal{T}, \omega_{\text{cot}})$ whose integral curves are precisely the γ_E 's.

Analogous to Moncrief's flow for CMC-foliations in constant curvature MGHC spacetimes of dimension 3.

HAMILTONIAN FLOW: PROOF

$$w_k(E) := V(E_k) - \frac{1}{4} \int_{\Sigma_k} H_k \, da_k + \frac{1}{2} L_\mu(m)$$

HAMILTONIAN FLOW: PROOF

$$w_k(E) := V(E_k) - \frac{1}{4} \int_{\Sigma_k} H_k \, da_k + \frac{1}{2} L_\mu(m)$$

$$dw_k = -\Phi_k^* \lambda + \hat{T}h^* \lambda \quad (\text{Schläfli formulae})$$

HAMILTONIAN FLOW: PROOF

$$w_k(E) := V(E_k) - \frac{1}{4} \int_{\Sigma_k} H_k \, da_k + \frac{1}{2} L_\mu(m)$$

$$dw_k = -\Phi_k^* \lambda + \hat{T}h^* \lambda \quad (\text{Schläfli formulae})$$

$$d\dot{w}_k = -(\Phi_k^* \lambda)' = -\iota_{X_k} \Phi_k^* \omega_{\text{cot}} - d(\iota_{X_k} \Phi_k^* \lambda) \quad (\text{Cartan's formula})$$

HAMILTONIAN FLOW: PROOF

$$w_k(E) := V(E_k) - \frac{1}{4} \int_{\Sigma_k} H_k \, da_k + \frac{1}{2} L_\mu(m)$$

$$dw_k = -\Phi_k^* \lambda + \hat{T}h^* \lambda \quad (\text{Schläfli formulae})$$

$$d\dot{w}_k = -(\Phi_k^* \lambda)' = -\iota_{X_k} \Phi_k^* \omega_{\text{cot}} - d(\iota_{X_k} \Phi_k^* \lambda) \quad (\text{Cartan's formula})$$

$$\iota_{X_k} \Phi_k^* \omega_{\text{cot}} = d(-\dot{w}_k - \iota_{X_k} \Phi_k^* \lambda) \doteq dm_k \quad (\text{variation of } W)$$

HYPERBOLIC ENDS

A component E of $M \setminus CM$ is a hyperbolic end.

HYPERBOLIC ENDS

A component E of $M \setminus CM$ is a hyperbolic end.

$\mathcal{E}(\Sigma)$:= space of hyp ends. Two classical parametrizations:

$$\begin{array}{lll} \text{Sch : } & \mathcal{E}(\Sigma) & \longrightarrow T^*\mathcal{T} \\ & E & \longmapsto (c_\infty, q_\infty) \end{array} \qquad \begin{array}{lll} \text{Th : } & \mathcal{E}(\Sigma) & \longrightarrow \mathcal{T} \times \mathcal{ML} \\ & E & \longmapsto (m, \mu) \end{array}$$

HYPERBOLIC ENDS

A component E of $M \setminus CM$ is a hyperbolic end.

$\mathcal{E}(\Sigma)$:= space of hyp ends. Two classical parametrizations:

$$\begin{array}{lll} \text{Sch : } & \mathcal{E}(\Sigma) & \longrightarrow T^*\mathcal{T} \\ & E & \longmapsto (c_\infty, q_\infty) \end{array} \quad \begin{array}{lll} \text{Th : } & \mathcal{E}(\Sigma) & \longrightarrow \mathcal{T} \times \mathcal{ML} \\ & E & \longmapsto (m, \mu) \end{array}$$

Using k -surfaces, Labourie defined

$$\begin{array}{lll} \Phi_k : & \mathcal{E}(\Sigma) & \longrightarrow T^*\mathcal{T} \\ & E & \longmapsto (c_k, q_k) \end{array} \quad \begin{array}{lll} \Psi_k : & \mathcal{E}(\Sigma) & \longrightarrow \mathcal{T} \times \mathcal{T} \\ & E & \longmapsto (I_k, III_k) \end{array}$$

Theorem (Labourie '92)

Ψ_k and Φ_k are homeos for every k .

A component E of $M \setminus CM$ is a hyperbolic end.

$\mathcal{E}(\Sigma)$:= space of hyp ends. Two classical parametrizations:

$$\begin{array}{lll} \text{Sch : } & \mathcal{E}(\Sigma) & \longrightarrow T^*\mathcal{T} \\ & E & \longmapsto (c_\infty, q_\infty) \end{array} \quad \begin{array}{lll} \text{Th : } & \mathcal{E}(\Sigma) & \longrightarrow \mathcal{T} \times \mathcal{ML} \\ & E & \longmapsto (m, \mu) \end{array}$$

Using k -surfaces, Labourie defined

$$\begin{array}{lll} \Phi_k : & \mathcal{E}(\Sigma) & \longrightarrow T^*\mathcal{T} \\ & E & \longmapsto (c_k, q_k) \end{array} \quad \begin{array}{lll} \Psi_k : & \mathcal{E}(\Sigma) & \longrightarrow \mathcal{T} \times \mathcal{T} \\ & E & \longmapsto (I_k, III_k) \end{array}$$

Theorem (Labourie '92)

Ψ_k and Φ_k are homeos for every k .

From the works of Quinn and Belraouti, Φ_k and Ψ_k "converge" to Sch and Th, resp.

CONSEQUENCES: SYMPLECTOMORPHISMS

We change Th and Ψ_k^* into

$$\begin{array}{ccc} \hat{\text{Th}} : & \mathcal{E}(\Sigma) & \longrightarrow & T^*\mathcal{T} \\ & E & \longmapsto & (m, d(L_\mu)_m) \end{array} \quad \begin{array}{ccc} \hat{\Psi}_k : & \mathcal{E}(\Sigma) & \longrightarrow & T^*\mathcal{T} \\ & E & \longmapsto & (I_k, d(L_{\mathcal{M}_k})_{I_k}) \end{array}$$

Corollary (M)

For every $k, k' \in (-1, 0)$

$$\text{Sch}^* \omega_{\text{cot}} = \Phi_k^* \omega_{\text{cot}} = \hat{\Psi}_{k'}^* \omega_{\text{cot}} = \hat{\text{Th}}^* \omega_{\text{cot}}.$$

Generalization of results of Krasnov-Schlenker '09, Bonsante-Mondello-Schlenker '15.

CONSEQUENCES: McMULLEN'S RECIPROCITY

Theorem (M)

The image of the map

$$\begin{aligned} F_k : \quad Q\mathcal{F}(\Sigma) &\longrightarrow T^*(\mathcal{T} \times \overline{\mathcal{T}}) \\ M &\longmapsto (c_k^+, c_k^-; q_k^+, q_k^-) \end{aligned}$$

is a Lagrangian submanifold of $(T^*(\mathcal{T} \times \overline{\mathcal{T}}), \omega_{\text{cot}})$.

CONSEQUENCES: McMULLEN'S RECIPROCITY

Theorem (M)

The image of the map

$$\begin{aligned} F_k : \quad Q\mathcal{F}(\Sigma) &\longrightarrow T^*(\mathcal{T} \times \overline{\mathcal{T}}) \\ M &\longmapsto (c_k^+, c_k^-; q_k^+, q_k^-) \end{aligned}$$

is a Lagrangian submanifold of $(T^*(\mathcal{T} \times \overline{\mathcal{T}}), \omega_{\text{cot}})$.


Proof : the pullback of the Liouville form λ on $T^*(\mathcal{T} \times \overline{\mathcal{T}})$ by F_k is equal to dW_k (Schläfli formulae).

$$F_k^* \omega_{\text{cot}} = d(F_k^* \lambda) = d(dW_k) = 0.$$


THANK YOU FOR YOUR ATTENTION!

POLYHEDRA VS CONVEX CORES

POLYHEDRON




CONVEX CORE




POLYHEDRA VS CONVEX CORES

POLYHEDRON




CONVEX CORE




hyp metric on $\partial P \setminus V$

POLYHEDRA VS CONVEX CORES

POLYHEDRON




CONVEX CORE




hyp metric on $\partial P \setminus V$
bending $\sum_i \vartheta_i \cdot e_i$

POLYHEDRA VS CONVEX CORES

POLYHEDRON



CONVEX CORE




hyp metric on $\partial P \setminus V$

bending $\sum_i \vartheta_i \cdot e_i$

length $\sum_i \vartheta_i \cdot L_{e_i}$


POLYHEDRA VS CONVEX CORES

POLYHEDRON



hyp metric on $\partial P \setminus V$
bending $\sum_i \vartheta_i \cdot e_i$
length $\sum_i \vartheta_i \cdot L_{e_i}$


CONVEX CORE



hyp metric $m \in \mathcal{T}(\partial CM)$


POLYHEDRA VS CONVEX CORES

POLYHEDRON



hyp metric on $\partial P \setminus V$
bending $\sum_i \vartheta_i \cdot e_i$
length $\sum_i \vartheta_i \cdot L_{e_i}$


CONVEX CORE




hyp metric $m \in \mathcal{T}(\partial CM)$
bending $\mu := \sum_i \vartheta_i \cdot \gamma_i \in \mathcal{ML}$

POLYHEDRA VS CONVEX CORES

POLYHEDRON



CONVEX CORE




hyp metric on $\partial P \setminus V$
bending $\sum_i \vartheta_i \cdot e_i$
length $\sum_i \vartheta_i \cdot L_{e_i}$

hyp metric $m \in \mathcal{T}(\partial CM)$
bending $\mu := \sum_i \vartheta_i \cdot \gamma_i \in \mathcal{ML}$
length $L_\mu(m) := \sum_i \vartheta_i \cdot L_{\gamma_i}(m)$


DUAL VOLUME

Let P be a compact polyhedron.



DUAL VOLUME


Let P be a compact polyhedron.



$$V(P) + V^*(P) = \frac{1}{2} \sum_i \vartheta_i \cdot L_{e_i}$$

DUAL VOLUME

Let P be a compact polyhedron.




$$V(P) + V^*(P) = \frac{1}{2} \sum_i \boldsymbol{\vartheta}_i \cdot \mathbf{L}_{e_i}$$

$$dV(\dot{P}) = \frac{1}{2} \sum_i \dot{\boldsymbol{\vartheta}}_i \cdot L_{e_i} \quad (\text{Schläfli 1858})$$

DUAL VOLUME

Let P be a compact polyhedron.




$$V(P) + V^*(P) = \frac{1}{2} \sum_i \vartheta_i \cdot L_{e_i}$$

$$dV(\dot{P}) = \frac{1}{2} \sum_i \dot{\vartheta}_i \cdot L_{e_i} \quad (\text{Schläfli 1858})$$

$$dV^*(\dot{P}) = \frac{1}{2} \sum_i \vartheta_i \cdot \dot{L}_{e_i} \quad (\text{Schläfli}^*)$$

DUAL VOLUME (IN M)


Let $M \in Q\mathcal{F}(\Sigma)$.



$$V_C^*(M) := V_C(M) - \frac{1}{2}L_\mu(m)$$

DUAL VOLUME (IN M)

Let $M \in Q\mathcal{F}(\Sigma)$.




$$V_C^*(M) := V_C(M) - \frac{1}{2}L_\mu(m)$$

$$dV_C(\dot{M}) = \frac{1}{2}L_{\dot{\mu}}(m) \quad (\text{Bonahon-Schl\"afli '98})$$

DUAL VOLUME (IN M)

Let $M \in Q\mathcal{F}(\Sigma)$.



$$V_C^*(M) := V_C(M) - \frac{1}{2}L_\mu(m)$$

$$dV_C(\dot{M}) = \frac{1}{2}L_{\dot{\mu}}(m) \quad (\text{Bonahon-Schl\"afli '98})$$

$$dV_C^*(\dot{M}) = -\frac{1}{2}dL_\mu(\dot{m}) \quad (\text{dual B-S, Krasnov-Schlenker '09})$$

SCHLÄFLI FORMULAE

$$\delta V(N) = \frac{1}{2} \int_{\partial N} \left(\delta H + \frac{1}{2} (\delta I, II) \right) da . \quad (\text{Rivin-Schlenker '99})$$

Lemma (M)

$$\delta W(N) = \frac{1}{4} \int_{\partial N} \left(\left(\delta II, III - \frac{H}{2} II \right)_II + \frac{\delta K^e}{2K^e} H \right) da .$$

$$dW_k(\dot{M}) = \frac{1}{8} \int_{\partial M_k} \left(II_k, -2 \operatorname{Re} q_k \right)_{II_k} da_{II_k} = - \operatorname{Re} \langle q_k, \dot{c}_k \rangle$$