


# Shear-bend coordinates for pleated surfaces in $PSL_d \mathbb{C}$

Filippo Mazzoli (University of Virginia),  
joint work with



Sara Maloni,



Giuseppe Martone, & Tengren Zhang



AMS-SMF-EMS Meeting, Grenoble, July 20<sup>th</sup>, 2022.

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $X(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$(d=2 : \text{Thurston, Bonahon '96}$   
 $d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17})$

$(d=2 : \text{Thurston, Bonahon '96})$   
 $d>2 : \text{MMMZ}$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow Q\mathcal{T}(S) \subset \left\{ [p] \in X(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} p \text{ holonomy of} \\ \text{a pleated surface} \end{array} \right\} \not\subset X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $X(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$(d=2 : \text{Thurston, Bonahon '96}$   
 $d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17})$

$(d=2 : \text{Thurston, Bonahon '96}$   
 $d>2 : \text{MMMZ})$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow Q\mathcal{T}(S) \subset \left\{ [p] \in X(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} p \text{ holonomy of} \\ \text{a pleated surface} \end{array} \right\} \not\subset X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $X(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17} \end{array})$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{MMMZ} \end{array})$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow Q\mathcal{T}(S) \subset \left\{ [p] \in X(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} \text{pholonomy of} \\ \text{a pleated surface} \end{array} \right\} \not\subset X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $X(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

$$\begin{cases} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17} \end{cases}$$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$$\begin{cases} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{MMMZ} \end{cases}$$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow Q\mathcal{T}(S) \subset \left\{ [p] \in X(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} \text{pholonomy of} \\ \text{a pleated surface} \end{array} \right\} \not\subset X_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $\chi(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17} \end{array})$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{MMMZ} \end{array})$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow \mathcal{QF}(S) \subset \left\{ [p] \in \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} \text{pholonomy of} \\ \text{a pleated surface} \end{array} \right\} \not\subset \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Motivation

Goal : Study  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$ , with

- $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ ,
- $\Gamma := \pi_1(S)$ ,  $S$  oriented closed surface,  $\chi(S) < 0$ .

Focus : Coord's on "large" open subsets of  $\text{Hom}(\Gamma, \text{PSL}_d \mathbb{K})$  generalizing techniques from 2 & 3-dimensional hyperbolic geometry

I. shear coord's in  $\text{PSL}_d \mathbb{R}$

II. shear-bend coord's in  $\text{PSL}_d \mathbb{C}$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{Fock-Goncharov '06, Bonahon-Dreyer '17} \end{array})$

$(\begin{array}{l} d=2 : \text{Thurston, Bonahon '96} \\ d>2 : \text{MMMZ} \end{array})$

When  $d=2$  :

- $\text{PSL}_2 \mathbb{R} = \text{Isom}^+(\mathbb{H}^2) \rightsquigarrow \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{R}) = \text{Teich}(S) \cup \text{Teich}(\bar{S})$
- $\text{PSL}_2 \mathbb{C} = \text{Isom}^+(\mathbb{H}^3) \rightsquigarrow \mathcal{QF}(S) \subset \left\{ [p] \in \mathcal{X}(\Gamma, \text{PSL}_2 \mathbb{C}) \mid \begin{array}{l} p \text{ holonomy of} \\ \text{a pleated surface} \end{array} \right\} \subset \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$

## Hitchin representations

Hitchin :  $\text{Hit}_d(S) := \text{conn. comp. of } \text{Lirr}(\mathcal{K}_{d,f.}(\Gamma, \text{PSL}_2 \mathbb{R})) \subset \mathcal{K}(\Gamma, \text{PSL}_d \mathbb{R})$

$$\cong_{\text{diff}} \mathbb{R}^{(d^2-1)|X(S)|} \quad (\text{if } d=2, \text{ Hit}_d(S) = \text{Teich}(S))$$

Labourie : i)  $[p] \in \text{Hit}_d(S) \Rightarrow \exists \xi : \partial \Gamma \rightarrow \text{Flags}(\mathbb{R}^d)$   $p$ -equivariant  
 (Guichard :  $\Leftarrow$ )  $\quad$  (Hölder) continuous, hyperconvex Frenet curve  
 Gromov  $\partial$  of Cayley( $\Gamma$ )

$$\left( \begin{array}{l} \xi \text{ is transverse \& generic, i.e.} \\ \text{hyperconvex Frenet} \Rightarrow \forall x, y, z \in \Gamma \quad \forall i, j \geq 1 : i+j=d \quad \forall k, h, l \geq 1 : k+h+l=d \\ \xi(x)^i + \xi(y)^j = \xi(x)^k + \xi(y)^h + \xi(z)^l = \mathbb{R}^d \end{array} \right)$$

ii)  $p \in \text{Hitin} \Rightarrow p \text{ Borel-Anosov, i.e. the (lift of) the geodesic flow on}$

$$T^*S \times_p \mathbb{R}^d := T^*\tilde{S} \times \mathbb{R}^d / (v_*x) \sim (rv, p(r)x)$$

uniformly expands/contracts in the directions determined by  $\xi$ .

## Hitchin representations

Hitchin :  $\text{Hit}_d(S) := \text{conn. comp. of } \text{Lirr}(\mathcal{K}_{d,f.}(\Gamma, \text{PSL}_2 \mathbb{R})) \subset \mathcal{K}(\Gamma, \text{PSL}_d \mathbb{R})$

$$\cong_{\text{diff}} \mathbb{R}^{(d^2-1)|X(S)|} \quad (\text{if } d=2, \text{ Hit}_d(S) = \text{Teich}(S))$$

Labourie : i)  $[\rho] \in \text{Hit}_d(S) \Rightarrow \exists \xi : \partial \Gamma \rightarrow \text{Flags}(\mathbb{R}^d)$   $\rho$ -equivariant  
Gromov  $\partial$  of Cayley( $\Gamma$ )  
 (Guichard :  $\Leftarrow$ )  $(\text{H\"older})$  continuous, hyperconvex Frenet curve

$$\left( \begin{array}{l} \xi \text{ is transverse \& generic, i.e.} \\ \text{hyperconvex Frenet} \Rightarrow \forall x, y, z \in \partial \Gamma \quad \forall i, j \geq 1 : i+j=d \quad \forall k, h, l \geq 1 : k+h+l=d \\ \xi(x)^i + \xi(y)^j = \xi(x)^k + \xi(y)^h + \xi(z)^l = \mathbb{R}^d \end{array} \right)$$

ii)  $\rho$  Hitchin  $\Rightarrow$   $\rho$  ~~Borel-Anosov~~, i.e. the (lift of) the geodesic flow on

$$T^*S \times_{\rho} \mathbb{R}^d := T^*\tilde{S} \times \mathbb{R}^d / (v_* X) \sim (r_v, \rho(r) X)$$

uniformly expands/contracts in the directions determined by  $\xi$ .

## Hitchin representations

Hitchin :  $\text{Hit}_d(S) := \text{conn. comp. of } \text{Lirr}(\mathcal{K}_{d,f.}(\Gamma, \text{PSL}_2 \mathbb{R})) \subset \mathcal{K}(\Gamma, \text{PSL}_d \mathbb{R})$

$$\cong_{\text{diff}} \mathbb{R}^{(d^2-1)|X(S)|} \quad (\text{if } d=2, \text{ Hit}_d(S) = \text{Teich}(S))$$

Labourie : i)  $[p] \in \text{Hit}_d(S) \Rightarrow \exists \xi : \partial \Gamma \rightarrow \text{Flags}(\mathbb{R}^d)$   $p$ -equivariant  
 (Guichard :  $\Leftarrow$ ) (Hölder) continuous, hyperconvex Frenet curve

$$\left( \begin{array}{l} \xi \text{ is transverse \& generic, i.e.} \\ \text{hyperconvex Frenet} \Rightarrow \forall x, y, z \in \partial \Gamma \quad \forall i, j \geq 1 : i+j=d \text{ \& } \forall k, h, l \geq 1 : k+h+l=d \\ \xi(x)^i + \xi(y)^j = \xi(x)^k + \xi(y)^h + \xi(z)^l = \mathbb{R}^d \end{array} \right)$$

ii)  $p \text{ Hitchin} \Rightarrow p \text{ Borel-Anosov}$ , i.e. the (lift of) the geodesic flow on

$$T^*S \times_p \mathbb{R}^d := T^*\tilde{S} \times \mathbb{R}^d / (v_*x) \sim (rv, p(r)x)$$

uniformly expands/contracts in the directions determined by  $\xi$ .

## Hitchin representations

Hitchin :  $\text{Hit}_d(S) := \text{conn. comp. of } \text{Lirr}(\mathcal{K}_{d,f.}(\Gamma, \text{PSL}_2 \mathbb{R})) \subset \mathcal{K}(\Gamma, \text{PSL}_d \mathbb{R})$

$$\cong_{\text{diff}} \mathbb{R}^{(d^2-1)|X(S)|} \quad (\text{if } d=2, \text{ Hit}_d(S) = \text{Teich}(S))$$

Labourie : i)  $[p] \in \text{Hit}_d(S) \Rightarrow \exists \xi : \partial \Gamma \rightarrow \text{Flags}(\mathbb{R}^d)$   $p$ -equivariant  
 (Guichard :  $\Leftarrow$ ) (Hölder) continuous, hyperconvex Frenet curve

$$\left( \begin{array}{l} \xi \text{ is transverse \& generic, i.e.} \\ \text{hyperconvex Frenet} \Rightarrow \forall x, y, z \in \partial \Gamma \quad \forall i, j \geq 1 : i+j=d \text{ \& } \forall k, h, l \geq 1 : k+h+l=d \\ \xi(x)^i + \xi(y)^j = \xi(x)^k + \xi(y)^h + \xi(z)^l = \mathbb{R}^d \end{array} \right)$$

ii)  $p \in \text{Hitin} \Rightarrow p$  **Borel-Anosov**, i.e. the (lift of) the geodesic flow on

$$T^*S \times_p \mathbb{R}^d := T^*\tilde{S} \times \mathbb{R}^d / (v_* X) \sim (r_v, p(r)X)$$

uniformly expands/contracts in the directions determined by  $\xi$ .

## Geodesic laminations

Let  $\tilde{S} \rightarrow S$  univ. cover.

(oriented) geodesic  $g$  of  $\tilde{S}$  := pair  $(g^+, g^-) \in \partial\Gamma \times \partial\Gamma \setminus \{\text{diagonals}\}$


(oriented) ideal triangle  $T$  of  $\tilde{S}$  := triple  $(x_T, y_T, z_T) \in (\partial\Gamma)^3 \setminus \{\text{diagonals}\}$

(Every reHom.d.f( $\Gamma, PSL_2\mathbb{R}$ ) induces  $\phi_\circ : \partial\Gamma \rightarrow \partial\mathbb{H}^2$   $\tau$ -equiv homeo)



$\Gamma$ -invariant closed subset

(geodesic) lamination of  $S$ :  $\tilde{\lambda} \subset \{\text{geodesics of } \tilde{S}\}$   
s.t.  $\forall g \neq h \in \tilde{\lambda}$   $g, h$  disjoint



$\lambda := \tilde{\lambda}/\Gamma$  is maximal if  $S \setminus \lambda = \bigsqcup$  interior of ideal triangles


## Geodesic laminations

Let  $\tilde{S} \rightarrow S$  univ. cover.

(oriented) geodesic  $g$  of  $\tilde{S}$  := pair  $(g^+, g^-) \in \partial\Gamma \times \partial\Gamma \setminus \{\text{diagonals}\}$


(oriented) ideal triangle  $T$  of  $\tilde{S}$  := triple  $(x_T, y_T, z_T) \in (\partial\Gamma)^3 \setminus \{\text{diagonals}\}$

(Every  $r \in \text{Hom}_d(\Gamma, \text{PSL}_2 \mathbb{R})$  induces  $\phi_r : \partial\Gamma \rightarrow \partial\mathbb{H}^2$   $r$ -equiv homeo)



$\Gamma$ -invariant closed subset

(geodesic) lamination of  $S$ :  $\tilde{\lambda} \subset \{\text{geodesics of } \tilde{S}\}$   
s.t.  $\forall g \neq h \in \tilde{\lambda}$   $g, h$  disjoint



$\lambda := \tilde{\lambda}/\Gamma$  is maximal if  $S \setminus \lambda = \bigsqcup$  interior of ideal triangles


## Geodesic laminations

Let  $\tilde{S} \rightarrow S$  univ. cover.

(oriented) geodesic  $g$  of  $\tilde{S}$  := pair  $(g^+, g^-) \in \partial\Gamma \times \partial\Gamma \setminus \{\text{diagonals}\}$


(oriented) ideal triangle  $T$  of  $\tilde{S}$  := triple  $(x_T, y_T, z_T) \in (\partial\Gamma)^3 \setminus \{\text{diagonals}\}$

(Every  $r \in \text{Homeo}_+(\Gamma, \text{PSL}_2 \mathbb{R})$  induces  $\phi_r : \partial\Gamma \rightarrow \partial\mathbb{H}^2$   $r$ -equiv homeo)



$\Gamma$ -invariant closed subset

(geodesic) lamination of  $S$ :  $\tilde{\lambda} \subset \{\text{geodesics of } \tilde{S}\}$   
 s.t.  $\forall g \neq h \in \tilde{\lambda} \quad g, h \text{ disjoint}$



$\lambda := \tilde{\lambda}/\rho$  is maximal if  $S \setminus \lambda = \bigsqcup$  interior of ideal triangles


## Geodesic laminations

Let  $\tilde{S} \rightarrow S$  univ. cover.

(oriented) geodesic  $g$  of  $\tilde{S}$  := pair  $(g^+, g^-) \in \partial\Gamma \times \partial\Gamma \setminus \{\text{diagonals}\}$


(oriented) ideal triangle  $T$  of  $\tilde{S}$  := triple  $(x_T, y_T, z_T) \in (\partial\Gamma)^3 \setminus \{\text{diagonals}\}$

(Every  $r \in \text{Homeo}_+(\Gamma, \text{PSL}_2 \mathbb{R})$  induces  $\phi_r : \partial\Gamma \rightarrow \partial\mathbb{H}^2$   $r$ -equiv homeo)



$\Gamma$ -invariant closed subset

(geodesic) lamination of  $S$ :  $\tilde{\lambda} \subset \{\text{geodesics of } \tilde{S}\}$   
 s.t.  $\forall g \neq h \in \tilde{\lambda}$   $g, h$  disjoint



$\lambda := \tilde{\lambda}/\Gamma$  is maximal if  $S \setminus \lambda = \bigsqcup$  interior of ideal triangles

# Fock-Goncharov '06, Bonahon-Dreyer '17 (for PSLd IR)

Fix  $\lambda$  max lam.  $\nabla T$  triangle of  $\tilde{\lambda}$ ,  $\forall k, h, l \geq 1 : k+h+l=d$ ,  
define **triangle invariants**

$$\tau_{khl}(x_T, y_T, z_T) := \log \left( \frac{\xi(x_T)^{h+1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l-1} \dots}{\xi(x_T)^{h-1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l+1} \dots} \right) \in \mathbb{R}$$

and **shears between triangles  $T, T'$**   $\forall i \in \{1, \dots, d-1\}$

$$\sigma_i(T, T') := \log \left( - \frac{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_T)^1}{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_{T'})^1} \dots \right) \in \mathbb{R}$$

$\Rightarrow (\sigma, \tau)$  are a  **$\lambda$ -cocyclic pair**, i.e.



holds only if adjacent

i) symmetries:  $\sigma_i(T, T') = \sigma_{d-i}(T', T)$ ,  $\tau_{khl}(x_T, y_T, z_T) = \tau_{k' l' h'}(\underbrace{y_T, z_T, x_T}_{\text{inverted}})$ ,

ii)  $\Gamma$ -invariance:  $\sigma_i(\gamma T, \gamma T') = \sigma_i(T, T')$ ,  $\tau_{khl}(r x_T, r y_T, r z_T) = \tau_{khl}(x_T, y_T, z_T)$ ,

iii) cocycle prop:  $\sigma_i(T, T'') = \sigma_i(T, T') + \sigma_i(T', T'') + \left( \begin{array}{c} \text{sum of some} \\ \text{triangle invariants} \quad (=0 \text{ when } d=2) \\ \text{of } T \end{array} \right)$

# Fock-Goncharov '06, Bonahon-Dreyer '17 (for PSLd IR)


Fix  $\lambda$  max lam. &  $T$  triangle of  $\tilde{\lambda}$ ,  $\forall k, h, l \geq 1 : k+h+l=d$ ,  
define triangle invariants

$$\tau_{khl}(x_T, y_T, z_T) := \log \left( \frac{\xi(x_T)^{h+1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l-1} \dots}{\xi(x_T)^{h-1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l+1} \dots} \right) \in \mathbb{R}$$

and shears between triangles  $T, T'$   $\forall i \in \{1, \dots, d-1\}$

$$\sigma_i(T, T') := \log \left( - \frac{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_T)^1 \dots}{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_{T'})^1 \dots} \right) \in \mathbb{R}$$

$\Rightarrow (\sigma, \tau)$  are a  $\lambda$ -cocyclic pair, i.e.



holds only if adjacent

i) symmetries:  $\sigma_i(T, T') = \sigma_{d-i}(T', T)$ ,  $\tau_{khl}(x_T, y_T, z_T) = \tau_{k'l'h}(y_T, z_T, x_T)$ ,

ii)  $\Gamma$ -invariance:  $\sigma_i(\gamma T, \gamma T') = \sigma_i(T, T')$ ,  $\tau_{khl}(\gamma x_T, \gamma y_T, \gamma z_T) = \tau_{khl}(x_T, y_T, z_T)$ ,

iii) cocycle prop:  $\sigma_i(T, T'') = \sigma_i(T, T') + \sigma_i(T', T'') + \left( \begin{array}{c} \text{sum of some} \\ \text{triangle invariants} \quad (=0 \text{ when } \\ \text{of } T) \end{array} \right)$

Fock-Goncharov '06, Bonahon-Dreyer '17 (for PSLd IR)

Fix  $\lambda$  max lam. &  $T$  triangle of  $\tilde{\lambda}$ ,  $\forall k, h, l \geq 1 : k+h+l=d$ ,  
define triangle invariants

$$\tau_{khl}(x_T, y_T, z_T) := \log \left( \frac{\xi(x_T)^{h+1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l-1} \dots}{\xi(x_T)^{h-1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l+1} \dots} \right) \in \mathbb{R}$$

and shears between triangles  $T, T' \quad \forall i \in \{1, \dots, d-1\}$


$$\sigma_i(T, T') := \log \left( - \frac{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_T)^1 \dots}{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_{T'})^1 \dots} \right) \in \mathbb{R}$$

$\Rightarrow (\sigma, \tau)$  are a  $\lambda$ -cocyclic pair, i.e.

i) symmetries:  $\sigma_i(T, T') = \sigma_{d-i}(T', T)$ ,  $\tau_{khl}(x_T, y_T, z_T) = \tau_{khl}(y_T, z_T, x_T)$ ,

ii)  $\Gamma$ -invariance:  $\sigma_i(\gamma T, \gamma T') = \sigma_i(T, T')$ ,  $\tau_{khl}(\gamma x_T, \gamma y_T, \gamma z_T) = \tau_{khl}(x_T, y_T, z_T)$ ,

iii) cocycle prop:  $\sigma_i(T, T'') = \sigma_i(T, T') + \sigma_i(T', T'') + \left( \begin{array}{c} \text{sum of some} \\ \text{triangle invariants} \left( \begin{array}{c} =0 \text{ when} \\ d=2 \end{array} \right) \end{array} \right)$



# Fock-Goncharov '06, Bonahon-Dreyer '17 (for PSLd IR)

Fix  $\lambda$  max lam.  $\nabla T$  triangle of  $\tilde{\lambda}$ ,  $\forall k, h, l \geq 1 : k+h+l=d$ ,  
define triangle invariants

$$\tau_{khl}(x_T, y_T, z_T) := \log \left( \frac{\xi(x_T)^{h+1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l-1} \dots}{\xi(x_T)^{h-1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l+1} \dots} \right) \in \mathbb{R}$$

and shears between triangles  $T, T'$   $\forall i \in \{1, \dots, d-1\}$


$$\sigma_i(T, T') := \log \left( - \frac{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_T)^1 \dots}{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_{T'})^1 \dots} \right) \in \mathbb{R}$$

$\Rightarrow (\sigma, \tau)$  are a  $\lambda$ -cocyclic pair, i.e.

i) symmetries:  $\sigma_i(T, T') = \sigma_{d-i}(T', T)$ ,  $\tau_{khl}(x_T, y_T, z_T) = \tau_{k'l'h}(y_T, z_T, x_T)$ ,

ii)  $\Gamma$ -invariance:  $\sigma_i(\gamma T, \gamma T') = \sigma_i(T, T')$ ,  $\tau_{khl}(\gamma x_T, \gamma y_T, \gamma z_T) = \tau_{khl}(x_T, y_T, z_T)$ ,

iii) cocycle prop:  $\sigma_i(T, T'') = \sigma_i(T, T') + \sigma_i(T', T'') + \left( \begin{array}{c} \text{sum of some} \\ \text{triangle invariants} \\ \text{of } T \\ (=0 \text{ when } d=2) \end{array} \right)$



# Fock-Goncharov '06, Bonahon-Dreyer '17 (for PSLd IR)

Fix  $\lambda$  max lam.  $\nabla T$  triangle of  $\tilde{\lambda}$ ,  $\forall k, h, l \geq 1 : k+h+l=d$ ,  
define triangle invariants

$$\tau_{khl}(x_T, y_T, z_T) := \log \left( \frac{\xi(x_T)^{h+1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l-1} \dots}{\xi(x_T)^{h-1} \wedge \xi(y_T)^k \wedge \xi(z_T)^{l+1} \dots} \right) \in \mathbb{R}$$

and shears between triangles  $T, T' \quad \forall i \in \{1, \dots, d-1\}$


$$\sigma_i(T, T') := \log \left( - \frac{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_T)^1}{\xi(x_T)^i \wedge \xi(y_T)^{d-i-1} \wedge \xi(z_{T'})^1} \dots \right) \in \mathbb{R}$$

$\Rightarrow (\sigma, \tau)$  are a  $\lambda$ -cocyclic pair, i.e.

i) symmetries:  $\sigma_i(T, T') = \sigma_{d-i}(T', T)$ ,  $\tau_{khl}(x_T, y_T, z_T) = \tau_{k'l'h}(y_T, z_T, x_T)$ ,

ii)  $\Gamma$ -invariance:  $\sigma_i(\gamma T, \gamma T') = \sigma_i(T, T')$ ,  $\tau_{khl}(\gamma x_T, \gamma y_T, \gamma z_T) = \tau_{khl}(x_T, y_T, z_T)$ ,

iii) cocycle prop:  $\sigma_i(T, T'') = \sigma_i(T, T') + \sigma_i(T', T'') + \left( \begin{array}{c} \text{sum of some} \\ \text{triangle invariants} \left( \begin{array}{c} = 0 \text{ when} \\ d=2 \end{array} \right) \end{array} \right)$



## Shear coord's for $\text{Hit}_d(S)$

Thm (Bonahon-Dreyer '17, Bonahon '96 if  $d=2$ )  $\forall d \geq 2 \quad \forall \lambda \max \text{lam}$  the map

$$\varphi_d^\lambda : \text{Hit}_d(S) \longrightarrow \mathcal{C}(\lambda, d, \text{IR}) \subset \left\{ \begin{array}{l} \text{R-valued} \\ \lambda\text{-cocyclic pairs} \end{array} \right\}$$

$$[\rho] \mapsto (\sigma^\rho, \tau^\rho)$$

is a real analytic diffeomorphism, whose image  $= \mathcal{C}(\lambda, d, \text{IR}) = \bigcap_{k<\infty}$  half-spaces is determined by suitable length functions.

Q: What about  $\text{PSL}_2 \mathbb{C}$ ?

$G = \text{PSL}_2 \mathbb{C}$ : Bonahon '96 constructed shear-bend coord's on  $\mathcal{R}_\lambda$ , with  $\text{QF}(S) \subset \mathcal{R}_\lambda \not\subset \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$ , via equivariant pleated surfaces

## Shear coord's for $\text{Hit}_d(S)$

Thm (Bonahon-Dreyer '17, Bonahon '96 if  $d=2$ )  $\forall d \geq 2 \quad \forall \lambda \max \text{lam}$  the map

$$\varphi_d^\lambda : \text{Hit}_d(S) \longrightarrow \mathcal{C}(\lambda, d, \text{IR}) \subset \left\{ \begin{array}{l} \text{R-valued} \\ \lambda\text{-cocyclic pairs} \end{array} \right\}$$

$$[\rho] \mapsto (\sigma^\rho, \tau^\rho)$$

is a real analytic diffeomorphism, whose image  $= \mathcal{C}(\lambda, d, \text{IR}) = \bigcap_{k < \infty}$  half-spaces is determined by suitable length functions.

Q: What about  $\text{PSL}_2 \mathbb{C}$ ?

$G = \text{PSL}_2 \mathbb{C}$ : Bonahon '96 constructed shear-bend coord's on  $\mathcal{R}_\lambda$ , with  $\text{QF}(S) \subset \mathcal{R}_\lambda \not\subset \mathcal{X}_{\text{d.f.}}(\Gamma, \text{PSL}_2 \mathbb{C})$ , via equivariant pleated surfaces



Credits: Jeffrey Brock, Mickey Mouse example

## Pleated surfaces in higher rank

Dfn (MMMZ) Let  $d \geq 2$ ,  $\lambda$  max lam. A  $d$ -pleated surface with pleating locus  $\lambda$

is a pair  $(\rho: \Gamma \rightarrow \mathrm{PSL}_d \mathbb{C}, \xi: \tilde{\lambda} \subset \partial \Gamma \rightarrow \mathrm{Flags}(\mathbb{C}^d))$  satisfying:

- $\xi$   $\rho$ -equivariant,  $\xi_{\tilde{\lambda}}: \tilde{\lambda} \rightarrow \mathrm{Flags}(\mathbb{C}^d)^2$  (locally Hölder) continuous,
- $\xi$   $\lambda$ -transverse and  $\lambda$ -generic,
- $\rho$  is  $\lambda$ -Borel Anosov, i.e. the lift of the geodesic flow  $(\psi_t)_t$  over

$$T^1 \lambda \times_{\rho} \mathbb{C}^d := T^1 \tilde{\lambda} \times \mathbb{C}^d /_{(v, X) \sim (\gamma v, \rho(\gamma)X)} \rightarrow T^1 \lambda$$

uniformly contracts/expands in the directions prescribed by  $\xi$ .

Let

$$\tilde{\mathcal{R}}_d^\lambda := \left\{ \rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_d \mathbb{C}) \mid \begin{array}{l} \text{$\rho$ holonomy of a $d$-pleated} \\ \text{surface with pleating locus $\lambda$} \end{array} \right\}$$

## Pleated surfaces in higher rank

Dfn (MMMZ) Let  $d \geq 2$ ,  $\lambda$  max lam. A  $d$ -pleated surface with pleating locus  $\lambda$

is a pair  $(\rho: \Gamma \rightarrow \mathrm{PSL}_d \mathbb{C}, \xi: \tilde{\lambda} \subset \partial \Gamma \rightarrow \mathrm{Flags}(\mathbb{C}^d))$  satisfying:

- $\xi$   $\rho$ -equivariant,  $\xi_{\tilde{\lambda}}: \tilde{\lambda}^{\text{or}} \rightarrow \mathrm{Flags}(\mathbb{C}^d)^2$  (locally Hölder) continuous,
- $\xi$   $\lambda$ -transverse and  $\lambda$ -generic,
- $\rho$  is  $\lambda$ -Borel Anosov, i.e. the lift of the geodesic flow  $(\psi_t)_t$  over

$$T^1 \lambda \times_{\rho} \mathbb{C}^d := T^1 \tilde{\lambda} \times \mathbb{C}^d /_{(v, X) \sim (\gamma v, \rho(\gamma)X)} \rightarrow T^1 \lambda$$

uniformly contracts/expands in the directions prescribed by  $\xi$ .

Let

$$\tilde{\mathcal{R}}_d^\lambda := \left\{ \rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_d \mathbb{C}) \mid \begin{array}{l} \rho \text{ holonomy of a } d\text{-pleated} \\ \text{surface with pleating locus } \lambda \end{array} \right\}$$

## Pleated surfaces in higher rank

Dfn (MMMZ) Let  $d \geq 2$ ,  $\lambda$  max lam. A  $d$ -pleated surface with pleating locus  $\lambda$

is a pair  $(\rho: \Gamma \rightarrow \mathrm{PSL}_d \mathbb{C}, \xi: \partial \tilde{\lambda} \subset \partial \Gamma \rightarrow \mathrm{Flags}(\mathbb{C}^d))$  satisfying:

- $\xi$   $\rho$ -equivariant,  $\xi_{\tilde{\lambda}}: \tilde{\lambda}^{\text{or}} \rightarrow \mathrm{Flags}(\mathbb{C}^d)^2$  (locally Hölder) continuous,
- $\xi$   $\lambda$ -transverse and  $\lambda$ -generic,
- $\rho$  is  $\lambda$ -Borel Anosov, i.e. the lift of the geodesic flow  $(\psi_t)_t$  over

$$T^1 \lambda \times_{\rho} \mathbb{C}^d := T^1 \tilde{\lambda} \times \mathbb{C}^d /_{(v, X) \sim (\gamma_v, \rho(\gamma)X)} \rightarrow T^1 \lambda$$

uniformly contracts/expands in the directions prescribed by  $\xi$ .

Let

$$\tilde{\mathcal{R}}_d^\lambda := \left\{ \rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_d \mathbb{C}) \mid \begin{array}{l} \rho \text{ holonomy of a } d\text{-pleated} \\ \text{surface with pleating locus } \lambda \end{array} \right\}$$

## Pleated surfaces in higher rank

Dfn (MMMZ) Let  $d \geq 2$ ,  $\lambda$  max lam. A  $d$ -pleated surface with pleating locus  $\lambda$

is a pair  $(\rho: \Gamma \rightarrow \mathrm{PSL}_d \mathbb{C}, \xi: \partial \tilde{\lambda} \subset \partial \Gamma \rightarrow \mathrm{Flags}(\mathbb{C}^d))$  satisfying:

- $\xi$   $\rho$ -equivariant,  $\xi_{\tilde{\lambda}}: \tilde{\lambda}^{\text{or}} \rightarrow \mathrm{Flags}(\mathbb{C}^d)^2$  (locally Hölder) continuous,
- $\xi$   $\lambda$ -transverse and  $\lambda$ -generic,
- $\rho$  is  $\lambda$ -Borel Anosov, i.e. the lift of the geodesic flow  $(\psi_t)_t$  over

$$T^1 \lambda \times_{\rho} \mathbb{C}^d := T^1 \tilde{\lambda} \times \mathbb{C}^d /_{(v, X) \sim (\gamma v, \rho(\gamma)X)} \rightarrow T^1 \lambda$$

uniformly contracts/expands in the directions prescribed by  $\xi$ .

Let

$$\tilde{\mathcal{R}}_d^\lambda := \left\{ \rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_d \mathbb{C}) \mid \begin{array}{l} \rho \text{ holonomy of a } d\text{-pleated} \\ \text{surface with pleating locus } \lambda \end{array} \right\}$$

## Pleated surfaces in higher rank

Dfn (MMMZ) Let  $d \geq 2$ ,  $\lambda$  max lam. A  $d$ -pleated surface with pleating locus  $\lambda$

is a pair  $(\rho: \Gamma \rightarrow \mathrm{PSL}_d \mathbb{C}, \xi: \partial \tilde{\lambda} \subset \partial \Gamma \rightarrow \mathrm{Flags}(\mathbb{C}^d))$  satisfying:

- $\xi$   $\rho$ -equivariant,  $\xi_{\tilde{\lambda}}: \tilde{\lambda}^{\text{or}} \rightarrow \mathrm{Flags}(\mathbb{C}^d)^2$  (locally Hölder) continuous,
- $\xi$   $\lambda$ -transverse and  $\lambda$ -generic,
- $\rho$  is  $\lambda$ -Borel Anosov, i.e. the lift of the geodesic flow  $(\psi_t)_t$  over

$$T^1 \lambda \times_{\rho} \mathbb{C}^d := T^1 \tilde{\lambda} \times \mathbb{C}^d /_{(v, X) \sim (\gamma v, \rho(\gamma)X)} \rightarrow T^1 \lambda$$

uniformly contracts/expands in the directions prescribed by  $\xi$ .

Let  $\tilde{\mathcal{R}}_d^\lambda := \left\{ \rho \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_d \mathbb{C}) \mid \begin{array}{l} \rho \text{ holonomy of a } d\text{-pleated} \\ \text{surface with pleating locus } \lambda \end{array} \right\}$

## Main result

Given  $(\rho, \xi)$   $d$ -pleated surface, let

$$\sigma_i(T, T') = \log \left( \frac{\text{same expression}}{\text{as Bonahon-Dreyer}} \right), \quad \tau_{hkl}(x_T, y_T, z_T) = \log \left( \frac{\text{same expression}}{\text{as Bonahon-Dreyer}} \right) \in \mathbb{C}/2\pi i \mathbb{Z}$$

Rmk  $(\sigma, \tau)$  well defined since  $\xi$  is  $\lambda$ -transverse &  $\lambda$ -generic

$$C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) := \{(\sigma, \tau) \in \mathcal{Y}(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \mid (\operatorname{Re} \sigma, \operatorname{Re} \tau) \in \varphi_d^\lambda(\operatorname{Hit}_d(S))\}$$

Thm (MMMZ, Bonahon '96 for  $d=2$ ) If  $d \geq 2$ , if  $\lambda$  max fam, the map

$$\begin{aligned} \psi_d^\lambda: \tilde{\mathcal{R}}_d^\lambda &\longrightarrow C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \times \mathbb{X}_d \leftarrow \text{affine space over } \operatorname{PSL} d \mathbb{C} \\ \rho &\mapsto (\sigma^\rho, \tau^\rho; \xi(x_{T_0}), \xi(y_{T_0}), \xi(z_{T_0})') \end{aligned}$$

is a biholomorphism.

## Main result

Given  $(\rho, \xi)$   $d$ -pleated surface, let

$$\sigma_i(T, T') = \log \left( \begin{array}{l} \text{same expression} \\ \text{as Bonahon-Dreyer} \end{array} \right), \quad \tau_{hkl}(x_T, y_T, z_T) = \log \left( \begin{array}{l} \text{same expression} \\ \text{as Bonahon-Dreyer} \end{array} \right) \in \mathbb{C}/2\pi i \mathbb{Z}$$

Rmk  $(\sigma, \tau)$  well defined since  $\xi$  is  $\lambda$ -transverse &  $\lambda$ -generic

$$C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) := \{(\sigma, \tau) \in Y(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \mid (\operatorname{Re} \sigma, \operatorname{Re} \tau) \in \varphi_d^{-1}(\operatorname{Hit}_d(S))\}$$

Thm (MMMZ, Bonahon '96 for  $d=2$ ) If  $d \geq 2$ , if  $\lambda$  max fam, the map

$$\begin{aligned} \psi_d^\lambda: \tilde{\mathcal{R}}_d^\lambda &\longrightarrow C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \times \mathbb{X}_d \leftarrow \text{affine space over } \operatorname{PSL} d \mathbb{C} \\ \rho &\mapsto (\sigma^\rho, \tau^\rho; \xi(x_{T_0}), \xi(y_{T_0}), \xi(z_{T_0})^*) \end{aligned}$$

is a biholomorphism.

## Main result

Given  $(\rho, \xi)$   $d$ -pleated surface, let

$$\sigma_i(T, T') = \log \left( \begin{array}{l} \text{same expression} \\ \text{as Bonahon-Dreyer} \end{array} \right), \quad \tau_{hkl}(x_T, y_T, z_T) = \log \left( \begin{array}{l} \text{same expression} \\ \text{as Bonahon-Dreyer} \end{array} \right) \in \mathbb{C}/2\pi i \mathbb{Z}$$

Rmk  $(\sigma, \tau)$  well defined since  $\xi$  is  $\lambda$ -transverse &  $\lambda$ -generic

$$C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) := \{(\sigma, \tau) \in Y(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \mid (\operatorname{Re} \sigma, \operatorname{Re} \tau) \in \varphi_d^1(\operatorname{Hit}_d(S))\}$$

Thm (MMMZ, Bonahon '96 for  $d=2$ ) If  $d \geq 2$ , if  $\lambda$  max fam, the map

$$\psi_d^\lambda: \tilde{\mathcal{R}}_d^\lambda \rightarrow C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z}) \times \mathbb{X}_d \quad \text{affine space over } \operatorname{PSL}_d \mathbb{C}$$

$$\rho \mapsto (\sigma^\rho, \tau^\rho; \xi(x_{T_0}), \xi(y_{T_0}), \xi(z_{T_0})')$$

is a biholomorphism.

## Steps of the proof :

i)  $\forall (\xi, p)$  d-pleated surface, its shear-bend coord's  $(\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$

( extend length functions  $l_i^R$  from Dreyer, Bonahon-Dreyer to  $l_i^C$  +  
homological interpretation of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$  )

ii)  $\psi_d^\lambda$  surjective (hard!) :  $\forall (\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ , pick  $p_0 \in \text{Hit}_n(S)$  such that  
 $\psi_d^\lambda(p_0) = (\text{Re}\sigma, \text{Re}\tau)$ . Then "bend"  $(p_0, \xi_0)$  to obtain  $(p, \xi)$  s.t.

$$\psi_d^\lambda(p, \xi) = (\text{Re}\sigma + i \text{Im}\sigma, \text{Re}\tau + i \text{Im}\tau)$$

(Generalizes the case  $d=2$ , requires notion of eruption from Wienhard - Zhang - Zhu)

iii)  $\psi_d^\lambda$  injective + holomorphic : The key ingredient is

$$\forall r \in \Gamma \quad p(r) = \lim_{n \rightarrow \infty} \left( \begin{array}{l} \text{explicit sequence of matrices that} \\ \text{depend holomorphically on } (\sigma, \tau) \end{array} \right)$$

## Steps of the proof :

i)  $\forall (\xi, p)$  d-pleated surface, its shear-bend coord's  $(\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$

(extend length functions  $l_i^R$  from Dreyer, Bonahon-Dreyer to  $l_i^C$  +  
homological interpretation of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ )

ii)  $\psi_d^\lambda$  surjective (hard!) :  $\forall (\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ , pick  $p_0 \in \text{Hit}_n(S)$  such that  
 $\psi_d^\lambda(p_0) = (\text{Re}\sigma, \text{Re}\tau)$ . Then "bend"  $(p_0, \xi_0)$  to obtain  $(p, \xi)$  s.t.

$$\psi_d^\lambda(p, \xi) = (\text{Re}\sigma + i \text{Im}\sigma, \text{Re}\tau + i \text{Im}\tau)$$

(Generalizes the case  $d=2$ , requires notion of eruption from Wienhard-Zhang-Zhu)

iii)  $\psi_d^\lambda$  injective + holomorphic : The key ingredient is

$$\forall r \in \Gamma \quad p(r) = \lim_{n \rightarrow \infty} \left( \begin{array}{l} \text{explicit sequence of matrices that} \\ \text{depend holomorphically on } (\sigma, \tau) \end{array} \right)$$

## Steps of the proof :

i)  $\forall (\xi, p)$  d-pleated surface, its shear-bend coord's  $(\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$

(extend length functions  $l_i^R$  from Dreyer, Bonahon-Dreyer to  $l_i^C$  +  
homological interpretation of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ )

ii)  $\psi_d^\lambda$  surjective (hard!) :  $\forall (\sigma, \tau) \in C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ , pick  $p_0 \in \text{Hit}_n(S)$  such that  
 $\psi_d^\lambda(p_0) = (\text{Re}\sigma, \text{Re}\tau)$ . Then "bend"  $(p_0, \xi_0)$  to obtain  $(p, \xi)$  s.t.

$$\psi_d^\lambda(p, \xi) = (\text{Re}\sigma + i \text{Im}\sigma, \text{Re}\tau + i \text{Im}\tau)$$

(Generalizes the case  $d=2$ , requires notion of eruption from Wienhard-Zhang-Zhu)

iii)  $\psi_d^\lambda$  injective + holomorphic : The key ingredient is

$$\forall r \in \Gamma \quad p(r) = \lim_{n \rightarrow \infty} \left( \begin{array}{l} \text{explicit sequence of matrices that} \\ \text{depend holomorphically on } (\sigma, \tau) \end{array} \right)$$

Thank you for your  
attention !

## Remarks/questions

- $\forall d \geq 2$  and  $\forall \lambda$  (finite) max lam,  $\exists [\rho] \in \tilde{\mathcal{R}}_d^\lambda$  non-discrete.
- if  $d > 2$  and  $\lambda$  finite max lam, then  $\tilde{\mathcal{R}}_d^\lambda \subset \text{Hom}(\Gamma, \text{PSL}_d \mathbb{C})$  contains reducible representations, and

$$\mathcal{R}_d^\lambda := \tilde{\mathcal{R}}_d^\lambda / \text{PSL}_d \mathbb{C} \subset \mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$$

contains non-Hausdorff points of  $\mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$ . Can we characterize them in terms of shear-bend coord's?

- What is the topology of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ ? It is an open subset of a cplx Abelian Lie group. Is it connected? (When  $d=2$  it has 2 connected components, see Bonahon '96)
- already when  $d=2$ , not clear how to characterize  $QF(S) \subset \mathcal{R}_2^\lambda \forall \lambda$  in terms of shear-bend coord's.

What properties does  $\rho \in \bigcap_{\lambda \text{ max lam}} \mathcal{R}_d^\lambda$  satisfy?

## Remarks/questions

- $\forall d \geq 2$  and  $\forall \lambda$  (finite) max lam,  $\exists [\rho] \in \tilde{\mathcal{R}}_d^\lambda$  non-discrete.
- if  $d > 2$  and  $\lambda$  finite max lam, then  $\tilde{\mathcal{R}}_d^\lambda \subset \text{Hom}(\Gamma, \text{PSL}_d \mathbb{C})$  contains reducible representations, and

$$\mathcal{R}_d^\lambda := \tilde{\mathcal{R}}_d^\lambda / \text{PSL}_d \mathbb{C} \subset \mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$$

contains non-Hausdorff points of  $\mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$ . Can we characterize them in terms of shear-bend coord's?

- What is the topology of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ ? It is an open subset of a cplx Abelian Lie group. Is it connected? (When  $d=2$  it has 2 connected components, see Bonahon '96)
- already when  $d=2$ , not clear how to characterize  $QF(S) \cap \mathcal{R}_2^\lambda \forall \lambda$  in terms of shear-bend coord's.

What properties does  $\rho \in \bigcap_{\lambda \text{ max lam}} \mathcal{R}_d^\lambda$  satisfy?

## Remarks/questions

- $\forall d \geq 2$  and  $\forall \lambda$  (finite) max lam,  $\exists [\rho] \in \tilde{\mathcal{R}}_d^\lambda$  non-discrete.
- if  $d > 2$  and  $\lambda$  finite max lam, then  $\tilde{\mathcal{R}}_d^\lambda \subset \text{Hom}(\Gamma, \text{PSL}_d \mathbb{C})$  contains reducible representations, and

$$\mathcal{R}_d^\lambda := \tilde{\mathcal{R}}_d^\lambda / \text{PSL}_d \mathbb{C} \subset \mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$$

contains non-Hausdorff points of  $\mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$ . Can we characterize them in terms of shear-bend coord's?

- What is the topology of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ ? It is an open subset of a cplx Abelian Lie group. Is it connected? (When  $d=2$  it has 2 connected components, see Bonahon '96)
- already when  $d=2$ , not clear how to characterize  $QF(S) \cap \mathcal{R}_2^\lambda \forall \lambda$  in terms of shear-bend coord's.

What properties does  $\rho \in \bigcap_{\lambda \text{ max lam}} \mathcal{R}_d^\lambda$  satisfy?

## Remarks/questions

- $\forall d \geq 2$  and  $\forall \lambda$  (finite) max lam,  $\exists [\rho] \in \tilde{\mathcal{R}}_d^\lambda$  non-discrete.
- if  $d > 2$  and  $\lambda$  finite max lam, then  $\tilde{\mathcal{R}}_d^\lambda \subset \text{Hom}(\Gamma, \text{PSL}_d \mathbb{C})$  contains reducible representations, and

$$\mathcal{R}_d^\lambda := \tilde{\mathcal{R}}_d^\lambda / \text{PSL}_d \mathbb{C} \subset \mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$$

contains non-Hausdorff points of  $\mathcal{X}(\Gamma, \text{PSL}_d \mathbb{C})$ . Can we characterize them in terms of shear-bend coord's?

- What is the topology of  $C(\lambda, d, \mathbb{C}/2\pi i \mathbb{Z})$ ? It is an open subset of a cplx Abelian Lie group. Is it connected? (When  $d=2$  it has 2 connected components, see Bonahon '96)
- already when  $d=2$ , not clear how to characterize  $QF(S) \cap \mathcal{R}_2^\lambda \forall \lambda$  in terms of shear-bend coord's.

What properties does  $\rho \in \bigcap_{\lambda \text{ max lam}} \mathcal{R}_d^\lambda$  satisfy?