

Pleated Surfaces for $SO_0(2, n)$ -maximal representations

Filippo Mazzoli (University of Virginia),
joint work with Gabriele Viaggi (Heidelberg University).




Geometry Seminar, University of Virginia

April 5, 2022.

Outline

0. Motivation
1. The geometry of $H^{2,n}$
2. $SO_0(2,n+1)$ -maximal representations
3. Pleated surfaces in $H^{2,n}$
4. Some applications




Pleated surfaces (in hyp 3-mfld N)

Def $f: (\Sigma, h) \rightarrow N$ path-isometry such that

- i) (Σ, h) hyp surface, N hyp 3-mfld
- ii) \exists max geodesic lamination λ (closed subset foliated by geodesics) s.t. $\forall g \in \lambda$
 $f(g)$ geodesic in N

and $\forall T \subset S \setminus \lambda$

$f|_T$ totally geodesic immersion in N




Pleated surfaces (in hyp 3-mfld N)

Def $f: (\Sigma, h) \rightarrow N$ path-isometry such that

- i) (Σ, h) hyp surface, N hyp 3-mfld
- ii) \exists max geodesic lamination λ (closed subset foliated by geodesics) s.t. $\forall g \in \lambda$
 $f(g)$ geodesic in N

and $\forall T \subset \Sigma \setminus \lambda$

$f|_T$ totally geodesic immersion in N



Pleated surfaces (in hyp 3-mfld N)


Def $f: (\Sigma, h) \rightarrow N$ path-isometry such that

- i) (Σ, h) hyp surface, N hyp 3-mfld
- ii) \exists max geodesic lamination λ (closed subset foliated by geodesics) s.t. $\forall g \in \lambda$

$f(g)$ geodesic in N

and $\forall T \subset S \setminus \lambda$

$f|_T$ totally geodesic immersion in N



Thm (Thurston) Let $u: \Sigma \rightarrow M$ map s.t. $\Gamma \cong u_*(\Gamma)$ has no parabolics,
then for (almost every) max geod lamination λ \exists pleated surface
homotopic to u with pleating locum $\subseteq \lambda$.

Many applications to Thurston's geometrization, Ending lamination,
quasi-Fuchsian mflds & their convex core, ...

Q: Is there a good notion of pleated surfaces for $(\mathbb{H}^{2,n}, SO_0(2,n+1))$ -manifolds
with maximal holonomy?
?

Thm (Thurston) Let $u: \Sigma \rightarrow M$ map s.t. $\Gamma \cong u_*(\Gamma)$ has no parabolics,
then for (almost every) max geod lamination $\lambda \exists$ pleated surface
homotopic to u with pleating locum $\subseteq \lambda$.

Many applications to Thurston's geometrization, Ending lamination,
quasi-Fuchsian mflds & their convex core, ...

Q: Is there a good notion of pleated surfaces for $(\mathbb{H}^{2,n}, SO_0(2,n+1))$ -manifolds
with maximal holonomy?
?

Thm (Thurston) Let $u: \Sigma \rightarrow M$ map s.t. $\Gamma \cong u_*(\Gamma)$ has no parabolics,
then for (almost every) max geod lamination $\lambda \exists$ pleated surface
homotopic to u with pleating locum $\subseteq \lambda$.

Many applications to Thurston's geometrization, Ending lamination,
quasi-Fuchsian mflds & their convex core, ...

Q: Is there a good notion of pleated surfaces for $(\mathbb{H}^{2,n}, SO_0(2,n+1))$ -manifolds
with maximal holonomy?
?


Geometry in $\mathbb{H}^{2,n}$

$$\langle x, x \rangle_{(2,n+1)} := x_1^2 + x_2^2 - x_3^2 - \cdots - x_{n+3}^2$$

$$\mathbb{H}^{2,n} := P(\{x \in \mathbb{R}^{2,n+1} \mid \langle x, x \rangle_{(2,n+1)} < 0\}) \subset \mathbb{R}P^{n+2}$$

Pseudo-Riemannian mfd of signature $(2,n)$

$$Ein^{1,n} := \partial \mathbb{H}^{2,n} = P(\{x \in \mathbb{R}^{2,n+1} \mid \langle x, x \rangle_{(2,n+1)} = 0\})$$



($n=1$ in affine chart)

Ideal boundary

$$Isom_0(\mathbb{H}^{2,n}) = SO_0(2,n+1) \curvearrowright \mathbb{H}^{2,n} \text{ & } Ein^{1,n} \quad (\text{homogeneous spaces})$$


Geometry in $H^{2,n}$

$$\langle x, x \rangle_{(2,n+1)} := x_1^2 + x_2^2 - x_3^2 - \cdots - x_{n+1}^2$$

$$H^{2,n} := P(\{x \in \mathbb{R}^{2,n+1} \mid \langle x, x \rangle_{(2,n+1)} < 0\}) \subset \mathbb{R}P^{n+2}$$

Pseudo-Riemannian mfd of signature $(2,n)$

$$Ein^{1,n} := \partial H^{2,n} = P(\{x \in \mathbb{R}^{2,n+1} \mid \langle x, x \rangle_{(2,n+1)} = 0\})$$



($n=1$ in affine chart)

Ideal boundary

$$Isom_0(H^{2,n}) = SO_0(2,n+1) \cap H^{2,n} \text{ & } Ein^{1,n} \quad (\text{homogeneous spaces})$$

1-dim: geodesics $\ell = PV$, for some $V \in \mathbb{R}^{2,n+1}$
with $\dim = 2$

- * ℓ is space-like if $\langle \ell'(t), \ell'(t) \rangle > 0$
- * ℓ is light-like if $\langle \ell'(t), \ell'(t) \rangle = 0$
- * ℓ is timelike if $\langle \ell'(t), \ell'(t) \rangle < 0$


2-dim: totally geodesic space-like planar $H = PW$, $\dim W = 3$
and $\langle \cdot, \cdot \rangle|_H > 0$

$$\Rightarrow H \cong_{\text{isom}} \mathbb{H}^2$$

If $x, y \in \mathbb{H}^{2,n}$ joined by a space-like geodesic $[x, y]$, then

$$d_{\mathbb{H}^{2,n}}(x, y) := L([x, y]) \quad \leftarrow \text{Alert! No triangle inequality!!}$$

Subspaces



1-dim: geodesics $\ell = PV$, for some $V \in \mathbb{R}^{2,n+1}$
with $\dim = 2$

- * ℓ is space-like if $\langle \ell'(t), \ell'(t) \rangle > 0$
- * ℓ is light-like if $\langle \ell'(t), \ell'(t) \rangle = 0$
- * ℓ is timelike if $\langle \ell'(t), \ell'(t) \rangle < 0$


2-dim: totally geodesic space-like planar $H = PW$, $\dim W = 3$
and $\langle \cdot, \cdot \rangle|_H > 0$

$$\Rightarrow H \simeq_{\text{isom}} \mathbb{H}^2$$

If $x, y \in \mathbb{H}^{2,n}$ joined by a space-like geodesic $[x, y]$, then

$$d_{\mathbb{H}^{2,n}}(x, y) := L([x, y]) \quad \leftarrow \text{Alert! No triangle inequality!!}$$

Subspaces



1-dim: geodesics $\ell = PV$, for some $V \in \mathbb{R}^{2,n+1}$
with $\dim = 2$

- * ℓ is space-like if $\langle \ell'(t), \ell'(t) \rangle > 0$
- * ℓ is light-like if $\langle \ell'(t), \ell'(t) \rangle = 0$
- * ℓ is timelike if $\langle \ell'(t), \ell'(t) \rangle < 0$


2-dim: totally geodesic space-like planar $H = PW$, $\dim W = 3$
and $\langle \cdot, \cdot \rangle|_H > 0$

$$\Rightarrow H \simeq_{\text{isom}} \mathbb{H}^2$$

If $x, y \in \mathbb{H}^{2,n}$ joined by a space-like geodesic $[x, y]$, then

$$d_{\mathbb{H}^{2,n}}(x, y) := L([x, y]) \quad \leftarrow \text{Alert! No triang ineq!!}$$

Subspaces



Maximal representations


Wide class of surface groups repr's in G Lie group of Hermitian type

Here $G := SO_0(2, n+1)$, $\Gamma := \pi_1(\Sigma_g)$, $g \geq 2$

Thm (Burger - Iozzi - Labourie - Wienhard) $\rho: \Gamma \rightarrow SO_0(2, n+1)$ max

iff $\exists \xi: \partial\Gamma \rightarrow \text{Ein}^{1,n}$ ρ -equiv homeo + spacelike, i.e.

$\forall x, y, z \in \partial\Gamma$ distinct $\exists H$ spacelike plane in $H^{2,n}$
s.t. $\xi(x), \xi(y), \xi(z) \in \partial H$



Maximal representations


Wide class of surface groups repr's in G Lie group of Hermitian type

Here $G := SO_0(2, n+1)$, $\Gamma := \pi_1(\Sigma_g)$, $g \geq 2$

Thm (Burger - Iozzi - Labourie - Wienhard) $\rho: \Gamma \rightarrow SO_0(2, n+1)$ max

iff $\exists \xi: \partial\Gamma \rightarrow \text{Ein}^{1,n}$ ρ -equiv homeo + space-like, i.e.


$\forall x, y, z \in \partial\Gamma$ distinct $\exists H$ space-like plane in $H^{2,n}$
s.t. $\xi(x), \xi(y), \xi(z) \in \partial H$



Example:

$n=0$: $(\mathbb{H}^2, SO_0(2,1))$ $\rho: \Gamma \rightarrow \text{Isom}^+(\mathbb{H}^2)$ maximal iff

ρ Fuchsian, i.e. $\rho(\Gamma) \backslash \mathbb{H}^2$ hyperbolic surface



$n=1$: $(\mathbb{H}^{2,1}, SO_0(2,2)) = (\text{AdS}^3, \text{Isom}_0(\text{AdS}^3) \cong \text{PSL}_2\mathbb{R} \times \text{PSL}_2\mathbb{R})$

$\rho = (\rho_e, \rho_r): \Gamma \rightarrow \text{PSL}_2\mathbb{R} \times \text{PSL}_2\mathbb{R}$ maximal iff

both ρ_e & ρ_r are Fuchsian




$\forall n$: $\exists C_p$ closed ρ -equiv convex s.t. $\Gamma \xrightarrow{\rho} C_p$ cocompact, $M_\rho := C_p / \rho(\Gamma)$

Let's go back to the original def...

Example:

$n=0$: $(\mathbb{H}^2, SO_0(2,1))$ $\rho: \Gamma \rightarrow \text{Isom}^+(\mathbb{H}^2)$ maximal iff


ρ Fuchsian, i.e. $\rho(\Gamma) \backslash \mathbb{H}^2$ hyperbolic surface



$n=1$: $(\mathbb{H}^{2,1}, SO_0(2,2)) = (AdS^3, \text{Isom}_0(AdS^3) \cong PSL_2\mathbb{R} \times PSL_2\mathbb{R})$

$\rho = (\rho_e, \rho_r): \Gamma \rightarrow PSL_2\mathbb{R} \times PSL_2\mathbb{R}$ maximal iff

both ρ_e & ρ_r are Fuchsian




$\forall n$: $\exists C_\rho$ closed ρ -equiv convex s.t. $\Gamma \xrightarrow{\rho} C_\rho$ cocompact, $M_\rho := C_\rho / \rho(\Gamma)$

Let's go back to the original def...

Example:

$n=0$: $(\mathbb{H}^2, SO_0(2,1))$ $\rho: \Gamma \rightarrow \text{Isom}^+(\mathbb{H}^2)$ maximal iff


ρ Fuchsian, i.e. $\rho(\Gamma) \backslash \mathbb{H}^2$ hyperbolic surface



$n=1$: $(\mathbb{H}^{2,1}, SO_0(2,2)) = (AdS^3, \text{Isom}_0(AdS^3) \cong PSL_2\mathbb{R} \times PSL_2\mathbb{R})$

$\rho = (\rho_e, \rho_r): \Gamma \rightarrow PSL_2\mathbb{R} \times PSL_2\mathbb{R}$ maximal iff

both ρ_e & ρ_r are Fuchsian



$\forall n$: $\exists C_p$ closed ρ -equiv convex s.t. $\Gamma \xrightarrow{\rho} C_p$ cocompact, $M_\rho := C_p / \rho(\Gamma)$

Let's go back to the original def...

Pleated surfaces (in hyp 3-mflds)

Def $f: (\Sigma, h) \rightarrow M$ path-isometry such that \leftarrow main problems!!

i) (Σ, h) hyp surface, M hyp 3-mfd $M = C\rho/\rho(\Gamma)$


ii) \exists max geodesic lamination λ (closed subset

foliated by geodesics) s.t. $\forall g \in \lambda$

$f(g)$ geodesic in M spacelike

and $\forall T \subset S \setminus \lambda$

$f(T)$ totally geodesic immersion spacelike



Pleated surfaces (in hyp 3-mflds)

Def $f: (\Sigma, h) \rightarrow M$ path-isometry such that \leftarrow main problems!!

i) (Σ, h) hyp surface, M hyp 3-mfd $M = C\rho/\rho(\Gamma)$


ii) \exists max geodesic lamination λ (closed subset

foliated by geodesics) s.t. $\forall g \in \lambda$

$f(g)$ geodesic in M spacelike

and $\forall T \in S \setminus \lambda$

$f(T)$ totally geodesic immersion spacelike



Pleated surfaces (in hyp 3-mflds)

Def $f: (\Sigma, h) \rightarrow M$ | path-isometry such that \leftarrow main problems!!

i) (Σ, h) hyp surface, M hyp 3-mfd $M = C\rho/\rho(\Gamma)$


ii) \exists max geodesic lamination λ (closed subset

foliated by geodesics) s.t. $\forall g \in \lambda$

$f(g)$ geodesic in M spacelike

and $\forall T \in S \setminus \lambda$

$f(T)$ totally geodesic immersion spacelike



No path metric $c \Rightarrow$ how to get a hyp structure on Σ ?

Main tool: p max repr, ξ limit map $\rightsquigarrow \beta^P$ cross ratio

$$\beta^P(u, v, w, z) = \left(\frac{\langle \xi u, \xi w \rangle_{(z, n+1)}}{\langle \xi u, \xi z \rangle_{(z, n+1)}} \frac{\langle \xi v, \xi z \rangle_{(z, n+1)}}{\langle \xi v, \xi w \rangle_{(z, n+1)}} \right)^{1/2}$$

satisfier:

- $\beta(u, v, w, z) = 0 \Leftrightarrow u = w \text{ or } v = z$
- $\beta(u, u, w, z) = \beta(u, v, w, w) = 1$
- $\beta(u, v, w, z) = \beta(w, z, u, v)$
- $\beta(u, v, w, z) = \beta(u, v, x, z) \beta(u, v, w, x)$
- $|\beta(u, v, w, z) \beta(w, u, v, z) \beta(v, w, u, z)| = 1$

$\left. \begin{array}{l} \beta : \partial P^{(4)} \rightarrow \mathbb{R} \text{ is a cross ratio} \\ \text{if it satisfier these symmetries} \end{array} \right\}$

No path metric $c \Rightarrow$ how to get a hyp structure on Σ ?

Main tool: p max repr, ξ limit map $\rightsquigarrow \beta^p$ cross ratio

$$\beta^p(u, v, w, z) = \left(\frac{\langle \xi u, \xi w \rangle_{(z, n+1)}}{\langle \xi u, \xi z \rangle_{(z, n+1)}} \frac{\langle \xi v, \xi z \rangle_{(z, n+1)}}{\langle \xi v, \xi w \rangle_{(z, n+1)}} \right)^{1/2}$$

satisfier:

- $\beta(u, v, w, z) = 0 \Leftrightarrow u = w \text{ or } v = z$
- $\beta(u, u, w, z) = \beta(u, v, w, w) = 1$
- $\beta(u, v, w, z) = \beta(w, z, u, v)$
- $\beta(u, v, w, z) = \beta(u, v, x, z) \beta(u, v, w, x)$
- $|\beta(u, v, w, z) \beta(w, u, v, z) \beta(v, w, u, z)| = 1$

$\left. \begin{array}{l} \beta : \partial P^{(4)} \rightarrow \mathbb{R} \text{ is a cross ratio} \\ \text{if it satisfier these symmetries} \end{array} \right\}$

No path metric $c \Rightarrow$ how to get a hyp structure on Σ ?

Main tool: p max repr, ξ limit map $\rightsquigarrow \beta^p$ cross ratio

$$\beta^p(u, v, w, z) = \left(\frac{\langle \xi u, \xi w \rangle_{(z, n+1)}}{\langle \xi u, \xi z \rangle_{(z, n+1)}} \frac{\langle \xi v, \xi z \rangle_{(z, n+1)}}{\langle \xi v, \xi w \rangle_{(z, n+1)}} \right)^{1/2}$$

satisfier:

- $\beta(u, v, w, z) = 0 \iff u = w \text{ or } v = z$
- $\beta(u, u, w, z) = \beta(u, v, w, w) = 1$
- $\beta(u, v, w, z) = \beta(w, z, u, v)$
- $\beta(u, v, w, z) = \beta(u, v, x, z) \beta(u, v, w, x)$
- $|\beta(u, v, w, z) \beta(w, u, v, z) \beta(v, w, u, z)| = 1$


$\left. \begin{array}{l} \beta : \partial \Gamma^{(4)} \rightarrow \mathbb{R} \text{ is a cross ratio} \\ \text{if it satisfies these symmetries} \end{array} \right\}$

Why are cross ratios useful?

If λ max lamination and β_0 is the cross ratio on $\partial\tilde{\Sigma} \cong_h \partial\mathbb{H}^2$,

shear between ideal triangles

$$\sigma_\lambda(T, T') := \log |\beta_0(\xi_u, \xi_v, \xi_w, \xi_z)|$$



σ_λ extends to a function $(T, T') \mapsto \sigma_\lambda(T, T')$ on pairs of distinct triangles of λ satisfying:

- $\sigma_\lambda(T, T') = \sigma_\lambda(T', T)$
- $\sigma_\lambda(T, T') = \sigma_\lambda(\gamma T, \gamma T') \quad \forall \gamma \in \pi_1(\Sigma)$
- $\sigma_\lambda(T, T'') = \sigma_\lambda(T, T') + \sigma_\lambda(T', T'') \quad \forall T' \text{ separating } T \text{ & } T''$


σ_λ is a Hölder cocycle!
 $\sigma_\lambda \in \mathcal{H}(\lambda)$

Why are cross ratios useful?

If λ max lamination and β_0 is the cross ratio on $\partial\tilde{\Sigma} \cong_h \partial\mathbb{H}^2$,

shear between ideal triangles

$$\sigma_\lambda(T, T') := \log |\beta_0(\xi_u, \xi_v, \xi_w, \xi_z)|$$



σ_λ extends to a function $(T, T') \mapsto \sigma_\lambda(T, T')$ on pairs of distinct triangles of λ satisfying:

- $\sigma_\lambda(T, T') = \sigma_\lambda(T', T)$
- $\sigma_\lambda(T, T') = \sigma_\lambda(\gamma T, \gamma T') \quad \forall \gamma \in \pi_1(\Sigma)$
- $\sigma_\lambda(T, T'') = \sigma_\lambda(T, T') + \sigma_\lambda(T', T'') \quad \forall T' \text{ separating } T \text{ & } T''$

σ_λ is a Hölder cocycle!
 $\sigma_\lambda \in \mathcal{H}(\lambda)$

Thm(Bonahon '96) If the map

$$\begin{array}{ccc} \{\text{hyp str's on } \Sigma\} / \text{isotopy} & \longrightarrow & G(\lambda) \subset \mathcal{H}(\lambda) \\ (\Sigma, h) & \longmapsto & \sigma_\lambda = \sigma_\lambda^h \end{array}$$

is a diffeo onto its image.

We can generalize the shear cocycle machinery $\beta, \lambda \mapsto \sigma_\lambda^\beta \in \mathcal{H}(\lambda)$ obtaining:

Thm(M-Viaggi) Let β be a strictly positive and locally bounded cross ratio.

Then for all max lam's λ $\exists! X$ hyp surface with shear coord's σ_λ^β

Thm(Bonahon '96) At the map

$$\left\{ \text{hyp str's on } \Sigma \right\} / \text{isotopy} \longrightarrow G(\lambda) \subset \mathcal{H}(\lambda)$$

$$(\Sigma, h) \xrightarrow{\quad} \sigma_\lambda = \sigma_\lambda^h \quad \text{shear coord's}$$

is a diffeo onto its image.

We can generalize the shear cocycle machinery $\beta, \lambda \mapsto \sigma_\lambda^\beta \in \mathcal{H}(\lambda)$ obtaining:


Thm(M-Viaggi) Let β be a strictly positive and locally bounded cross ratio.

Then for all max lam's λ $\exists! X = X_\lambda^\beta$ hyp surface with shear coord's σ_λ^β .


Thm 2 (M-Viaggi) Let $\rho: \Gamma \rightarrow SO_0(2, n+1)$ be a max repr. Then

$\forall \lambda$ max geod lam there exist unique

- $S_\lambda \hookrightarrow C\rho/\rho(\Gamma)$ embedded spacelike surface
- X_λ hyp structure, \leftarrow intrinsic description using cross ratio
- homeo $u: S_\lambda \rightarrow X_\lambda$ such that
 - $u^{-1}(g)$ spacelike geo $\forall g \subset \lambda$ &
 - $u^{-1}(\gamma)$ spacelike triangle $\forall \gamma \subset X_\lambda \cap \lambda$




Moreover $L_{X_\lambda}(\gamma) \leq L_\rho(\gamma) \quad \forall \gamma \in \Gamma \setminus \{e\}$, with = iff $\gamma \subset \lambda$.




Thm 2 (M-Viaggi) Let $\rho: \Gamma \rightarrow SO_0(2, n+1)$ be a max repr. Then

$\forall \lambda$ max geod lam there exist unique

- o $S_\lambda \hookrightarrow C\rho/\rho(\Gamma)$ embedded spacelike surface
- o X_λ hyp structure, $\xleftarrow{\text{intrinsic description using cross ratio}}$
- o homeo $u: S_\lambda \rightarrow X_\lambda$ such that
 - i) $u^{-1}(g)$ spacelike geo $\forall g \subset \lambda$ &
 - ii) $d_{\mathbb{H}^2}(\tilde{u}(x), \tilde{u}(y)) \leq d_{\mathbb{H}^{2,n}}(x, y)$



Moreover $L_{X_\lambda}(\gamma) \leq L_\rho(\gamma) \quad \forall \gamma \in \Gamma \setminus \{e\}$, with = iff $\gamma \subset \lambda$.




Thm 2 (M-Viaggi) Let $\rho: \Gamma \rightarrow SO_0(2, n+1)$ be a max repr. Then


$\forall \lambda$ max geod lam there exist unique

- o $S_\lambda \hookrightarrow C\rho/\rho(\Gamma)$ embedded spacelike surface
- o X_λ hyp structure, $\xleftarrow{\text{intrinsic description using cross ratio}}$
- o homeo $u: S_\lambda \rightarrow X_\lambda$ such that
 - i) $u^{-1}(g)$ spacelike geo $\forall g \subset \lambda$ &
 - $u^{-1}(\gamma)$ spacelike triangle $\forall \gamma \subset X_\lambda \setminus \lambda$
 - ii) $d_{\mathbb{H}^{n+1}}(\tilde{u}(x), \tilde{u}(y)) \leq d_{\mathbb{H}^{n+1}}(x, y)$

Moreover $L_{X_\lambda}(\gamma) \leq L_\rho(\gamma) \quad \forall \gamma \in \Gamma \setminus \{e\}$, with = iff $\gamma \subset \lambda$.



$$\begin{array}{c} u \\ \uparrow \end{array}$$




Thm 2 (M-Viaggi) Let $\rho: \Gamma \rightarrow SO_0(2, n+1)$ be a max repr. Then


$\forall \lambda$ max geod lam there exist unique

- o $S_\lambda \hookrightarrow C\rho/\rho(\Gamma)$ embedded spacelike surface
- o X_λ hyp structure, $\xleftarrow{\text{intrinsic description using cross ratio}}$
- o homeo $u: S_\lambda \rightarrow X_\lambda$ such that
 - i) $u^{-1}(g)$ spacelike geo $\forall g \subset \lambda$ &
 - $u^{-1}(\gamma)$ spacelike triangle $\forall \gamma \subset X_\lambda \setminus \lambda$
 - ii) $d_{\mathbb{H}^{n+1}}(\tilde{u}(x), \tilde{u}(y)) \leq d_{\mathbb{H}^{n+1}}(x, y)$

Moreover $L_{X_\lambda}(\gamma) \leq L_\rho(\gamma) \quad \forall \gamma \in \Gamma \setminus \{e\}$, with =
iff $\gamma \subset \lambda$.



$\uparrow u$



Applications:

- In AdS^3 , $\forall \lambda \in \text{max}\ \text{lam}$ we find

$$\varphi_\lambda : X_{\max}(\Gamma, SO_0(2,2)) \longrightarrow \text{Im } \varphi_\lambda \subset \mathcal{U}(\lambda; \mathbb{R} + z\mathbb{R})$$

$$p \longmapsto \hat{\delta}_\lambda^p \quad \leftarrow \text{shear-bend cocycle}$$

para-holomorphic & symplectic parametrization (analogue of the work of Bonahon for repr's in $\text{Isom}^+(\mathbb{H}^3) = \text{PSL}_2 \mathbb{C}$)

(if λ finite, then there is a similar result using $\text{Pho}^{2,n}$ -structures ($n \geq 2$), compare with Collier - Tholozan - Toulisse)

Applications:

- In AdS^3 , $\forall \lambda \in \text{max}\ \text{lam}$ we find

$$\varphi_\lambda : X_{\max}(\Gamma, SO_0(2,2)) \longrightarrow \text{Im } \varphi_\lambda \subset \mathcal{U}(\lambda; \mathbb{R} + z\mathbb{R})$$

$$p \longmapsto \hat{\delta}_\lambda^p \quad \leftarrow \text{shear-bend cocycle}$$

para-holomorphic & symplectic parametrization (analogue of the work of Bonahon for repr's in $\text{Isom}^+(\mathbb{H}^3) = \text{PSL}_2 \mathbb{C}$)

(if λ finite, then there is a similar result using $\text{Phi}^{2,n}$ -structures ($n \geq 2$), compare with Collier - Tholozan - Toulisse)

- $\forall \lambda \max \quad \ell_p(\gamma) \geq \ell_{X_\lambda}(\gamma) \quad \forall \gamma \in \Gamma \Leftarrow$ Pseudo-Riemannian geometry of $\mathbb{H}^{2,n}$

When $n=1$, $p=(p_X, p_Y)$ satisfies $L_\gamma(p) = \frac{1}{2}(L_\gamma(X) + L_\gamma(Y))$, and $\sigma_\lambda^B = \frac{1}{2}(\sigma_\lambda^X + \sigma_\lambda^Y)$

$$\Rightarrow L_\gamma\left(\frac{1}{2}(\sigma_\lambda^X + \sigma_\lambda^Y)\right) \leq \frac{1}{2}(L_\gamma(X) + L_\gamma(Y)) \quad (\text{Bestvina-Bromberg-Fujiwara-Souto, Théret})$$

lengths are convex along shear paths

- recover Collier-Tholozan-Toulisse: $\forall p: \Gamma \rightarrow SD_0(2, n+1) \max$

$$\boxed{\exists X \text{ hyp str} \& \exists k \geq 1 \text{ s.t. } \ell_p(\gamma) \geq k \ell_X(\gamma) \quad \forall \gamma \in \Gamma}$$

and $k=1 \Leftrightarrow p$ Fuchsian (but w/o Higgs bundles!)

$(\Rightarrow \text{entropy}(p) \leq 1 \& = 1 \Leftrightarrow p$ Fuchsian)

- $\forall \lambda \max \quad \ell_p(\gamma) \geq \ell_{X_\lambda}(\gamma) \quad \forall \gamma \in \Gamma \Leftarrow$ Pseudo-Riemannian geometry of $\mathbb{H}^{2,n}$

When $n=1$, $p=(p_X, p_Y)$ satisfies $L_\gamma(p) = \frac{1}{2}(L_\gamma(X) + L_\gamma(Y))$, and $\sigma_\lambda^\beta = \frac{1}{2}(\sigma_\lambda^X + \sigma_\lambda^Y)$

$$\Rightarrow L_\gamma\left(\frac{1}{2}(\sigma_\lambda^X + \sigma_\lambda^Y)\right) \leq \frac{1}{2}(L_\gamma(X) + L_\gamma(Y)) \quad (\text{Bestvina-Bromberg-Fujiwara-Souto, Théret})$$

lengths are convex along shear paths

- recover Collier-Tholozan-Toulisse: $\forall \rho: \Gamma \rightarrow SO_0(2, n+1)$ max

$$\exists X \text{ hyp str } \& \exists k \geq 1 \text{ s.t. } \ell_\rho(\gamma) \geq k \ell_X(\gamma) \quad \forall \gamma \in \Gamma$$

and $k=1 \Leftrightarrow \rho$ Fuchsian (but w/o Higgs bundles!)

$(\Rightarrow \text{entropy}(\rho) \leq 1 \& = 1 \Leftrightarrow \rho \text{ Fuchsian})$

Questions

i) CROSS RATIOS: Thm 1 applies to a wide class of cross ratios

(other Anosov repr's come with good cross ratios, e.g. Hitchin repr's, or max repr's in $Sp(2n, \mathbb{R})$, see Martone-Zhang, Labourie)

Q: Does σ_λ^β have nice geom meaning for other types of ρ ?

ii) CONVEX CORE GEOMETRY: every $\rho: \Gamma \rightarrow SO_0(2, n+1)$ max has a cpt convex core $CC(\rho) \supset$ all pleated surfaces

Q: Can pleated surfaces help us to understand geometry of convex core? Already for $d=1$ not much is known about the interior of the convex core

Questions

i) CROSS RATIOS: Thm 1 applies to a wide class of cross ratios

(other Anosov repr's come with good cross ratios, e.g. Hitchin repr's, or max repr's in $Sp(2n, \mathbb{R})$, see Martone-Zhang, Labourie)

Q: Does σ_λ^β have nice geom meaning for other types of ρ ?

ii) CONVEX CORE GEOMETRY: every $\rho: \Gamma \rightarrow SO_0(2, n+1)$ max has a cpt convex core $CC(\rho) \supset$ all pleated surfaces

Q: Can pleated surfaces help us to understand geometry of convex core? Already for $d=1$ not much is known about the interior of the convex core

Thank you for your attention !