

# Improved Statistical Inference for Expectile-Based Risk Measures

An Optimal Pooling Approach in the Heavy-Tailed Regime

Filippo Gombac

Supervisor: Simone Padoan

Department of Decision Sciences

Bocconi University

June 6, 2026

# Abstract

[TODO: ]

# Contents

|          |                                                            |           |
|----------|------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                        | <b>1</b>  |
| <b>2</b> | <b>Background</b>                                          | <b>2</b>  |
| 2.1      | Regular variation and the second-order condition . . . . . | 2         |
| 2.2      | Tail-index and extreme-quantile estimation . . . . .       | 3         |
| 2.2.1    | The Hill estimator . . . . .                               | 3         |
| 2.2.2    | Weissman extrapolation to extreme quantiles . . . . .      | 4         |
| 2.3      | Expectiles . . . . .                                       | 4         |
| 2.3.1    | Definition . . . . .                                       | 4         |
| 2.3.2    | Basic properties . . . . .                                 | 5         |
| 2.3.3    | Coherence and elicibility . . . . .                        | 5         |
| 2.4      | The high-expectile/high-quantile bridge . . . . .          | 6         |
| 2.4.1    | The expectile-quantile asymptotic equivalence . . . . .    | 6         |
| 2.4.2    | Relation to earlier expectile estimators . . . . .         | 7         |
| 2.5      | The optimal pooling framework . . . . .                    | 7         |
| 2.5.1    | Setup . . . . .                                            | 7         |
| 2.5.2    | The pooled Hill estimator . . . . .                        | 8         |
| 2.5.3    | Variance- and AMSE-optimal weights . . . . .               | 9         |
| 2.5.4    | Weighted geometric pooling of extreme quantiles . . . . .  | 10        |
| 2.5.5    | Estimated weights and plug-in objects . . . . .            | 11        |
| 2.5.6    | Source diagnostics and intervals . . . . .                 | 12        |
| 2.5.7    | Distributed inference as a special case . . . . .          | 12        |
| <b>3</b> | <b>Pooled extreme expectiles</b>                           | <b>14</b> |
| 3.1      | Estimator and exact decomposition . . . . .                | 14        |
| 3.2      | A deterministic two-weight theorem . . . . .               | 16        |
| 3.3      | Weight consequences . . . . .                              | 19        |
| 3.4      | Estimated pooled-Weissman weights and inference . . . . .  | 20        |
|          | <b>Bibliography</b>                                        | <b>22</b> |

# Chapter 1

## Introduction

[TODO: ]

## Chapter 2

# Background

### 2.1 Regular variation and the second-order condition

Throughout the thesis we work with a real-valued random loss  $X$  that has continuous distribution function  $F$  and survival function  $\bar{F} = 1 - F$ . Given an iid sample  $X_1, \dots, X_n$  from  $F$ , the corresponding order statistics are denoted by  $X_{1:n} \leq X_{2:n} \leq \dots \leq X_{n:n}$ .

Heavy-tailed asymptotics are governed by the language of regular variation; the reference is Haan and Ferreira (2006, Appendix B.1).

**Definition 2.1** (Regular variation). A Lebesgue measurable function  $f : (0, \infty) \rightarrow \mathbb{R}$  that is eventually positive is *regularly varying at infinity with index*  $\alpha \in \mathbb{R}$ , written  $f \in \text{RV}_\alpha$ , if for every  $y > 0$ ,

$$\lim_{t \rightarrow \infty} \frac{f(ty)}{f(t)} = y^\alpha.$$

The functions in  $\text{RV}_0$  are called *slowly varying*.

The first-order condition under which all of our asymptotics are taken asserts that  $F$  has a Pareto-type tail.

**Condition 2.2** (First-order condition). There exists  $\gamma > 0$  such that  $\bar{F} \in \text{RV}_{-1/\gamma}$ , or equivalently the *tail quantile function*

$$U(t) := \inf \left\{ x \in \mathbb{R} : \frac{1}{\bar{F}(x)} \geq t \right\}, \quad t > 1,$$

satisfies  $U \in \text{RV}_\gamma$ .

Throughout, for a continuous distribution function  $F$ , we write

$$Q_X(\tau) := F^{\leftarrow}(\tau) = \inf \{ x \in \mathbb{R} : F(x) \geq \tau \}, \quad 0 < \tau < 1.$$

With the tail quantile  $U(t) = F^{\leftarrow}(1 - 1/t)$ ,  $t > 1$ , this gives

$$Q_X(\tau) = U((1 - \tau)^{-1}), \quad Q_X(1 - p) = U(1/p).$$

**Condition 2.2** is the Fréchet max-domain condition with extreme-value index  $\gamma$  (Haan and Ferreira 2006, Theorem 1.2.1); standard examples include Pareto, Burr, Student- $t$ , Fréchet and  $F$ -distribution tails (Haan and Ferreira 2006, §2.6). For normal limits we use the standard second-order refinement.

**Condition 2.3** (Second-order condition  $\mathcal{C}_2(\gamma, \rho, A)$ ). There exist  $\gamma > 0$ , a second-order parameter  $\rho \leq 0$ , and a measurable auxiliary function  $A$  of constant sign with  $A(t) \rightarrow 0$  as  $t \rightarrow \infty$ , such that for every  $y > 0$ ,

$$\lim_{t \rightarrow \infty} \frac{1}{A(t)} \left[ \frac{U(ty)}{U(t)} - y^\gamma \right] = y^\gamma \frac{y^\rho - 1}{\rho},$$

where the right-hand side is read as  $y^\gamma \log y$  when  $\rho = 0$ .

The function  $|A|$  is itself regularly varying with index  $\rho$  (Haan and Ferreira 2006, Theorem 2.3.9). Since  $A$  has constant sign eventually, signed ratios  $A(ty)/A(t)$  converge to  $y^\rho$ ; the rate  $A$  is the source of the second-order bias terms below.

**Condition 2.4** (Finite first-moment condition). The extreme-value index satisfies  $\gamma \in (0, 1)$  and the negative part  $X^- = \max(-X, 0)$  satisfies

$$\mathbb{E} X^- < \infty.$$

Under **Condition 2.2**,  $\gamma < 1$  gives  $\mathbb{E} X^+ < \infty$ ; together with the displayed lower-tail condition this is the finite-mean regime for the quantile-based expectile approximation.

## 2.2 Tail-index and extreme-quantile estimation

We collect the two classical estimators used throughout the thesis: the Hill estimator of  $\gamma$  and the Weissman extrapolation for extreme quantiles. The textbook reference is Haan and Ferreira (2006, Ch. 3–4).

### 2.2.1 The Hill estimator

Given an iid sample  $X_1, \dots, X_n$  from  $F$  and an integer  $k = k_n \in \{1, \dots, n-1\}$ , the *Hill estimator* (Hill 1975) of  $\gamma$  based on the  $k+1$  top order statistics is

$$\hat{\gamma}_n(k) = \frac{1}{k} \sum_{i=1}^k \log \left( \frac{X_{n-i+1:n}}{X_{n-k:n}} \right). \quad (2.1)$$

Since  $U(t) \rightarrow \infty$ ,  $X_{n-k:n} > 0$  with probability tending to one for intermediate  $k$ ; log-based estimators are understood on this event. For tail-index estimation one may also shift the data, but shifts must be undone for quantile or expectile estimation. The consistency result below is Haan and Ferreira (2006, Theorem 3.2.2); the normal limit is Haan and Ferreira (2006, Theorem 3.2.5) when  $\rho < 0$ , with the  $\rho = 0$  endpoint covered by the  $m = 1$  specialisation of Daouia, Padoan, et al. (2024, Theorem 1).

**Theorem 2.5.** Assume  $\mathcal{C}_2(\gamma, \rho, A)$  with  $\gamma > 0$ . Let  $k = k_n \rightarrow \infty$  and  $k/n \rightarrow 0$  as  $n \rightarrow \infty$ . Then

$\hat{\gamma}_n(k) \xrightarrow{\mathbb{P}} \gamma$ . If, in addition,  $\sqrt{k} A(n/k) \rightarrow \lambda \in \mathbb{R}$ , then

$$\sqrt{k} (\hat{\gamma}_n(k) - \gamma) \xrightarrow{d} \mathcal{N}\left(\frac{\lambda}{1-\rho}, \gamma^2\right).$$

The choice of  $k$  controls the usual bias–variance trade-off; here it is treated as fixed by the data analyst.

### 2.2.2 Weissman extrapolation to extreme quantiles

The *Weissman extrapolation* (Weissman 1978) extrapolates from the intermediate-order statistic  $X_{n-k:n}$  — which estimates the quantile  $Q_X(1 - k/n)$  at the intermediate level  $1 - k/n$  — to a quantile at a very-extreme level  $1 - p$  with  $p = p_n \rightarrow 0$  as  $n \rightarrow \infty$ . Plugging the Hill estimator into the first-order identity  $U(ty)/U(t) \rightarrow y^\gamma$  gives the Weissman estimator

$$\hat{q}_n^*(1 - p \mid k) = \left(\frac{k}{np}\right)^{\hat{\gamma}_n(k)} X_{n-k:n}. \quad (2.2)$$

The theorem below, the Hill-estimator specialisation of Haan and Ferreira (2006, Theorem 4.3.8), covers the far-tail regime  $np = o(k)$ ; the case  $np = O(1)$  is very-extreme in the sense that only finitely many observations are expected beyond the target.

**Theorem 2.6.** Assume  $\mathcal{C}_2(\gamma, \rho, A)$  with  $\gamma > 0$  and  $\rho < 0$ . Let  $k = k_n \rightarrow \infty$ ,  $k/n \rightarrow 0$ ,  $p = p_n \rightarrow 0$  with  $k/(np) \rightarrow \infty$  and  $\sqrt{k}/\log(k/(np)) \rightarrow \infty$ . Suppose  $\sqrt{k} A(n/k) \rightarrow \lambda \in \mathbb{R}$ . Then

$$\frac{\sqrt{k}}{\log(k/(np))} \left( \frac{\hat{q}_n^*(1 - p \mid k)}{Q_X(1 - p)} - 1 \right) \xrightarrow{d} \mathcal{N}\left(\frac{\lambda}{1-\rho}, \gamma^2\right).$$

The log-denominator is the cost of extrapolation. Up to that factor the limit is the Hill limit, which is why Daouia, Padoan, et al. (2024, §2.3) pool Weissman estimators geometrically. This pooled-Weissman law is the source result against which the rebuilt expectile estimator will be compared.

## 2.3 Expectiles

Expectiles were introduced by Newey and Powell (1987) as asymmetric-least-squares analogues of quantiles. Their role as coherent and elicitable risk measures is developed in Bellini et al. (2014), Ziegel (2016), Gneiting (2011), and Ehm et al. (2016).

### 2.3.1 Definition

Let  $X$  be a real-valued random variable with  $\mathbb{E}|X| < \infty$ , and fix a level  $\tau \in (0, 1)$ . The  $\tau$ -*expectile* of  $X$  is defined as

$$\xi_\tau(X) := \arg \min_{\theta \in \mathbb{R}} \mathbb{E}[\eta_\tau(X - \theta) - \eta_\tau(X)], \quad \eta_\tau(u) := |\tau - \mathbb{1}\{u \leq 0\}| u^2. \quad (2.3)$$

The subtraction of  $\eta_\tau(X)$  makes the criterion finite under  $\mathbb{E}|X| < \infty$  and does not affect the minimiser; at  $\tau = 1/2$  one recovers the mean.

The convex objective (2.3) has a unique minimiser, and its first-order condition reduces to the implicit equation

$$\tau \mathbb{E}[(X - \xi_\tau(X))_+] = (1 - \tau) \mathbb{E}[(\xi_\tau(X) - X)_+], \quad (2.4)$$

where  $u_+ = \max(u, 0)$ . Thus expectiles balance upper and lower first partial moments, unlike quantiles, which balance tail probabilities.

### 2.3.2 Basic properties

The following properties are immediate from (2.3) or (2.4); we collect them without proof and refer to Newey and Powell (1987, §2) and Bellini et al. (2014, Sections 2–3) for details.

**Proposition 2.7.** *Let  $X$  and  $Y$  be real-valued random variables with  $\mathbb{E}|X|, \mathbb{E}|Y| < \infty$ . For each  $\tau \in (0, 1)$ :*

- (i) **Existence and uniqueness.**  $\xi_\tau(X)$  exists and is uniquely defined.
- (ii) **Law invariance.** If  $X$  and  $Y$  have the same distribution, then  $\xi_\tau(X) = \xi_\tau(Y)$ .
- (iii) **Mean recovery.**  $\xi_{1/2}(X) = \mathbb{E}X$ .
- (iv) **Location equivariance.**  $\xi_\tau(X + c) = \xi_\tau(X) + c$  for  $c \in \mathbb{R}$ .
- (v) **Positive homogeneity.**  $\xi_\tau(\lambda X) = \lambda \xi_\tau(X)$  for  $\lambda \geq 0$ .
- (vi) **Sign reflection.**  $\xi_\tau(-X) = -\xi_{1-\tau}(X)$ .
- (vii) **Monotonicity in  $\tau$ .** The map  $\tau \mapsto \xi_\tau(X)$  is nondecreasing on  $(0, 1)$ , and is strictly increasing when  $X$  is nondegenerate, with the boundary limits  $\xi_\tau(X) \rightarrow \text{ess inf}(X)$  as  $\tau \rightarrow 0$  and  $\xi_\tau(X) \rightarrow \text{ess sup}(X)$  as  $\tau \rightarrow 1$  (infinite limits allowed).
- (viii) **Monotonicity in the argument.** If  $X \leq Y$  almost surely, then  $\xi_\tau(X) \leq \xi_\tau(Y)$ .

For losses, expectiles are translation equivariant, positively homogeneous and monotone. The remaining coherence issue is subadditivity.

### 2.3.3 Coherence and elicibility

In the rest of the thesis  $X$  denotes a loss. The coherence literature is often written for financial positions, so in this subsection only write  $Y$  for a position and define the capital requirement

$$\rho_\tau(Y) = \xi_\tau(-Y).$$

**Theorem 2.8** (Bellini et al. (2014)). *The functional  $\rho_\tau(Y) = \xi_\tau(-Y)$  is a coherent risk measure if and only if  $\tau \geq 1/2$ .*



Recall that a functional is *elicitable* if it is the unique minimiser of an expected strictly consistent score (Gneiting 2011). Expectiles are elicitable on classes with finite first moment; the asymmetric squared score is strictly consistent under finite second moments, and the general scoring class has Bregman-type and Choquet-type representations (Gneiting 2011; Ehm et al. 2016). This scoring structure motivates forecast comparison, but the asymptotic theory below uses only (2.4) and the extreme expectile–quantile equivalence.

## 2.4 The high-expectile/high-quantile bridge

The contribution chapter uses expectile theory through the population relationship between a high expectile and the quantile of the same level. The stochastic extreme-quantile part is supplied by the Daouia–Padoan–Stupfler pooled Weissman estimator recalled in Section 2.5; the expectile-specific work is to control the deterministic bridge between  $\xi_\tau$  and  $Q_X(\tau)$ . This section therefore records only the bridge statements needed later.

### 2.4.1 The expectile–quantile asymptotic equivalence

The single most important fact for extreme expectile inference is that, in the heavy-tailed regime, the expectile is asymptotically proportional to the quantile of the same level.

**Proposition 2.9** (Asymptotic equivalence; Bellini et al. (2014), Daouia, Girard, et al. (2019), Corollary 1 with  $p = 2$ ). *Suppose  $X$  satisfies Condition 2.2 with index  $\gamma \in (0, 1)$  and  $\mathbb{E} X^- < \infty$ ; equivalently, suppose Condition 2.4 holds. Then*

$$\frac{\xi_\tau(X)}{Q_X(\tau)} \xrightarrow{\tau \uparrow 1} \psi(\gamma) := (\gamma^{-1} - 1)^{-\gamma}. \quad (2.5)$$

The bridge requires only the finite-mean range  $\gamma < 1$  and lower-tail integrability; stronger restrictions used for direct empirical expectile estimation are not part of the quantile-based bridge route. The function  $\psi$  is continuous on  $(0, 1)$ , with  $\psi(\gamma) \rightarrow 1$  as  $\gamma \downarrow 0$  and  $\psi(\gamma) \rightarrow \infty$  as  $\gamma \uparrow 1$ . It is *not* monotone: its logarithmic derivative

$$m(\gamma) := \frac{d}{d\gamma} \log \psi(\gamma) = \frac{1}{1 - \gamma} - \log(\gamma^{-1} - 1) \quad (2.6)$$

changes sign at  $\gamma^* \approx 0.2178$ , where  $\psi$  reaches its minimum. In the pooled extreme-expectile theorem below,  $m(\gamma)$  appears in the Taylor expansion of the bridge-Hill term; when  $m(\gamma) = 0$ , that term is even smaller at first order.

Under the second-order condition  $\mathcal{C}_2(\gamma, \rho, A)$  with  $\gamma \in (0, 1)$  and  $\mathbb{E} X^- < \infty$ , the first-order equivalence (2.5) sharpens to the second-order expansion (Daouia, Girard, et al. 2020, Proposi-

tion 1(i))

$$\begin{aligned} \frac{\xi_\tau(X)}{Q_X(\tau)} &= \psi(\gamma) \left[ 1 + c(\gamma, \rho) A((1 - \tau)^{-1}) + \gamma(\gamma^{-1} - 1)^\gamma \frac{\mathbb{E} X}{Q_X(\tau)} \right. \\ &\quad \left. + o(A((1 - \tau)^{-1})) + o(Q_X(\tau)^{-1}) \right], \\ c(\gamma, \rho) &:= \frac{(\gamma^{-1} - 1)^{-\rho}}{1 - \gamma - \rho} + \frac{(\gamma^{-1} - 1)^{-\rho} - 1}{\rho}. \end{aligned} \quad (2.7)$$

At  $\rho = 0$  the second term in  $c(\gamma, \rho)$  is read by continuity, so  $c(\gamma, 0) = m(\gamma)$ . The  $A$ -term is the second-order expectile–quantile bias used later to control the population bridge term. The displayed first-moment term is  $O(Q_X(\tau)^{-1})$ ; in [Chapter 3](#) it is controlled on the pooled-Weissman scale by the explicit bridge-rate condition.

### 2.4.2 Relation to earlier expectile estimators

The iid extreme-expectile literature also studies direct asymmetric-least-squares estimators, quantile-based intermediate plug-in estimators, and Weissman-type expectile extrapolations from an intermediate expectile anchor (Daouia, Girard, et al. [2018](#); Daouia, Girard, et al. [2020](#); Davison et al. [2023](#)). Those results explain why the expectile–quantile bridge is useful, but they are not theorem inputs for the contribution chapter. The estimator analysed below performs the extreme extrapolation in the pooled-quantile component and then applies the population bridge at the target level. Accordingly, the needed expectile background is the first- and second-order bridge above, rather than a generic CLT for an intermediate expectile estimator.

## 2.5 The optimal pooling framework

We now recall the pooling framework of Daouia, Padoan, et al. ([2024](#)). The rebuilt thesis uses this framework as the source theory for pooled tail-index and extreme-quantile inference before applying the expectile–quantile bridge. The framework pools tail-index and extreme-quantile estimators from  $m$  samples, allowing heterogeneous sizes, second-order properties and cross-sample tail dependence.

### 2.5.1 Setup

Throughout [Section 2.5](#),  $m$  is fixed while  $n = \sum_{j=1}^m n_j \rightarrow \infty$ . Let  $\mathbf{X}_i = (X_{i,1}, \dots, X_{i,m})^\top$ ,  $i \geq 1$ , be iid copies of  $\mathbf{X} = (X_1, \dots, X_m)^\top$ . Margin  $j$  is observed for  $i = 1, \dots, n_j$  and has continuous right-heavy-tailed distribution  $F_j$ , tail quantile function  $U_j = (1/\bar{F}_j)^\leftarrow$  and extreme-value index  $\gamma_j > 0$ ; its order statistics are denoted  $X_{1:n_j,j} \leq \dots \leq X_{n_j:n_j,j}$ .

Tail inference in sample  $j$  uses its top  $k_j + 1$  order statistics, where the effective sample size  $k_j = k_j(n)$  satisfies  $k_j \rightarrow \infty$  and  $k_j/n_j \rightarrow 0$  as  $n \rightarrow \infty$ . The total effective sample size is  $k = \sum_{j=1}^m k_j$ . Assume the asymptotic-proportionality conditions

$$\frac{n_1}{n_j} \rightarrow b_j \in (0, \infty), \quad \frac{k_1}{k_j} \rightarrow c_j \in (0, \infty), \quad (2.8)$$

with  $b_1 = c_1 = 1$ . The marginal  $F_j$  satisfies  $\mathcal{C}_2(\gamma_j, \rho_j, A_j)$  ([Condition 2.3](#)). The thesis works

under the *common-tail-index* assumption

$$(H_\gamma) \quad \gamma_1 = \dots = \gamma_m = \gamma.$$

Under this assumption, all common- $\gamma$  vector limits below are expressed on the aggregate scale  $\sqrt{k}$ , with the constants  $c_j$  translating the local  $\sqrt{k_j}$  rates.

The cross-sample asymptotic dependence is captured by a pairwise *tail-copula* structure (Daouia, Padoan, et al. 2024, §2.1).

**Condition  $\mathcal{J}(\mathbf{R})$  (tail-copula condition).** For each pair  $(j, \ell)$  with  $j \neq \ell$ , there exists a function  $R_{j,\ell} : [0, \infty]^2 \setminus \{(\infty, \infty)\} \rightarrow [0, \infty)$  such that, for every  $(x_j, x_\ell) \in [0, \infty]^2 \setminus \{(\infty, \infty)\}$ ,

$$\lim_{s \rightarrow \infty} s \mathbb{P}(1 - F_j(X_j) \leq x_j/s, 1 - F_\ell(X_\ell) \leq x_\ell/s) = R_{j,\ell}(x_j, x_\ell).$$

The case  $R_{j,\ell} \equiv 0$  is asymptotic tail independence, implied by the independent samples used later in the thesis.

### 2.5.2 The pooled Hill estimator

Each sample contributes a Hill estimator  $\hat{\gamma}_j(k_j)$  as in (2.1); we collect these into the random vector

$$\hat{\gamma}_n = (\hat{\gamma}_1(k_1), \dots, \hat{\gamma}_m(k_m))^\top.$$

Given a weight vector  $\omega \in \mathbb{R}^m$  with  $\omega^\top \mathbf{1} = 1$ , the *pooled Hill estimator* is the weighted arithmetic combination

$$\hat{\gamma}_n(\omega) := \omega^\top \hat{\gamma}_n = \sum_{j=1}^m \omega_j \hat{\gamma}_j(k_j). \quad (2.9)$$

The affine constraint  $\omega^\top \mathbf{1} = 1$  ensures consistency for the common  $\gamma$  under  $(H_\gamma)$ .

The fundamental asymptotic result is the joint CLT for the marginal Hill estimators.

**Theorem 2.10** (Daouia, Padoan, et al. (2024), Theorem 1). *Assume  $\mathcal{C}_2(\gamma_j, \rho_j, A_j)$  for every  $j$ , the tail-copula condition  $\mathcal{J}(\mathbf{R})$ , the proportionality conditions (2.8), and  $\sqrt{k_j} A_j(n_j/k_j) \rightarrow \lambda_j \in \mathbb{R}$  for every  $j$ . Then*

$$(\sqrt{k_1}(\hat{\gamma}_1(k_1) - \gamma_1), \dots, \sqrt{k_m}(\hat{\gamma}_m(k_m) - \gamma_m))^\top \xrightarrow{d} \mathcal{N}(\mathbf{B}, \mathbf{V}),$$

with  $\mathbf{B} = (\lambda_j/(1 - \rho_j))_{j=1}^m$  and  $\mathbf{V}$  the symmetric matrix whose entries are

$$V_{j,\ell} = \begin{cases} \gamma_j^2, & j = \ell, \\ \frac{\gamma_j \gamma_\ell}{\max(b_j, b_\ell) \sqrt{c_j c_\ell}} R_{j,\ell}(b_j c_\ell, b_\ell c_j), & j < \ell. \end{cases}$$

The matrix is symmetric, so  $V_{\ell,j} = V_{j,\ell}$  for  $\ell > j$ . Suppose, in addition, the common- $\gamma$  condition

$(H_\gamma)$ . Then the same marginal estimators, now viewed on the common  $\sqrt{k}$  scale, satisfy

$$\sqrt{k}(\hat{\gamma}_n - \gamma \mathbf{1}) \xrightarrow{d} \mathcal{N}(\mathbf{B}_c, \mathbf{V}_c),$$

where

$$(\mathbf{B}_c)_j = \sqrt{c_j \sum_{i=1}^m c_i^{-1} \frac{\lambda_j}{1 - \rho_j}} \quad j = 1, \dots, m.$$

The covariance matrix  $\mathbf{V}_c$  is given by

$$(\mathbf{V}_c)_{j,\ell} = \left( \sum_{i=1}^m c_i^{-1} \right) \begin{cases} \gamma^2 c_j, & j = \ell, \\ \frac{\gamma^2}{\max(b_j, b_\ell)} R_{j,\ell}(b_j c_\ell, b_\ell c_j), & j < \ell, \end{cases}$$

with  $(\mathbf{V}_c)_{\ell,j} = (\mathbf{V}_c)_{j,\ell}$  for  $\ell > j$ . Consequently, for any  $\boldsymbol{\omega}$  with  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ ,

$$\sqrt{k}(\hat{\gamma}_n(\boldsymbol{\omega}) - \gamma) \xrightarrow{d} \mathcal{N}(\boldsymbol{\omega}^\top \mathbf{B}_c, \boldsymbol{\omega}^\top \mathbf{V}_c \boldsymbol{\omega}).$$

For independent samples, all off-diagonal entries vanish and  $\mathbf{V}_c$  becomes the diagonal matrix

$$\mathbf{V}_c = \gamma^2 \left( \sum_{i=1}^m c_i^{-1} \right) \text{diag}(c_1, \dots, c_m). \quad (2.10)$$

Equation (2.10) is the covariance form to be used when the rebuilt main theorem is specialised to independent samples.

### 2.5.3 Variance- and AMSE-optimal weights

The relevant weights solve quadratic programmes under the affine constraint  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ ; unless stated otherwise, they need not be nonnegative. In Chapter 3, these formulas are applied only to the pooled-Weissman weights; they do not define or optimise the separate bridge-Hill weights introduced there.

**Proposition 2.11** (Optimal weights; Daouia, Padoan, et al. (2024), Theorem 1; estimated-weight implementation in §2.2 and Corollary 1). *Suppose  $\mathbf{V}_c$  is positive definite. Under the common- $\gamma$  condition  $(H_\gamma)$ :*

(i) *The variance-optimal weights, minimising  $\boldsymbol{\omega}^\top \mathbf{V}_c \boldsymbol{\omega}$  subject to  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ , are*

$$\boldsymbol{\omega}^{\text{Var}} = \frac{\mathbf{V}_c^{-1} \mathbf{1}}{\mathbf{1}^\top \mathbf{V}_c^{-1} \mathbf{1}},$$

*with corresponding asymptotic variance  $\boldsymbol{\omega}^{\text{Var},\top} \mathbf{V}_c \boldsymbol{\omega}^{\text{Var}} = (\mathbf{1}^\top \mathbf{V}_c^{-1} \mathbf{1})^{-1}$ .*

(ii) *The AMSE-optimal weights, minimising  $(\boldsymbol{\omega}^\top \mathbf{B}_c)^2 + \boldsymbol{\omega}^\top \mathbf{V}_c \boldsymbol{\omega}$  subject to  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ , are*

$$\boldsymbol{\omega}^{\text{AMSE}} = \frac{(1 + \mathbf{B}_c^\top \mathbf{V}_c^{-1} \mathbf{B}_c) \mathbf{V}_c^{-1} \mathbf{1} - (\mathbf{1}^\top \mathbf{V}_c^{-1} \mathbf{B}_c) \mathbf{V}_c^{-1} \mathbf{B}_c}{(1 + \mathbf{B}_c^\top \mathbf{V}_c^{-1} \mathbf{B}_c)(\mathbf{1}^\top \mathbf{V}_c^{-1} \mathbf{1}) - (\mathbf{1}^\top \mathbf{V}_c^{-1} \mathbf{B}_c)^2}.$$

In the independent-samples case,  $\mathbf{V}_c$  is diagonal (Equation (2.10)) and the variance-optimal weights reduce to

$$\omega_j^{\text{Var}} = \left( \sum_{i=1}^m c_i^{-1} \right)^{-1} c_j^{-1} \sim \frac{k_j}{k}, \quad (2.11)$$

where the equivalence uses  $c_i^{-1} = \lim k_i/k_1$ . The finite-sample distributed version is

$$\tilde{\omega}_n^{\text{Var}} = (k_1/k, \dots, k_m/k)^\top,$$

which is exactly normalised and converges to  $\omega^{\text{Var}}$ .

Feasible weights replace  $\mathbf{V}_c$  and  $\mathbf{B}_c$  by consistent plug-ins (Daouia, Padoan, et al. 2024, §2.2). If  $\hat{\omega}^\top \mathbf{1} = 1$  exactly and  $\hat{\omega} \xrightarrow{\mathbb{P}} \omega$ , then  $\hat{\gamma}_n(\hat{\omega})$  is  $\sqrt{k}$ -equivalent to  $\hat{\gamma}_n(\omega)$ ; the difficult practical step is stable estimation of  $\mathbf{V}_c$  and the bias quantities.

### 2.5.4 Weighted geometric pooling of extreme quantiles

For estimation of an extreme quantile  $q(1-p)$  with  $p = p(n) \rightarrow 0$ , each sample contributes a Weissman estimator

$$\hat{q}_j^*(1-p \mid k_j) = \left( \frac{k_j}{n_j p} \right)^{\hat{\gamma}_j(k_j)} X_{n_j - k_j : n_j, j}.$$

Because Weissman extrapolation is multiplicative, pooling is geometric on the original scale, equivalently linear on the log scale. It requires asymptotic equivalence of the marginal extreme quantiles:

**Condition 2.12** (Tail homoskedasticity ( $\mathcal{H}$ )). For every  $j \neq \ell$ ,  $U_j(t)/U_\ell(t) \rightarrow 1$  as  $t \rightarrow \infty$ .

Equivalently,  $Q_j(1-p)/Q_\ell(1-p) \rightarrow 1$  as  $p \downarrow 0$ . The pooled extreme-quantile estimator of Daouia, Padoan, et al. (2024, §2.3) is

$$\hat{q}_n^*(1-p \mid \omega) := \prod_{j=1}^m [\hat{q}_j^*(1-p \mid k_j)]^{\omega_j}. \quad (2.12)$$

Taking logs reduces pooling to weighted sums of the local identities  $\log \hat{q}_j^*(1-p \mid k_j) = \log(k_j/(n_j p)) \hat{\gamma}_j(k_j) + \log X_{n_j - k_j : n_j, j}$ ; so the Hill part is handled by Theorem 2.10; (2.8) gives  $\log(k_j/(n_j p)) = \log(k/(np)) + O(1)$ .

**Theorem 2.13** (Daouia, Padoan, et al. (2024), Theorem 2). Assume the hypotheses of Theorem 2.10, the common- $\gamma$  condition ( $H_\gamma$ ), the tail-homoskedasticity condition ( $\mathcal{H}$ ), and, in addition,  $\rho_j < 0$  for every  $j = 1, \dots, m$ . Let  $p = p(n) \rightarrow 0$  with  $k/(np) \rightarrow \infty$  and  $\sqrt{k}/\log(k/(np)) \rightarrow \infty$ . Then for any  $\omega$  with  $\omega^\top \mathbf{1} = 1$  and any  $j \in \{1, \dots, m\}$ ,

$$\frac{\sqrt{k}}{\log(k/(np))} \left( \frac{\hat{q}_n^*(1-p \mid \omega)}{Q_j(1-p)} - 1 \right) \xrightarrow{d} \mathcal{N}(\omega^\top \mathbf{B}_c, \omega^\top \mathbf{V}_c \omega).$$

The strict  $\rho_j < 0$  assumption is part of the Weissman theorem. It also supplies the rate needed to recenter by any marginal  $Q_j(1-p)$  under ( $\mathcal{H}$ ) (Daouia, Padoan, et al. 2024, Supplement,

Lemma B.2): for any  $j, \ell$ ,

$$\log Q_j(1-p) - \log Q_\ell(1-p) = O(|A_j(1/p)|) + O(|A_\ell(1/p)|) = o(k^{-1/2}),$$

along the Weissman sequence.

Thus the pooled Weissman and pooled Hill limits share the same mean, variance and optimal weights up to the log-rate factor. This shared structure is the reason the contribution chapter can import the Daouia–Padoan–Stupfler weight criteria for the pooled-Weissman component.

### 2.5.5 Estimated weights and plug-in objects

The optimal weights above are oracle quantities because they depend on the asymptotic bias and covariance objects. For inference they are replaced either by deterministic allocation weights, such as  $(k_1/k, \dots, k_m/k)^\top$  in the independent distributed case, or by estimated weights based on plug-in estimates of the same objects. The following result records the stability property used in the contribution chapter. There, estimated weights are used only in the pooled-Weissman role; the bridge-Hill vector remains deterministic.

**Proposition 2.14** (Estimated pooling weights; Daouia, Padoan, et al. (2024), Supplement, Theorem A.1; Theorem 2 and Corollary 8). *Assume the common- $\gamma$  conditions of Theorem 2.10. Let  $\hat{\omega}_n$  be a random weight vector satisfying*

$$\hat{\omega}_n^\top \mathbf{1} = 1, \quad \hat{\omega}_n \xrightarrow{\mathbb{P}} \omega,$$

for some deterministic  $\omega$  with  $\omega^\top \mathbf{1} = 1$ . Then

$$\sqrt{k} (\hat{\gamma}_n(\hat{\omega}_n) - \gamma) \xrightarrow{d} \mathcal{N}(\omega^\top \mathbf{B}_c, \omega^\top \mathbf{V}_c \omega).$$

If, in addition, the assumptions of Theorem 2.13 hold, then

$$\frac{\sqrt{k}}{\log(k/(np))} \left( \frac{\hat{q}_n^*(1-p | \hat{\omega}_n)}{Q_j(1-p)} - 1 \right) \xrightarrow{d} \mathcal{N}(\omega^\top \mathbf{B}_c, \omega^\top \mathbf{V}_c \omega), \quad j = 1, \dots, m.$$

In the independent common-distribution distributed case, the same statement holds with the diagonal covariance matrix (2.10) and the distributed bias constants of Daouia, Padoan, et al. (2024, Corollary 5).

The supplement’s Theorem A.1 is the generic estimated-weight transfer: exact affine normalisation and convergence in probability of the weights make the random-weight pooled estimator asymptotically equivalent to the pooled estimator with limiting deterministic weights. Applied to Theorem 2.10, this gives the Hill statement. The Weissman statement is the corresponding random-weight version of Theorem 2.13; in the distributed common-marginal case it is the published Corollary 8 of Daouia, Padoan, et al. (2024).

The next proposition separates the probabilistic consistency of the estimated bias and variance objects from the algebra of applying a chosen weight vector.

**Proposition 2.15** (Plug-in bias and variance). *Let  $\widehat{\mathbf{B}}_{cn}$  and  $\widehat{\mathbf{V}}_{cn}$  be estimators of  $\mathbf{B}_c$  and  $\mathbf{V}_c$  satisfying*

$$\widehat{\mathbf{B}}_{cn} \xrightarrow{\mathbb{P}} \mathbf{B}_c, \quad \widehat{\mathbf{V}}_{cn} \xrightarrow{\mathbb{P}} \mathbf{V}_c,$$

*and let  $\widehat{\boldsymbol{\omega}}_n \xrightarrow{\mathbb{P}} \boldsymbol{\omega}$ . Define*

$$\widehat{B}_{\widehat{\boldsymbol{\omega}},n} = \widehat{\boldsymbol{\omega}}_n^\top \widehat{\mathbf{B}}_{cn}, \quad \widehat{V}_{\widehat{\boldsymbol{\omega}},n} = \widehat{\boldsymbol{\omega}}_n^\top \widehat{\mathbf{V}}_{cn} \widehat{\boldsymbol{\omega}}_n.$$

*Then*

$$\widehat{B}_{\widehat{\boldsymbol{\omega}},n} \xrightarrow{\mathbb{P}} \boldsymbol{\omega}^\top \mathbf{B}_c, \quad \widehat{V}_{\widehat{\boldsymbol{\omega}},n} \xrightarrow{\mathbb{P}} \boldsymbol{\omega}^\top \mathbf{V}_c \boldsymbol{\omega}.$$

All vectors and matrices have fixed dimension. The maps  $(w, b) \mapsto w^\top b$  and  $(w, V) \mapsto w^\top V w$  are continuous, so the result follows from the continuous mapping theorem.

Daouia, Padoan, et al. (2024, §2.2 and Corollary 1) give a general route to the consistency assumptions in Proposition 2.15 through estimated second-order and tail-dependence quantities. In the common-distribution distributed case, their Corollaries 6–7 give the corresponding deterministic variance allocation and plug-in AMSE constructions under the stated second-order parameter consistency assumptions.

### 2.5.6 Source diagnostics and intervals

Daouia, Padoan, et al. (2024) also develop tail-homogeneity tests for the common- $\gamma$  condition, confidence intervals for the common tail index, and log-scale intervals for pooled Weissman quantiles. These tools are useful diagnostics in the source pooling framework. The expectile interval in Chapter 3, however, is not imported as a direct restatement of a DPS quantile interval; it is obtained by Slutsky’s theorem and inversion after the exact expectile decomposition.

### 2.5.7 Distributed inference as a special case

In the independent common-distribution distributed setting, the samples are iid subdivisions of one common population. Then:

- the common- $\gamma$  condition ( $H_\gamma$ ) and the tail-homoskedasticity condition ( $\mathcal{H}$ ) are trivially satisfied;
- the tail copula  $R_{j,\ell}$  between any pair of distinct samples vanishes identically (independence implies tail independence);
- $\mathbf{V}_c$  in Theorem 2.10 becomes the diagonal matrix (2.10);
- the variance-optimal weights of Proposition 2.11 reduce to  $\omega_j^{\text{var}} \propto k_j$  as in (2.11).

Under the stronger balanced-allocation assumptions studied by Daouia, Padoan, et al. (2024, Corollary 6, Theorem 3 and Corollary 8), the variance-optimally pooled Hill and pooled Weissman estimators match the corresponding unfeasible estimators built from the entire combined sample at first order. This oracle equivalence is useful benchmark context for distributed inference. The main theorem in Chapter 3 uses the more direct consequence needed for the expectile problem:

in the independent common-distribution case, the pooled-Weissman bias and variance objects are the diagonal distributed objects displayed above.



## Chapter 3

# Pooled extreme expectiles

This chapter develops the pooled extreme-expectile estimator studied in the thesis. We work in the finite- $m$  distributed iid setting of [Section 2.5.7](#): each machine contains observations from the same continuous heavy-tailed loss distribution, and the target is the common marginal expectile  $\xi_{\tau_n}$ . In this setting the population target is unambiguous, and the pooled extreme-quantile component can be handled through the distributed pooled-Weissman theorem of Daouia, Padoan, et al. ([2024](#), Corollary 8).

The construction combines the geometrically pooled Weissman estimator with the asymptotic bridge between high expectiles and high quantiles. The role of the chapter is to show, from an exact log decomposition, which terms remain visible on the pooled-Weissman scale. Under the bridge-rate condition imposed below, the first-order law is inherited from the pooled extreme-quantile component. Accordingly, first-order weight optimisation concerns the Daouia–Padoan–Stupfler weights in the pooled-Weissman part.

### 3.1 Estimator and exact decomposition

Let  $p_n = 1 - \tau_n$ ,  $n = \sum_{j=1}^m n_j$ ,  $k = \sum_{j=1}^m k_j$ , and

$$\ell_n = \log\left(\frac{k}{np_n}\right).$$

For each machine  $j$ , let  $\hat{\gamma}_j(k_j)$  be the local Hill estimator and

$$\hat{q}_j^*(1 - p_n \mid k_j) = \left(\frac{k_j}{n_j p_n}\right)^{\hat{\gamma}_j(k_j)} X_{n_j - k_j : n_j, j}$$

be the local Weissman estimator. For an affine weight vector  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ , the geometrically pooled Weissman estimator is

$$\hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega}) = \prod_{j=1}^m [\hat{q}_j^*(1 - p_n \mid k_j)]^{\omega_j}.$$

For another affine weight vector  $\boldsymbol{\nu}^\top \mathbf{1} = 1$ , define the pooled Hill estimator used in the outer bridge by

$$\hat{\gamma}_n(\boldsymbol{\nu}) = \sum_{j=1}^m \nu_j \hat{\gamma}_j(k_j).$$

The two-weight pooled extreme-expectile estimator is

$$\hat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega}) = \psi(\hat{\gamma}_n(\boldsymbol{\nu})) \hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega}), \quad \psi(\gamma) = (\gamma^{-1} - 1)^{-\gamma}. \quad (3.1)$$

The vector  $\boldsymbol{\omega}$  weights the pooled-Weissman quantile estimator. The vector  $\boldsymbol{\nu}$  weights only the Hill estimator appearing in the outer expectile bridge. The diagonal construction  $\boldsymbol{\nu} = \boldsymbol{\omega}$  is a special case, not a first-order necessity.

**Lemma 3.1** (Exact log decomposition). *Let  $Q(1 - p_n)$  denote the common marginal quantile and let  $\xi_{\tau_n}$  denote the corresponding expectile. Then*

$$\log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}} = A_n(\boldsymbol{\omega}) + B_n(\boldsymbol{\nu}) - C_n, \quad (3.2)$$

where

$$A_n(\boldsymbol{\omega}) = \log \frac{\hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega})}{Q(1 - p_n)}, \quad B_n(\boldsymbol{\nu}) = \log \psi(\hat{\gamma}_n(\boldsymbol{\nu})) - \log \psi(\gamma),$$

and

$$C_n = \log \frac{\xi_{\tau_n}}{\psi(\gamma)Q(1 - p_n)}.$$

*Proof.* By the definition of the estimator,

$$\begin{aligned} \log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}} &= \log \frac{\psi(\hat{\gamma}_n(\boldsymbol{\nu})) \hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega})}{\xi_{\tau_n}} \\ &= \log \psi(\hat{\gamma}_n(\boldsymbol{\nu})) + \log \hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega}) - \log \xi_{\tau_n}. \end{aligned}$$

Insert and subtract the two deterministic quantities  $\log Q(1 - p_n)$  and  $\log \psi(\gamma)$ . This gives

$$\begin{aligned} \log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}} &= \{\log \hat{q}_n^*(1 - p_n \mid \boldsymbol{\omega}) - \log Q(1 - p_n)\} \\ &\quad + \{\log \psi(\hat{\gamma}_n(\boldsymbol{\nu})) - \log \psi(\gamma)\} \\ &\quad + \{\log \psi(\gamma) + \log Q(1 - p_n) - \log \xi_{\tau_n}\}. \end{aligned}$$

The first bracket is  $A_n(\boldsymbol{\omega})$ , the second bracket is  $B_n(\boldsymbol{\nu})$ , and the third bracket equals

$$\log \frac{\psi(\gamma)Q(1 - p_n)}{\xi_{\tau_n}} = -\log \frac{\xi_{\tau_n}}{\psi(\gamma)Q(1 - p_n)} = -C_n.$$

Combining the three brackets yields (3.2).  $\square$

### 3.2 A deterministic two-weight theorem

The following theorem gives the first-order limit under a rate condition that makes the deterministic population gap between expectiles and quantiles negligible on the pooled-Weissman scale.

**Theorem 3.2** (Pooled extreme expectile, deterministic weights). *Let  $m$  be fixed. Suppose the observations across and within machines are iid copies of a common continuous loss variable  $X$ . Let  $U(t) = Q(1 - 1/t)$  be the common tail quantile, and assume  $C_2(\gamma, \rho, A)$  with  $0 < \gamma < 1$  and  $\rho < 0$ . Assume  $\mathbb{E} X^- < \infty$  and  $U(t) > 0$  for all sufficiently large  $t$ .*

Let  $n = \sum_{j=1}^m n_j$ ,  $k = \sum_{j=1}^m k_j$ , and assume

$$\frac{n_1}{n_j} \rightarrow b_j \in (0, \infty), \quad \frac{k_1}{k_j} \rightarrow c_j \in (0, \infty),$$

with  $b_1 = c_1 = 1$ . Suppose

$$k \rightarrow \infty, \quad \frac{k}{n} \rightarrow 0, \quad \sqrt{k} A(n/k) \rightarrow \lambda \in \mathbb{R}.$$

Let  $p_n = 1 - \tau_n$ ,  $\ell_n = \log\{k/(np_n)\}$ , and assume the extreme Weissman regime

$$p_n \rightarrow 0, \quad \frac{k}{np_n} \rightarrow \infty, \quad \frac{\sqrt{k}}{\ell_n} \rightarrow \infty.$$

Finally impose the expectile-bridge rate condition

$$\eta_n = \frac{\sqrt{k}}{\ell_n Q(1 - p_n)} \rightarrow 0. \quad (3.3)$$

For deterministic affine weights  $\boldsymbol{\nu}^\top \mathbf{1} = 1$  and  $\boldsymbol{\omega}^\top \mathbf{1} = 1$ , the estimator (3.1) satisfies

$$\frac{\sqrt{k}}{\ell_n} \log \frac{\widehat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}} \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega), \quad (3.4)$$

where

$$B_\omega = \frac{\lambda}{1 - \rho} \sum_{j=1}^m d_j^\rho \omega_j, \quad V_\omega = \gamma^2 \left( \sum_{j=1}^m c_j^{-1} \right) \left( \sum_{j=1}^m c_j \omega_j^2 \right),$$

and

$$d_j = \frac{c_j}{b_j} \frac{\sum_{i=1}^m c_i^{-1}}{\sum_{i=1}^m b_i^{-1}}.$$

Moreover,

$$\frac{\sqrt{k}}{\ell_n} \left( \frac{\widehat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}} - 1 \right) \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega). \quad (3.5)$$

*Proof.* Start from the exact identity (3.2). We treat the three terms separately on the scale  $s_n = \sqrt{k}/\ell_n$ .

For  $A_n(\boldsymbol{\omega})$ , the common-marginal pooled-Weissman theorem of Daouia, Padoan, et al. (2024,

Corollary 8) gives

$$\frac{\sqrt{k}}{\ell_n} \left( \frac{\hat{q}_n^*(1 - p_n \mid \omega)}{Q(1 - p_n)} - 1 \right) \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega).$$

Set

$$x_n = \frac{\hat{q}_n^*(1 - p_n \mid \omega)}{Q(1 - p_n)} - 1.$$

The condition  $\sqrt{k}/\ell_n \rightarrow \infty$  implies  $x_n = O_P(\ell_n/\sqrt{k}) = o_P(1)$ . Hence

$$\log(1 + x_n) = x_n + O_P(x_n^2)$$

and

$$\frac{\sqrt{k}}{\ell_n} \{\log(1 + x_n) - x_n\} = O_P\left(\frac{\ell_n}{\sqrt{k}}\right) = o_P(1).$$

Therefore

$$\frac{\sqrt{k}}{\ell_n} A_n(\omega) \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega). \quad (3.6)$$

For  $B_n(\nu)$ , write

$$g(x) = \log \psi(x) = -x \log(x^{-1} - 1).$$

On every compact subinterval of  $(0, 1)$ ,  $g$  is twice continuously differentiable, with

$$g'(x) = m(x) = \frac{1}{1-x} - \log(x^{-1} - 1), \quad g''(x) = \frac{1}{(1-x)^2} + \frac{1}{1-x} + \frac{1}{x}.$$

By the pooled Hill CLT in [Theorem 2.10](#),  $\hat{\gamma}_n(\nu) - \gamma = O_P(k^{-1/2})$  and  $\hat{\gamma}_n(\nu) \xrightarrow{\mathbb{P}} \gamma$ . Hence, with probability tending to one, the line segment between  $\hat{\gamma}_n(\nu)$  and  $\gamma$  lies in a fixed compact subset of  $(0, 1)$ . Taylor's theorem on that event gives

$$B_n(\nu) = m(\gamma)\{\hat{\gamma}_n(\nu) - \gamma\} + O_P\left(\{\hat{\gamma}_n(\nu) - \gamma\}^2\right) = m(\gamma)\{\hat{\gamma}_n(\nu) - \gamma\} + O_P(k^{-1}).$$

Consequently

$$\frac{\sqrt{k}}{\ell_n} B_n(\nu) = \frac{m(\gamma)}{\ell_n} \sqrt{k}\{\hat{\gamma}_n(\nu) - \gamma\} + O_P\left(\frac{1}{\ell_n \sqrt{k}}\right) = o_P(1), \quad (3.7)$$

because  $\ell_n \rightarrow \infty$ . If  $m(\gamma) = 0$ , the displayed conclusion follows from the quadratic remainder alone.

It remains to control  $C_n$ . Put

$$a_n = A(1/p_n), \quad b_n = Q(1 - p_n)^{-1}, \quad \alpha_\gamma = \gamma(\gamma^{-1} - 1)^\gamma, \quad \mu = \mathbb{E} X.$$

The second-order expectile-quantile expansion [\(2.7\)](#) gives, after dividing by  $\psi(\gamma)$ ,

$$\Delta_n := \frac{\xi_{\tau_n}}{\psi(\gamma)Q(1 - p_n)} - 1 = c(\gamma, \rho)a_n + \alpha_\gamma \mu b_n + o(|a_n|) + o(b_n).$$

Since  $\Delta_n \rightarrow 0$ ,

$$C_n = \log(1 + \Delta_n) = \Delta_n + O(\Delta_n^2).$$

The regular variation of  $A$ , the strict inequality  $\rho < 0$ , and

$$\frac{1/p_n}{n/k} = \frac{k}{np_n} \rightarrow \infty$$

imply

$$\frac{A(1/p_n)}{A(n/k)} = \left( \frac{k}{np_n} \right)^\rho \{1 + o(1)\} \rightarrow 0.$$

Together with  $\sqrt{k}A(n/k) \rightarrow \lambda$ , this yields

$$\frac{\sqrt{k}}{\ell_n} a_n \rightarrow 0.$$

The imposed bridge-rate condition (3.3) is exactly

$$\frac{\sqrt{k}}{\ell_n} b_n \rightarrow 0.$$

Therefore all linear terms in  $C_n$  are  $o(1)$  after multiplication by  $\sqrt{k}/\ell_n$ . The logarithmic square remainder is also negligible, since

$$\frac{\sqrt{k}}{\ell_n} (|a_n| + b_n)^2 \rightarrow 0$$

follows from  $(\sqrt{k}/\ell_n)|a_n| \rightarrow 0$ ,  $(\sqrt{k}/\ell_n)b_n \rightarrow 0$ , and  $|a_n| + b_n \rightarrow 0$ . Hence

$$\frac{\sqrt{k}}{\ell_n} C_n \rightarrow 0. \tag{3.8}$$

Combining (3.6), (3.7), and (3.8) in the exact decomposition gives (3.4) by Slutsky's theorem.

For the relative-error statement, define

$$L_n = \log \frac{\hat{\xi}_{\tau_n}^{\text{pool}, \star}(\boldsymbol{\nu}, \boldsymbol{\omega})}{\xi_{\tau_n}}.$$

The log limit implies  $L_n = O_P(\ell_n/\sqrt{k}) = o_P(1)$ . Therefore

$$e^{L_n} - 1 = L_n + O_P(L_n^2)$$

and

$$\frac{\sqrt{k}}{\ell_n} \{e^{L_n} - 1 - L_n\} = O_P\left(\frac{\ell_n}{\sqrt{k}}\right) = o_P(1).$$

This proves (3.5). □

*Remark 3.3* (What the theorem does and does not identify). The limit in Theorem 3.2 depends on  $\boldsymbol{\omega}$ , the geometric pooled-Weissman weights, but not on  $\boldsymbol{\nu}$ , the bridge-Hill weights. Thus  $\boldsymbol{\nu}$  is first-order unidentified in the present regime. The diagonal estimator  $\boldsymbol{\nu} = \boldsymbol{\omega}$  is a permissible

deterministic convention, and the practical convention used for plug-in inference below is

$$\boldsymbol{\nu}_n^k = \left( \frac{k_1}{k}, \dots, \frac{k_m}{k} \right)^\top.$$

Neither convention is an optimality result for  $\boldsymbol{\nu}$ . Any optimal choice of bridge-Hill weights would require a lower-order theory not developed here.

### 3.3 Weight consequences

The preceding theorem determines the relevant first-order weight criterion. The leading bias and variance coincide with those of the Daouia–Padoan–Stupfler pooled-Weissman component in the distributed common-marginal case. Define

$$\mathbf{B}_c^{\text{dist}} = \frac{\lambda}{1-\rho} (d_1^\rho, \dots, d_m^\rho)^\top, \quad \mathbf{V}_c^{\text{dist}} = \gamma^2 \left( \sum_{i=1}^m c_i^{-1} \right) \text{diag}(c_1, \dots, c_m).$$

Then  $B_\omega = \omega^\top \mathbf{B}_c^{\text{dist}}$  and  $V_\omega = \omega^\top \mathbf{V}_c^{\text{dist}} \omega$ .

**Corollary 3.4** (First-order criteria for  $\omega$ ). *Under the assumptions of [Theorem 3.2](#), the first-order variance criterion for the pooled extreme-expectile estimator is*

$$\omega^\top \mathbf{V}_c^{\text{dist}} \omega,$$

and the first-order AMSE criterion is

$$(\omega^\top \mathbf{B}_c^{\text{dist}})^2 + \omega^\top \mathbf{V}_c^{\text{dist}} \omega,$$

both under the affine constraint  $\omega^\top \mathbf{1} = 1$ . Hence the deterministic/oracle variance- and AMSE-optimal formulas of Daouia, Padoan, et al. (2024) apply to  $\omega$ . In particular,

$$\omega^{\text{Var}} = \frac{(\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{1}}{\mathbf{1}^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{1}},$$

so, in the independent distributed case,

$$\omega_j^{\text{Var}} = \frac{c_j^{-1}}{\sum_{i=1}^m c_i^{-1}}, \quad j = 1, \dots, m.$$

If  $\mathbf{V}_c^{\text{dist}}$  is positive definite, the AMSE-optimal weights are

$$\omega^{\text{AMSE}} = \frac{(1 + (\mathbf{B}_c^{\text{dist}})^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{B}_c^{\text{dist}}) (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{1} - (\mathbf{1}^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{B}_c^{\text{dist}}) (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{B}_c^{\text{dist}}}{(1 + (\mathbf{B}_c^{\text{dist}})^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{B}_c^{\text{dist}}) (\mathbf{1}^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{1}) - (\mathbf{1}^\top (\mathbf{V}_c^{\text{dist}})^{-1} \mathbf{B}_c^{\text{dist}})^2}.$$

No first-order variance or AMSE criterion for  $\boldsymbol{\nu}$  is identified by [Theorem 3.2](#).

*Proof.* By [Theorem 3.2](#),

$$\frac{\sqrt{k}}{\ell_n} \log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\boldsymbol{\nu}, \omega)}{\xi_{\tau_n}} \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega).$$

Thus the scaled first-order mean-square criterion is  $B_\omega^2 + V_\omega$ , and the unscaled criterion differs only by the common factor  $\ell_n^2/k$ . The same criterion applies to relative error by (3.5). Substituting  $B_\omega = \omega^\top B_c^{\text{dist}}$  and  $V_\omega = \omega^\top V_c^{\text{dist}} \omega$  gives exactly the quadratic criteria recalled in Proposition 2.11. The variance and AMSE formulas follow from that result.  $\square$

### 3.4 Estimated pooled-Weissman weights and inference

The deterministic theorem is the base result. Estimated weights are introduced only in the pooled-Weissman role  $\omega$ , where Daouia, Padoan, et al. (2024) already provide random-weight quantile limits under their own consistency conditions. The bridge-Hill vector remains deterministic in this chapter.

**Corollary 3.5** (Estimated- $\omega$  transfer). *Assume the conditions of Theorem 3.2, with the deterministic  $\omega$  in the estimator replaced by a random  $\hat{\omega}_n$  satisfying the estimated-weight conditions of Proposition 2.14, with limiting vector  $\omega$ . Let  $\nu$  be deterministic and affine, or let  $\nu = \nu_n^k = (k_1/k, \dots, k_m/k)^\top$ . Then*

$$\frac{\sqrt{k}}{\ell_n} \log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\nu, \hat{\omega}_n)}{\xi_{\tau_n}} \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega).$$

The same limit holds on relative-error scale.

*Proof.* The exact decomposition becomes

$$\log \frac{\hat{\xi}_{\tau_n}^{\text{pool},*}(\nu, \hat{\omega}_n)}{\xi_{\tau_n}} = A_n(\hat{\omega}_n) + B_n(\nu) - C_n.$$

Proposition 2.14 gives the relative-error limit for the pooled-Weissman component. The same Taylor conversion used in the proof of Theorem 3.2 gives

$$\frac{\sqrt{k}}{\ell_n} A_n(\hat{\omega}_n) \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega).$$

The bridge-Hill term is unchanged. If  $\nu$  is fixed, the proof of (3.7) applies verbatim. If  $\nu = \nu_n^k$ , then  $\nu_n^k$  is deterministic, affine, bounded, and convergent under the threshold-proportion assumptions; the vector pooled Hill CLT again gives  $\hat{\gamma}_n(\nu_n^k) - \gamma = O_P(k^{-1/2})$ , so the same Taylor bound applies. The population bridge term  $C_n$  is deterministic and is still controlled by (3.8). Slutsky's theorem gives the log limit, and the exponential Taylor step gives the relative-error version.  $\square$

**Corollary 3.6** (Plug-in studentisation and log-scale interval). *Let*

$$\hat{\xi}_{\tau_n}^{\text{plug}} = \hat{\xi}_{\tau_n}^{\text{pool},*}(\nu_n^k, \hat{\omega}_n), \quad \nu_n^k = \left( \frac{k_1}{k}, \dots, \frac{k_m}{k} \right)^\top.$$

Under the assumptions of [Corollary 3.5](#), let  $\widehat{\mathbf{B}}_{c_n}^{\text{dist}}$  and  $\widehat{\mathbf{V}}_{c_n}^{\text{dist}}$  be plug-in estimators satisfying

$$\widehat{\mathbf{B}}_{c_n}^{\text{dist}} \xrightarrow{\mathbb{P}} \mathbf{B}_c^{\text{dist}}, \quad \widehat{\mathbf{V}}_{c_n}^{\text{dist}} \xrightarrow{\mathbb{P}} \mathbf{V}_c^{\text{dist}}.$$

Define

$$\widehat{B}_{\widehat{\omega},n} = \widehat{\omega}_n^\top \widehat{\mathbf{B}}_{c_n}^{\text{dist}}, \quad \widehat{V}_{\widehat{\omega},n} = \widehat{\omega}_n^\top \widehat{\mathbf{V}}_{c_n}^{\text{dist}} \widehat{\omega}_n,$$

and assume  $V_\omega > 0$ . Then

$$T_n^{\text{plug}} = \frac{\frac{\sqrt{k}}{\ell_n} \log\left(\frac{\widehat{\xi}_{\tau_n}^{\text{plug}}}{\xi_{\tau_n}}\right) - \widehat{B}_{\widehat{\omega},n}}{\sqrt{\widehat{V}_{\widehat{\omega},n}}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Let  $z_{1-\alpha/2}$  be the standard normal quantile. On the event  $\widehat{V}_{\widehat{\omega},n} > 0$ , define

$$I_{n,1-\alpha}^{\text{plug}} = \left[ \widehat{\xi}_{\tau_n}^{\text{plug}} \exp\left\{-\frac{\ell_n}{\sqrt{k}} \left(\widehat{B}_{\widehat{\omega},n} + z_{1-\alpha/2} \sqrt{\widehat{V}_{\widehat{\omega},n}}\right)\right\}, \right. \\ \left. \widehat{\xi}_{\tau_n}^{\text{plug}} \exp\left\{-\frac{\ell_n}{\sqrt{k}} \left(\widehat{B}_{\widehat{\omega},n} - z_{1-\alpha/2} \sqrt{\widehat{V}_{\widehat{\omega},n}}\right)\right\} \right].$$

Then

$$\mathbb{P}\left\{\xi_{\tau_n} \in I_{n,1-\alpha}^{\text{plug}}\right\} \rightarrow 1 - \alpha.$$

*Proof.* [Corollary 3.5](#) gives

$$\frac{\sqrt{k}}{\ell_n} \log\left(\frac{\widehat{\xi}_{\tau_n}^{\text{plug}}}{\xi_{\tau_n}}\right) \xrightarrow{d} \mathcal{N}(B_\omega, V_\omega).$$

[Proposition 2.15](#), applied with  $\mathbf{B}_c = \mathbf{B}_c^{\text{dist}}$  and  $\mathbf{V}_c = \mathbf{V}_c^{\text{dist}}$ , gives

$$\widehat{B}_{\widehat{\omega},n} \xrightarrow{\mathbb{P}} B_\omega, \quad \widehat{V}_{\widehat{\omega},n} \xrightarrow{\mathbb{P}} V_\omega.$$

Together with  $V_\omega > 0$ , these convergences give the standard normal limit by Slutsky's theorem. Since  $\widehat{V}_{\widehat{\omega},n} \xrightarrow{\mathbb{P}} V_\omega > 0$ , the event  $\widehat{V}_{\widehat{\omega},n} > 0$  has probability tending to one. Solving

$$\left|T_n^{\text{plug}}\right| \leq z_{1-\alpha/2}$$

for the unknown  $\xi_{\tau_n}$  gives the displayed multiplicative interval, so its asymptotic coverage follows from the studentised limit.  $\square$

*Remark 3.7* (Scope of the inference corollary). The interval in [Corollary 3.6](#) is obtained by inverting the plug-in-centred expectile statistic above. It is therefore a consequence of the transferred expectile limit, rather than a direct restatement of the quantile interval in Daouia, Padoan, et al. (2024). The centering and variance objects remain those of the pooled-Weissman component because, under  $\eta_n \rightarrow 0$ , the additional expectile-bridge terms are lower order on that scale.



# Bibliography

- Bellini, F., B. Klar, A. Müller, and E. Rosazza Gianin (2014). “Generalized quantiles as risk measures”. In: *Insurance: Mathematics and Economics* 54, pp. 41–48.
- Daouia, A., S. Girard, and G. Stupfler (2018). “Estimation of tail risk based on extreme expectiles”. In: *Journal of the Royal Statistical Society. Series B* 80.2, pp. 263–292. DOI: [10.1111/rssb.12254](https://doi.org/10.1111/rssb.12254).
- (2019). “Extreme M-quantiles as risk measures: From  $L^1$  to  $L^p$  optimization”. In: *Bernoulli* 25.1, pp. 264–309. DOI: [10.3150/17-BEJ987](https://doi.org/10.3150/17-BEJ987).
- (2020). “Tail expectile process and risk assessment”. In: *Bernoulli* 26.1, pp. 531–556. DOI: [10.3150/19-BEJ1137](https://doi.org/10.3150/19-BEJ1137).
- Daouia, A., S. A. Padoan, and G. Stupfler (2024). “Optimal weighted pooling for inference about the tail index and extreme quantiles”. In: *Bernoulli* 30.2, pp. 1287–1312. DOI: [10.3150/23-BEJ1632](https://doi.org/10.3150/23-BEJ1632). arXiv: [2111.03173](https://arxiv.org/abs/2111.03173).
- Davison, A. C., S. A. Padoan, and G. Stupfler (2023). “Tail risk inference via expectiles in heavy-tailed time series”. In: *Journal of Business & Economic Statistics* 41.3, pp. 876–889. DOI: [10.1080/07350015.2022.2078332](https://doi.org/10.1080/07350015.2022.2078332).
- Ehm, W., T. Gneiting, A. Jordan, and F. Krüger (2016). “Of quantiles and expectiles: Consistent scoring functions, Choquet representations and forecast rankings”. In: *Journal of the Royal Statistical Society. Series B* 78.3, pp. 505–562.
- Gneiting, T. (2011). “Making and evaluating point forecasts”. In: *Journal of the American Statistical Association* 106.494, pp. 746–762.
- Haan, L. de and A. Ferreira (2006). *Extreme Value Theory: An Introduction*. Springer Series in Operations Research and Financial Engineering. New York: Springer.
- Hill, B. M. (1975). “A simple general approach to inference about the tail of a distribution”. In: *The Annals of Statistics* 3.5, pp. 1163–1174.
- Newey, W. K. and J. L. Powell (1987). “Asymmetric least squares estimation and testing”. In: *Econometrica* 55.4, pp. 819–847.
- Weissman, I. (1978). “Estimation of parameters and large quantiles based on the  $k$  largest observations”. In: *Journal of the American Statistical Association* 73.364, pp. 812–815.
- Ziegel, J. F. (2016). “Coherence and elicibility”. In: *Mathematical Finance* 26.4, pp. 901–918. DOI: [10.1111/mafi.12080](https://doi.org/10.1111/mafi.12080).