

Alessandro Frasca, Roderik Bruce, Francesco Cerutti, CERN, Geneva, Switzerland
 Alessia Ciccottelli, University of Manchester, Manchester, UK and CERN, Geneva, Switzerland
 Marcin Patecki, Warsaw University of Technology, Poland

LHC heavy ion operation

- fully-stripped lead ions $^{208}\text{Pb}^{+82}$
- up to 7 Z TeV beam energy
- Pb-Pb collisions at the four experiments ATLAS, CMS, ALICE and LHCb
- electromagnetic interactions dominate over nuclear inelastic ones


Bound-free pair production (BFPP)



- highest cross section
 - small transverse momentum recoil
 - secondary beams with well-defined magnetic rigidity change
 - localised power deposition downstream the IP
- $$P_p = \mathcal{L} \sigma_p E_b$$
- risk of beam dumps or magnet quenches
 - luminosity limitations

BFPP at LHCb

- so far luminosity levelling to a safe value of $10^{27} \text{cm}^{-2} \text{s}^{-1}$
- recent request for an **higher integrated luminosity** until the end of LHC Run 4
- mitigation strategy for BFPP** to increase LHCb peak luminosity
- TCLD collimators, as adopted for ALICE, presently not foreseen




- B1 BFPP peak loss at LHCb measured by beam loss monitors
- physics operation in 2018

SixTrack simulation goals

- tracking of BFPP particles from LHCb
- study of loss patterns
- investigation of orbit bumps as mitigation strategy


BFPP beam tracking from LHCb

- 2018 LHC heavy-ion optics
- both 2018 and design beam parameters
- dispersive trajectory of $^{208}\text{Pb}^{+81}$ ions mimicked by tracking $^{208}\text{Pb}^{+82}$ ions with a fractional momentum offset $\delta = 1/81$
- initially distributed as the colliding ions
- BFPP beams hit the superconductive dipoles in cell 10 MB.B10R8.B1 and MB.B10L8.B2



Orbit bumps as alleviation strategy for LHCb

- horizontal orbit bumps** of -5.3 mm on MQML.10R8.B1 and +5 mm on MQML.10L8.B2
- BFPP losses shifted from cell 10 to cell 12
- simulation of new loss patterns show new impact locations on superconductive dipoles MB.C12R8.B1 and MB.C12L8.B2
- in cell 12 **larger β -function** and hence larger BFPP beam transverse size
- BFPP power deposition more spread out



- a full energy deposition study with FLUKA using these results as input predicts a peak power load **lower by a factor of 4** in cell 12
- the levelled luminosity at LHCb could be safely increased by a factor 2-3** in future Pb-Pb runs