Uniform Convergence of a Sequence of Functions

Note 1

1bf1a79b9eba47cf852e1a9c7468c5f7

Let (f_n) be well a sequence of function on a set A. We say we say we converges pointwise on A to a function f if we for all $x \in A$

$$\left(f_n(x)\right) \underset{n \to \infty}{\longrightarrow} f(x).$$

,,

Note 2

11dc20a5619424cafc97ab1b4d64b5f

Let (f_n) be a sequence of function on a set A. If (f_n) converges pointwise on A to f, we write

$$\text{ (cl::} f_n \to f \text{)} \quad \text{or} \quad \text{ (cl::} \lim_{n \to \infty} f_n = f. \text{)}$$

Note 3

6f3f051b9e0741dcbd85037d47c4fd19

Let
$$f_n(x) = \frac{x^2 + nx}{n}$$
.

$$\lim_{n\to\infty}f_n(x)=\text{\{c1::}x.\text{\}}$$

Note 4

3c7731c6b70a4c28972a5ea2e88a1e5f

Let
$$f_n(x) = x^n$$
, $f_n : [0,1] \to \mathbb{R}$.

$$\lim_{n o \infty} f_n(x) = \sup \left\{ egin{aligned} 0 & ext{for } 0 \leq x < 1, \ 1 & ext{for } x = 1. \end{aligned}
ight.$$

Note 5

7218c9c8b0f04d4887dc2345da75c6c6

Let (f_n) be a sequence of function on a set A. We say $\{(c^2)^n (f_n)\}$ converges uniformly on A to a function f_n if $\{(c^2)^n (f_n)\}$

$$\forall \epsilon > 0 \quad \exists N \in \mathbf{N} \quad \forall n \ge N$$

 $|f_n - f| < \epsilon.$

}}

Let (f_n) be a sequence of function on a set A. If (f_n) converges uniformly on A to f, we write (f_n)

$$f_n \rightrightarrows f$$
.

}}

Note 7

77ef924775b2453cb303b726f3081917

What is the key distinction between the definitions of pointwise and uniform convergences of a sequence of functions?

The dependence of N on x.

Note 8

42d2e1017eac4382878c195aa5a4c54d

What is the visual behind the uniform convergence of a sequence of functions?

Eventually every f_n is completely contained in the ϵ -strip.

Note 9

0c853e2f4ed04acf9dae0b00c1a751f3

Which is stronger, uniform or pointwise convergence?

Uniform convergence is stronger.

Note 10

ed7804cf8d4d48d5b0efb426d130fb52

Uniform convergence implies (convergence.)

Note 11

c9h4c187h4d54a78a9500289aa5899d

Let (f_n) be a sequence of function on a set A.

$$\text{((c2::} f_n \Longrightarrow f \text{))} \quad \text{((c3::} \Longleftrightarrow \text{))} \quad \text{((c1::} \sup \left| f_n - f \right| \underset{n \to \infty}{\longrightarrow} 0.\text{))}$$

(in terms of sup)

Let (f_n) be a sequence of function on a set A. (case Then (f_n) converges uniformly on A)) (case if and only if)

$$\{\{\text{c1::} \forall \varepsilon > 0 \quad \exists N \in \mathbf{N} \quad \forall m, n \geq N\}\}$$

Note 13

b9e4671775a43e9aa4a6b4d581b1658

Let (f_n) be a sequence of function on a set A. Then $f_n \rightrightarrows f$ if and only if

$$\forall \varepsilon > 0 \quad \exists N \in \mathbf{N} \quad \forall m, n \ge N$$

$$|f_n - f_m| < \varepsilon.$$

«{{c1::Cauchy Criterion}}»

Note 14

3fa98b94397f4cc2b2d766dd41934f67

What is the key idea in the proof of necessity of the Cauchy Criterion for uniform convergence?

Follows immediately from the definition.

Note 15

f2d15c9af98f4b82956e48ed7df71fc9

What is the key idea in the proof of sufficiency of the Cauchy Criterion for uniform convergence?

Define a candidate for the limit and prove by definition.

Note 16

1525b27207e74da186a95d7656e895da

In the proof of sufficiency of the Cauchy Criterion for uniform convergence, how do you define a candidate for the limit?

Use the pointwise limit.

In the proof of sufficiency of the Cauchy Criterion for uniform convergence, how do we know the pointwise limit exists?

Due to the Cauchy Criterion for sequences.

Note 18

a1ccae80d31b4f38a4fc876e1ffe4ae7

In the proof of sufficiency of the Cauchy Criterion for uniform convergence, we have $f_n \to f$. How do you show that $f_n \rightrightarrows f$?

Take the limit of the inequality from the Cauchy Criterion.

Note 19

baab958475694fc08316e2031a57fa58

Let $f_n \to f$ on a set A and $c \in A$. If (can the convergence is uniform)) and (can all f_n are continuous at c.)) then (can f is continuous at c.))

Note 20

026cf3ddb2f4d5b9a94b36b2bc20ef9

Let $f_n \to f$ on a set A and $c \in A$. If the convergence is uniform and all f_n are continuous at c, then f is continuous at c.

«{{c1::Continuous Limit Theorem}}»

Note 21

5fd08fca82504ff0af82d320da351ff7

What is the key idea in the proof of the Continuous Limit Theorem for a sequence of functions?

Triple triangle inequality after adding and subtracting f_N .

Note 22

06425162bee447479d3a4f5c71c9cf2a

Let $f_n \to f$ on a set A and $c \in A$. If we the convergence is uniform and all f_n are continuous at c, then

$$\lim_{x \to c} \lim_{n \to \infty} f_n(x) = \lim_{x \to c} \lim_{n \to \infty} \lim_{x \to c} f_n(x).$$

Let $f_n \to f$ on a set A. If each f_n is continuous, but f is discontinuous, then {convergence is not uniform.}

Note 24

5ee2f3836bd4545afde8c2d7ecda40e

Give an example of a sequence of functions $f_n \to f$ such that

- each f_n is continuous almost everywhere; and
- *f* is nowhere continuous.
- Step-by-step construction of the Dirichlet's function.

Note 25

31c5e1a2081241d1973bb2cacde92627

Assume $f_n \to f$ on a set A and each f_n is uniformly continuous. If $\{(c) = f_n \rightrightarrows f_n\}$ then $\{(c) = f \in f \text{ is uniformly continuous.}\}$

Note 26

f819f1c60074468ba1e718298059ade4

Assume $f_n \to f$ on a set A and each f_n is bounded. If $\{\{e^2\}: f_n \rightrightarrows f, \}$ then $\{\{e^1\}: f \text{ is bounded.}\}$

Note 27

b1fded6e729d40ba99a9d087781866dd

Assume $f_n \to f$ on a set A and each f_n has a finite number of discontinuities. If $f_n \rightrightarrows f$, then (c) f has at most a countable number of discontinuities.

Note 28

a010908ba95d473ea734442288757314

Assume $f_n \rightrightarrows f$ on a set A and $c \in A$. If $\{c \in F\}$ is discontinuous at c, then $\{c \in A\}$ are eventually discontinuous at c.

Note 29

5ca0ebc56cc947d1bc6a5ed00cd1617l

Assume $f_n \rightrightarrows f$ on a set A and $c \in A$. If f is discontinuous at c, then all f_n are eventually discontinuous at c. What is the key idea in the proof?

By contradiction + choose a subsequence continuous at c.

Note 30

4c8d50b955be4fa0a3ba792c5699174f

Let f be (c2::continuous) on all of ${f R}$. Then $f(x+{1\over n})$ (c1::converges to f.)

Note 31

59f59d25a40a4e72afdd62a2dd24bd1

Let f be {{c2:}uniformly continuous}} on all of ${\bf R}$. Then $f(x+\frac{1}{n})$ {{c1:}converges uniformly to f.}

Uniform Convergence and Differentiation

Note 1

7f46dbb00f54423a835a842d402aa10

What sequence is considered in the Differentiable Limit Theorem?

A sequence of differentiable functions that converges pointwise on a closed interval.

Note 2

19574e41800e43678628e78581f801ce

When applying the Differentiable Limit Theorem, is it necessary for the limit to be differentiable?

No, this is one of the implications.

Note 3

5ef400e26d2541e589faa672492059bf

When do we conclude something form the Differentiable Limit Theorem?

When the derivatives converges uniformly.

Note 4

f7da48c586d2457baad72d900c07defd

What do we conclude from The Differentiable Limit Theorem?

The limit f is differentiable and $f' = \lim f'_n$.

Note 5

61acf9aeed834980a9dbaa77746b89e0

Let $f_n \to f$ on [a, b] and each f_n is differentiable. What do we know about f if $f'_n \to g$?

Nothing special.

Note 6

63a1ccb4818a4cd281f9b4d9513500a0

Let $f_n \to f$ on [a, b] and each f_n is differentiable. What do we know about f if $f'_n \rightrightarrows g$?

f is differentiable and f' = g.

Note 7

a720c08c553f46a0b0423c46f4c19a2e

What is the key idea in the proof of the Differentiable Limit Theorem?

Rewrite the limit's derivative by definition.

Note 8

31222913007d4ceda945e1a21642c876

In the proof of the Differentiable Limit Theorem, how do you find an upper bound for

$$\left| \frac{f(x+h) - f(x)}{h} - g(x) \right| ?$$

Expand it using the triple triangle inequality involving f_N .

Note 9

d239aa3eedd346a69139a5a8b1d94ce7

In the proof of the Differentiable Limit Theorem, how do you choose N?

By the Cauchy Criterion for $f'_n \rightrightarrows g$.

Note 10

70bbcff5bceb49c7b0abb25a8ab9be35

In the proof of the Differentiable Limit Theorem, how do you find an upper bound for

$$|f_N'(x) - g(x)|?$$

Take the limit of the inequality from the Cauchy Criterion.

Note 11

ee6b7ee23d0a48a8a32afe978be50a7d

In the proof of the Differentiable Limit Theorem, how do you find an upper bound for

$$\left| \frac{f_N(x+h) - f_N(x)}{h} - f_N'(x) \right| ?$$

Pick δ by the definition of differentiability of f_N .

Note 12

cdb10b03a9254c5abfe796106c1d3e9b

In the proof of the Differentiable Limit Theorem, how do you find an upper bound for

$$\left| \frac{f(x+h) - f(x)}{h} - \frac{f_N(x+h) - f_N(x)}{h} \right| ?$$

The Mean Value Theorem for $f_N - f_m$ and make $m \to \infty$.

Note 13

o4b2753226ff4d839269bbf795c0230

Let (f_n) be noted a sequence of differentiable functions on [a,b] and noted are uniformly. If noted $f_n(x_0)$ exists for some x_0 , then noted converges uniformly.

Note 14

8c542d7e30524e129805ce26973b0925

How can we weaken the hypothesis of the Differentiable Limit Theorem?

 (f_n) converges at a single point.

Series of Functions

Note 1

h2a393ededc84241bh594417273dea7e

Let (f_n) be (case a sequence of functions on a set A.) (case A functional series) is (case a formal expression of the form

$$\sum_{n=1}^{\infty} f_n(x).$$

Note 2

6291bcd4e0274102bfe4090eebac24e

Let (f_n) be a sequence of functions on a set A. We say $\sum_n f_n(x)$ we converges pointwise on A to a function f(x) if we have sequence of partial sums converges pointwise on A to f.

Note 3

084d4603478b4dc48c0d1837ff30dfd8

Let (f_n) be a sequence of functions on a set A. If $\{c^2 = \sum_n f_n(x) \}$ converges pointwise to f(x), we write $\{c^2 = \sum_n f_n(x) \}$

$$f(x) = \sum_{n} f_n(x).$$

}}

Note 4

2922cd6ac8ff42fabe5bc630fa320169

Let (f_n) be a sequence of functions on a set A. We say $\sum f_n(x)$ (converges uniformly on A to a function f(x)) if (contact the sequence of partial sums converges uniformly on A to f.)

Note 5

2b28ab51bc7f45ca934cc405e7de388f

Let $\sum_n f_n(x)$ be a functional series. (CLE) A series

$$\sum_{n=k+1}^{\infty} f_n(x) \quad \text{for } k \in \mathbf{N},$$

)} is called {{c2::a tail of $\sum_n f_n(x)$.}}

A series $\sum_n f_n(x)$ (converges pointwise) (confidential converges pointwise to 0.1)

(in terms of the tail)

Note 7

16325daa37b14ddebc3939e1d2ea063b

A series $\sum_n f_n(x)$ (converges uniformly) (converges uniformly to 0.)

(in terms of the tail)

Note 8

891381b2ecd44c2cb160d114479f0b20

A series $\sum_n f_n(x)$ (c2::converges pointwise) (c3::only if) (c1:: $f_n o 0$.

Note 9

767a398cce7c40b781b0c39db5f9b9a

A series $\sum_n f_n(x)$ ((c2::converges uniformly)) ((c3::only if)) ((c1:: $f_n
ightharpoonup 0.$

Note 10

296676411bf5475eacdde73dc1c2b008

What series is considered in the Weierstrass M-Test?

• A series of bounded functions.

Note 11

5c393b177b724cf69790bafcf0ff7b23

When do we conclude something from the Weierstrass M-Test?

When the series of "absolute" bounds converges.

Note 12

f964f11937374d53be121d3893daeef6

Which bounds are considered in the Weierstrass M-Test?

The upper bound for the absolute value of $f_n(x)$.

Note 13

c48d5e20f5d24ca58a0c3bd71ab7b256

What do we conclude from the Weierstrass M-Test?

The functional series converges.

Note 14

f9827fda17c4670b0d2bd4728303ae

What is the key idea in the proof of the Weierstrass M-Test?

A corollary of the Cauchy Criterion.