

Basics

Note 1

21e64c2f0430467f8a36481045e172b3

$\{\{c2::\mathbb{Z}^+, \mathbb{Q}^+ \text{ and } \mathbb{R}^+\}\}$ denote $\{\{c1::\text{the positive (nonzero) elements in } \mathbb{Z}, \mathbb{Q} \text{ and } \mathbb{R}, \text{ respectively.}\}\}$

Note 2

ad32571de5d04cafb2a7d6a27aee4b14

Given a function $f : A \rightarrow B$, $\{\{c1::\text{the set } B\}\}$ is called the codomain of f .

Note 3

0cd492ad876a4dfbbb22c9210039fcc1

Given a function $f : A \rightarrow B$ and $\{\{c3::a \in B\}\}$ $\{\{c2::\text{the preimage of } \{b\} \text{ under } f\}\}$ is called $\{\{c1::\text{the fiber of } f \text{ over } b.\}\}$

Note 4

b2a02c66209140a591b43dede69ffbf1

If $f : A \rightarrow B$ and $g : B \rightarrow C$, then the $\{\{c1::\text{composite map}\}\}$

$$g \circ f : A \rightarrow C$$

is defined by

$$(g \circ f)(a) = g(f(a)).$$

Note 5

1b2bf2fe79dc4063a151a960f45698d9

A function $f : A \rightarrow B$ $\{\{c3::\text{has a left inverse}\}\}$ if there is a function $g : \{\{c2::B \rightarrow A\}\}$, such that $\{\{c1::$

$$g \circ f = id_A.$$

$\}\}$

Note 6

d9a63bd7866e44ab83172cf9189e9b9a

A function $f : A \rightarrow B$ $\{\{c3::\text{has a right inverse}\}\}$ if there is a function $g : \{\{c2::B \rightarrow A\}\}$, such that $\{\{c1::$

$$f \circ g = id_B.$$

$\}\}$

Note 7

8098015b57774529a721663e39eabb18

A map f is $\{\{c1::\text{injective}\}\}$ if and only if f has a $\{\{c2::\text{left}\}\}$ inverse.

Note 8

205100b0fd6447a9bcc94e4d7711a606

A map f is $\{\{c1::\text{surjective}\}\}$ if and only if f has a $\{\{c2::\text{right}\}\}$ inverse.

Note 9

8e4ddf27550f4aa9a1daf0b67cd2f7e4

A $\{\{c2::\text{permutation}\}\}$ of a set A is $\{\{c1::\text{a bijection from } A \text{ to itself.}\}\}$

Note 10

1feef80fbcd48618084ce93c88df83b

If $A \subseteq B$ and $f : B \rightarrow C$, $\{\{c1::\text{the restriction of } f \text{ to } A\}\}$ is denoted $\{\{c1::f|_A\}\}$

Note 11

01a5a3b0e5f24f6782e689090b17c437

If $A \subseteq B$ and $g : A \rightarrow C$ and there is a function $f : B \rightarrow C$ such that $\{\{c2::f|_A = g\}\}$ we shall say f is $\{\{c1::\text{an extension of } g \text{ to } B.\}\}$

Note 12

6ca7e478954c4c898718ce116219822f

$\{\{c2::\text{A binary relation on a set } A\}\}$ is $\{\{c1::\text{a subset } R \text{ of } A \times A.\}\}$

Note 13

50bb82cd97cb40bf8621065845545d18

Let R be a binary relation on a set A . We write $\{\{c2::A \sim B\}\}$ if $\{\{c1::(a, b) \in R.\}\}$

Note 14

65287096376a47f399a0048c0d8092d0

A binary relation R on A is said to be $\{\{c2::\text{reflexive}\}\}$ if $\{\{c1::$

$$a \sim a, \text{ for all } a \in A.$$

$\}\}$

Note 15

71b961a1f8f347dcbf7b9c7c8dec303c

A binary relation R on A is said to be $\{\{c2::\text{symmetric}\}\}$ if $\{\{c1::$

$$a \sim b \text{ implies } b \sim a \text{ for all } a, b \in A.$$

$\}\}$

Note 16

40964931d9594b2997437cc9e3e150cc

A binary relation R on A is said to be $\{\{c2::\text{transitive}\}\}$ if $\{\{c1::$

$a \sim b$ and $b \sim c$ implies $a \sim c$ for all $a, bc \in A$.

$\}\}$

Note 17

54a959a8e36045c1aea2d838ce8998b8

A binary relation is $\{\{c2::\text{an equivalence relation}\}\}$ if $\{\{c1::\text{it is reflexive, symmetric and transitive.}\}\}$

Note 18

7c28d643ddd74509b88cfe2f75e6d743

If \sim defines an $\{\{c3::\text{equivalence}\}\}$ relation on A , then $\{\{c2::\text{the equivalence class}\}\}$ of $a \in A$ is defined to be $\{\{c1::$

$$\{x \in A \mid x \sim a\}.$$

$\}\}$

Note 19

323fae73cd4b47ddb8c19cb515ffd4cf

If C is an equivalence class, $\{\{c2::\text{any element of } C\}\}$ is called $\{\{c1::\text{a representative of the class } C.\}\}$

Note 20

3a597e1d5c48420490d792b972a38fe6

$\{\{c2::\text{A partition of } A\}\}$ is $\{\{c3::\text{any collection } \{A_i \mid i \in I\} \text{ of nonempty subsets of } A\}\}$ such that $\{\{c1::A \text{ is the disjoint union of all } A_i.\}\}$

Note 21

c2216701429649b7a262afdd5c85a72d

If \sim defines an equivalence relation on A then $\{\{c2::\text{the set of equivalence classes of } \sim\}\}$ form $\{\{c1::\text{a partition of } A.\}\}$

Properties of the Integers

Note 1

f535d29c343f494fa35bccefc9d6988

Let $a, b \in \mathbb{Z}$. We write $\{\{c2:a \mid b\}\}$ if $\{\{c1:a \text{ divides } b\}\}$

Note 2

96293ae3b76348d8ba9f0b02c8b49a94

Let $a, b \in \mathbb{Z}$ with $a \neq 0$. We write $\{\{c2:a \nmid b\}\}$ if $\{\{c1:a \text{ does not divide } b\}\}$

Note 3

533403fe830341a39cee216314b861e8

Let $a, b \in \{\{c3:\mathbb{Z} - \{0\}\}\}$. $\{\{c2:\text{The greatest common divisor of } a \text{ and } b\}\}$ is denoted by $\{\{c1:(a, b)\}\}$

Note 4

20b204b896884b6b9d07ca3023b7cf4a

Let $a, b \in \{\{c3:\mathbb{Z} - \{0\}\}\}$. If $\{\{c2:(a, b) = 1\}\}$ we say that a and b are $\{\{c1:\text{relatively prime}\}\}$

Note 5

69adfe8820204997a5aa44c50b353a40

If $a, b \in \mathbb{Z} - \{0\}$, then there exists unique $q, r \in \mathbb{Z}$ such that

$$a = qb + r \text{ and } 0 \leq r < |b|,$$

where q is $\{\{c1:\text{the quotient}\}\}$ and r $\{\{c1:\text{the remainder}\}\}$

« $\{\{c2:\text{Division Algorithm}\}\}$ »

Note 6

6267be99c4884a09b1282d041ac05e18

If $a, b \in \mathbb{Z} - \{0\}$, then there exist $x, y \in \{\{c3:\mathbb{Z}\}\}$ such that

$$\{\{c2:(a, b)\}\} = \{\{c1:xa + yb\}\}$$

Note 7

e30ea564f2ce479391e71512867aea51

If p is prime and $p \mid ab$, for some $a, b \in \mathbb{Z}$, then $\{\{c1:$

either $p \mid a$ or $p \mid b$.

$\}\}$

Note 8

3cc931ead0ec4cddae21114d84f1de0c

The Euler φ -function is defined as follows: for $n \in \mathbb{Z}^+$ let $\varphi(n)$ be the number of positive integers $a \leq n$ with a relatively prime to n .

Note 9

03f37e11eb9d40d29ca92031ac27d9ed

Let φ stand for the Euler φ -function. If p is prime and $a \geq 1$, then

$$\varphi(p^a) = p^a - p^{a-1}.$$

Note 10

7dc766a783c04a309951678711bd8317

Let φ stand for the Euler φ -function. Then

$$\varphi(ab) = \varphi(a)\varphi(b) \quad \text{if } (a, b) = 1.$$