

# Basics

## Note 1

21e64c2f0430467f8a36481045e172b3

$\{\{c2::\mathbb{Z}^+, \mathbb{Q}^+ \text{ and } \mathbb{R}^+\}\}$  denote  $\{\{c1::\text{the positive (nonzero) elements in } \mathbb{Z}, \mathbb{Q} \text{ and } \mathbb{R}, \text{ respectively.}\}\}$

## Note 2

ad32571de5d04cafb2a7d6a27aee4b14

Given a function  $f : A \rightarrow B$ ,  $\{\{c1::\text{the set } B\}\}$  is called the codomain of  $f$ .

## Note 3

0cd492ad876a4dfbbb22c9210039fcc1

Given a function  $f : A \rightarrow B$  and  $\{\{c3::a \in B\}\}$   $\{\{c2::\text{the preimage of } \{b\} \text{ under } f\}\}$  is called  $\{\{c1::\text{the fiber of } f \text{ over } b.\}\}$

## Note 4

b2a02c66209140a591b43dede69ffb1

If  $f : A \rightarrow B$  and  $g : B \rightarrow C$ , then the  $\{\{c1::\text{composite map}\}\}$

$$g \circ f : A \rightarrow C$$

is defined by

$$(g \circ f)(a) = g(f(a)).$$

## Note 5

1b2bf2fe79dc4063a151a960f45698d9

A function  $f : A \rightarrow B$   $\{\{c3::\text{has a left inverse}\}\}$  if there is a function  $g : \{\{c2::B \rightarrow A\}\}$ , such that  $\{\{c1::$

$$g \circ f = id_A.$$

$\}\}$

## Note 6

d9a63bd7866e44ab83172cf9189e9b9a

A function  $f : A \rightarrow B$   $\{\{c3::\text{has a right inverse}\}\}$  if there is a function  $g : \{\{c2::B \rightarrow A\}\}$ , such that  $\{\{c1::$

$$f \circ g = id_B.$$

$\}\}$

**Note 7**

8098015b57774529a721663e39eabb18

A map  $f$  is  $\{\{c1::\text{injective}\}\}$  if and only if  $f$  has a  $\{\{c2::\text{left}\}\}$  inverse.

**Note 8**

205100b0fd6447a9bcc94e4d7711a606

A map  $f$  is  $\{\{c1::\text{surjective}\}\}$  if and only if  $f$  has a  $\{\{c2::\text{right}\}\}$  inverse.

**Note 9**

8e4ddf27550f4aa9a1daf0b67cd2f7e4

A  $\{\{c2::\text{permutation}\}\}$  of a set  $A$  is  $\{\{c1::\text{a bijection from } A \text{ to itself.}\}\}$

**Note 10**

1feef80fbcd48618084ce93c88df83b

If  $A \subseteq B$  and  $f : B \rightarrow C$ ,  $\{\{c2::\text{the restriction of } f \text{ to } A\}\}$  is denoted  $\{\{c1::f|_A\}\}$

**Note 11**

01a5a3b0e5f24f6782e689090b17c437

If  $A \subseteq B$  and  $g : A \rightarrow C$  and there is a function  $f : B \rightarrow C$  such that  $\{\{c2::f|_A = g\}\}$  we shall say  $f$  is  $\{\{c1::\text{an extension of } g \text{ to } B.\}\}$

**Note 12**

6ca7e478954c4c898718ce116219822f

$\{\{c2::\text{A binary relation on a set } A\}\}$  is  $\{\{c1::\text{a subset } R \text{ of } A \times A.\}\}$

**Note 13**

50bb82cd97cb40bf8621065845545d18

Let  $R$  be a binary relation on a set  $A$ . We write  $\{\{c2::a \sim b\}\}$  if  $\{\{c1::(a, b) \in R.\}\}$

**Note 14**

65287096376a47f399a0048c0d8092d0

A binary relation  $R$  on  $A$  is said to be  $\{\{c2::\text{reflexive}\}\}$  if  $\{\{c1::$

$$a \sim a, \text{ for all } a \in A.$$

$\}\}$

**Note 15**

71b961a1f8f347dcbf7b9c7c8dec303c

A binary relation  $R$  on  $A$  is said to be  $\{\{c2::\text{symmetric}\}\}$  if  $\{\{c1::$

$$a \sim b \text{ implies } b \sim a \text{ for all } a, b \in A.$$

$\}\}$

### Note 16

40964931d9594b2997437cc9e3e150cc

A binary relation  $R$  on  $A$  is said to be  $\{\{c2::\text{transitive}\}\}$  if  $\{\{c1::$

$a \sim b$  and  $b \sim c$  implies  $a \sim c$  for all  $a, bc \in A$ .

$\}\}$

### Note 17

54a959a8e36045c1aea2d838ce8998b8

A binary relation is  $\{\{c2::\text{an equivalence relation}\}\}$  if  $\{\{c1::\text{it is reflexive, symmetric and transitive.}\}\}$

### Note 18

7c28d643ddd74509b88cfe2f75e6d743

If  $\sim$  defines an  $\{\{c3::\text{equivalence}\}\}$  relation on  $A$ , then  $\{\{c2::\text{the equivalence class}\}\}$  of  $a \in A$  is defined to be  $\{\{c1::$

$$\{x \in A \mid x \sim a\}.$$

$\}\}$

### Note 19

323fae73cd4b47ddb8c19cb515ffd4cf

If  $C$  is an equivalence class,  $\{\{c2::\text{any element of } C\}\}$  is called  $\{\{c1::\text{a representative of the class } C.\}\}$

### Note 20

3a597e1d5c48420490d792b972a38fe6

$\{\{c2::\text{A partition of a set } A_i\}\}$  is  $\{\{c3::\text{any collection } \{A_i \mid i \in I\}$  of nonempty subsets of  $A\}\}$  such that  $\{\{c1::A$  is the disjoint union of all  $A_i.\}\}$

### Note 21

c2216701429649b7a262afdd5c85a72d

If  $\sim$  defines an equivalence relation on  $A$  then  $\{\{c2::\text{the set of equivalence classes of } \sim\}\}$  form  $\{\{c1::\text{a partition of } A.\}\}$

# Properties of the Integers

## Note 1

f535d29c343f494fa35bccefc9d6988

Let  $a, b \in \mathbb{Z}$ . We write  $\{\{c2:a \mid b\}\}$  if  $\{\{c1:a \text{ divides } b\}\}$

## Note 2

96293ae3b76348d8ba9f0b02c8b49a94

Let  $a, b \in \mathbb{Z}$  with  $a \neq 0$ . We write  $\{\{c2:a \nmid b\}\}$  if  $\{\{c1:a \text{ does not divide } b\}\}$

## Note 3

533403fe830341a39cee216314b861e8

Let  $a, b \in \{\{c3:\mathbb{Z} - \{0\}\}\}$ .  $\{\{c2:\text{The greatest common divisor of } a \text{ and } b\}\}$  is denoted by  $\{\{c1:(a, b)\}\}$

## Note 4

20b204b896884b6b9d07ca3023b7cf4a

Let  $a, b \in \{\{c3:\mathbb{Z} - \{0\}\}\}$ . If  $\{\{c2:(a, b) = 1\}\}$  we say that  $a$  and  $b$  are  $\{\{c1:\text{relatively prime}\}\}$

## Note 5

69adfe8820204997a5aa44c50b353a40

If  $a, b \in \mathbb{Z} - \{0\}$ , then there exists unique  $q, r \in \mathbb{Z}$  such that

$$a = qb + r \text{ and } 0 \leq r < |b|,$$

where  $q$  is  $\{\{c1:\text{the quotient}\}\}$  and  $r$   $\{\{c1:\text{the remainder}\}\}$

« $\{\{c2:\text{Division Algorithm}\}\}$ »

## Note 6

6267be99c4884a09b1282d041ac05e18

If  $a, b \in \mathbb{Z} - \{0\}$ , then there exist  $x, y \in \{\{c3:\mathbb{Z}\}\}$  such that

$$\{\{c2:(a, b)\}\} = \{\{c1:xa + yb\}\}$$

## Note 7

e30ea564f2ce479391e71512867aea51

If  $p$  is prime and  $p \mid ab$ , for some  $a, b \in \mathbb{Z}$ , then  $\{\{c1:$

either  $p \mid a$  or  $p \mid b$ .

$\}\}$

### Note 8

3cc931ead0ec4cddae21114d84f1de0c

The Euler  $\varphi$ -function is defined as follows: for  $n \in \mathbb{Z}^+$  let  $\varphi(n)$  be the number of positive integers  $a \leq n$  with  $a$  relatively prime to  $n$ .

### Note 9

03f37e11eb9d40d29ca92031ac27d9ed

Let  $\varphi$  stand for the Euler  $\varphi$ -function. If  $p$  is prime and  $a \geq 1$ , then

$$\varphi(p^a) = p^a - p^{a-1}.$$

### Note 10

7dc766a783c04a309951678711bd8317

Let  $\varphi$  stand for the Euler  $\varphi$ -function. Then

$$\varphi(ab) = \varphi(a)\varphi(b) \quad \text{if } (a, b) = 1.$$