On the complete integrability of the geodesic flow on pseudo-H type Lie groups

Wolfram Bauer

Leibniz Universität Hannover 30167 Hannover, Germany bauer@math.uni-hannover.de

Pseudo-H type groups form a subclass of all 2-step nilpotent Lie groups. They generalize the notion of Heisenberg and H-type groups and were introduced by P. Ciatti in 2000. More recently, the classification problem for such groups and the question whether they admit integral lattices have gained attention (see [2,3]). In a natural way such groups can be equipped with a left-invariant (pseudo)-Riemannian metric.

In this talk we discuss the problem of complete integrability of the geodesic flow on pseudo-H type groups. There are several sources for the explicit construction of sets of Poisson commuting first integrals. In particular, we explain how to obtain a maximal set which then proves the complete integrability. We mention that in the case of the Heisenberg group this problem has been previously solved in [4]. The problem of how to descend the first integrals to the quotient of the group by a standard lattice constructed in [3] is discussed. Finally, we address the question whether "one can hear the complete integrability of the geodesic flow". More precisely, we ask whether the complete integrability of the geodesic flow of a closed Riemannian manifold is determined by the spectral data. In this framework two examples (not of pseudo-H-type) by D. Schueth are presented.

This talk is based on the joint work [1] with Daisuke Tarama (Ritsumeikan University, Japan).

REFERENCES

- [1] W. Bauer, D. Tarama, On the complete integrability of the geodesic flow of pseudo-H-type Lie groups, to appear in: Analysis and Math. Phys.
- [2] K. Furutani, I. Markina, *Complete classification of pseudo H-type Lie algebras: I*, Geom. Dedicata 190 (2017), 23 51.
- [3] K. Furutani, I. Markina, *Existence of lattice on general H-type groups*, J. Lie Theory 24, 979 1011.
- [4] A. Kocsard, G.P. Ovando, S. Reggiani, *The geodesic flow on nilmanifolds*, Differential Geom. Appl. 49 (2016), 496 509.