THE ROOT FUNCTOR

FRANCESCA PRATALI

ABSTRACT. In this paper, we show that any ∞-operad is equivalent to the localization of a discrete Σ-free operad; this result extend Joyal's delocalization theorem for categories to the operadic setting. Along the way, we pursue a systematic study of ∞-operadic localization in the dendroidal context and its compatibility with un/straightening equivalences, deducing another description of algebras over ∞-operads.

CONTENTS

Introduction and main results 1
1. Dendroidal recollections 4
2. Localization of dendroidal ∞-operads 9
3. The delocalization theorem 11
4. Localization and un/straightening equivalences 18

INTRODUCTION AND MAIN RESULTS

Introduction. Operads were introduced by May [May06] and Boardman-Vogt [BV73] in order to describe the up-to-homotopy algebraic structures on iterated loop spaces. In modern homotopy theory, one works with topological spaces up to weak homotopy equivalence, and for this purpose the language of ∞-*operads* offers a tractable way to articulate homotopy-coherence. The theory of ∞-operads has led to advancements in mathematical questions that predate its formulation ([GW99], [Wei99], [BdBW13], [KK24], [Lur17]).

Intuitively, a (colored) ∞-operad $\mathcal{P}$ is given by a set of objects and, for any choice of objects $x_1, \dots, x_n, y$, a space of operations $\mathcal{P}(x_1, \dots, x_n; y)$, together with operadic partial composition laws

$$\circ_{x_i} \colon P(x_1, \dots, x_k; y) \times P(z_1, \dots, z_m; x_i) \longrightarrow P(x_1, \dots, x_{i-1}, z_1, \dots, z_m, x_{i+1}, \dots, x_k; y)$$

equivariant with respect to some specified action of the symmetric groups given by 'permuting the variables'.



FIGURE 1. A graphical representation of the operadic composition $\circ_{x_2}$.

A $\mathcal{P}$-algebra is a family of spaces $F = \{F(x)\}_{x \in \mathsf{Ob}(P)}$ on which the operad acts via homotopy coherent maps

$$P(x_1, \dots, x_n; y) \longrightarrow \mathsf{Map}(F(x_1) \times \cdots \times F(x_n); F(y)).$$

Key words and phrases. Dendroidal sets, ∞-operads, localization, covariant model structure.

We say that $\mathcal{P}$ is *discrete* if it is 0-truncated, that is, if its spaces of operations are sets.

Given an ∞-operad $\mathcal{P}$ and a class of unary morphisms $\mathcal{W}$, the *localization* of $\mathcal{P}$ at $\mathcal{W}$ is the ∞-operad $\mathcal{L}_{\mathcal{W}}\mathcal{P}$ obtained by freely inverting the arrows in $\mathcal{W}$. Expressing an ∞-operad $\mathcal{Q}$ as a localization $\mathcal{L}_{\mathcal{W}}\mathcal{P}$ can allow to infer properties of $\mathcal{Q}$ from those of $\mathcal{P}$, notably about the ∞-categories of algebras over the operads.

Localization of ∞-operads can be approached from two sides. First, it can be seen as a generalization of localization of ∞-categories, in light of the fully faithful inclusion $\mathsf{Cat}_\infty \hookrightarrow \mathsf{Op}_\infty$. Second, it can be used to deduce descriptions of the ∞-category of algebras over an ∞-operads as some 'locally constant' one over another. In this regard, localization plays for instance an important role in the theory of $\mathbb{E}_n$-algebras, factorization algebras and factorization homology. In its simplest form, operadic localization appears in the relationship between the little n-disks operad $\mathbb{E}_n$ and the discrete operad of disks Disk_n. By [Lur17, Theorem 5.4.5.9], the natural map of ∞-operads $\mathcal{N}\mathsf{Disk}_n^{\otimes} \to \mathbb{E}_n$ is a localization at the preimage $\mathcal{J}$ of equivalences, inducing an equivalence $\mathsf{Alg}_{\mathbb{E}_n} \simeq \mathsf{Alg}^{\mathcal{J}}_{\mathsf{Disk}_n}$; analogous statements hold for disk operads on n-(stratified, framed) manifolds (see [AFT17], [KSW24]) and are crucial for identifying certain colimits and operadic Kan extensions.

It is natural to ask whether this behaviour is specific to little disks operads or holds more generally for *every* ∞-operad. To answer to this question, we go back to the perspective of ∞-operads as a generalization of ∞-categories. By *Joyal's delocalization theorem*, formulated in [Joy07, §13.6] by Joyal (a first construction of the map appearing in [Wal85, §1.6]) and proven by Stevenson in [Ste17], we know that it always happens that an ∞-category is equivalent to the localization of a discrete one.

move this to later

In this paper, we study localization of ∞-operads through the formalism of dendroidal sets. Our main result consists in extending Joyal's delocalization theorem to a delocalization theorem for every ∞-operad. This will allow us to give statements also for symmetric monoidal ∞-categories, and to deduce a 'locally constant' description for algebras over ∞-operads. Let us give a more precise overview of our main results in the next section.

to add only if I expand on this

Main results. As already anticipated, the starting point for our operadic delocalization theorem is Joyal's delocalization theorem for ∞-categories, which we recall in the following

Theorem (Joyal delocalization theorem)**.** *Let X be a simplicial set, and consider its category of elements Δ/X. The objects are strings of composable morphisms in X, i.e. a map $\alpha\colon \Delta^n \to X$. We denote such an object by $([n],\alpha)$ and represent it as $\alpha(0) \to \cdots \to \alpha(n)$. There is a morphism from the nerve of this category back to X,*

$$\mathcal{N}(\Delta/X) \longrightarrow X \quad ([n],\alpha) \mapsto \alpha(n)\ ;$$

which sends a string to its last vertex. This map is a weak categorical equivalence upon localizing $\mathcal{N}(\Delta/X)$ at the preimage of the identities.

For the operadic generalization, we pass to the category of *dendroidal sets*, which we denote by $\mathbf{dSets}$. These are presheaves on a category $\mathbf{\Omega}$ whose objects are trees and with fully faithful inclusions $\Delta \subset \mathbf{\Omega} \subset \mathsf{Op}$. Canonically, one may form the dendroidal nerve functor

$$\mathcal{N}_d\colon \mathsf{Op} \to \mathbf{dSets}$$

from the category of discrete operads to dendroidal sets. There is a model structure on dendroidal sets where fibrant objects are *quasioperads* ([CM11]), and it extends Joyal's model structure on simplicial sets for quasicategories.

In this paper, we construct a natural transformation $(\mathbf{\Omega}/-)^{\otimes}\colon \mathbf{dSets} \to \mathsf{Op}$ fitting into the commutative diagram

$$\begin{tikzcd}
\mathbf{dSets} \arrow[r, "(\mathbf{\Omega}/-)^{\otimes}"] & \mathsf{Op} \arrow[r, "\mathcal{N}_d"] & \mathbf{dSets} \\
\mathbf{sSets} \arrow[u, hook] \arrow[r, "\Delta/"] & \mathsf{Cat} \arrow[u, hook] \arrow[r, "\mathcal{N}"] & \mathbf{sSets}. \arrow[u, hook]
\end{tikzcd}$$

The operad $(\mathbf{\Omega}/X)^{\otimes}$ is defined in Theorem 3.6. An object of $(\mathbf{\Omega}/X)^{\otimes}$ is given by (T, α), with T a tree in $\mathbf{\Omega}$ and $\alpha\colon T \to X$ a map of dendroidal sets, that is, a labelling of T by objects (the edges) and operations (the vertices) of X, together with all the higher coherences. A k-ary operation $((T_1, \alpha_1), \dots (T_n, \alpha_n)) \longrightarrow (S, \beta)$ is a map of dendroidal sets $T_1 \sqcup \dots T_n \to S$ over X satisfying certain *wideness* and *independency* conditions.

The *root* of T is a canonical edge, and evaluating an object (T, α) at the root yields an assignemt $\mathsf{Ob}((\mathbf{\Omega}/X)^{\otimes}) \to X_0$. In some sense, $(\mathbf{\Omega}/X)^{\otimes}$ can be characterized as the maximal suboperad of the symmetric monoidal category $(\mathbf{\Omega}/X)^{\sqcup}$ for which the assignment $(*)$ extends to a functor of dendroidal set

$$\mathscr{r}_X : \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}) \longrightarrow X$$

cocontinuously in X as a natural transformation of functors. We call this map the *root functor* (Theorem 3.12). When X is a simplicial set, the operad $(\mathbf{\Omega}/X)^{\otimes}$ is just the category of elements of X, and the root functor is the last vertex map $\mathcal{N}(\Delta/X) \to X$. Our main result is the following.

Theorem (3.14)**.** *Let X be a normal*[1] *dendroidal set, and let $\mathcal{R}$ be the set of morphisms of $(\mathbf{\Omega}/X)^{\otimes}$ sent to identities by $\mathscr{r}_X$. The root functor $\mathscr{r}_X$ induces an operadic weak equivalence of dendroidal sets*

$$\overline{\mathscr{r}_X} : \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})[\mathcal{R}^{-1}] \xrightarrow{\sim} X$$

between the localization of $\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})$ at $\mathcal{R}$ and X.

Let us refer to the above as the *(operadic) delocalization theorem*.

When X is a simplicial set, the root functor coincides with the last vertex functor and one recovers Joyal delocalization theorem for ∞-categories.

From the delocalization theorem we deduce that in the stable model structure on dendroidal sets ([BN14]), any ∞-operad X is weakly equivalent to the infinite loop space of the discrete operad $(\mathbf{\Omega}/X)^{\otimes}$ (Corollary 3.15)[2]. When specialized to an ∞-category, it gives another proof of the well-known fact that every ∞- category has the weak homotopy type of the nerve of a category (used for instance in [Wal85]).

In the last Section 4, we study how operadic localization interacts with the ∞-categories involved in the un/straightening equivalence: that of algebras over an ∞-operad and that of operadic left fibrations ([Heu11], [BdBM20], [Ram22], [Ker23], [Pra25b]). We use the formalism of dendroidal sets and the notion of dendroidal left fibrations, establishing Quillen equivalences of (semi)model categories.

More precisely, we consider a localization of dendroidal sets $X \to X[S^{-1}]$ and study the model category structure on $\mathbf{dSets}/X$ obtained as a left Bousfield localization of the covariant model structure (Theorem 4.1). Its fibrant objects are those dendroidal left fibrations $E \to X$ for which a morphism $f\colon x \to y$ in S induces a weak homotopy equivalence of Kan complexes $f_!\colon E_x \to E_y$ between the fibres. We denote it by $\mathcal{L}_S(\mathbf{dSets}/X)$. The first result we prove is the following

Proposition (4.4)**.** *For every localization map of dendroidal sets $\lambda\colon X \to X[S^{-1}]$, the pullback along λ induces a Quillen equivalence of model categories*

$$\lambda_!\colon \mathcal{L}_S(\mathbf{dSets}/X) \rightleftarrows \mathbf{dSets}/X[S^{-1}] : \lambda^*.$$

By un/straightening, the following result follows.

Theorem (4.10)**.** *Let X be a normal dendroidal ∞-operad. Let $\mathsf{Alg}_X(\mathbf{sSets})$ be the category of algebras over X with the projective model structure.* [3] *There is a zig-zag of Quillen equivalences*

[1] It means cofibrant in the operadic model structure. It corresponds to an operad being Σ-free.

[2] The stable model structure is a left Bousfield localization of the operadic model structure. It is Quillen equivalent to that of group-like $\mathbb{E}_\infty$-spaces ([BdBM20]), or equivalently that of infinite loop spaces. Under the equivalence $\mathbf{dSets}/\eta \simeq \mathbf{sSets}$, one recovers Kan-Quillen model structure.

[3] The ∞-category of algebras over a dendroidal ∞-operad X is modeled by the projective model structure on $\mathsf{Alg}_{W_!(X)}(\mathbf{sSets})$, where $W_!(X)$ is the simplicial operad given by the Boardman-Vogt resolution of X ([CM13]).

of semi-model categories

$$\mathsf{Alg}_X(\mathbf{sSets}) \xleftarrow{\sim} \bullet \xrightarrow{\sim} \mathcal{L}_{\mathcal{R}_X}\left(\mathsf{Alg}_{(\mathbf{\Omega}/X)^{\otimes}}(\mathbf{sSets})\right),$$

where on the right the Bousfield localization has as fibrant objects $\mathcal{R}_X$-locally constant algebras.

Relation with other works. There is no canonical way to extend categorical results to operadic ones, and sometimes such extensions may not exist. The core of this result relies in the construction of the operad of elements $(\mathbf{\Omega}/X)^{\otimes}$. Once the construction is fixed, the formal steps of Theorem 3.14 follow Stevenson's approach in the proof of Joyal's delocalization theorem [Ste17]. A new contracting homotopy for trees ($\mathfrak{l}_T$) must be built, and care is needed when extending simplicial arguments to dendroidal sets, e.g., assuming X is normal.

It is natural to ask what happens when X is a symmetric monoidal ∞-category, also in light of [Ara26]. We address this in Remark 3.17.

Outline.

(1) In Section 1 we recall the necessary preliminary notions and set up notation.
(2) In Section 2, we set up the theory of dendroidal localization. Our definition extends Lurie's simplicial localization for quasicategories in [Lur17] to quasioperads.
(3) Section 3 is the heart of the article: here we define the operad of elements (Theorem 3.6), construct the root functor (Theorem 3.12) and prove the delocalization theorem (Theorem 3.14).
(4) In Section 4, we study interaction of dendroidal localization with operadic un/straightening equivalences, and then specify to the case of the operad of elements to deduce local un/straightening equivalences.

Acknowledgements. I thank Gijs Heuts for suggesting this problem to me and for many valuable discussions. I am also grateful to Miguel Barata,Victor Carmona, Denis-Charles Cisinski and Ieke Moerdijk for several fruitful exchanges. I would like to thank my advisor, Eric Hoffbeck, for his constant support and interest in my projects, and for carefully reading my drafts.

This project was realized during my PhD in Paris13 and a visiting in Utrecht. I hence thank the EOLE grant of French-Dutch Network (RFN) and the European Union's Horizon 2020 research and innovation programme, which funded my PhD under the Marie Skłodowska-Curie grant agreement No 945332.

1. Dendroidal recollections

For this section, complete proofs and definitions can be found in [HM22].

1.1. **Discrete operads.** From now on, we will call *operad* a discrete, strict operad P. We write Op for the category of operads and morphisms between them. A morphism of operads $f\colon P \to Q$ is the datum of a map of sets $\mathsf{Ob}(P) \to \mathsf{Ob}(Q)$ together with maps $P(c_1,\dots,c_n;d) \to Q(f(c_1),\dots,f(c_n);f(d))$ for any choice of objects $c_1,\dots,c_n,d$, compatible with the operadic composition and the symmetric group action.

Remark 1.1. The set $\mathsf{Ob}(P)$ has the structure of a poset, where $c \leqslant d$ if and only if P has an operation with target d and set of inputs containing c. Observe that a morphism of operads $f\colon P \to Q$ induces a morphism of posets $f\colon \mathsf{Ob}(P) \to \mathsf{Ob}(Q)$.

There is a fully faithful functor $j_!\colon \mathsf{Cat} \to \mathsf{Op}$, where $j_!$ acts by 'extension by zero', in the sense that

$$j_!C(c_1,\dots,c_n;d) = \begin{cases} \varnothing & \text{if } n \neq 1 \\ C(c_1;d) & \text{if } n = 1. \end{cases} \tag{1.1}$$

Its right adjoint $j^*\colon \mathsf{Op} \to \mathsf{Cat}$ is called the *underlying category* functor.

Definition 1.2. An operad P is Σ-*free* if, for any natural number n and any choice of objects $c_1, \dots, c_n, c$, the symmetric group on n-elements Σ_n acts freely on the set $\bigcup_{\sigma \in \Sigma_n} P(c_{\sigma(1)}, \dots, c_{\sigma(n)}; c)$.

Given any (small) symmetric monoidal category $(\mathbb{V}, \otimes)$, one may form the operad $\mathbb{V}^{\otimes}$, with objects the objects of $\mathbb{V}$ and operations $\mathbb{V}^{\otimes}(x_1, \dots, x_n; y) = \mathbb{V}(x_1 \otimes \cdots \otimes x_n; y)$, where composition is that of multivariable functions. A P-algebra in $\mathbb{V}$ is a morphism of operads $F\colon P \to \mathbb{V}^{\otimes}$. In this paper, we use $\mathbb{V} = \mathbf{sSets}$ and the subcategory $\mathbb{V} = \mathbf{Sets}$, symmetric monoidal with the cartesian product.

Denote by $\mathsf{Alg}_P(\mathbb{V})$ the category of P-algebras in $\mathbb{V}$ and morphisms between them. Observe that for a category C, we have a natural identification $\mathsf{Alg}_{j_!C}(\mathbb{V}) = \mathsf{Fun}(C, \mathbb{V})$.

1.2. **The dendroidal category.** Let Δ be the category of finite linear orders $[n] = \{0 < 1 < \cdots < n\}$, $n \geqslant 0$, and morphisms the maps of posets. Simplicial sets $\mathbf{sSets}$ are the category of presheaves over Δ. There is a fully faithful functor $\Delta \to \mathsf{Cat}$, which by restricted Yoneda embedding induces the nerve functor $\mathcal{N}\colon \mathsf{Cat} \to \mathbf{sSets}$.

The dendroidal category $\mathbf{\Omega}$ has objects non planar finite rooted trees.



FIGURE 2. Some typical trees in $\mathbf{\Omega}$.

Any such tree T yields an operad $\Omega(T)$[4], so morphisms are defined by declaring $\Omega(-)\colon \mathbf{\Omega} \to \mathsf{Op}$ a fully faithful embedding[5].

There is a fully faithful embedding $\Delta \to \mathbf{\Omega}$, which looks like this:



FIGURE 3. The ordinal $[2]$ is realized as the linear tree with 3 edges and 2 unary vertices.

It is customary, and we adopt this convention here as well, to denote by η the image of the ordinal $[0]$ under the above inclusion. It is the unique tree with no vertex.

Dendroidal sets $\mathbf{dSets}$ are the category of presheaves on $\mathbf{\Omega}$. The restricted Yoneda embedding yields the fully faithful functor $\mathcal{N}_d\colon \mathsf{Op} \to \mathbf{dSets}$, called the *dendroidal nerve functor*. Explicitly, we have $\mathcal{N}_d(P) := \mathsf{Op}(\Omega(-), P)$.

The diagram on the left commutes, making the one the right commute as well.

$$\begin{tikzcd} \mathsf{Cat} \arrow[d] & \Delta \arrow[l] \arrow[d] \\ \mathsf{Op} & \mathbf{\Omega} \arrow[l] \end{tikzcd} \qquad \begin{tikzcd} \mathsf{Cat} \arrow[r, "\mathcal{N}"] \arrow[d] & \mathbf{sSets} \arrow[d, "i_!"] \\ \mathsf{Op} \arrow[r, "\mathcal{N}_d"'] & \mathbf{dSets} \end{tikzcd},$$

[4]The set of objects of $\Omega(T)$ is the set of edges of T, and for any choice of edges $e_1, \dots, e_n, e$, one has $\Omega(T)(e_1, \dots, e_n; e) = \{*\}$ if and only if there exists a subtree of T with leaves $\{e_1, \dots, e_n\}$ and root e (necessarily unique), and $\Omega(T)(e_1, \dots, e_n; e) = \varnothing$ otherwise. The operadic composition corresponds to *grafting* of subtrees, which means successive identifications of the root of a subtree with a leaf of another.

[5]That is, $\mathbf{\Omega}(S, T) = \mathsf{Op}(\Omega(S), \Omega(T))$.

where $i_!$ is the left Kan extension of $i\colon \Delta \to \mathbf{\Omega}$ along the Yoneda embedding.

There is a natural isomorphism $\Delta \simeq \mathbf{\Omega}/\eta$, which yields the isomorphism $\mathbf{dSets}/\eta \simeq \mathbf{sSets}$, under which the inclusion $i_!$ is the forgetful functor $\mathbf{dSets}/\eta \to \mathbf{dSets}$. More generally, for any simplicial set M, one has $\mathbf{dSets}/i_!M \simeq \mathbf{sSets}/M$.

1.2.1. *Dendroidal notions.* In a tree T of $\mathbf{\Omega}$, every vertex has a single output edge and n input edges for some $n \geqslant 0$ (the *arity* of v). The root is the unique edge without an input vertex, and we denote it by r_T. The *leaves* of T are those edges which have no output vertex. A vertex is *external* if its input edges are contained in the leaves of T or if it is a *stump*, that is, a vertex with no output edge. We denote by C_n the the essentially unique tree with one vertex and n leaves, and call it the *n-corolla*. Let η be the unique tree with no vertex. Denote by $E(T)$ the set of edges of T.

We rephrase the poset structure on $E(T)$ (see Remark 1.1) as follows: for two edges e, f, one has $e \leqslant f$ if and only if the (unique) path from e to the root of T contains f. In particular, the root r_T is the unique maximal element, while the minimal elements are the leaves of T and the input edges of stumps.

We say that S is a subtree of T if it can be obtained from T by successively pruning away external vertices and the outer edges attached to them from T.

Given two trees R, S and a leaf ℓ of S, the *grafting* of R onto S along ℓ is the tree $S \cup_\ell R$ given by the pushout

$$\begin{tikzcd} \eta \arrow[r, "\ell"] \arrow[d, "r_R"'] & S \arrow[d] \\ R \arrow[r] & S \cup_\ell R \end{tikzcd}$$

where both arrows $R \to S \cup_\ell R \leftarrow S$ are inclusions of subtrees.

Any tree T is ismorphic to a grafting of subtrees, the indecomposable elements being the n-corollas and the trivial tree η.

For the scope of this section, we denote by $\Omega[T]$ the representable dendroidal set associated to a tree T. Observe that there is a natural isomorphism of dendroidal sets

$$\Omega[T] \simeq \mathcal{N}_d(\Omega(T)).$$

1.3. **Tensor product of dendroidal sets.** An homotopy equivalence of maps of dendroidal sets $f, g\colon X \to Y$ is modeled by considering the dendroidal tensor product with the interval (i.e. the 1 corolla $C_1 \simeq [1]$) $h\colon X \otimes C_1 \to Y$ with the usual restrictions $h_0 = f$, $h_1 = g$.

Tensor product of dendroidal sets is the bifunctor

$$\otimes\colon \mathbf{dSets} \times \mathbf{dSets} \longrightarrow \mathbf{dSets}$$

defined as the two variable left Kan extension along the Yoneda embeddings under the bifunctor $\mathbf{\Omega} \times \mathbf{\Omega} \longrightarrow \mathbf{dSets}$, $\quad (T, S) \mapsto \mathcal{N}_d(\Omega(T) \otimes_{BV} \Omega(S))$, where $\otimes_{BV}$ denotes the Boardman-Vogt tensor product of discrete operads ([BV73]). Restricted to simplicial sets, the tensor product is just the cartesian product.

The BV tensor product of discrete operads is characterized by the universal property that are equivalences of categories

$$\mathsf{Alg}_Q(\mathsf{Alg}_P(\mathbf{sSets})) \simeq \mathsf{Alg}_{P \otimes_{BV} Q}(\mathbf{sSets}) \simeq \mathsf{Alg}_P(\mathsf{Alg}_Q(\mathbf{sSets})),$$

for any two operads P, Q, and the product for algebras is the objectwise cartesian product. The operad $P \otimes Q$ is a quotient of $P \times Q$ under some relations, notably the *interchange relation*:

$$(p \otimes y) \circ (c_1 \otimes q, \ldots, c_n \otimes q) = \sigma^*_{n,m}(d \otimes q) \circ (p \otimes d_1, \ldots, p \otimes d_m),$$

where $\circ$ denotes the total operadic composition and the permutation $\sigma_{n,m}$ is the unique element of Σ_{nm} making sense of the above formula.

Let us now illustrate with an example our case of interest, the tensoring with the 1-corolla.

Example 1.3. *Consider the operads*



The Boardman-Vogt interchange relation for $\Omega(C_3) \otimes \Omega(C_1)$ *yields the identification:*



FIGURE 4. The Boardman-Vogt interchange relation for $\Omega(C_3) \otimes \Omega(C_1)$.

On dendroidal sets, the tensor product is commutative, but it is not, in fact, associative up to coherent isomorphisms, hence it does not make **dSets** symmetric monoidal. It is however associative up to coherent homotopies [HM22, §4.4]. However, it is so up to homotopy (see Section 1.4) [6], and is the good homotopical notion of product to consider on dendroidal sets, thanks also to the behaviour of its internal hom object bifunctor. We denote by $\hom(-,-)$ the underlying simplicial set, that is

$$\hom\colon \mathbf{dSets}^{\mathrm{op}} \times \mathbf{dSets} \longrightarrow \mathbf{sSets} \quad (X, Y) \mapsto \hom(X, Y)_\bullet \simeq \mathbf{dSets}(i_!(\Delta^\bullet) \otimes X, Y).$$

1.4. **Dendroidal homotopy theory.** The homotopy theory of dendroidal ∞-operads was defined in [MW07] and [CM11] by extending Joyal's theory of ∞-categories as quasicategories. We will not need explicit definition of the following fundamental notions, which we will treat syntetically:

- Analogously to the description of the morphisms in Δ as generated by faces and degeneracies under the simplicial identities, morphisms in $\mathbf{\Omega}$ are also generated ([HM22, §3.3.4]) by dendroidal faces and degeneracies, together with isomorphisms of trees, which satisfy the *dendroidal identities*.
- Some of the notions of the homotopy theory of simplicial sets can be formulated in the context of dendroidal sets. In particular, one can talk about the *boundary* of a tree and inner and external horns of a tree. Inner horns are relative to inner edges, while external horns divide into to classes, relative respectively to leaf vertices or root vertices [7].

Definition 1.4. A dendroidal set X is a *dendroidal ∞-operad* if it has the right lifting property against inner horn inclusions of trees.

A dendroidal set X is *normal* if, for any tree T, the action of $\mathrm{Aut}(T)$ on the set X_T is free.

Observe that, if $X \simeq \mathcal{N}_d P$, then X is normal if and only if P is Σ-free.

Theorem 1.5 ([CM11])**.** *There exists a model structure on the category* **dSets** *of dendroidal sets, called the* operadic model structure*, with the following properties:*

[6] This tensor product endows the ∞-category of dendroidal ∞-operads with a symmetric monoidal structure, see [HM24, §5].

[7] They extend, respectively, left and right fibrations of simplicial sets. Contrarily to these latter, they are not dual to each other as $\mathbf{\Omega}$ does not admit a self duality.

(1) The cofibrations are the normal monomorphisms.
(2) The fibrant objects are the dendroidal ∞-operads.
(3) A map between normal dendroidal sets is a weak equivalence if and only if for every dendroidal ∞-operad X, the map

$$\hom(B, X) \longrightarrow \hom(A, X)$$

is a categorical equivalence of ∞-categories.

Moreover, this model structure is left proper and cofibrantly generated.

The homotopy theory of dendroidal left fibrations, which will appear in Section 4, was first defined in [Heu11] and generalizes that for left fibration of simplicial sets. It models ∞-operads cofibred in groupoids.

Definition 1.6. A morphism of dendroidal sets $p\colon Y \to X$ is a *dendroidal left fibration* if it has the right lifting properties against inner and leaf horn inclusions of trees.

Notation 1.7. For a morphism of dendroidal sets $p\colon Y \to X$ and an object x of X, we denote by Y_x the fibre over x, that is,

$$Y_x \simeq p^{-1}(x).$$

If p is a dendroidal left fibration, Y_x is a Kan complex ([HM22, Remark 9.60]).

Theorem 1.8 ([Heu11])**.** *Let X be a dendroidal set. The category* $\mathbf{dSets}/X$ *carries a left proper cofibrantly generated model structure, called the* covariant model structure*, with the following properties:*

(1) The cofibrations are the normal monomorphisms over X.
(2) The fibrant objects are the dendroidal left fibrations over X.
(3) A map $(A, u) \to (B, v)$ between normal objects over X is a weak equivalence if and only if for any dendroidal left fibration (Y, p), the map

$$\hom_X(B, Y) \longrightarrow \hom_X(A, Y)$$

is a weak homotopy equivalence of Kan complexes, where

$$\hom_X(A, Y) = \hom(A, Y) \times_{\hom(A,X)} \{u\}.$$

(4) A map $(Y, p) \to (Z, q)$ between dendroidal left fibrations is a weak equivalence if and only if, for any object x of X, the induced map $Y_x \to Z_x$ between the fibres over x is a weak homotopy equivalence of Kan complexes.

Moreover, if $f\colon X \to Y$ is a map of dendroidal sets, the adjunction

$$f_!\colon \mathbf{dSets}/X \rightleftarrows \mathbf{dSets}/Y : f^*$$

is a Quillen adjunction with respect to the covariant model structure, which is a Quillen equivalence when f is an operadic weak equivalence.

Slicing over the point, that is under the equivalence $\mathbf{sSets} \simeq \mathbf{dSets}/\eta$, the operadic model structure yields the Joyal model structure, and the covariant model structure yields the homonimous one for left fibrations over a simplicial set.

Coming back to the discussion after Example 1.3, let us observe that $\hom(-,-)$ homotopy-enriches $\mathbf{dSets}$ with the operadic model structure in simplicial sets with the Joyal model structure ([HM22, Proposition 9.28]). When localizing for the covariant model structure on $\mathbf{dSets}/X$, we find a simplicial enrichment with respect to the Kan-Quillen model structure ([HM22, Proposition 9.66]). In particular, $\hom(-,-)$, resp. $\hom_X(-,-)$, gives a model for the mapping spaces in the operadic, resp. covariant, model structure when computed on bifibrant objects.

1.5. **Last conventions.** When in presence of the canonical fully faithful functors $i\colon \Delta \to \mathbf{\Omega}$, $i_!\colon \mathbf{sSets} \to \mathbf{dSets}$, $\mathbf{\Omega} \to \mathsf{Op} \to \mathbf{dSets}$. In particular, given a tree T, we still write T for the operad $\Omega(T)$ and for the dendroidal set $\Omega[T]$. Observe that there are isomorphisms of dendroidal sets $\eta \simeq \Delta^0$ and $C_1 \simeq \Delta^1$, but that the analogy stops here, as for $n \neq 1$, there is not even a map $C_n \to \Delta^n$. We will freely use any of these functors explicitly whenever it is convenient or helps to emphasize a point.

2. Localization of dendroidal ∞-operads

We now define derived localization of dendroidal sets, in a way that extends that of quasi-categories in the sense of Joyal and Lurie ([Lur17]). We then construct a model for normal dendroidal sets in Theorem 2.4. For compatibility of localization of dendroidal sets with the covariant model structure we direct the reader to Section 4.1.

2.1. **The definition of localization.** Let us start by some preliminary

Definition 2.1. A *normalization* of a dendroidal set X is a trivial fibration $X' \xrightarrow{\sim} X$ in the operadic model structure, with X' normal.

Given a morphism of dendroidal sets $f\colon X \to Y$, a *normalization* of f is a commutative diagram

$$\begin{tikzcd} X' \arrow[r, "f'"] \arrow[d, "\wr"'] & Y' \arrow[d, "\wr"] \\ X \arrow[r, "f"] & Y \end{tikzcd}$$

where both vertical arrows are normalizations.

Normalizations exist and are unique up to operadic weak equivalence. As explained in [HM22, Remark 9.21], an explicit construction of a normalization with contractible fibres is given by the projection $X \times W^*\mathcal{E} \to X$, where $\mathcal{E}$ is the simplicial Barratt-Eccles operad and W^* is the operadic homotopy coherent nerve functor, right adjoint to $W_!$.

Definition 2.2. Let $\lambda\colon X \to Y$ be a morphism between normal dendroidal sets and let $S \subseteq X_{C_1}$ be a subset of 1-morphisms. The map λ is a *localization of X at the set of morphisms S* if, for any dendroidal ∞-operad Z, the morphism between the simplicial hom objects

$$\hom(Y, Z) \longrightarrow \hom(X, Z)$$

is fully faithful, with essential image given by those maps $X \to Z$ sending S to equivalences.

If $\lambda\colon X \to Y$ is a morphism between non-necessarily normal dendroidal sets, we say that λ is a localization if any, or equivalently one, normalization λ' of λ is.

Localization is unique up to operadic weak equivalence, and we denote by $X[S^{-1}]$ 'the' localization.

Observe that, if we denote by $\hom_S(X, Z)$ the full sub simplicial set of $\hom(X, Z)$ spanned by those maps $X \to Z$ sending S to equivalences in Z, the universal property of the localization allows to identify $\hom(X[S^{-1}], Z)$ with $\hom_S(X, Z)$.

Remark 2.3. It is immediate to see that the definition applied to a morphism of dendroidal sets recovers the localization of quasi-categories in the sense of [Lur17, Definition 1.3.4.1].

We can construct an explicit model for the localization of a normal dendroidal set.

Proposition 2.4. *Denote by J be the nerve of the connected groupoid on two objects. Given a normal dendroidal set X and a subset $S \subseteq X_{C_1}$ of 1-morphisms in X, the localization of X at S is realized by the map $\lambda\colon X \to \mathcal{L}(X, S)$ defined by the pushout diagram*

$$\begin{tikzcd} \coprod_{s\in S} C_1 \arrow[r] \arrow[d] & X \arrow[d, "\lambda"] \\ \coprod_{s\in S} J \arrow[r] & \mathcal{L}(X,S)\,. \end{tikzcd} \tag{2.1}$$

Proof. Observe that, by left properness of the operadic model structure, the pushout in Equation (2.1) is a homotopy pushout. Let Z be a dendroidal ∞-operad and consider the map

$$\lambda^*\colon\ \hom(\mathcal{L}(X,S),Z) \longrightarrow \hom(X,Z).$$

The essential image of λ^* consists of all functors sending S to equivalences, so we just need to show λ is also fully faithful. To this end, it is sufficient to show that the diagram

$$\begin{tikzcd} \hom(\mathcal{L}(X,S)\otimes C_1, Z) \arrow[r] \arrow[d] & \hom(X\otimes C_1, Z) \arrow[d] \\ \hom(\mathcal{L}(X,S)\otimes \partial C_1, Z) \arrow[r] & \hom(X\otimes \partial C_1, Z) \end{tikzcd}$$

is a homotopy pullback for the Joyal model structure, which happens if the diagram

$$\begin{tikzcd} X\otimes \partial C_1 \arrow[r] \arrow[d] & X\otimes C_1 \arrow[d] \\ \mathcal{L}(X,S)\otimes \partial C_1 \arrow[r] & \mathcal{L}(X,S)\otimes C_1 \end{tikzcd} \tag{2.2}$$

is a homotopy pushout for the operadic model structure. Since the diagram in (2.1) is a transfinite composition of homotopy pushouts along the morphisms $\{C_1 \xrightarrow{s} J\}_{s\in S}$, it is sufficient to show that it is an homotopy pushout just in the case when $S=\{s\}$, where it appears as the front face of the commutative cube

$$\begin{tikzcd} C_1\otimes\partial C_1 \arrow[rr] \arrow[dr, "s\otimes\mathrm{id}"] \arrow[dd] & & C_1\otimes C_1 \arrow[dr, "s\otimes\mathrm{id}"] \arrow[dd] & \\ & X\otimes\partial C_1 \arrow[rr] \arrow[dd] & & X\otimes C_1 \arrow[dd] \\ J\otimes\partial C_1 \arrow[rr] \arrow[dr] & & J\otimes C_1 \arrow[dr] & \\ & \mathcal{L}(X,S)\otimes\partial C_1 \arrow[rr] & & \mathcal{L}(X,S)\otimes C_1 \end{tikzcd}$$

The front face of the cube is a homotopy pushout because all the others are, and this concludes the proof. □

The above construction essentially means that to localize a dendroidal set at a set of 1-morphisms, we can first localize its underlying simplicial set and then glue it to the original dendroidal set X, as we explain in the next

Remark 2.5. Let X be a normal dendroidal set, and write $M := i^*X$ for its underlying simplicial set. Of course, we have $S \subseteq X_{C_1} = M_1$, so we can localize M at S. Let $\overline{\lambda}\colon M \to \mathcal{L}(M,S)$ be the localization map. The localization of X at S can be realized as the following homotopy pushout

$$\begin{tikzcd} M = i^*X \arrow[r, "\epsilon_X"] \arrow[d, "\bar{\lambda}"] & X \arrow[d, "\lambda"] \\ \mathcal{L}(M,S) \arrow[r] & \mathcal{L}(X,S) \end{tikzcd}$$

where the top horizontal map is the counit of the adjunction $(i_!, i^*)$.

Given two endofunctors of X which preserve S, there is an easy way to check when they are homotopy equivalent as endofunctors of the localization $X[S^{-1}]$. Recall the following

Definition 2.6. Let $f, g\colon X \to Y$ be two maps of dendroidal sets. An *homotopy* between f and g is a morphism $h\colon X \otimes C_1 \to Y$ in **dSets** with the property that $h_0 = f$ and $h_1 = g$, where h_i is the restriction of h along the leaf, resp. root, inclusions $\eta \xrightarrow{\{i\}} C_1$, $i = 0$, resp. $i = 1$.

For $x \in X_\eta$, the arrow $h_x\colon f(x) \to g(x)$ in Y_{C_1} is called a *component* of h.

Lemma 2.7. *Let X be normal, $S \subseteq X_{C_1}$ a set of 1-morphisms. Let $f, g\colon X \to X$ be two maps sending S to itself. If f and g are homotopic via a homotopy whose components are in S, then f and g are homotopy equivalent as maps $f, g\colon X[S^{-1}] \to X[S^{-1}]$.*

Proof. Let $h\colon X \otimes C_1 \to X$ be the homotopy between f and g whose components lie in S. The transpose of $h^*\colon \hom(X,Z) \to \hom(X \otimes C_1, Z) \simeq \hom(C_1, \hom(X,Z))$ induces a morphism of simplicial sets $\hom_S(X,Z) \times \Delta^1 \to \hom(X,Z)$. By the hypotheses on f and g and fullness of $\hom_S(X,Z)$ in $\hom(X,Z)$, there is an induced homotopy $\mathfrak{h}\colon \hom_S(X,Z) \times \Delta^1 \to \hom_S(X,Z)$. To prove that $\mathfrak{h}$ is a natural equivalence, it suffices to see that, for any object ϕ of $\hom_S(X,Z)$, the arrow $\mathfrak{h}_\phi\colon f^*(\phi) \to g^*(\phi)$ is an equivalence in $\hom_S(X,Z)$, that is in $\hom(X,Z)$. So we only need to check that for any $x \in X_\eta$, the 1-morphism $(\mathfrak{h}_\phi)_x\colon \phi(f(x)) \to \phi(g(x))$ is an equivalence in Z. As $(\mathfrak{h}_\phi)_x = \phi(h_x)$, and since by hypothesis $\phi(h_x)$ is a weak equivalence as h_x is an arrow in S, we have that $\mathfrak{h}$ establishes an equivalence between f^* and g^*. This concludes the proof. □

The homotopy theory of algebras over a dendroidal ∞-operad X is governed by the covariant model structure on the over-category $\mathbf{dSets}/X$, so it is a natural question to investigate compatibility of the localization of dendroidal ∞-operads with the covariant model structure for dendroidal left fibrations. We study this in Section 4.1 (Theorem 4.4).

3. The delocalization theorem

We start Section 3.1 by introducing the formalism of forests and wide independent maps between them. We define the operad of elements and the root functor of a dendroidal set in Section 3.3, and prove the localization result, namely Theorem 3.14, in Section 3.4.

3.1. The category of forests.

A *forest* is a finite disjoint union of trees. We denote by Φ be the *category of forests* obtained from $\mathbf{\Omega}$ by formally adjoining finite coproducts. By extending the inclusion $\mathbf{\Omega} \hookrightarrow \mathsf{Op}$ to forests under finite-coproduct, one has $\Omega\colon \Phi \hookrightarrow \mathsf{Op}$, where, given a forest $F = \bigsqcup_{i=1}^n T_i$, the operad $\Omega(F)$ is defined as $\Omega(F) := \bigsqcup_{i=1}^n \Omega(T_i)$. For any such F, every tree T_i is called a *constituent* of F, and $\mathbf{\Omega}$ embeds into Φ by seeing a tree to a forest with one constituent. Let ϕ be the subcategory of Φ whose objects are non-empty forests and where morphisms are the *independent* maps of forests, where

Definition 3.1. A morphism of forests $f\colon F \longrightarrow G$ is called *independent* if F and G are non-empty and, writing $F = \bigsqcup_{i=1}^n T_i$ and $G = \bigsqcup_{j=1}^m S_j$, f is specified by a map of sets $\alpha\colon \{1,\dots,n\} \to \{1,\dots,m\}$ and morphisms of trees $f_i\colon T_i \to S_{\alpha(i)}$, for $i = 1,\dots,n$, such that, whenever $i \neq i'$ and $\alpha(i) = \alpha(i') = j$, for all edges $e \in E(T_i)$, $e' \in E(T_{i'})$ the edges $f_i(e)$ and $f_{i'}(e')$ are incomparable in the poset $E(S_j)$.

By the independency condition, it suffices to check that the edges $f(r_{T_i})$ and $f(r_{T_{i'}})$ are independent.

The restriction of the disjoint union to ϕ endows it with a non-unital symmetric monoidal structure; we write $\oplus := \bigsqcup_{|\phi}$ and call it direct sum. In the category ϕ, every morphism can be written as a direct sum of maps whose target has only one constituent. Observe also that ϕ is no longer the coproduct in ϕ, as a candidate for the codiagonal $F \oplus F \to F$ would not satisfy the independence property.

Remark 3.2. The two categories of forests presented here are the ones appearing in various comparisons of the dendroidal and Lurie's model for ∞-operads: the category Φ is the one considered in [HM24], while the category ϕ is the one appearing in [HHM16].

3.2. **Wide maps of forests.** These were first introduced in [HM22].

Definition 3.3. A map of forests $f\colon F \to G$ is called *wide* if, for any constituent tree S of the forest G, any maximal monotonic path in the poset $E(S)$, that is from a minimal element to the root r_S, contains one element of the form $f(r_T)$ for some T constituent tree of F.

Observe that if f is also independent, if there exists such a constituent T, then it is unique.

When the target of a map of forests has only one constituent, we can reformulate the wideness condition in the following useful way.

Lemma 3.4. *Let $T_1, \dots, T_n, S$ be trees and $f\colon T_1 \oplus \cdots \oplus T_n \to S$ a map of forests. If f is independent, then f is wide if and only if $S(f(r_{T_1}), \dots, f(r_{T_n}); r_S) \neq \varnothing$.*

Proof. It suffices to observe that, given edges $e_1, \dots, e_n$ of S, the only obstruction to the existence of a subtree with leaves $e_1, \dots, e_n$ and root r_S is the existence of a maximal monotonic path $\mathfrak{p} = \{l < a_1 < \cdots < a_m < r_S\}$ in the poset of edges of S such that $\{e_1, \dots, e_n\} \cap \mathfrak{p} = \varnothing$. □

Every wide morphism is the direct sum of morphisms of the above type, and with Theorem 3.4 it is immediate to see that wide maps are closed under composition. We introduce some classes of wide maps in the following

Example 3.5.

(1) All maps of linear trees $[n] \to [m]$ are wide and independent.
(2) All maps of trees $T \to S$ are independent maps of forests.
(3) A morphism of trees $f\colon T \to S$ is root preserving *if $f(r_T) = r_S$. Any root preserving map is wide, and the converse is true whenever the root vertex of S is not unary.*
(4) Given a forest F, one can construct the tree $\overline{F}$ as follows. Write $F = \oplus_{i=1}^n T_i$ and let $\overline{F}$ be the tree obtained as the grafting of the trees $(T_1, \dots, T_n)$ on the leaves $\{l_1, \dots, l_n\}$ of an n-corolla C_n,

$$\overline{F} = C_n \circ (T_1, \dots, T_n).$$

The forest root face *of $\overline{F}$ is the natural inclusion*

$$F = T_1 \oplus \cdots \oplus T_n \longrightarrow C_n \circ (T_1, \dots, T_n) = \overline{F}.$$



FIGURE 5. A forest root face.

In fact, it is easy to see that wide independent maps of forests are generated, under composition and direct sum, by forest root face inclusions and root preserving morphisms.

3.3. **The operad of elements and the root functor.** We want to functorially associate to every dendroidal set X a discrete operad $(\mathbf{\Omega}/X)^{\otimes}$, which we call the *operad of elements* of X, with the requirement that, when X is a simplicial set, $(\mathbf{\Omega}/X)^{\otimes}$ is a category and coincides with the category of elements of X. Also, the nerve of the constructed operad will need to map back to X.

Recall that, for every dendroidal set X, the overcategory $\mathbf{dSets}_{/X}$ is a symmetric monoidal category with the disjoint union $\sqcup$. Let $(\mathbf{dSets}_{/X})^{\sqcup}$ be the operad associated to it. A first, naive way would be to define the operad of elements of X as the suboperad $(\mathbf{\Omega}/X)^{\sqcup}$ of $(\mathbf{dSets}_{/X})^{\sqcup}$ spanned by the representable objects. However, it turns out this is not the good definition (see later Remark 3.11), so we need to give a more elaborate one.

Definition 3.6. Let X be a dendroidal set. Its *operad of elements* $(\mathbf{\Omega}/X)^{\otimes}$ is the sub operad of $(\mathbf{dSets}_{/X})^{\sqcup}$ specified by the following data:

- the set of objects of $(\mathbf{\Omega}/X)^{\otimes}$ is the set of elements of X as a presheaf, that is,
$$\mathsf{Ob}((\mathbf{\Omega}/X)^{\otimes}) = \{(T,\alpha) \mid T \in \mathbf{\Omega} \text{ and } \alpha\colon T \to X\};$$
- for objects $(S_1,\alpha_1),\dots,(S_n,\alpha_n)$, (R,β), the set of operations
$$(\mathbf{\Omega}/X)^{\otimes}((S_1,\alpha_1),\dots,(S_n,\alpha_n);(R,\beta))$$
is given by the wide and independent maps of forests $f\colon S_1 \oplus \cdots \oplus S_n \to R$ such that
$$\beta \circ f = (\alpha_1,\dots,\alpha_n)\colon S_1 \bigsqcup \dots \bigsqcup S_n \to X$$
as maps of dendroidal sets.

To be explicit, the partial composition of f as above with $g \in (\mathbf{\Omega}/X)^{\otimes}((Q_1,\beta_1),\dots,(Q_m,\beta_m);(S_i,\alpha_i))$ along (S_i,α_i) is given by the wide map of forests
$$(S_1 \oplus \cdots \oplus S_{i-1}) \oplus (\oplus_{j=1}^m Q_j) \oplus S_{i+1} \cdots \oplus S_n \xrightarrow{(\mathrm{id}^{\oplus i-1},g,\mathrm{id}^{\oplus n-i})} \oplus_{h=1}^n S_h \xrightarrow{f} R.$$

Remark 3.7. By the independency condition on forest morphisms, the operad of elements $(\mathbf{\Omega}/X)^{\otimes}$ is Σ-free.

Example 3.8.

- *For a simplicial set M, the operad of elements $(\mathbf{\Omega}/M)^{\otimes}$ coincides with the category of elements of M $(\mathbf{\Omega}/M)^{\otimes} = \Delta/M$. Its objects are the pairs $([n],f)$, with $n \geqslant 0$ and $f\colon \Delta^n \to M$; a morphism $F\colon ([n],f) \to ([m],g)$ is just a map of representable simplicial sets $F\colon \Delta^n \to \Delta^m$ over M.*
- *If $X \simeq \mathcal{N}_d P$ for an operad P, an object of $\mathbf{\Omega}/X$ can be written as a pair (T,α), with α in $\mathrm{Hom}_{\mathsf{Op}}(T,P)$. Similarly, if $X \simeq W^*\mathbb{P}$, where W^* is the homotopy coherent nerve of a simplicial operad $\mathbb{P}$, then for an object (S,β) of $(\mathbf{\Omega}/X)^{\otimes}$ β is an element in $\mathrm{Hom}_{\mathsf{sOp}}(W(S),\mathbb{P})$, where $W(S)$ is the simplicial operad given by the Boardman-Vogt resolution of $\Omega(S)$.*

The construction $X \mapsto (\mathbf{\Omega}/X^{\otimes}$ is functorial in X via postcomposition, and we get an endofunctor
$$\mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})\colon \mathbf{dSets} \to \mathbf{dSets}.$$
Let us postpone to later the proof of the next central

Proposition 3.9. *The functor $\mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})$ is cocontinuous. Moreover, it preserves normal monomorphisms of dendroidal sets.*

In other words, it means that $\mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})$ is equivalent to the left Kan extension of its restriction to representables, and moreover in an homotopy-coherent way

Now, let T be a tree and (S,α) an object of $(\mathbf{\Omega}/T)^{\otimes}$. Evaluation of α at the root of S yields an assignment
$$\mathsf{Ob}((\mathbf{\Omega}/T)^{\otimes}) \ni (S,\alpha) \mapsto \alpha(r_S) \in \mathsf{Ob}(T) = E(T). \tag{3.1}$$

Proposition 3.10. *The assignment in Equation* (3.1) *extends to a map of operads*

$$\mathfrak{r}_T\colon (\mathbf{\Omega}/T)^{\otimes} \longrightarrow T$$

that we call the root functor *for* T.

Proof. As the set of operations of T is either empty or a singleton, we just need to check that, if there exists a wide independent map of forests $f\colon (S_1,\alpha_1),\dots,(S_n,\alpha_n) \to (R,\beta)$, then $T(\alpha_1(r_{S_1}),\dots,\alpha_n(r_{S_n});\beta(r_R)) \neq \varnothing$. By Lemma 3.4, $R(f(r_{S_1}),\dots,f(r_{S_n});r_R) \neq \varnothing$, and since β is a map of operads and $\beta\circ f = (\alpha_1,\dots,\alpha_n)$ we have the thesis, as wanted. □

Remark 3.11. The assignment in Equation (3.1) makes sense even for the *full* suboperad $(\mathbf{\Omega}/X)^{\sqcup} \subseteq (\mathbf{dSets}/X)^{\sqcup}$, where one considers all maps of forests $T_1\oplus\cdots\oplus T_n \to S$. However, Proposition 3.10 does *not* hold for $(\mathbf{\Omega}/X)^{\sqcup}$. An easy way to see this is taking $X = C_n$ any n-corolla, and the binary operation $(\mathrm{id},\mathrm{id})\colon (C_n,\mathrm{id}),(C_n,\mathrm{id}) \to (C_n,\mathrm{id})$. A candidate image for this latter has to be a subtree of C_n with leaves (r_{C_n},r_{C_n}) and root r_{C_n}, but such subtree does not exist. In this sense, Proposition 3.10 characterizes $\mathbf{\Omega}/X$ as the maximal sub operad of $(\mathbf{\Omega}/X)^{\sqcup}$ such that Proposition 3.10 holds.

Because of cocontinuity of $\mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})$ (Theorem 3.9), we can extend the root functor to every dendroidal set.

Definition 3.12. Let X be a dendroidal set. The *root functor* of X is the morphism of dendroidal sets

$$\mathfrak{r}_X\colon \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}) \longrightarrow X$$

defined as the colimit

$$\mathfrak{r}_X := \operatorname*{colim}_{T\to X} \mathfrak{r}_T\colon \mathcal{N}_d((\mathbf{\Omega}/T)^{\otimes}) \longrightarrow T.$$

We can interpret an object (S,α) in $\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})$ as a decoration of the tree S with objects and operations in X, together with all higher coherences governing operadic composition (which are not visible in the picture). In particular, by contracting all inner edges of S one obtains an operation of X. The root functor then sends (S,α) to the target of this operation. Visually



FIGURE 6. The root functor for $S \simeq C_3$ and a map $\alpha\colon C_3 \to X$.

Observe that, for any tree T and morphism $\alpha\colon T \to X$, the following diagram commutes:

$$\begin{tikzcd}
(\mathbf{\Omega}/T)^{\otimes} \arrow[r, "(\mathbf{\Omega}/\alpha)^{\otimes}"] \arrow[d, "\mathfrak{r}_T"'] & (\mathbf{\Omega}/X)^{\otimes} \arrow[d, "\mathfrak{r}_X"] \\
T \arrow[r, "\alpha"] & X
\end{tikzcd}$$

Remark 3.13. If M is a simplicial set, the root functor coincides with the *last vertex functor*

$$\mathfrak{r}_M\colon \Delta/M \to M, \quad \mathfrak{r}_M([n],f) = f(n).$$

We conclude this section with the promised proof of cocontinuity of the nerve operad of elements construction.

Proof (of Proposition 3.9). We denote by θ the restriction of $\mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})$ to the representables, that is $\theta = \mathcal{N}_d((\mathbf{\Omega}/-)^{\otimes})_{|\mathbf{\Omega}}$, and we write $\widehat{\theta}$ for its left Kan extension along the Yoneda embedding.

Given a dendroidal set X and a tree T, the set $\widehat{\theta}(X)_T$ may be described as

$$\widehat{\theta}(X)_T = \{((Q,x),(T,u)) \mid Q \in \mathbf{\Omega}, x\colon Q \to X, u\colon T \to \theta(Q)\}/\sim,$$

where, for any $\alpha\colon S \to Q$, one identifies

$$((Q,x),(R,\alpha_* u)) \sim ((S,x\alpha),(R,u)),$$

where $\alpha_* = \mathcal{N}_d((\mathbf{\Omega}/\alpha)^{\otimes})$. Functoriality in T is given by letting faces and degeneracies act on the second component, while functoriality in X is obtained via the first component.

We construct a natural equivalence

$$\psi_X\colon \widehat{\theta}(X) \to \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})$$

by defining its components $(\psi_X)_T$ by induction on the number of vertices of the tree T. When $T \simeq \eta$, we set

$$\psi_X((Q,x),(\eta,u)) \coloneqq x_* \circ u\colon \eta \to \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}).$$

The assignment respects the equivalence relation, as (with the same notations as above) one has

$$\psi_X((Q,x),(\eta,\alpha_* u)) = x_*(\alpha_* u) = (x\alpha)_* u = \psi_X((S,x\alpha),(\eta,u)),$$

and it is straightforward to see that it induces a bijection.

Similarly, given a n-corolla C_n, $n \geqslant 0$, and an element $((S,x),(C_n,u))$, we set

$$\psi_X((S,x),(C_n,u)) \coloneqq x_* u\colon C_n \to \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}),$$

and it is the same calculation which shows that it descends to the quotient and is a bijection.

For the inductive step, let us consider a tree T with at least two vertices and decompose it as the grafting $T = R \cup_a S$, with $R, S \neq \eta$. For any dendroidal set Y, there is a natural isomorphism

$$\mathcal{N}_d((\mathbf{\Omega}/Y)^{\otimes})_T \simeq \mathcal{N}_d((\mathbf{\Omega}/Y)^{\otimes})_R \underset{\mathcal{N}_d((\mathbf{\Omega}/Y)^{\otimes})_\eta}{\times} \mathcal{N}_d((\mathbf{\Omega}/Y)^{\otimes})_S.$$

The isomorphism is compatible with the equivalence relation defining $\widehat{\theta}(X)$, which means that $\widehat{\theta}(X)$ satisfies the same strict Segal condition, that is, there is a natural isomorphism

$$\widehat{\theta}(X)_T \simeq \widehat{\theta}(X)_R \underset{\widehat{\theta}(X)_\eta}{\times} \widehat{\theta}(X)_S.$$

One defines the map $(\psi_X)_T$ as

$$(\psi_X)_T \coloneqq (\psi_X)_R \underset{(\psi_X)_\eta}{\times} (\psi_X)_S.$$

It respects the equivalence relation and is a bijection, so we only need to check that ψ_X is well defined. This follows from the fact that any tree T decomposes as the grafting of corollas, and the decomposition is unique up to isomorphism and operadic associativity relations, and the Segal isomorphism is compatible with these latter, which shows that the definition of $(\psi_X)_T$ does not depend on the decomposition of T, as wanted. □

3.4. **The delocalization result.** In Theorem 3.5 we introduced root preserving morphisms of trees, which are in particular wide independent maps of forests. Given a dendroidal set X, we say that a morphism $f\colon (S,\alpha) \to (R,\beta)$ is *root preserving* if $f\colon S \to R$ is a root preserving map of trees. Denote by $\mathcal{R}_X$ the set of root preserving morphisms of $(\mathbf{\Omega}/X)^{\otimes}$.

If $f\colon (S,\alpha) \to (R,\beta)$ is root preserving, then $\wr(f) = \mathrm{id}_{f(r_S)}$, so the root functor factors via the localization of $\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})$ at $\mathcal{R}_X$, as

$$\begin{tikzcd} \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}) \arrow[rr, "\mathfrak{r}_X"] \arrow[dr, "\lambda"'] & & X \\ & \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})[\mathcal{R}_X^{-1}] \, . \arrow[ur, "\overline{\mathfrak{r}_X}"'] & \end{tikzcd}$$

Theorem 3.14. *For any normal dendroidal set X, the root functor induces an operadic weak equivalence of dendroidal sets*

$$\overline{\mathfrak{r}_X} \colon \mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})[\mathcal{R}_X^{-1}] \xrightarrow{\sim} X.$$

Proof. Let us use Theorem 2.4 and choose $\mathcal{L}(\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes}), \mathcal{R}_X)$ for the model of the localization of $\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})$ at $\mathcal{R}_X$. By Theorem 3.9, the functor $\mathcal{L}(-, \mathcal{R}_-) \colon \mathbf{dSets} \to \mathbf{dSets}$ is cocontinuous and preserves normal monomorphisms, so we proceed by skeletal filtration, reducing the proof to the case of $X \simeq T$ a tree, that is a representable dendroidal set. The root functor is the nerve of the map of operads $\mathfrak{r} \colon (\mathbf{\Omega}/T)^{\otimes} \to T$, and we construct a section $\mathfrak{l}_T$ of $\mathfrak{r}_T$,

$$\mathfrak{l}_T \colon T \to (\mathbf{\Omega}/T)^{\otimes}$$

and a homotopy

$$h \colon \mathcal{N}_d((\mathbf{\Omega}/T)^{\otimes}) \otimes C_1 \to \mathcal{N}_d((\mathbf{\Omega}/T)^{\otimes})$$

between $\mathfrak{l}_T \circ \mathfrak{r}_T$ and $\mathrm{id}_{(\mathbf{\Omega}/T)^{\otimes}}$, such that its components are root preserving morphisms. After Theorem 2.7, this implies that h becomes an equivalence between $\mathfrak{l}_T \circ \mathfrak{r}_T$ and id_T after localizing at $\mathcal{R}_T$, which means that $\mathfrak{l}_T$ and $\mathfrak{r}_T$ are homotopy inverses of the other once localized.

To construct h, consider an edge e of T, and let $T_e^{\uparrow}$ be the biggest subtree of T having e as root. We denote by $\iota_e \colon T_e^{\uparrow} \hookrightarrow T$ the associated subtree inclusion. Observe that, if $e \leqslant f$, then T_e is a subtree of T_f.

We define the morphism $\mathfrak{l}_T$ on an edge e of T as

$$\mathfrak{l}_T(e) := (T_e^{\uparrow}, \iota_e) \in \mathsf{Ob}((\mathbf{\Omega}/T)^{\otimes}).$$

Given edges $e_1, \dots, e_n, e$ such that $T(e_1, \dots, e_n; e) = \{*\}$, the map

$$\mathfrak{l}_T \colon \{*\} \longrightarrow (\mathbf{\Omega}/T)^{\otimes}((T_{e_1}^{\uparrow}, \iota_{e_1}), \dots, (T_{e_n}^{\uparrow}, \iota_{e_n}); (T_e^{\uparrow}, \iota_e))$$

selects the operation of $(\mathbf{\Omega}/T)^{\otimes}$ given by the forest inclusion

$$\oplus_{i=1}^n T_{e_i}^{\uparrow} \hookrightarrow T_e^{\uparrow},$$

defined by the composition of the forest root face $\oplus_{i=1}^n T_{e_i}^{\uparrow} \to \overline{\oplus_{i=1}^n T_{e_i}^{\uparrow}}$ and the inner face $\partial \colon \overline{\oplus_{i=1}^n T_{e_i}^{\uparrow}} \to T_e^{\uparrow}$ sending each $T_{e_i}^{\uparrow}$ to itself and the new root vertex to the subtree T_e^{e}.

As the operadic composition for T is the grafting of subtrees, we can check that it is a well defined map of operads $\mathfrak{l}_T \colon T \to (\mathbf{\Omega}/T)^{\otimes}$. One checks that $\mathfrak{r}_T \circ \mathfrak{l}_T = \mathrm{id}_T$; for the other composition $\mathfrak{l}_T \circ \mathfrak{r}_T$, consider an object (S, α) in $(\mathbf{\Omega}/T)^{\otimes}$. We have that

$$\mathfrak{l}_T \circ \mathfrak{r}_T(S, \alpha) = \mathfrak{l}_T(\alpha(r_S))) = (T_{\alpha(r_S)}^{\uparrow}, \iota_{\alpha(r_S)}) \, .$$

Since the image of α is contained in the subtree $T_{\alpha(r_S)}^{\uparrow}$, we can always write α as a composition

$$\begin{tikzcd} S \arrow[rr, "\alpha"] \arrow[dr, "h_{(S,\alpha)}"'] & & T \\ & T_{\alpha(r_S)}^{\uparrow} \arrow[ur, hook, "\iota_{\alpha(r_S)}"'] & \end{tikzcd}$$

for an unique root preserving morphism $h_{(S,\alpha)}$. In particular, $h_{(S,\alpha)}$ is a morphism in $(\mathbf{\Omega}/T)^{\otimes}$ and we obtain the collection of morphisms

$$h := \{h_{(S,\alpha)} \colon (S,\alpha) \to (T_{\alpha(r_S)}, \iota_{\alpha(r_S)})\}_{(S,\alpha)\in \mathsf{Ob}(\mathbf{\Omega}/T)}.$$

Let us show that h defines an homotopy between $\mathrm{id}_{(\mathbf{\Omega}/T)^{\otimes}}$ and $\mathfrak{l}_T \circ \mathfrak{r}_T$. For this purpose, we need to check the Boardman-Vogt interchange relation, so consider objects $(S_1,\beta_1),\dots,(S_n,\beta_n),(R,\gamma)$ and an operation $f \in (\mathbf{\Omega}/T)^{\otimes}((S_1,\beta_1),\dots,(S_n,\beta_n);(R,\gamma))$. Since $\gamma \circ f = (\beta_1,\dots,\beta_n)$, we have that

$$\gamma \circ f = (\iota_{\mathfrak{r}_T(\beta_1)},\dots,\iota_{\mathfrak{r}_T(\beta_n)}) \circ (h_{(S_1,\beta_1)},\dots,h_{(S_n,\beta_n)}),$$

which is precisely the wanted relation. Consider the functor of dendroidal set given by the dendroidal nerve of h; precomposing $\mathcal{N}_d(h)$ with the natural map $\mathcal{N}_d((\mathbf{\Omega}/T)^{\otimes}) \otimes C_1 \to \mathcal{N}_d((\mathbf{\Omega}/T)^{\otimes} \otimes C_1)$, we obtain an homotopy between between the identity of $(\mathbf{\Omega}/T)^{\otimes}$ and the composition $\mathfrak{l}_T \circ \mathfrak{r}_T$. As the components of this homotopy are root preserving morphisms, it becomes an equivalence after localizing at $\mathcal{R}_T$, and this concludes the argument. □

The operadic model structure on dSets admits a left Bousfield localization whose homotopy category is equivalent to that of group-like $\mathbb{E}_\infty$-algebras, in turn equivalent to infinite loop spaces. It is called the *stable model structure*, and was first introduced by Bašić-Nikolaus in [BN14]. It has the property that the induced model structure on the overcategory sSets $\simeq$ dSets$/\eta$ coincides with the Kan-Quillen model structure. In particular, one localizes also by the arrow $C_1 \to J$, which becomes an equivalence, so from Theorem 3.14 we deduce the following

Corollary 3.15. *For any normal dendroidal set X, the root functor $\mathfrak{r}_X \colon \mathcal{N}((\mathbf{\Omega}/X)^{\otimes}) \to X$ is a weak equivalence in the stable model structure for* dSets.

Specializing the above results to simplicial sets, we hence obtain the following well-known results.

Corollary 3.16 (Joyal [Joy07], Stevenson [Ste17])**.** *For any simplicial set M, the last vertex functor induces a categorical weak equivalence*

$$\Delta/M[\mathcal{R}_M^{-1}] \xrightarrow{\sim} M,$$

where $\mathcal{R}_M$ is the class of morphisms $f \colon ([n],\alpha) \to ([m],\beta)$ such that $f(n) = m$.
In particular, the last vertex functor $\Delta/M \to M$ is a weak homotopy equivalence.

Remark 3.17. It is natural to ask what happens when X is a symmetric monoidal ∞-category. In the dendroidal formalism, this corresponds to being a coCartesian fibration over the terminal object, see [Heu11]. By [Ara26], we abstractly know that for every such X there exists a relative discrete symmetric monoidal category which localizes at X. The question becomes whether this equivalence can be realized by means of the root functor.

The operad $(\mathbf{\Omega}/X)^{\otimes}$ does not have a symmetric monoidal structure when X is symmetric monoidal. One can then consider the map induced by adjunction $\mathsf{Env}((\mathbf{\Omega}/X)^{\otimes}) \to X$ from the envelope of $(\mathbf{\Omega}/X)^{\otimes}$ (still discrete) into X, and ask whether this is a localization of symmetric monoidal categories. However, it is easy to see that this holds if and only if X is itself equivalent to its envelope, $X \simeq \mathsf{Env}(UX)$ [8]. However, when X is symmetric monoidal, $(\mathbf{\Omega}/X)^{\otimes}$ has a partial monoidal structure: it is a *pre-coCartesian operad*, in the sense of [KSW24]. It would be interesting to explore further this direction.

here expand: we know the result!

[8] Observe here we need to change model and consider Lurie ∞-operads, as there is no homotopical notion of dendroidal envelope yet (see [Bar24]).

4. Localization and un/straightening equivalences

In this section, we describe the homotopy theory of algebras over a dendroidal ∞-operad in terms of locally constant algebras over its operad of elements. To this end, we study the compatibility of dendroidal localization with the covariant model structure for dendroidal left fibrations, appearing in the un/straightening equivalence.

4.1. **Localization and the covariant model structure.** For a dendroidal set X and any subset of morphisms $S \subseteq X_{C_1}$, the localization map $\lambda\colon X \to X[S^{-1}]$ induces an adjunction of over-categories

$$\lambda_!\colon \mathbf{dSets}/X \rightleftarrows \mathbf{dSets}/X[S^{-1}] : \lambda^*,$$

which is a Quillen adjunction with respect to the covariant model structure for dendroidal left fibrations ([Heu11, Proposition 2.4]). It is natural to expect dendroidal left fibrations over the localization to be weakly equivalent to those left fibrations over X for which all the maps of fibres $f_!\colon Y_a \to Y_b$ induced by the morphisms $f\colon a \to b$ in S are weak homotopy equivalences of spaces. This is indeed the case, as we show in Theorem 4.4. Let us introduce some constructions first.

Construction 4.1. Let X be a dendroidal set and S a subset of X_{C_1}. For any $f\colon a \to b$ in S, we define the $r_f\colon (\eta, \{b\}) \longrightarrow (C_1, f)$ in $\mathbf{dSets}/X$ as the following commutative triangle

$$\begin{tikzcd} \eta \arrow[rr, hook, "r"] \arrow[dr, "b"'] & & C_1 \arrow[dl, "f"] \\ & X & \end{tikzcd}$$

where the arrow $r\colon \eta \to C_1$ is the inclusion of the edge η into the root of C_1, or more familiary the map $\{1\}\colon [0] \to [1]$. We write $S_{/X}$ for the set of morphisms of the form r_f, for $f \in S$.

Given a model category $\mathcal{M}$ and some set $\mathcal{S}$ of morphisms in it, we can talk about *$\mathcal{S}$-local* objects in $\mathcal{M}$: there are the fibrant objects M for which, for any morphism $s\colon A \to B$ in $\mathcal{S}$, the morphism of mapping spaces

$$s_*\colon \mathsf{Map}_{\mathcal{M}}(B, M) \longrightarrow \mathsf{Map}_{\mathcal{M}}(A, M)$$

is a weak homotopy equivalence of spaces. For $\mathcal{M}$ given by the covariant model structure for dendroidal left fibrations over a dendroidal set X, we have an explicit way of computing mapping spaces between fibrant-cofibrant objects:

Remark 4.2. If (A, u) is a normal dendroidal set over X and (E, p) a dendroidal left fibration over X, we have an equivalence of spaces

$$\mathsf{Map}_{\mathbf{dSets}/X}((A, u), (E, p)) \simeq \hom_X(A, E) = \hom(A, E) \times_{\hom(A,X)} \{u\}.$$

Maps in $S_{/X}$ have both cofibrant source and taget; we can rephrase $S_{/X}$-locality as follows.

Definition 4.3. A (E, p) dendroidal left fibration over X is $S_{/X}$-local if, for any f in S, the morphism

$$(r_f)^*\colon \hom(C_1, E) \times_{\hom(C_1,X)} \{f\} \longrightarrow \hom(\eta, E) \times_{\hom(\eta,X)} \{b\} \simeq E_b = p^{-1}(b),$$

induced by the root inclusion $\eta \hookrightarrow C_1$[9] is a weak homotopy equivalence of Kan complexes.

We write $\mathcal{L}_S(\mathbf{dSets}/X)$ for the left Bousfield localization of the covariant model structure on $\mathbf{dSets}/X$ at the set of arrows $S_{/X}$, which, recall, is left proper and cofibrantly generated. In this model structure, the fibrant objects are the $S_{/X}$-local dendroidal left fibrations, while the cofibrations are the normal monomorphisms over X. In particular, a morphism between $S_{/X}$-local left fibrations $\varphi\colon (E, p) \to (E', p')$ is a weak equivalence in the localization if and

[9] Again, this is just the map $[0] \xrightarrow{\{1\}} [1]$.

only if it is a covariant weak equivalence, hence a fibrewise weak homotopy equivalence. We call the model structure

$$\mathcal{L}_S\left(\mathbf{dSets}/X\right)$$

the $S_{/X}$*-local covariant model structure*. It is precisely this localization that encodes the compatibility of localization with the covariant model structure:

Proposition 4.4. *Let X be a normal dendroidal set and $S \subseteq X_{C_1}$ a subset of morphisms. The localization map $\lambda\colon X \to X[S^{-1}]$ induces a Quillen adjunction*

$$\lambda_!\colon \mathcal{L}_S\left(\mathbf{dSets}/X\right) \rightleftarrows \mathbf{dSets}/X[S^{-1}] : \lambda^*$$

between the $S_{/X}$-local covariant model structure on $\mathbf{dSets}/X$ and the covariant model structure on $\mathbf{dSets}/X[S^{-1}]$. Moreover, the adjunction is a Quillen equivalence.

Proof. We know that the adjunction

$$\lambda_!\colon \mathbf{dSets}/X \rightleftarrows \mathbf{dSets}/X[S^{-1}] : \lambda^*$$

is a Quillen adjunction between covariant model structures ([HM22, Proposition 9.62]). To prove that it is still a Quillen adjunction after localizing on the left, it suffices to show that, given any dendroidal left fibration (E,p) over X, the element (E,p) is $S_{/X}$-local if and only if there exists a dendroidal left fibration (Y,p') over $X[S^{-1}]$ and a covariant weak equivalence $(E,p) \xrightarrow{\sim} \lambda^*(Y,p)$ in $\mathbf{dSets}/X$.

Fix any such dendroidal left fibration (E,p). Write $S \times C_1$, resp. $S \times J$, for the simplicial set given by the union $S \times C_1 = \bigsqcup_{s\in S} C_1$, resp. $S \times J = \bigsqcup_{s\in S} J$, where J is the nerve of the connected groupoid on two objects.

The pullback

$$E' := (S \times C_1) \times_X E \longrightarrow S \times C_1$$

is a left fibration of simplicial sets, and after [Lur09, Proposition 2.1.3.1] it is a Kan fibration. As such, it can be factored as the composition $E' \to E'' \to S \times C_1$, where $E' \to E''$ is a trivial Kan fibration and $E'' \to S \times C_1$ is a minimal Kan fibration. As explained in [GZ67, §5.4], there is an isomorphism $E'' \simeq S \times C_1 \times M$ for some minimal Kan complex M, and the map $E'' \to S \times C_1$ is the projection. In particular, the map $S \times J \times M \to S \times J$ is a minimal Kan fibration, fitting into the pullback diagram

$$\begin{tikzcd} E'' \arrow[r, "\phi"] \arrow[d] & S \times J \times M \arrow[d] \\ S \times C_1 \arrow[r] & S \times J \end{tikzcd}$$

An argument due to Joyal (see [KL21, Lemma 2.2.5]) shows that we can find a trivial Kan fibration $Z \to S \times J \times M$ and an isomorphism $E' \simeq \phi^*(Z)$ over E''. Thus we obtain a commutative diagram

$$\begin{tikzcd} E \arrow[d] & E' \arrow[l] \arrow[r] \arrow[d] & Z \arrow[d] \\ X & S \times C_1 \arrow[l] \arrow[r] & S \times J \end{tikzcd}$$

where the right hand square is a pullback and the map $Z \to S \times J$ is a Kan fibration. Define

$$Y := E \bigcup_{E'} Z$$

there is a canonical map $p'\colon Y \to X[S^{-1}]$, and its pullback along the surjection $(S \times J) \sqcup X \to X[S^{-1}]$ consists in the dendroidal left fibration $Z \sqcup E \to (S \times J) \sqcup X$. In particular, p' is a dendroidal left fibration as well. Moreover, for any object x of X, there is a weak homotopy equivalence of fibres $(E,p)_x \to \lambda^*(Y,p')_x$, hence a covariant weak equivalence

$(E,p) \to \lambda^*(Y,p')$ over X, as wanted. This concludes the proof that there is an induced Quillen adjunction. It is straightforward to check that $\mathbb{R}\lambda^*$ is fully faithful, hence we conclude that the induced adjunction is a Quillen equivalence, as wanted. □

Remark 4.5. The above proof essentially extends to the dendroidal context Stevenson's proof of [Ste17, Proposition 5.11] for simplicial sets.

4.2. **Locally constant algebras.** We now perform an analogous localization of the projective model structure on the category $\mathsf{Alg}_P(\mathbf{sSets})$ of algebras over a (discrete) operad P. In this context, one must work in the framework of semi-model categories (see [Hov98], [Bar10], [WY18]; cf. [Car23, §2] for a concise account).

Recall that a semi-model category consists of classes of weak equivalences, cofibrations, and fibrations satisfying the usual axioms, except that the lifting property for trivial fibrations and the factorization axioms are required only for maps with cofibrant source.

This weakening is necessary because left Bousfield localization of a model category at a set of maps yields a model structure under left properness, a condition that frequently fails for categories of operadic algebras. Nevertheless, such a localization still produces a semi-model structure, and it is in this sense that we proceed.

Let us give the following definition. This makes sense for any simplicial operad P.

Definition 4.6. A simplicial P-algebra A is *S-locally constant* if A sends S to equivalences, that is for any $f\colon a \to b$ in S, the map $A(f)\colon A(a) \to A(b)$ is a weak homotopy equivalence of simplicial sets.

Let us consider the operadic un/straightening equivalence of [Pra25a, Theorem 5.9], which consists, for any Σ-free operad P, in a Quillen equivalence

$$\rho_!^P\colon \mathbf{dSets}/\mathcal{N}_d(P) \rightleftarrows \mathsf{Alg}_P(\mathbf{sSets}) : \rho_P^* \tag{4.1}$$

between the covariant model structure on the left and the projective model structure on the right. The left adjoint $\rho_!^P$ is defined as follows.

Definition 4.7. The functor $\rho_!^P\colon \mathbf{dSets}/\mathcal{N}_d P \to \mathsf{Alg}_P(\mathbf{sSets})$ is the essentially unique cocontinuous functor characterized by the fact that, for any tree T and morphism $\alpha\colon T \to \mathcal{N}_d(P)$, the P-algebra $\rho_!^P(T,\alpha)$ is defined as

$$\mathsf{Ob}(P) \ni c \mapsto \rho_!^P(T,\alpha)(c) \simeq \mathsf{Env}(T) \underset{\mathsf{Env}(P)}{\times} \mathsf{Env}(P)_{/c},$$

where Env denotes the symmetric monoidal envelope [Lur17, §2.2.4].

We can now prove the following

Proposition 4.8. *Let P be a discrete Σ-free operad. The operadic un/straightening adjunction $(\rho_!^P, \rho_P^*)$ induces a Quillen equivalence*

$$\rho_!^P\colon \mathcal{L}_S\left(\mathbf{dSets}/\mathcal{N}_d(P)\right) \rightleftarrows \mathcal{L}_S\left(\mathsf{Alg}_P(\mathbf{sSets})\right) : \rho_P^*$$

between the covariant model structure for S-local left fibrations and the left Bousfield localization of the projective model structure whose fibrant objects are S-locally constant P-algebras.

Proof. In general, given a Quillen adjunction of model categories $F\colon \mathcal{M} \rightleftarrows \mathcal{N} : G$ and a choice of (cofibrant) arrows $\mathcal{S}$ in $\mathcal{M}$, one can consider the left Bousfield localizations $\mathcal{L}_{\mathcal{S}}\mathcal{M}$ on $\mathcal{M}$ and the transfered localization on $\mathcal{N}$, resp. $\mathcal{L}_{\mathbb{L}F(\mathcal{S})}\mathcal{N}$, of $\mathcal{M}$ at $\mathcal{S}$, resp. of $\mathcal{N}$ at $\mathbb{L}F(\mathcal{S})$. It is a standard result of model categories ([Hir03]) that, if (F,G) is a Quillen equivalence, then there is an induced Quillen equivalence $F\colon \mathcal{L}_{\mathcal{S}}\mathcal{M} \rightleftarrows \mathcal{L}_{\mathbb{L}F(\mathcal{S})}\mathcal{N} : G$, where $\mathcal{L}_{\mathbb{L}F(\mathcal{S})}\mathcal{N}$ is the left Bousfield localization of $\mathcal{N}$ at $\mathbb{L}F(\mathcal{S})$. This implies that we only need to characterize the fibrant objects of $\mathcal{L}_{\rho_!^P(S_{/\mathcal{N}_d(P)})}\mathsf{Alg}_P(\mathbf{sSets})$ as the locally constant fibrant P-algebras.

An object A in $\mathcal{L}_{\rho_!^P(\mathcal{S})}(\mathsf{Alg}_P(\mathbf{sSets}))$ is fibrant if and only if it is projectively fibrant and $\rho_!^P(\mathcal{S}_{/\mathcal{N}_d P})$-local. As the pair $(\rho_!^P, \rho_P^*)$ is a Quillen adjunction, this is equivalent to asking $\rho_P^*(A)$ to be $S_{/\mathcal{N}_d P}$-local, that is, that

$$\mathsf{Map}_{\mathbf{dSets}/\mathcal{N}_d(P)}((C_1, f), \rho_P^*(A)) \longrightarrow \mathsf{Map}_{\mathbf{dSets}/\mathcal{N}_d(P)}((\eta, \{b\}), \rho_P^*(A))$$

is a weak homotopy equivalence. As A is projectively fibrant, $\rho_P^*(A)$ is a dendroidal left fibration, so in particular local with respect to the map $(\eta, \{a\}) \to (C_1, f)$ induced by the leaf inclusion $\ell\colon \eta \hookrightarrow C_1$, so the above is an equivalence if and only if the map of spaces

$$\mathsf{Map}_{\mathbf{dSets}/\mathcal{N}_d(P)}((\eta, \{a\}), \rho_P^*(A)) \longrightarrow \mathsf{Map}_{\mathbf{dSets}/\mathcal{N}_d(P)}((\eta, \{b\}), \rho_P^*(A))$$

is a weak homotopy equivalence. After Theorem 4.2, there is an equivalence of spaces

$$\mathsf{Map}_{\mathbf{dSets}/\mathcal{N}_d(P)}((\eta, \{x\}) \simeq \rho_P^*(A)_x.$$

As the right adjoint ρ_P^* enjoys the property that for any P-algebra A and any object x of P, there is an equivalence

$$\rho_P^*(A)_c \simeq A(c)$$

between the fibre of $\rho_P^*(A)$ over c and the value of A on c ([Pra25a, Lemma 2.9]), we conclude that $\rho_P^*(A)$ is $S_{/\mathcal{N}_d P}$-local if and only if the map

$$A(f)\colon A(a) \longrightarrow A(b)$$

is a weak homotopy equivalence of spaces, which means that A is S-locally constant. This concludes the proof. □

Remark 4.9. The above result only depends on the fact that $\rho_!^P$ preserves and detects fibrant S-locally constant algebras, which follows from the equivalence $\rho_!^P(F)_a \simeq F(a)$ between the fiber over an object c of P of the unstraightening of an algebra F and evaluation of F at c.

As a corollary, by instantiating Proposition 4.8 with $P = (\mathbf{\Omega}/X)^{\otimes}$, which is Σ-free (Remark 3.7), we obtain a new version of the un/straightening equivalence for a normal dendroidal ∞-operad X, that is an equivalence of semi-model categories

$$\rho_!^{(\mathbf{\Omega}/X)^{\otimes}}\colon \mathcal{L}_{\mathcal{R}_X}\left(\mathbf{dSets}/\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})\right) \rightleftarrows \mathcal{L}_{\mathcal{R}_X}\left(\mathsf{Alg}_{(\mathbf{\Omega}/X)^{\otimes}}(\mathbf{sSets})\right) : \rho^*_{(\mathbf{\Omega}/X)^{\otimes}}$$

between the covariant model structure for $\mathcal{R}_X$-local dendroidal left fibrations over $\mathcal{N}_d\mathbf{\Omega}/X$ and the projective semi-model structure for $\mathcal{R}_X$-locally constant $(\mathbf{\Omega}/X)^{\otimes}$-algebras.

We may interpret this as a 'local' un/straightening equivalence. As a corollary, we deduce the following local description for algebras.

Corollary 4.10. *Let X be a normal dendroidal ∞-operad. There is a zig-zag of Quillen equivalences*

$$\mathsf{Alg}_{W_!(X)}(\mathbf{sSets}) \xleftarrow{\sim} \bullet \xrightarrow{\sim} \mathcal{L}_{\mathcal{R}_X}\left(\mathsf{Alg}_{(\mathbf{\Omega}/X)^{\otimes}}(\mathbf{sSets})\right)$$

between the projective model structures for simplicial $W_!(X)$-algebras and for $\mathcal{R}_X$-locally constant algebras on $(\mathbf{\Omega}/X)^{\otimes}$.

Proof. By [Heu11, Proposition 2.4], if $f\colon X \to Y$ is an operadic weak equivalence between normal dendroidal sets, the induced adjunction $(f_!, f^*)$ on over categories is a Quillen equivalence between covariant model structures. As this is natural in the dendroidal sets, applying Theorem 4.8 we obtain that the root functor yields a Quillen equivalence

$$(\mathscr{r}_X)_!\colon \mathcal{L}_{\mathcal{R}_X}\left(\mathbf{dSets}/\mathcal{N}_d((\mathbf{\Omega}/X)^{\otimes})\right) \rightleftarrows \mathbf{dSets}/X : \mathscr{r}_X^*.$$

Combining this with Proposition 4.8 and the classical [Heu11, Theorem 2.7], we get the wanted zig-zag of Quillen equivalences of semi-model categories. □

References

[AFT17] David Ayala, John Francis, and Hiro Lee Tanaka. Factorization homology of stratified spaces. *Selecta Mathematica*, 23(1):293–362, 2017.

[Ara26] Kensuke Arakawa. Monoidal relative categories model monoidal ∞-categories. *Journal of Pure and Applied Algebra*, page 108183, 2026.

[Bar10] Kan Barwick. On left and right model categories and left and right bousfield localizations. *Homology Homotopy and Applications*, 12(2):245–320, 2010.

[Bar24] Miguel Barata. The right cancellation property for certain classes of dendroidal anodynes, 2024.

[BdBM20] Pedro Boavida de Brito and Ieke Moerdijk. Dendroidal spaces, Γ-spaces and the special Barratt–Priddy–Quillen theorem. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2020(760):229–265, 2020.

[BdBW13] Pedro Boavida de Brito and Michael S Weiss. Manifold calculus and homotopy sheaves. *Homology, Homotopy and Applications*, 15(2):361–383, 2013.

[BN14] Matija Bašić and Thomas Nikolaus. Dendroidal sets as models for connective spectra. *Journal of K-Theory*, 14(3):387–421, 2014.

[BV73] John M. Boardman and Rainer M. Vogt. *Homotopy invariant algebraic structures on topological spaces*, volume 347. Lecture Notes in Mathematics, Vol. 347, Springer-Verlag, 1973.

[Car23] Víctor Carmona. When bousfield localizations and homotopy idempotent functors meet again. *Homology, Homotopy and Applications*, 25:187–218, 01 2023.

[CM11] Denis-Charles Cisinski and Ieke Moerdijk. Dendroidal sets as models for homotopy operads. *J. Topol.*, 4(2):257–299, 2011.

[CM13] Denis-Charles Cisinski and Ieke Moerdijk. Dendroidal sets and simplicial operads. *Journal of Topology*, 6(3):705–756, 04 2013.

[GW99] Thomas G Goodwillie and Michael Weiss. Embeddings from the point of view of immersion theory: Part ii. *Geometry & Topology*, 3(1):103–118, 1999.

[GZ67] Peter Gabriel and Michel Zisman. *Calculus of Fractions and Homotopy Theory*. Springer-Verlag New York, 1967.

[Heu11] Gijs Heuts. Algebras over ∞-operads. *arXiv:1110.1776*, 2011.

[HHM16] Gijs Heuts, Vladimir Hinich, and Ieke Moerdijk. On the equivalence between Lurie's model and the dendroidal model for ∞-operads. *Advances in Mathematics*, 302:869–1043, 2016.

[Hir03] Philip S. Hirschhorn. *Model Categories and Their Localizations*. Mathematical surveys and monographs. American Mathematical Society, 2003.

[HM22] Gijs Heuts and Ieke Moerdijk. *Simplicial and dendroidal homotopy theory*, volume 75 of *Ergeb. Math. Grenzgeb., 3. Folge*. Cham: Springer, 2022.

[HM24] Vladimir Hinich and Ieke Moerdijk. On the equivalence of the Lurie's ∞-operads and dendroidal ∞-operads. *https://arxiv.org/abs/2206.14033*, 2024.

[Hov98] Mark Hovey. Monoidal model categories. *arXiv preprint math/9803002*, 1998.

[Joy07] André Joyal. Notes on quasi-categories. available for download at *http://www.fields.utoronto.ca/av/slides/06-07/crs-quasibasic/joyal/download.pdf*, 2007.

[Ker23] David Kern. Monoidal envelopes and Grothendieck construction for dendroidal Segal objects. *arXiv preprint arXiv:2301.10751*, 2023.

[KK24] Manuel Krannich and Alexander Kupers. ∞-operadic foundations for embedding calculus. *arXiv preprint arXiv:2409.10991*, 2024.

[KL21] Krzysztof Kapulkin and Peter LeFanu Lumsdaine. The simplicial model of univalent foundations (after Voevodsky). *Journal of the European Mathematical Society*, 23(6):2071–2126, 2021.

[KSW24] Eilind Karlsson, Claudia I Scheimbauer, and Tashi Walde. Assembly of constructible factorization algebras. *arXiv preprint arXiv:2403.19472*, 2024.

[Lur09] Jacob Lurie. *Higher Topos Theory*. Princeton University Press, 2009.

[Lur17] Jacob Lurie. *Higher Algebra*. Available at *http://math.ias.edu/ lurie/*., 2017.

[May06] J Peter May. *The geometry of iterated loop spaces*, volume 271. Springer, 2006.

[MW07] Ieke Moerdijk and Ittay Weiss. Dendroidal sets. *Algebraic & Geometric Topology*, 7:1441–1470, 2007.

[Pra25a] Francesca Pratali. Rectification of dendroidal left fibrations. *arXiv:2502.17415*, 2025.

[Pra25b] Francesca Pratali. A straightening-unstraightening equivalence for ∞-operads, 2025.

[Ram22] Maxime Ramzi. A monoidal Grothendieck construction for ∞-categories. *arXiv:2209.12569*, 2022.

[Ste17] Danny Stevenson. Covariant model structures and simplicial localization. *North-West. Eur. J. Math.*, 3:141–202, 2017.

[Wal85] Friedhelm Waldhausen. Algebraic K-theory of spaces. In Andrew Ranicki, Norman Levitt, and Frank Quinn, editors, *Algebraic and Geometric Topology*, pages 318–419, Berlin, Heidelberg, 1985. Springer Berlin Heidelberg.

[Wei99] Michael Weiss. Embeddings from the point of view of immersion theory: Part I. *Geometry & Topology*, 3(1):67–101, 1999.

[WY18] David White and Donald Yau. Bousfield localization and algebras over colored operads. *Applied Categorical Structures*, 26, 02 2018.