


# RESPIRATION

All living organisms need energy. Respiration: the life process by which organisms convert the chemical energy stored in food to a form of energy more easily utilized by the cell. Process of Cell Respiration: a biochemical process used by cells to release energy from organic molecules (food) such as glucose:

- ~ This energy is stored in the molecule ATP
- ~ ATP = adenosine triphosphate or A~P~P~P
- ~ The process is controlled by enzymes



Life on Earth first evolved without free oxygen ( $O_2$ ) in atmosphere and energy had to be captured from organic molecules in absence of  $O_2$ , so organisms that evolved glycolysis are ancestors of all modern life.

Cellular respiration is a chain of pathway, through which the carbohydrate molecule and other molecule is oxidized into  $CO_2$  and  $H_2O$  and release of energy which is used in reactions that needs energy, as well as obtaining carbon skeletons needed for growth and development of the cell. Higher plants are aerobic organisms, that gets its energy and carbon by respiration



Oxidation of hexose sugar into  $CO_2$  and  $H_2O$  takes place in three separate stages:

1. Glycolysis.
2. Citric acid cycle (or Krebs cycle).
3. Electron transport chain.


Glycolysis

Glycolysis is breaking down glucose □ “glyco – lysis” (splitting sugar).


It is the most ancient form of energy capture and the starting point for all cellular respiration □ inefficient □ generate only 2 ATP for every 1 glucose □ in cytosol.

It is also known as EMP (or Eembden - Myerhofparnas).


It represents the first stage of respiration. It consists of a chain of anaerobic reactions which takes place in the cytoplasm. The six-carbon sugar molecule is divided and partially oxidized into 3-carbon inorganic acid called pyruvic acid or pyruvate.



The overall balance (or balanced reaction) of glycolysis reaction:






Further transformation of pyA depends on the presence or absence of oxygen. In aerobic conditions the pyA is transferred into mitochondria, where it is oxidized into CO<sub>2</sub> and H<sub>2</sub>O. Under anaerobic conditions (e.g. in the roots when the soil is saturated with water).fermentation takes place, and here pyA will either be transformed into ethyl alcohol (which is the result of fermentation in most of the plants) or lactic acid or other organic acids depending on microorganism.



## Mitochondria

- Double membrane
  - ◆ outer membrane
  - ◆ inner membrane
    - highly folded cristae\*
    - fluid-filled space between membranes = intermembrane space
- ◆ matrix
  - central fluid-filled space






Citric acid cycle:

It is known as Krebs cycle or Tricarboxylic acid cycle (TCA).

## Oxidation of pyruvate

### Pyruvate enters mitochondria




- ◆ 3 step oxidation process
- ◆ releases 1  $\text{CO}_2$  (count the carbons!)
- ◆ reduces  $\text{NAD} \rightarrow \text{NADH}$  (stores energy)
- ◆ produces acetyl CoA

### Acetyl CoA enters Krebs cycle

In the presence of oxygen, pyA is oxidized into  $\text{CO}_2$  and  $\text{H}_2\text{O}$  in the cycle. It is the second stage in respiration and takes place in the mitochondria.

Pyruvic acid is transferred to the mitochondria and passes into a series of reactions (5 reactions), that ends up with the oxidation of pyA molecule and produces a 2-Carbon compound called Acetyl Co-A.



## Pyruvate oxidized to Acetyl CoA



The Acetyl-Co-A enters Krebs cycle where it first react with a 4-carbon organic acid called OAA (Oxaloacetic acid) forming Citric acid which enters a chain of reactions through which energy is released in the form of ATP, or used for reduction of the soluble (respiration carries) NAD into NADH2 and FAD+(flavin adenine dinucleotide) into FADH2 .

Every NADH that its electron is transferred into the ETC will produce 3 molecules of ATP, and each FADH2 produces 2 ATP.


Overall energy (in the of ATP) that results from complete oxidation of one glucose molecule in respiration is as follow



Net gain = 2 ATP  
= 8 NADH + 2 FADH<sub>2</sub>

Pathway	NADH X3	FADH <sub>2</sub> X2	ATP	Total ATP
Glycolysis	2(6ATP)	0	2	8
PyA → acetyl Co-A	2(6ATP)	0	0	6
Krebs cycle	6(18ATP)	2(4ATP)	2	24
Total ATP	10x3=30	2x2=4	4	38






## Electron Transfer Chain

Redox reactions release energy when electrons “fall” from a hydrogen carrier to highly electronegative oxygen

- **NADH ( & FADH<sub>2</sub> )** delivers electrons to a series of electron carriers in an electron transport chain
  - As electrons move from carrier to carrier, their energy is released in small quantities




The number of ATP molecules that results from glycolysis and Krebs cycle is low (i.e. four molecules only). Most of potential energy in the sugar (e.g. glucose molecule) is present in the form of electron pairs (12 pairs), 10 pairs in the form of NADH and 2 pairs in the form of FADH<sub>2</sub>. The stored energy in these reduced carriers is liberated in the form of ATP in the third stage of respiration. In this stage the electrons from NADH and FADH<sub>2</sub> are transferred to the oxygen, coupled with transformation of energy in ATP in the electron transfer system which takes place in the inner membrane of the mitochondria.

## Electron Transport Chain

### ■ NADH passes electrons to ETC

- ◆ H cleaved off NADH & FADH<sub>2</sub>
- ◆ electrons stripped from H atoms → **H<sup>+</sup>** (H ions)
- ◆ electrons passed from one electron carrier to next in mitochondrial membrane (ETC)
- ◆ transport proteins in membrane pump **H<sup>+</sup>** across inner membrane to intermembrane space



’ Biology

2005-20

The electron transfer chain consists of a chain of carriers of NAD, FMN, CoQ and cytochromes b, c, a and a<sub>3</sub> with gradual reduction in energy (i.e. each carrier has an energy less than the previous one). The energy released in the different stages of the electron transfer chain is used for ATP building.

The carriers are enzyme & coenzymes located on the inner membrane of the mitochondria & it resembles the photosynthesis carriers that found in the thylakoids of the chloroplast.




## Why the build up H<sup>+</sup>?

### ■ ATP synthase

- ♦ enzyme in inner membrane of mitochondria




- ♦ only channel permeable to H<sup>+</sup>
- ♦ H<sup>+</sup> flow down concentration gradient = provides energy for ATP synthesis
  - molecular power generator!
  - flow like water over water wheel
  - flowing H<sup>+</sup> cause change in shape of ATP synthase enzyme
  - powers bonding of P<sub>i</sub> to ADP
  - "proton-motive" force



## ATP synthesis


### ■ Chemiosmosis couples ETC to ATP synthesis

- ♦ build up of H<sup>+</sup> gradient just so H<sup>+</sup> could flow through ATP synthase enzyme to build ATP




## Electrons flow downhill

- Electrons move in steps from carrier to carrier downhill to  $O_2$ 
  - ♦ each carrier more electronegative
  - ♦ controlled oxidation
  - ♦ controlled release of energy



AP Biology


## Cellular respiration



The Pentose Phosphate Pathway  
(or Hexose Monophosphate Shunt).

In higher plant, as well as animals, this pathway for glucose metabolism is also found. It takes place in the cytoplasm and requires oxygen. The first step is oxidation of glucose -6-p to 6-phosphogluconate, which loses a  $CO_2$  to produce a phosphorylated 5-

carbon sugar called Ribulose-5-phosphate, which can be converted into other compound, or becomes a part of nucleotides of DNA and RNA or other compound.




1. Production of reduction power NADPH (instead of NADH) which is used as a donor for electron in metabolic reactions.
2. Production of pentose sugar, which is the source for Ribose and deoxyribose needed for nucleic acid synthesis (DNA and RNA).
3. Production of erythrose-4-P, which is the source of aromatic amino acids, lignin and flavonoids.

Relationship between respiration and building blocks:-

The intermediate compound of respiration usually enters other metabolic pathways to build up building blocks for the cell before complete oxidation into CO<sub>2</sub> and water.

This means that the main functions of Respiration are:-


1. Liberation of energy needed for metabolism.
2. Building of cell building blocks or units, regardless of their type.



## Respiratory Quotient (RQ)

It is the ratio of liberated  $\text{CO}_2$  to the absorbed  $\text{O}_2$  when substrate (i.e. the matter of respiration) is oxidized.

In aerobic Respiration: This ratio is almost 1 always because:



Therefore for Carbohydrates:

$$RQ = \frac{6(\text{CO}_2)}{6(\text{O}_2)} = 1.0$$

For lipids:

RQ depends on the chemical structure of the molecule. It is usually between 0.7-0.8



$$RQ = \frac{18(\text{CO}_2)}{26(\text{O}_2)} = 0.7$$

For Protein

RQ usually between 0.8 and 1.0, depending on the oxidation degree

Measuring of RQ is necessary to determine the nature of the oxidized compound to produce the energy

- **Glycolysis accepts a wide range of carbohydrates fuels**


- ♦ polysaccharides  $\rightarrow \rightarrow \rightarrow$  glucose  
hydrolysis

- ex. starch, glycogen

- ♦ other 6C sugars  $\rightarrow \rightarrow \rightarrow$  glucose  
modified

- ex. galactose, fructose

- Proteins  $\rightarrow \rightarrow \rightarrow \rightarrow$  amino acids  
hydrolysis






■ Fats  $\rightarrow \rightarrow \rightarrow \rightarrow \rightarrow$  glycerol & fatty acids  
hydrolysis

◆ glycerol (3C)  $\rightarrow \rightarrow$  PGAL  $\rightarrow \rightarrow$  glycolysis

◆ fatty acids  $\rightarrow$  2C acetyl  $\rightarrow$  acetyl  $\rightarrow$  Krebs  
groups coA cycle



## Carbohydrates vs. Fats

■ Fat generates 2x ATP vs. carbohydrate

◆ more C in gram of fat

◆ more O in gram of carbohydrate

▪ so it's already partly oxidized

