

Power Point No. 4

Intracellular bacterial cell structure

- In comparison to [eukaryotes](#), the intracellular features of the bacterial cell are extremely simple. Bacteria do not contain [organelles](#) in the same sense as [eukaryotes](#). Instead, the [chromosome](#) and perhaps [ribosomes](#) are the only easily observable intracellular structures found in all [bacteria](#). They do exist, however, specialized groups of bacteria that contain more complex intracellular structures
- **The bacterial chromosome and plasmids**
- Unlike [eukaryotes](#), the bacterial [chromosome](#) is not enclosed inside of a membrane-bound [nucleus](#) but instead resides inside the bacterial [cytoplasm](#). This means that the transfer of cellular information through the processes of [translation](#), [transcription](#) and [DNA replication](#) all occur within the same compartment and can interact with other cytoplasmic structures, most notably [ribosomes](#).

- The bacterial chromosome is not packaged using [histones](#) to form [chromatin](#) as in [eukaryotes](#) but instead exists as a highly compact [supercoiled](#) structure and containing spermine and spyrmidine , the function of which remains unclear.
- Most bacterial chromosomes are [circular](#) although some examples of linear chromosomes exist (e.g. [Borrelia burgdorferi](#)). Along with chromosomal DNA, most bacteria also contain small independent pieces of DNA called [plasmids](#) that often encode for traits that are advantageous but not essential to their bacterial host. [Plasmids](#) can be easily gained or lost by a bacterium and can be transferred between bacteria as a form of [horizontal gene transfer](#)

Ribosomes and other multiprotein complexes

- In most [bacteria](#) the most numerous intracellular structure is the [ribosome](#), the site of [protein synthesis](#) in all living organisms. All [prokaryotes](#) have 70S ribosomes while [eukaryotes](#) contain larger 80S [ribosomes](#) in their [cytosol](#). The 70S [ribosome](#) is made up of a 50S and 30S subunits. The 50S subunit contains the 23S and 5S [rRNA](#) while the 30S subunit contains the 16S [rRNA](#). These [rRNA](#) molecules differ in size in [eukaryotes](#) and are complexed with a large number of ribosomal proteins, the number and type of which can vary slightly between organisms. While the [ribosome](#) is the most commonly observed intracellular multiprotein complex in [bacteria](#), other large complexes do occur and can sometimes be seen using [microscopy](#).

- **Intracellular membranes**

- While not typical of all [bacteria](#) some microbes contain intracellular membranes in addition to (or as extensions of) their cytoplasmic membranes. An early idea was that bacteria might contain membrane folds termed [mesosomes](#), but these were later shown to be artifacts produced by the chemicals used to prepare the cells for [electron microscopy](#). Examples of [bacteria](#) containing intracellular membranes are [phototrophs](#), [nitrifying bacteria](#) and [methane-oxidising](#) bacteria. Intracellular membranes are also found in [bacteria](#) belonging to the poorly studied [Planctomycetes](#) group, although these membranes more closely resemble organellar membranes in [eukaryotes](#) and are currently of unknown function.

- **Cytoskeleton**

- The prokaryotic cytoskeleton is the collective name for all structural [filaments](#) in [prokaryotes](#). It was once thought that prokaryotic cells did not possess [cytoskeletons](#), but recent advances in visualization technology and structure determination have shown that filaments indeed exist in these cells. In fact, [homologues](#) for all major cytoskeletal proteins in [eukaryotes](#) have been found in prokaryotes. Cytoskeletal elements play essential roles in cell division, protection, shape determination, and polarity determination in various prokaryotes.

- **Nutrient storage structures**

- Most Bacteria do not live in environments that contain large amounts of nutrients at all times. To accommodate these transient levels of nutrients. Bacteria contain several different methods of nutrient storage in times of plenty for use in times of want. For example, many bacteria store excess carbon in the form of [polyhydroxyalkanoates](#) or [glycogen](#). Some microbes store soluble nutrients such as [nitrate](#) in [vacuoles](#). Sulfide is most often stored as elemental (S^0) granules which can be deposited either intra- or extracellularly. Sulfur granules are especially common in [bacteria](#) that use [hydrogen sulfide](#) as an electron source.

- **Gas vesicles**
- Gas vesicles are spindle-shaped structures found in some [planktonic](#) bacteria that provides [buoyancy](#) to these cells by decreasing their overall cell density. They are made up of a protein coat that is very impermeable to solvents such as water but permeable to most gases. By adjusting the amount of gas present in their gas vesicles bacteria can increase or decrease their overall cell density and thereby move up or down within the water column to maintain their position in an environment optimal for growth.

- **Carboxysomes**

- Carboxysomes are intracellular structures found in many [autotrophic bacteria](#) such as *Cyanobacteria*, *Knallgasbacteria*, and *Nitrobacteria*. They are proteinaceous structures resembling phage heads in their [morphology](#) and contain the enzymes of carbon dioxide fixation in these organisms (especially ribulose biphosphate carboxylase/oxygenase (RuBisCO) and carbonic anhydrase). It is thought that the high local concentration of the enzymes along with the fast conversion of bicarbonate to carbon dioxide by carbonic anhydrase allows faster and more efficient carbon dioxide fixation than possible inside the cytoplasm. Similar structures are known to harbor the coenzyme B₁₂-containing glycerol dehydratase, the key enzyme of glycerol fermentation to 1,3-propanediol, in some Enterobacteriaceae (e. g. *Salmonella*).


- **Magnetosomes**

- Magnetosomes are intracellular organelles like structure found in magnetotactic bacteria that allow them to sense and align themselves along a magnetic field (magnetotaxis). The ecological role of magnetotaxis is unknown but it is hypothesized to be involved in the determination of optimal oxygen concentrations. Magnetosomes are composed of the mineral magnetite and are surrounded by a lipid bilayer membrane. The morphology of magnetosomes is species-specific.

- **Endospores**

- Perhaps the most well known bacterial adaptation to stress is the formation of [endospores](#). [Endospores](#) are bacterial survival structures that are highly resistant to many different types of chemical and environmental stresses and therefore enable the survival of [bacteria](#) in environments that would be lethal for these cells in their normal vegetative form. It has been proposed that [endospore](#) formation has allowed for the survival of some [bacteria](#) for hundreds of millions of years (e.g. in salt crystals). [Endospore](#) formation is limited to several genera of Gram-positive [bacteria](#) such as [Bacillus](#) and [Clostridium](#). It differs from reproductive spores in that only one spore is formed per cell resulting in no net gain in cell number upon [endospore](#) germination. The location of an [endospore](#) within a cell is species-specific and can be used to determine the identity of a [bacterium](#).

Steps of bacterial endospore formation



- **Intracellular Structures of Eukaryotic Microorganisms**
- **Microbodies:**
- Microbodies are a family of organelles bounded by single membrane. They carry out a range of functions in fungal cells. The major groups are peroxisomes that contain at least one oxidized enzyme that produce hydrogen peroxide (H_2O_2) together with peroxidase which decompose hydrogen peroxide to produce oxygen or oxidize substrates. Some peroxidase have another functions in the oxidation of fatty acids to give Acetyl-CoA as in the case of *S. cerevisiae* . Another function to provide site action of enzymes of glyoxylate cycle which required to covert carbon compounds to carbohydrate and this microbody named Glyoxysome

Peroxisomes plays an important roles in utilization of a wide range of carbon and nitrogen sources as in *Hansenula polymorpha* which contain enzymes for oxidation of methanol and methylamine. Other peroxisomes contain uricase enzyme which hydrolyze uric acid as in the case of *Candida lipolytica*. Peroxisomal enzymes allow a range of yeast species to grow on unusual substrate such as alkanes, alkylamine, fatty acids amino acids and uric acids.

- **Reserve Materials:**

- They also reserve glycogen 1-10 million molecular weight which is water soluble, but that of 100 million molecular weight are insoluble in water. Glycogen is a polymer of glucose molecules forming alfa-glucans linked together in identity $\alpha(1-4)$ glycosidic bonds forming long chains. After 10 glucose molecules it linked to third glucose molecules by $\alpha(1-6)$ glycosidic bond forming chain branching.
- Reserve materials in fungal cells also found as a soluble carbohydrates and the common one is trehalose as a disaccharide. Basic amino acids and poly phosphate were found in some fungal cells as a reserved materials.

- **Mitochondria and Mitochondrial DNA:**
- Mitochondria are the site of oxidative phosphorylation in eukaryotic cells such as fungi. The enzymes concerned with electron transport and ATP production located in the inner membrane of mitochondria. The area of which are increased by invagination named cristae.
- Mitochondria are derived from preexisting mitochondria and both parent contribute mitochondria. Fusion may occur between mitochondria and then fragmentation of large mitochondrion to yield several smaller mitochondrial organelles. Under anaerobic conditions the number of mitochondria decreased, become less complex smaller

in size and electron transport enzymes such as cytochromes disappear as in the case of *Saccharomyces cerevisiae*. The mitochondria are multiplied by simple fission in some fungi as in *Physirum polycephalum*. Mitochondria contain circular double stranded DNA (MitDNA). MitDNA constitutes the mitochondrial genome and contain gene for tRNA, rRNA and for some enzymes involved in oxidative phosphorylation as in *S. cerevisiae*.

- In *Neuorospora crassa* gene present in both nucleus and mitochondria involved in the production of these enzymes. The mitochondrial genome size in fungi is different from species to species and this size ranged between 26000bp in *Phycomyces* to about 8000bp in *S. cerevisiae*.

- **Lipid Particles: (Sphaerosomes):**
- Fungi like other organisms can accumulate lipids as a carbon source and the reserved oil droplets can be observed under light microscope in older cells. Function as storage vesicles which may serve as lipid reservoir for yeast membrane biosynthesis. They contain mainly sterol ester, but not triglycerides , and small amounts of phospholipids, protein and unsaturated fatty acids. The content of the latter are known to increase in backers yeast at the end of stationary growth phase in batch culture.

- **Proteasomes:**

- Are large multisubunit protease enzyme complexes found in cytoplasm of yeast cells with no apparent association with intracellular structures. In *S. cerevisiae*, 20S and 26S proteasomes exist whose function in regulating protein levels is essential for cell viability. The 26S particle is assembled from the 20S units which are cylindrical particles composed of four stacked rings. The 26S proteasome acts as an ATP dependent protease in the ubiquitin pathway which is responsible

- for the rapid degradation of short lived, abnormal proteins that are detrimental to the cell.
- Specific yeast physiology functions in which proteasomes are involved include cell cycle control, signal translocation , mating and adaptive stress responses. These multifunctional roles of proteasomes have postulated by Hilt and Wolf (1996).

- **Vacuoles:**

- Vacuoles are very abundant in fungal cells and most obvious organelle seen under light microscope. It is dynamic structure, playing multiple roles in hyphal and yeast cell growth rather than being inert depositories for storage of water and nutrients or waste materials.
- In growing fungi vacuole play an important in storage of metabolites, Regulator of pH, as an equivalent to an animal lysosomes, with range of lytic enzymes more than 60 enzymes such as proteases, nucleases, phosphatases, trahalase, lipases and glucosidases and secretion of extracellular polysaccharides. The vacuole surrounded by one layer membrane of lipoprotein named **Tonoplast** .

- The pH of the vacuole is about 4.5-5.5.
- Moreover, vacuole store amino acids as a source of nitrogen under the condition of nitrogen starvation and also store cations such K^+ , Ca^{+2} , Mg^{+2} , Mn^{+2} , Fe^{+2} and Zn^{+2} . The vacuolar hydrolytic enzymes may regulate the recycling of cell materials during growth, developments, lyses of dead cells and Liquefaction of tightly gills of fruit bodies of cup mushroom which allows the release of spores from the gills. In higher fungi such ascomycetes and basidiomycetes the vacuoles accumulate fluorescent dyes such as 6-carboxy-fluorecein which can be detected under fluorescence microscope. Finally the vacuoles can be transported across the fungal cell cytoplasm up to 60um distant by the help of microtubules.

- **6- Plasmid:**

- Plasmid are pieces of DNA can replicated independently of replication of chromosomes or mitochondrial DNA.
Saccharomyces cerevisiae contain circular plasmid of 2um double stranded DNA consist of about 6200base pair (bp) and have no any role in the life of yeast cells.
- The plasmid can be multiplied up to 100 copies providing important helps in the construction of vector for the genetic manipulation of the yeast.
- *Kluveromyces lactis* yeast also contain two linear plasmid of different length.

- One of them named killer plasmid producing an extracellular protein (toxin) that kill many yeasts such as *candida* and *Saccharomyces*. *Podospora anserine* contain plasmid originated from mitochondrial DNA causes fungal **senescence** in growth rate falls and vegetative propagation become impossible and this plasmid termed SecDNA which can be transmitted via parent materials to **ascospores**.