

BMS2004 Week 9

Liang Zhang

BMS

Liangzhang.28@cityu.edu.hk

Tel: 3442-4495

Questions to think about

- Why do we eat foods?
- What happens to the foods after we eat them?
- Why do we breathe oxygen all the time?
- Are other animals the same with us?

Metabolism and Energy

Metabolism


- The cost of living – energy.
- All the time, living organisms consumes and produces energy through chemical reactions.
- Collectively, we call these chemical reactions **metabolism**.
- What is true of E coli. is true of the elephant.

--- Jacques Monod

Counting the energy



- Ribonucleoside triphosphate are the currency of energy:

ATP, **GTP**, **UTP**, and **CTP**. The energy is stored in the multi-phosphate bonds.





The metabolic network

- A network of pathways.
- All inter-connected.
- Which direction to go?



The metabolic network

- A network of pathways.
- All inter-connected.
- Which direction to go?
--- Gibbs free energy and external regulation.



Direction of metabolic reactions – Gibbs free energy (G)



$$\Delta G = G_{(C + D)} - G_{(A + B)}$$

If $\Delta G < 0$, reaction goes left to right;

If $\Delta G = 0$, reaction is at equilibrium;

If $\Delta G > 0$, reaction goes right to left

What determines ΔG ?




Reactants

Products

$$\Delta G = \Delta G'_0 + RT \ln(\{\text{Products}\}/\{\text{Reactants}\})$$

What determines ΔG ?




$$\Delta G = \Delta G'_0 + RT \ln(\{\text{Products}\}/\{\text{Reactants}\})$$

$\Delta G'_0$: change of G at standard conditions.

For a given reaction, if $\Delta G'_0 < 0$, then that direction is energetically favorable, meaning the reaction tends to undergo in that direction.

If $\Delta G'_0 > 0$, then that direction is energetically unfavorable, meaning the reaction tends to undergo in the reverse direction.

What determines ΔG ?



$$\Delta G = \Delta G'_0 + RT \ln(\{\text{Products}\}/\{\text{Reactants}\})$$

But there is another part in the equation: $\{\text{Products}\}/\{\text{Reactants}\}$.



If the concentration of products is very low and the concentration of reactants is very high, even if $\Delta G'_0$ is positive (energetically unfavorable), ΔG may be negative and the reaction go as written – pushing and pulling.

Summary of objectives

- Metabolism is collection of chemical reactions in the body
- Central theme of metabolism: energy
- Determining the direction of reactions: ΔG , $\Delta G'_0$ and concentration of reactants/products (pushing and pulling).

Glycolysis

Metabolism of carbohydrates




http://www.genome.jp/kegg-bin/show_pathway?map01100

BMS2004



14

14


Central theme -- energy



The pathways are not isolated, and the reactions don't stop



Glycolysis – an introduction



Literally, glycolysis means the **splitting of sugar**.

It was the first metabolic pathway to be elucidated (yet it took almost 100 years!).

It is an almost **universal** central pathway of glucose metabolism, because it is found in mammals, plants, and microorganisms (especially anaerobic ones).

What is glycolysis?


Glycolysis is the sequence of reactions that metabolizes **one molecule of glucose** to **two molecules of pyruvate** with the concomitant net production of **TWO** molecules of ATP.

The net reaction of glycolysis is:




This process is **anaerobic** (i.e., it does not require O_2).


The reactions take place in the **cytosol**.



Some facts about glycolysis




Two ATP is generated, but is a major source of energy during intensive exercising, e.g. sprint.



A. Low O_2
(last seconds of a sprint)

B. Normal
(long slow run)

Biochemistry, 7th edition
© W.H. Freeman and company




Summary of objectives

- Glycolysis is the breakdown of simple sugars.
- Starting: 1 glucose; Ending: 2 pyruvates.
- ATP generated – 2.
- Location – cytosol.

The reactions


The reactions




10 steps in total

Can be divided into 2 phases

(a)




Step (1): Formation of glucose 6-phosphate




- (1) **Hexokinase** traps glucose in the cell because glucose 6-phosphate cannot diffuse through the membrane (as it is negatively charged).
- (2) The addition of the phosphoryl group begins to destabilize glucose, facilitating its further metabolism.

Step (2): Isomerization of glucose 6-phosphate to fructose 6-phosphate



In the open chain form, glucose has an **aldehyde group on C1**, whereas the fructose has a **keto group on C2**. Thus the isomerization of glucose 6-phosphate to fructose 6-phosphate is a conversion of an aldose to ketose. The purpose of this step is to make the C1 carrying a OH group for the next step.

Step (3): Formation of fructose 1,6-biphosphate




This reaction is catalyzed by *phosphofructokinase (PFK)*, an **allosteric enzyme** that sets the pace of glycolysis.

The reaction is almost irreversible under cellular condition, and it is the ***first committed step*** in the glycolytic pathway.

Thus fructose 1,6-bisphosphate is only targeted for glycolysis.

Step (4): Splitting of fructose 1,6-biphosphate

$\Delta G^{\circ} = 23.8 \text{ kJ/mol}$ (so how the cells drive this reaction?)




Fructose 1,6-bisphosphate

6C unit


A reversible reaction

Aldolase
 \rightleftharpoons



Dihydroxyacetone phosphate

3C unit




Glyceraldehyde 3-phosphate

3C unit

Glyceraldehyde 3-phosphate is on the direct pathway of glycolysis, while dihydroxyacetone phosphate is not (see next step)!


Step (5): Isomerization of dihydroxyacetone phosphate



N.B. At equilibrium, 96% of the triose phosphate is dihydroxyacetone phosphate. However, the reaction proceeds readily from dihydroxyacetone phosphate to glyceraldehyde 3-phosphate because of the efficient removal of this product by subsequent reactions.


At this point, *2 molecules of glyceraldehyde 3-phosphate* are formed from *1 glucose* molecule! *2 ATP* have been invested so far!

(a)




Step (6): Conversion of glyceraldehyde 3-phosphate into 1,3-bisphosphoglycerate

Start of payoff stage



1,3-Bisphosphoglycerate is an acyl phosphate, and one of its phosphoryl groups is transferred to ADP in the next step in glycolysis.

Step (7): Formation of ATP from 1,3-bisphosphoglycerate




ATP is formed!!!

The formation of ATP by phosphoryl group transfer from 1,3-bisphosphoglycerate is referred to as a **substrate-level phosphorylation**. It occurs in the cytoplasm and involves soluble enzymes and chemical intermediates.


Oxidative phosphorylation in mitochondria.

Photophosphorylation in plants.

Step (8): Formation of 2-phosphoglycerate




Step (9): Formation of phosphoenolpyruvate




PEP: one of the highest energy in the cells.

Step (10): Formation of Pyruvate




Phosphoenolpyruvate

The phosphoryl group traps the molecule in its unstable enol form. When the phosphoryl group has been donated to ATP, the enol molecule undergoes a conversion into the more stable ketone namely, pyruvate.



$$\Delta G'^{\circ} = -31.4 \text{ kJ/mol}$$

ATP is formed!!!




Enolpyruvate

enol form

The second substrate-level phosphorylation in glycolysis.


Tautomerization



Pyruvate

“Big bang” – almost enough energy to form a second ATP, but gets lost as heat.


Summary of objectives



1. Two phases: preparatory and payoff.
2. Not all reactions are energetically favorable.
3. 1 glucose molecule is converted into 2 molecules of pyruvate.
4. Investment of two ATP molecules generates 4 molecules of ATP (per glucose) and 2 molecules of NADH.


The fates of pyruvate

What happens to pyruvate?




A. Low O₂
(last seconds of a sprint)

B. Normal
(long slow run)




The fate of pyruvate – aerobic




Under aerobic conditions ***pyruvate is oxidized to acetate*** (or acetyl-CoA), which enters the citric acid cycle and is oxidized to CO_2 and H_2O , and NADH formed in glycolysis is reoxidized to NAD^+ by passage of its electrons to O_2 in mitochondria respiration, and this provides more energy for ATP synthesis (see coming sections).

Recycling NAD⁺ is important !




The fate of pyruvate in hypoxia



Under hypoxic conditions, **NADH generated** by glycolysis cannot be reoxidized by O₂. Failure to regenerate NAD⁺ would stop glycolysis and no energy can be produced. Most organisms have the ability to continually regenerate NAD⁺ during anaerobic glycolysis by transferring electrons from NADH to form lactate or ethanol.


Fermentation:

Formation of lactic acid



- ❑ The **lactate** formed by active skeletal muscles can be recycled: it is carried in the blood to the liver where it is **converted to glucose** during the recovery from strenuous muscular activity (see later sections).
- ❑ When lactate is produced in large amount, the acidification that results from ionization of lactic acid in muscles and blood causes pain and limits the period of vigorous activity.
- ❑ Erythrocytes have no mitochondria and thus cannot reoxidize pyruvate to carbon dioxide. These cells produce lactate even in the presence of oxygen.
- ❑ Thus through glycolysis and fermentation, the conversion of one molecule of glucose yields two lactate ions and two molecules of ATP.

Alcoholic Fermentation



- ❑ Yeast and other microorganisms ferment glucose to ethanol and CO_2 . Pyruvate is converted to ethanol and CO_2 in two steps.
- ❑ One molecule of glucose produces 2 molecules of ethanol, 2 molecules of carbon dioxide and 2 molecules of ATP.



Carbon dioxide release in fermentation

- Pyruvate decarboxylase is present in brewer's and **baker's yeast** (*Saccharomyces cerevisiae*), plants, and other microorganisms.


dough



- Carbon dioxide released in the reaction can be seen in the bottles of champagne and in the dough of bread.




What about animals?



- ☐ In animals including human, pyruvate decarboxylase is not present.
- ☐ Therefore, pyruvate needs to go down the lactate pathway.

But we have the other half to consume ethanol




So it is not always right:

What is true of E coli. is true of the elephant.

--- Jacques Monod

What is the product of ethanol consumption?



- ❑ In many organisms including human, alcohol dehydrogenase is present to convert ethanol back to acetaldehyde.
- ❑ Ethanol in the blood of human is converted back into acetaldehyde by **alcohol dehydrogenase (ADH)**, which is further oxidized to acetic acid by **aldehyde dehydrogenase (ALDH)**. High level of acetaldehyde (in alcoholics) exerts its toxic effect by inhibiting mitochondrial functions.
- ❑ Half of the east Asians have a genetic variant of the **aldehyde dehydrogenase (ALDH)**, which is less efficient at converting acetaldehyde into acetate. When acetaldehyde builds up, one would have flushing of their face, hot sensations, nausea, and palpitation.

Some common foods made by fermentation

Microorganism	Starting material	Fermentation product	Food Product
<i>Lactobacillus bulgaricus</i> (lactic acid bacteria)	milk	lactic acid	yogurt
<i>Propionibacterium freudenreichii</i>	milk	Propionic acid and CO ₂	Swiss cheese

The acids produced in fermentation can lower the pH to denature and precipitate the proteins in the process and make the food sour in taste.

Other foods made by fermentation: pickles, sauerkraut, sausage, soy sauce, kimchi.



BMS2004



Summary of objectives

1. The fate of pyruvate depends on the availability of oxygen.
2. With oxygen – citric acid cycle.
3. Low oxygen – fermentation.
4. Reason – recycle of NAD^+ .

