

Complex-Order and Fractional Derivatives: A First Exploration

Faisal Khalid

September 2025

Abstract

This paper presents an independent exploration of fractional and complex-order derivatives, building the framework from algebraic first principles using Gamma function extensions. Without prior exposure to existing fractional calculus literature, I derive formulas for D^α and D^z operators applied to elementary functions, prove fundamental properties (linearity, Index Law, product rules), and explore geometric interpretations through the " $D(i)$ plane." The work concludes with applications to cyclic derivatives and preliminary extensions to matrix orders.

Note to Readers: This represents independent rediscovery of classical fractional calculus concepts. The author has since learned this field has extensive existing literature (Riemann-Liouville, Caputo, etc.) and presents this work as a pedagogical exercise in mathematical exploration rather than novel research.

Part I

Algebraic foundation

1 The Generalized Operator for $f(x) = x^n$

1.1 From Integer to Fractional Order

We begin with the integer derivatives of $f(x) = x^n$:

$$D^k f(x) = n(n-1) \cdots (n-k+1) x^{n-k}$$

Using the identity $n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}$, we write:

$$D^k f(x) = \frac{n!}{(n-k)!} x^{n-k}$$

To generalize this for $k \in \mathbb{R}$, we substitute the factorial function with the continuous Gamma function, $\Gamma(z)$. We use the identities $n! = \Gamma(n+1)$ and

$\Gamma(z+1) = z\Gamma(z)$. The α -th derivative (where $\alpha \in \mathbb{R}$) is:

$$D^\alpha f(x) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha}$$

of course we can use this formula to get half-derivative of x^n

$$D^{\frac{1}{2}} = \frac{\Gamma(1+1)}{\Gamma(1-\frac{1}{2}+1)} x^{1-\frac{1}{2}} = \frac{1}{\Gamma(\frac{3}{2})} x^{\frac{1}{2}}$$

using the rule $\Gamma(n+1) = n\Gamma(n)$ and $\Gamma(\frac{1}{2}) = \sqrt{\pi}$

$$D^{\frac{1}{2}} = \frac{1}{\frac{1}{2}\Gamma(\frac{1}{2})} x^{\frac{1}{2}} = \frac{1}{\frac{\sqrt{\pi}}{2}} x^{\frac{1}{2}} = \frac{2}{\sqrt{\pi}} x^{\frac{1}{2}}$$

now if we take the half-derivative of that half-derivative it will be

$$\frac{2}{\sqrt{\pi}} D^{\frac{1}{2}} = \left(\frac{2}{\sqrt{\pi}}\right) \frac{\Gamma(1/2+1)}{\Gamma(\frac{1}{2}-\frac{1}{2}+1)} x^{\frac{1}{2}-\frac{1}{2}} = \left(\frac{2}{\sqrt{\pi}}\right) \frac{\Gamma(\frac{3}{2})}{1} x^0 = \left(\frac{2}{\sqrt{\pi}}\right) \left(\frac{\sqrt{\pi}}{2}\right) = 1$$

we talking the half-derivative twice to the same function gave what a one full derivative would give

of course such a proof is too simple and don't quite give the meaning of a proof, it's just a little confirmation for the time being, the full rigorous proof will be proven later with the properties of the D^z operator

Generalization to Complex Order $z = a + bi$

We now extend the derivative order to the complex number $z = a + bi$:

$$D^z f(x) = \frac{\Gamma(n+1)}{\Gamma(n-z+1)} x^{n-z}$$

To show the magnitude and phase components, we expand x^{n-z} using the property $x^{a+bi} = x^a e^{b \ln(x)i}$:

$$D^z f(x) = \frac{\Gamma(n+1)}{\Gamma(n-z+1)} x^{n-a} e^{-b \ln(x)i}$$

with these two formulas we can use them to find any \mathbb{C} or \mathbb{R} derivatives for x^n

Finding negative order derivatives

we can find the negative derivatives by putting -1 as the α and see what could happen

Putting -1 in the general formula gives the result

$$D^{-1} f(x) = \frac{\Gamma(n+1)}{\Gamma(n-(-1)+1)} x^{n-(-1)} = \frac{\Gamma(n+1)}{\Gamma(n+2)} x^{n+1}$$

and using the $\Gamma(z+1) = z\Gamma(z)$ we can say that

$$D^{-1} f(x) = \frac{\Gamma(n+1)}{(n+1)\Gamma(n+1)} x^{n+1} = \frac{x^{n+1}}{(n+1)}$$

which means that the negative order derivatives are the integrals a function
This result unifies the familiar integer derivative, the fractional derivative, and
the complex-order derivative into a single, elegant framework.

1.2 the x^{-n} problem

as we have seen, we can apply the past formula to any power of n weather it's
fractional or even complex but problems rise when we try to apply the past
formula to x^{-n}

$$D^\alpha(x^{-n}) = \frac{\Gamma(0)}{\Gamma(-\alpha)} x^{-n-\alpha}$$

not only do we have a **Gamma Pole** in the numerator but also for any value
 $\alpha \in \mathbb{Z}^+$ we also get a Gamma pole in the denominator which means that this
formula cant work and we need another formula
the m-th formula for x^{-n} is simply

$$\frac{d^m}{dx^m}(x^{-n}) = \frac{(-1)^m (n)^{(m)}}{x^{n+m}}$$

the $n^{(m)}$ here isn't a power but rather a rising factorial that can be expressed
as $n^{(m)} = \frac{(n+m-1)!}{(n-1)!}$ with this knowledge we can say

$$D^\alpha(x^{-n}) = \frac{(-1)^\alpha \frac{\Gamma(n+\alpha)}{\Gamma(n)}}{x^{n+\alpha}} = \frac{(-1)^\alpha \Gamma(n)}{x^{n+\alpha} \Gamma(n+\alpha)}$$

as simple as that, it didn't work with the original x^n formula , but this shows
something about fractional derivatives , if we went to find the first integral for
both x^n and x^{-n} we find that

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad \int x^{-n} dx = \ln|x| + C$$

Ignoring the integration constant we can find that x^n wasn't the original function
for x^{-n} but it actually transformed from $\ln(x)$

not only in complexity only but in dependency , the $D^z(x^n)$ is dependent on
the change of n more than x, while $\ln(x)$ is dependent on the change of x
this will be explained later in the **Transformative Functions** section , and
this is one of the few cases we use stanederd integration (other than the Gamma
function) in this research

2 Formulas for Other Algebraic Functions

2.1 The General Formula for a^x

Starting with the general integer rule for a^x :

$$D^n(x) = a^x \ln(a)^n$$

substituting α in the place of n gives us

$$D^\alpha f(x) = a^x \ln(a)^\alpha$$

we can see that this simple change was enough for the formula to work by taking the half-derivative twice and it gives us order one derivative

$$D^{1/2} f(x) = a^x \ln(a)^{1/2}$$

since $\ln(a)^{1/2}$ is a constant we can take it out simply when doing the derivative again

$$D^{1/2}(D^{1/2} f(x)) = \ln(a)^{1/2}(D^{1/2} f(x)) = \ln(a)^{1/2}(a^x \ln(a)^{1/2}) = a^x \ln(a)$$

which is true since our starting function was a^x and thus we can say this formula works

also from the same way we can find that $D^\alpha a^{-x} = (-1)^\alpha a^{-x} \ln(a)^\alpha$ which will be helpful for some serieses later

The Complex Generalization of this formula can be written like the D^α formula or like this

$$D^z f(x) = a^x \ln(a)^t e^{b \ln(a)i}$$

where $z = a + bi$

of course we can find the first Anti-derivative of this function by using -1 in the formula

$$D^{-1}(x) = a^x \ln(a)^{-1} = \frac{a^x}{\ln(a)}$$

and the first Complex derivative

$$D^i(x) = a^x \ln(a)^i = a^x e^{i \ln(a)}$$

2.2 The General Formula for e^x

The function e^x is known for it's "Unchanging Derivative" because it comes from the $D^n(a^x) = a^x \ln(a)^n$ and putting $a = e$ we get $D^n(e^x) = e^x$ so this also means there is no change affect the complex nor the fractional derivatives

$$D^\alpha f(x) = e^x \quad D^z f(x) = e^x$$

which means the Anti-derivative and the first complex derivative of the function

$$D^{-1}(x) = e^x \quad D^i(x) = e^x$$

2.3 The General Formula for e^{ax}

as we saw there isnot any change between e^x and any of it's derivatives , things change when we consider e^{ax} as we can see the rule of the first - second an derivative is

$$D^1 f(x) = a e^{ax} \quad D^2 f(x) = a^2 e^{ax} \quad D^3 f(x) = a^3 e^{ax}$$

so we can find the formula for the n-th derivative as

$$D^n f(x) = a^n e^{ax}$$

and changing the n to α we get

$$D^\alpha f(x) = a^\alpha e^{ax}$$

as simple as that we still have to test it to justify

$$D^{1/2} f(x) = a^{1/2} e^{ax}$$

since $a^{1/2}$ is a constant we can say that

$$D^{1/2}(D^{1/2} f(x)) = a^{1/2}(D^{1/2} f(x)) = (a^{1/2})(a^{1/2} e^{ax}) = a e^{ax}$$

this confirms that the formula work

and substituting z instead of α we get the same formula as above taht can also be written like that

$$D^z f(x) = a^z e^{bln(a)i+ax}$$

where $z = a + bi$

the first Anti-derivative for e^{ax} is

$$D^{-1} f(x) = a^{-1} e^{ax} = \frac{e^{ax}}{a}$$

and the first complex derivative is

$$D^i f(x) = a^i e^{ax} = e^{ln(a)i+ax}$$

2.4 The problem of $\log_a(x)$

$\log_a(x)$ The first derivative of $\log_a(x)$ is $\frac{1}{x \ln(a)}$ and the second derivative is $\frac{-1}{x^2 \ln(a)}$ the third derivative is $\frac{2}{x^3 \ln(a)}$ lastly the fourth derivative is $\frac{-6}{x^4 \ln(a)}$ is we can see the pattren of the n-th derivative

$$D^n f(x) = (-1)^{n+1} \frac{(n-1)!}{x^n \ln(a)}$$

and applying the gamma identity $n! = \Gamma(n+1)$ then changing n to α and reversing the x power in the denominator we get

$$D^\alpha f(x) = (-1)^{\alpha+1} \frac{\Gamma(\alpha)}{\ln(a)} x^{-\alpha}$$

But it fails, it doesnt work with fractions nor negative integares it only works positave integers, so what went wrong? my theory is that the problem is fairly simple, the $\log_a(x)$ can't be expressed as a Maclurin series and $\log_a(x+C)$ can only be expressed to a series with the condition $|x| < C$ or else it won't diverge

GDI theory : For Every function that isnt Function is analytic at $x=0$ and doesnt converge over \mathbb{R} the derivative formula $D^\alpha f(x)$ works only on \mathbb{Z}^+ for that function , we can also call it not full real differentiable

GDI hypothesis : the function is differentiable in all it's valid input values

These can also be rewritten like this:

Let $f(x)$ be an analytic function defined by its Maclaurin series

GDI theory: If $f(x)$ has a singularity at $x = 0$, then the generalized formula for $D^\alpha f(x)$ will contain a singularity at $\alpha = n$ for $n \in \mathbb{Z}^+$, preventing the generalized formula from equaling the expected integer fractional or anti-derivative.
 – Todo: change this to look better – this will be explained in detail in later sections

3 Properties of the D^z operator

these are properties to identify the nature of it that will help later with the formula and analysis of what i can call the D^z **plane** More on that later

3.1 General power series rule

power serieses are very important tools to analytically describe a function along the Real or Complex planes what is exactly what we need

The general power series definition for a function(Taylor series) is as following :

$$f(x) = f(a) + \frac{f^{(1)}(a)}{1!}(x-a) + \frac{f^{(2)}(a)}{2!}(x-a)^2 \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$$

taking the first derivative for both sides, the first term will cancel as it's a constant , the second term is linear so it will become constant, the third term is quadruple will become linear and $2!$ will cancel the 2 of the power and so on we can write is like this

$$D^1 f(x) = f^{(1)}(a) + \frac{f^{(2)}(a)}{1!}(x-a) + \frac{f^{(3)}(a)}{2!}(x-a)^2 \dots = \sum_{n=1}^{\infty} \frac{f^{(n+1)}(a)}{n!}(x-a)^n$$

the cancellation happen also because when we take a derivative the power of $(x-a)$ gets down by 1 to the numerator and we divide $n!$ by it leading to $(n-1)!$ which will lead to infinity in the denominator leading to the whole term to be zero so we can skip the first term and start from 1

taking the third and the forth derivative give the same result up to k-th derivative

$$D^k f(x) = f^{(k)}(a) + \frac{f^{(k+1)}(a)}{1!}(x-a) + \frac{f^{(k+2)}(a)}{2!}(x-a)^2 \dots = \sum_{n=k}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$$

that is of course for $k \in \mathbb{Z}$ now to make it full fractional we will put gamma instead of n and using the x^n general formula

$$D^\alpha f(x) = f(a)^{(\alpha)} + \frac{f^{(\alpha+1)}(a)}{\Gamma(2)}(x-a) + \frac{f^{(\alpha+2)}(a)}{\Gamma(3)}(x-a)^2 \dots = \sum_{n=0}^{\infty} \frac{f^{(n+\alpha)}(a)}{\Gamma(n+1)} \left[\frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} (x-a)^n \right]$$

in the brackets we can see the general derivative for x^n canceling the Gammas out we get

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n-\alpha+1)} (x-a)^{n-\alpha}$$

for powers of $(x-a)$ lesser than α it will get to ∞ in the denominator, at least when it's a negative integer other than that it will work normal, this happens because of the gamma pole, but taking the limit it will lead to 0 but still doesn't affect the sum, indeed helping us deleting the first $n < \alpha$ terms of course for negative integer derivative this works too since it will be positive

Note: because how much terms you take in the differentiation will always come terms that are replace to them because of the x^n differentiation and the infinite sum, **even if you differentiate it infinite amount of times** but you are still deleting values we are going to use that knowledge later in the integration and differential equations in later sections

we can now let $a = 0$ to get the important series we need, the Maclurin series

$$D^\alpha f(x) = f(0) + \frac{f^{(1)}(0)}{\Gamma(2)}(x) + \frac{f^{(2)}(0)}{\Gamma(3)}(x)^2 \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{\Gamma(n-\alpha+1)} x^{n-\alpha}$$

3.2 The linearity of D^z operator

for most of the operation with functions and real life applications we need to deal with linearity for D^z operator which is what we are going to prove in this small section **We Must Prove that:**

$$D^z(c_1 f(x) + c_2 g(x)) = c_1 D^z f(x) + c_2 D^z g(x)$$

let's begin with the simple x^n and let $f(x) = x^n, g(x) = x^m$ firstly we Differentiate them separately

$$D^z(c_1 f(x)) + D^z(c_2 g(x)) = c_1 D^z(f(x)) + c_2 D^z(g(x)) = c_1 \frac{\Gamma(n+1)}{\Gamma(n-z+1)} x^{n-z} + c_2 \frac{\Gamma(m+1)}{\Gamma(m-z+1)} x^{m-z}$$

Let this be statement 1

now let's differentiate them together we get

$$D^z(c_1 f(x) + c_2 g(x)) = c_1 \frac{\Gamma(n+1)}{\Gamma(n-z+1)} x^{n-z} + c_2 \frac{\Gamma(m+1)}{\Gamma(m-z+1)} x^{m-z}$$

Let this be statement 2

since **Statement 1 = Statement 2** we can say that

$$D^z(c_1f(x) + c_2g(x)) = c_1D^zf(x) + c_2D^zg(x)$$

Q.E.D

this is very useful , but to apply it to all functions , which we can do easily with infinite series

in other words

If D^z is linear on basis functions x^n , and $f = \sum c_n x^n$, then:

$$D^z(\sum c_n x^n) = \sum c_n D^z(x^n)$$

by uniform convergence of the series.

3.3 The Index law

The most important propriety for the formulas is the index law that is

$$D^{\alpha+\beta}f(x) = D^\alpha(D^\beta f(x))$$

we must prove this holds true for every case first we need to prove it for the simplest function we have which is x^n taking the D^z of the function we get

$$D^\alpha f(x) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha}$$

now let's apply the D^β with the knowledge that the Gamma functions are constants in the first derivative

$$D^\beta(D^\alpha f(x)) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} D^\beta(x^{n-\alpha}) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} \left[\frac{\Gamma(n-\alpha+1)}{\Gamma(n-\alpha-\beta+1)} x^{n-\alpha-\beta} \right]$$

the Gamma terms cancel out and we get

$$D^\beta(D^\alpha f(x)) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha-\beta+1)} x^{n-\alpha-\beta}$$

Let this be statement 1

now if we start from the beginning again but this time directly substitute $\alpha + \beta$ as O (stands for orders) we get

$$D^O f(x) = \frac{\Gamma(n+1)}{\Gamma(n-O+1)} x^{n-O}$$

if we substitute $O = \alpha + \beta$ back we get

$$D^{\alpha+\beta}f(x) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha-\beta+1)} x^{n-\alpha-\beta}$$

Let this be statement 2

if we Equal **statement 1** and **statement 2** we get

$$D^{\alpha+\beta}f(x) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha-\beta+1)}x^{n-\alpha-\beta} = D^{\beta}(D^{\alpha}f(x))$$

thus we can say that

$$D^{\alpha+\beta}f(x) = D^{\beta}(D^{\alpha}f(x))$$

Q.E.D

Note: this also works for imaginary numbers $z + w$

this by itself is a simple elegant proof but , it only works for x^n and applying the same method for each function will be very large waste of time

Instead we can use **Power Series** as they hold for every analytic function thus one proof will work for every function

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n+1)}(x-a)^n$$

we can immediately see that it's the simple x^n with everything else being a constant to the derivative

Since we've proven the Index Law for x^n , and the power series represents f as a sum of such terms, the Index Law extends to f by linearity of the operator and see that

$$D^{\alpha+\beta}f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n+1)} \left[\frac{\Gamma(n+1)}{\Gamma(n-\alpha-\beta+1)}(x-a)^{n-\alpha-\beta} \right] = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n-\alpha-\beta+1)}(x-a)^{n-\alpha-\beta}$$

which means that the Index law works for any analytic function **Q.E.D**

3.4 The Multiplication Law

A logical next step is to prove that there exists a law for multiplying orders of derivatives as it will help us later to prove and analysis a lot of topics in this research

let's take for example $f(x) = x^6$ we need (for now) to prove that multiplying two order derivatives for example 2 and 3 return the same result

for now we may write the multiplication as $M[D^{\alpha}f(x)]^{\beta}$ just as a placeholder for now

so **we need to prove that**

$$M[D^2f(x)]^3 = M[D^3f(x)]^2 = D^6f(x)$$

evaluating the first expression gives us $D^6f(x) = \frac{\Gamma(7)}{\Gamma(6-6+1)}x^{6-6} = \Gamma(7)$ a very logical step to do is to use the **Index Law** we just proved so we start with

$$D^2f(x) = \frac{\Gamma(7)}{\Gamma(6-2+1)}x^{6-2} = \frac{\Gamma(7)}{\Gamma(5)}x^4$$

$$\begin{aligned}
D^2(D^2 f(x)) &= \frac{\Gamma(7)}{\Gamma(5)} D^2 f(x) = \frac{\Gamma(7)}{\Gamma(5)} \times \frac{\Gamma(5)}{\Gamma(4-2+1)} x^{4-2} = \frac{\Gamma(7)}{\Gamma(5)} \times \frac{\Gamma(5)}{\Gamma(3)} x^2 \\
D^2(D^2(D^2 f(x))) &= \frac{\Gamma(7)}{\Gamma(5)} \times \frac{\Gamma(5)}{\Gamma(3)} D^2 f(x) = \\
&= \frac{\Gamma(7)}{\Gamma(5)} \times \frac{\Gamma(5)}{\Gamma(3)} \frac{\Gamma(3)}{\Gamma(2-2+1)} x^{2-2} = \frac{\Gamma(7)}{\Gamma(5)} \times \frac{\Gamma(5)}{\Gamma(3)} \times \frac{\Gamma(3)}{\Gamma(1)}
\end{aligned}$$

as we can see the orders cancel out perfectly leaving only $\Gamma(7)$

Let's generalize the idea with α and β following the same index law but with $f(x) = x^n$

$$\begin{aligned}
D^\alpha f(x) &= \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha} \\
D^\alpha(D^\alpha f(x)) &= \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} D^\alpha f(x) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} \frac{\Gamma(n-\alpha+1)}{\Gamma(n-\alpha-\alpha+1)} x^{n-\alpha-\alpha} \\
&= \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} \frac{\Gamma(n-\alpha+1)}{\Gamma(n-2\alpha+1)} x^{n-2\alpha} = \frac{\Gamma(n+1)}{\Gamma(n-2\alpha+1)} x^{n-2\alpha}
\end{aligned}$$

of course we can do it again

$$\begin{aligned}
D^\alpha(D^\alpha(D^\alpha f(x))) &= \frac{\Gamma(n+1)}{\Gamma(n-2\alpha+1)} D^\alpha(D^\alpha f(x)) = \frac{\Gamma(n+1)}{\Gamma(n-2\alpha+1)} \frac{\Gamma(n-2\alpha+1)}{\Gamma(n-2\alpha-\alpha+1)} x^{n-2\alpha-\alpha} \\
&= \frac{\Gamma(n+1)}{\Gamma(n-2\alpha+1)} \frac{\Gamma(n-2\alpha+1)}{\Gamma(n-3\alpha+1)} x^{n-3\alpha} = \frac{\Gamma(n+1)}{\Gamma(n-3\alpha+1)} x^{n-3\alpha}
\end{aligned}$$

we can continue like this up till the β -th multiplication and get the same result , so from this we can define

$$M[D^\alpha f(x)]^\beta = \frac{\Gamma(n+1)}{\Gamma(n-\alpha\beta+1)} x^{n-\alpha\beta}$$

and since $\alpha\beta = \gamma$ where γ is the original intended order , then

$$M[D^\alpha]^\beta = D^\gamma$$

Q.E.D

of course this works only when $\beta \in \mathbb{N}$ since it's the set we can apply Multiplication Law in terms of Index Law

I believe to ascend it to general multiplication we may first define it in a better mathematical language as

Definition : For $\beta \in \mathbb{N}$, define ${}^\beta D^\alpha$ inductively

$$1- {}^1 D^\alpha = D^\alpha$$

$$2- {}^{n+1} D^\alpha = D^\alpha \circ {}^n D^\alpha$$

By induction, we proved: ${}^n D^\alpha = D^{n\alpha}$ for $n \in \mathbb{N}$

And to extend this to \mathbb{R}/\mathbb{C} for general β , we define: ${}^\beta D^\alpha := D^{\alpha\beta}$

Then we need to verify this satisfies expected properties: ${}^{\beta_1}({}^{\beta_2} D^\alpha) = D^{\beta_1(\beta_2\alpha)} =$

$$D^{(\beta_1\beta_2)\alpha} = \beta_1\beta_2 D^\alpha$$

$$^1D^\alpha = D^\alpha$$

$$^0D^\alpha = D^0 \text{ identity}$$

This makes it a definition extended by continuity/analyticity.

Which also means that

$$M[D^\alpha]^\beta = M[D^\beta]^\alpha$$

of course that is when the operation itself is in a commutative ring like (\mathbb{R}, \mathbb{C}) ,
for other rings that isn't commutative this is False like \mathbb{H}

We adopt the notation $^\beta D^\alpha$ to denote the multiplicative action of order β on derivative D^α , distinguishing it from composition $D^\beta \circ D^\alpha$ (which gives $D^{\alpha+\beta}$ by Index Law) and the direct derivative $D^{\alpha\beta}$ but i recommend the first one as it shows we are going from α to β which may be used later without ruining the first shape of D^z operator

since the Index Law works well for Taylor series this will also work well

3.5 Operator inverses and identities

The inverse addition order:

from the Index law we can assume that there exists an equation in which

$$D^\gamma D^\alpha(f(x)) = f(x)$$

writing $f(x)$ with the geneal D^z operator we get

$$D^\gamma D^\alpha(f(x)) = D^0(f(x))$$

we can use the opposite of the operator in the other side which gives us

$$D^\alpha(f(x)) = D^{-\gamma} D^0(f(x)) = D^{-\gamma} f(x)$$

this can hold true if and only if $\gamma = -\alpha$ substituting this back we get

$$D^\alpha(f(x)) = D^{-(-\alpha)} f(x)$$

which is true thus $D^{-\alpha} D^\alpha f(x) = f(x)$

You can see this as a **generalization of the fundamental theorem of calculus** as it hold true for any set of numbers not only integers

The inverse Multiplication order:

from the multiplication law we can assume that there exist an equation where

$$^\gamma D^\alpha f(x) = D^1 f(x)$$

using the inverse of the multiplication rule we get

$$D^\alpha f(x) = {}^{\frac{1}{\gamma}} D^1 f(x)$$

which with the multiplication law we get

$$D^\alpha f(x) = D^{\frac{1}{\gamma}} f(x)$$

this only happens if and only if $\gamma = \frac{1}{\alpha}$ substituting this back we get

$$D^\alpha f(x) = D^{\frac{1}{\alpha}} f(x) = D^\alpha f(x)$$

which is true thus

$$D^{\frac{1}{\alpha}} D^\alpha f(x) = D^1 f(x) = D^\alpha f(x)$$

inverse Multiplication order theorem: For any analytic function that can be expressed as $\sum_{n=0}^{\infty} a_n x^n$, for the order derivative α there exists an order β such that ${}^\beta D^\alpha f(x) = D^1 f(x)$ and $\beta = \frac{1}{\alpha}$

of course there are some things that we can state :

The derivative order addition identity: D^0

The derivative order multiplication identity: D^1

The zeroth-derivative order multiplication propertie: ${}^0 D^\alpha$ and ${}^\beta D^0$ or in other words, the function itself is the zero of derivative order multiplication, going to or going from

Every derivative order has it's addition inverse and multiplication inverse that lead to the function itself and the first order derivative function

there are infinity many ways to get to a function from it's derivatives and integrals

Distribution of orders and Order of operations:

since the whole operation is linear we can distribute operations on the order, but to prove this we need to prove that ${}^\beta D^{(\alpha+\gamma)} = {}^\beta D^\alpha \circ {}^\beta D^\gamma$ on the left hand side we can let $\sigma = \alpha + \gamma$ so we go:

$${}^\beta D^\sigma = D^{\beta\sigma} = D^{\beta(\alpha+\gamma)} = D^{\beta\alpha+\beta\gamma}$$

with the right hand side we can go

$${}^\beta D^\alpha \circ {}^\beta D^\gamma = D^{\beta\alpha} \circ D^{\beta\gamma} = D^{\beta\alpha+\beta\gamma}$$

because both the right hand side and the left hand side are equal we can say that there is order for operations as multiplication comes before addition and there is distribution between the order operations let's take for example ${}^\beta D^\alpha \circ D^\gamma$, using what we know to write this as one derivative order we can write

$$D^{\beta\alpha} D^\gamma = D^{\beta\alpha+\gamma}$$

multiplication then addition

Derivative Order constant

we can from the integer derivative assume in function x^n that there exist a derivative order with constant, and that will also happen when $\alpha = n$ because then we have x^0 , so applying this we get

$$D^\alpha x^n = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha} = \frac{\Gamma(n+1)}{\Gamma(1)} x^0 = \Gamma(n+1)$$

Order conjugate

Order exponentiation

Order exponentiation(Logarithms of order)

Order Limits

4 Deriving the Rules of the Fractional and complex derivatives

most of the other functions need us to derive the rules of the fractional Differ-integar operator, the α -th derivative for functions like $\tan(x), \operatorname{arcsech}(x), \ln(x)$ etc.. can only be found using power serieses and product rule

4.1 General product rule

One of the most important and needed formulas in calculus in the product rule let $f(x) = g(x)h(x)$

$$f'(x) = g(x)h'(x) + g'(x)h(x)$$

$$f''(x) = g(x)h''(x) + 2 g'(x)h'(x) + g''(x)h(x)$$

$$f'''(x) = g(x)h'''(x) + 3 g'(x)h''(x) + 3 g''(x)h'(x) + g'''(x)h(x)$$

this is very similar to the binomial theorem , the only difference is it deals with derivatives instead of powers

the general product rule also known as the **General Leibniz rule** is

$$D^n(fg) = \sum_{k=0}^n \binom{n}{k} D^{n-k}(f) D^k(g)$$

of course whatever the formula for fractional derivatives turns out, this is the Integer Value Product Rule

this simple yet elegant formula is what we are going to use for the General product rule, before we try to do a simple substitution of α we need to use the generalized nC_k which means using $n! = \Gamma(n+1)$ in the formula $\frac{n!}{(n-k)!k!}$

$$D^n(fg) = \sum_{k=0}^n \frac{\Gamma(n+1)}{\Gamma(n-k+1)\Gamma(k+1)} D^{n-k}(f) D^k(g)$$

this is the same formula just and works the same for positive integers, but let's try to use fractions and ignore the sigma upper term and expand it , for example one half expansion will be

$$D^{1/2}(fg) = \frac{\Gamma(\frac{1}{2})}{\Gamma(1\frac{1}{2} - 0)\Gamma(1)} D^{1/2}(f)D^0(g) + \frac{\Gamma(\frac{1}{2})}{\Gamma(1\frac{1}{2} - 1)\Gamma(2)} D^{-1/2}(f)D^1(g) + \dots$$

$$D^{1/2}(fg) = \frac{\Gamma(\frac{1}{2})}{\Gamma(1\frac{1}{2})} D^{1/2}(f)g + D^{-1/2}(f)D^1(g) + \frac{\Gamma(\frac{1}{2})}{\Gamma(-\frac{1}{2})\Gamma(3)} D^{-3/2}(f)D^2(g) + \dots$$

as we can see , it expands to an infinite sum, it will never stop because the lower value never hit the upper value, the main reason the simple form works for integers is that even if the k value goes higher than the n value it will get a negative integer in a Gamma which is a pole and thus equal zero that means in order for a term to be it must not have the Gamma of integers so that means we can't find a non-positive integer product rule yet, but at anyway turning back the sum from what we know will be

$$D^\alpha(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha-k}(f)D^k(g)$$

we can mark this as the general power rule but we need to make our proof

we can do a simple proof by intuition

first we do it for 1:

$$D^1(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(2)}{\Gamma(-k + 2)\Gamma(k + 1)} D^{1-k}(f)D^k(g) = \frac{1}{1} D^1 f(x)g(x) + \frac{1}{1} f(x)D^1 g(x)$$

then we have the formula for α -th derivative

$$D^\alpha(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha-k}(f)D^k(g)$$

then we make two different statements

In the first we take the α -th derivative and derivative it again

$$D^1(D^\alpha(fg)) = D^1\left(\sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha-k}(f)D^k(g)\right) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^1(D^{\alpha-k}(f)D^k(g))$$

we can do this move since only the functions have x in them, like doing it to $\sum_n^\infty a_n x^n$, with the index law we can say that this equal to $D^{\alpha+1}$

$$D^{\alpha+1}(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} \left[D^{\alpha-k+1}(f)D^k(g) + D^{\alpha-k}f(x)D^{k+1}g(x) \right]$$

and we can split the sums

$$D^{\alpha+1}(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha-k+1}(f)D^k(g) + \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha-k}f(x)D^{k+1}g(x)$$

then in the second sum we can replace k with $k - 1$, which makes the sum start from 1

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - (k - 1) + 1)\Gamma((k - 1) + 1)} D^{\alpha - (k - 1)} f(x) D^{(k - 1) + 1} g(x) \\ = \sum_{k=1}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 2)\Gamma(k)} D^{\alpha + 1 - k} f(x) D^k g(x) \end{aligned}$$

but since the zero-th term vanishes because of $\Gamma(0)$ in the denominator, we can start from there, so the first statement change to

$$D^{\alpha + 1}(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} D^{\alpha - k + 1}(f) D^k(g) + \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 2)\Gamma(k)} D^{\alpha - k + 1} f(x) D^k g(x)$$

factoring both sums out we get

$$\begin{aligned} D^{\alpha + 1}(fg) &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)} + \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 2)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \\ &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha - k + 2)\Gamma(k)\Gamma(\alpha + 1) + \Gamma(\alpha - k + 1)\Gamma(k + 1)\Gamma(\alpha + 1)}{\Gamma(\alpha - k + 1)\Gamma(k + 1)\Gamma(\alpha - k + 2)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \end{aligned}$$

let's try to simplify things using the gamma identity $\Gamma(x + 1) = x\Gamma(x)$

$$\begin{aligned} &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)((\alpha - k + 1)\Gamma(\alpha - k + 1)\Gamma(k) + \Gamma(\alpha - k + 1)k\Gamma(k))}{\Gamma(\alpha - k + 1)k\Gamma(k)(\alpha - k + 1)\Gamma(\alpha - k + 1)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \\ &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)((\alpha - k + 1)\cancel{\Gamma(\alpha - k + 1)}\Gamma(k) + \cancel{\Gamma(\alpha - k + 1)}k\Gamma(k))}{\cancel{\Gamma(\alpha - k + 1)}k\Gamma(k)(\alpha - k + 1)\Gamma(\alpha - k + 1)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \\ &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)((\alpha - k + 1) + k)}{k(\alpha - k + 1)\Gamma(\alpha - k + 1)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \\ &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)(\alpha + 1)}{(\alpha - k + 1)\Gamma(\alpha - k + 1)k\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \end{aligned}$$

using the Gamma identity again but backwards gives us

$$\begin{aligned} &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 1)((\alpha - k + 1) + k)}{k(\alpha - k + 1)\Gamma(\alpha - k + 1)\Gamma(k)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \\ &= \sum_{k=0}^{\infty} \left[\frac{\Gamma(\alpha + 2)}{\Gamma(\alpha - k + 2)\Gamma(k + 1)} \right] D^{\alpha - k + 1} f(x) D^k g(x) \end{aligned}$$

We can let this be **statement 1**
then we do the $D^{\alpha+1}$ derivative from the beginning

$$D^{\alpha+1}(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha+2)}{\Gamma(\alpha-k+2)\Gamma(k+1)} D^{\alpha+1-k}(f) D^k(g)$$

we can let this be **statement 2**
since we can see that **statement 1** = **statement 2**, thus the formula works
Q.E.D

of course there will exist some functions that doesn't have closed form other than power series so we can get a simple series version of the α -th derivative
we can use the original formula but we replace every function with it's series

$$D^{\alpha}(fg) = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha+1)}{\Gamma(\alpha-k+1)\Gamma(k+1)} \sum_{j=0}^{\infty} \frac{f^{(j)}(0)}{\Gamma(j-\alpha+k+1)} x^{(j-\alpha+k)} \sum_{i=0}^{\infty} \frac{g^{(i)}(0)}{\Gamma(i-\alpha+k+1)} x^{(i-\alpha+k)}$$

4.2 General chain rule

The General Chain Rule is hard, we can try any of our normal methods, yet they fail
the easier way is to go with the series option we can get the Maclaurin series with ease

4.2.1 General $f(x^n)$ formula

first we define our functions for simpler solving

$$u(x) = x^n \quad g(x) = f(u)$$

then we differentiate for the first three derivatives for $u(x)$

$$u'(x) = nx^{n-1} \quad u''(x) = n(n-1)x^{n-2} \quad u'''(x) = n(n-1)(n-2)x^{n-3}$$

then we differentiate $g(x) = f(u)$ with respect to u

$$g'(x) = u'f'(u) \quad g''(x) = u''f'(u) + (u')^2f''(u)$$

$$g'''(x) = u'''f'(u) + u''u'f''(u) + 2u'u''f''(u) + (u')^3f'''(u)$$

then we evaluate with the Maclaurin series

$$g(x) = f(0) + u'(0)f'(0)x + \frac{u''(0)f'(0) + (u'(0))^2f''(0)}{2!} + \dots$$

using the chain rule, for the x_1 term we have $nx^{n-1}f'(x^n)$
evaluating at 0 gives us 0 for any term that isn't 1, evaluating at one gives us

$0^0 f'(0)$, so we need to approach it with limits
if we take the limit as x approach zero so we have

$$\lim_{x \rightarrow 0} x^0 f'(x^n) = f'(0)$$

so it works fine with the first term, we can do the same for the rest terms and we get different results based on different multiples of n , if we choose k to be 2 only even terms will show up as other terms equal 0, for $n = 3$ we get multiples of 3 so on so forth, and so we get the series

$$f(x^n) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^{kn}$$

we can give it a simple check by calculating random value for a random function like $\sin(x^3)$

the series for this function will be $\sin(x^3) = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} x^{6n+3}$ we let it equal to $f(x)$ so we can see the difference between both sides

$$\sin(0.365^2) = 0.0486079633377 \quad f(0.365) = 0.0486079633377$$

we can see that it works fine, now for the α -th derivative we get a simple

$$D^\alpha f(x^n) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{\Gamma(kn - \alpha + 1)} x^{kn - \alpha}$$

thus a simple formula but something closer to the general Macluarin series chain rule

4.2.2 General $f(ax)$ formula

Although not quite the difficult formula to get yet it's the easier with a closed form and have big importance in other fields like Fourier transforms
first let $g(x) = f(ax)$, for the first few derivatives we have

$$g'(x) = af'(ax) \quad g''(x) = a^2 f''(ax) \quad g'''(x) = a^3 f'''(ax)$$

you probably can guess the rule from the pattern, but to make sure let's continue with the Macluarin series

$$g(x) = f(0) + af'(0)x + \frac{a^2 f''(0)}{2!} x^2 + \dots = \sum_{n=0}^{\infty} \frac{a^n f^{(n)}(0)}{n!} x^n$$

and from this we can say that $D^n f(ax) = a^n f^{(n)}(ax)$ applying α instead of n will give us

$$D^\alpha f(ax) = a^\alpha f^{(\alpha)}(ax)$$

simple yet needed

4.2.3 General Series form for chain rule

first we find the first derivatives with $h(x) = f(g(x))$
 $h'(x) = f'(g(x))g'(x)$, $h''(x) = f'(g(x))(g'(x))^2 + g''(x)f'(g(x))$ then we do a simple series expansion

$$h(x) = h(0) + h'(0)x + \frac{h''(0)x^2}{2!} + \dots = \sum_{n=0}^{\infty} \frac{h^{(n)}(0)}{n!} x^n$$

then we substitute the derivatives

$$\begin{aligned} f(g(x)) &= f(g(0)) + f'(g(0))g'(0)x + \frac{(f'(g(0))(g'(0))^2 + g''(0)f'(g(0)))x^2}{2!} + \dots \\ &= \sum_{n=0}^{\infty} \frac{f^{(n)}(g(0))}{n!} (g(x) - g(0))^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(g(0))}{n!} \left(\sum_{k=0}^{\infty} \frac{g^{(k)}(0)}{k!} x^k - g(0) \right)^n \end{aligned}$$

4.2.4 General Closed form for chain rule

the general formula for integer chain rule is called Faà di Bruno's formula and it stats

$$\frac{d^n}{dx^n} f(g(x)) = \sum_{k=0}^n f^{(k)}(g(x)) \times {}^n B_k(g'(x), g''(x), \dots, g^{(n-k+1)}(x))$$

and as we can see there are bunch of problems here ,first if we tried to plug any fraction in place of the n we get an infinte series

5 Interpretation of fractional and complex derivative in other functions

5.1 Trigonometric Functions

Deriving the trigonometric functions can be quite tricky , as there exists n-th derivative formula for them but don't seem to work as intended

5.1.1 $\sin(x)$ and $\cos(x)$

for $\sin(x)$ there exists a formula which is

$$D^n \sin(x) = \sin\left(\frac{n\pi}{2} + x\right)$$

we need to first expand it to it's exponentiation formula

$$\sin(x) = \frac{e^{ix} - e^{-ix}}{2i}$$

then we here can differentiate with ease using our $D^\alpha e^{ax} = a^\alpha e^{ax}$ formula

$$D^\alpha \sin(x) = \frac{i^\alpha e^{ix} - (-i)^\alpha e^{-ix}}{2i}$$

to make it look simpler we can use $i = e^{\frac{i\pi}{2}}$ and $i = e^{\frac{i\pi}{2} + i\pi} = e^{\frac{-i\pi}{2}}$

$$\begin{aligned} D^\alpha \sin(x) &= \frac{(e^{\frac{i\pi}{2}})^\alpha e^{ix} - (e^{\frac{-i\pi}{2}})^\alpha e^{-ix}}{2i} = \frac{(e^{\frac{i\pi\alpha}{2}})e^{ix} - (e^{\frac{-i\pi\alpha}{2}})e^{-ix}}{2i} \\ &= \frac{e^{\frac{i\pi\alpha}{2} + ix} - e^{\frac{-i\pi\alpha}{2} - ix}}{2i} = \frac{e^{i(\frac{\pi\alpha}{2} + x)} - e^{-i(\frac{\pi\alpha}{2} + x)}}{2i} = \sin\left(\frac{\alpha\pi}{2} + x\right) \end{aligned}$$

and use the Index Law work here too so $D^\alpha(D^\beta \sin(x)) = \sin(\frac{(n+m)\pi}{2} + x)$ now we can test it for half derivative

$$D^{\frac{1}{2}} \sin(x) = \sin\left(\frac{\frac{1}{2}\pi}{2} + x\right) = \sin\left(\frac{\pi}{4} + x\right)$$

$$D^{\frac{1}{2}}(D^{\frac{1}{2}} \sin(x)) = \sin\left(\frac{(\frac{1}{2} + \frac{1}{2})\pi}{2} + x\right) = \sin\left(\frac{2\pi}{4} + x\right) = \sin\left(\frac{\pi}{2} + x\right)$$

the same also works for $\cos(x)$

$$D^\alpha \cos(x) = \frac{e^{i(\frac{\pi\alpha}{2} + x)} + e^{-i(\frac{\pi\alpha}{2} + x)}}{2i} = \cos\left(\frac{\alpha\pi}{2} + x\right)$$

but there is another proof that this works for all real numbers.

from the euler formula $e^{ix} = \cos(x) + i \sin(x)$ we can say that

$$\sin(x) = \text{Im}(e^{ix})$$

taking the alpha-th derivative of both sides

$$D^\alpha \sin(x) = D^\alpha \text{Im}(e^{ix}) = \text{Im}(i^\alpha e^{ix})$$

knowing that $i = e^{i\pi/2}$

$$D^\alpha \sin(x) = \text{Im}(e^{i\pi\alpha/2} e^{ix}) = \text{Im}(e^{i\pi\alpha/2 + ix}) = \text{Im}(e^{i(\alpha\pi/2 + x)})$$

turning this back to the euler formula will give us

$$D^\alpha \text{Im}(e^{ix}) = \sin\left(\frac{\alpha\pi}{2} + x\right)$$

which indeed proves it's true from the same formula we can also get the α -th for $\cos(x)$ with the same formula turning this back to the euler formula will give us

$$D^\alpha \text{Re}(e^{ix}) = \cos\left(\frac{\alpha\pi}{2} + x\right)$$

now we can write them as

$$D^\alpha \sin(x) = \text{Im}(e^{i(\alpha\pi/2+x)}) \quad D^\alpha \cos(x) = \text{Re}(e^{i(\alpha\pi/2+x)})$$

or

$$D^\alpha \sin(x) = \sin\left(\frac{\alpha\pi}{2} + x\right) \quad D^\alpha \cos(x) = \cos\left(\frac{\alpha\pi}{2} + x\right)$$

and to make it to the complex plane we can use these formulas also the negative derivative of these is

$$D^{-1} \sin(x) = \sin\left(\frac{-\pi}{2} + x\right) = -\cos(x) \quad D^{-1} \cos(x) = \cos\left(\frac{-\pi}{2} + x\right) = \sin(x)$$

and the first complex derivative of these is

$$D^i \sin(x) = \sin\left(\frac{i\pi}{2} + x\right) = \sin\left(\frac{\ln(-1)}{2} + x\right) \quad D^i \cos(x) = \cos\left(\frac{i\pi}{2} + x\right) = \cos\left(\frac{\ln(-1)}{2} + x\right)$$

5.1.2 $\tan(x)$ and $\sec(x)$

finding the alpha-th derivative for $\tan(x)$ is quite hard since we didn't get any direct formulas for quotients, and there is no direct integer derivative formula we can plug in and generalize to the Real numbers, we can try to change it a little with some algebra

$$\tan(x) = \sin(x) (\cos(x))^{-1}$$

and then use the general product rule, but quickly we can see the problem

$$D^\alpha (\sin(x) (\cos(x))^{-1}) = \sum_{k=0}^{\infty} \frac{\Gamma(0)}{\Gamma(\alpha - k + 1) \Gamma(k + 1)} D^{\alpha-k}(\sin(x)) D^k(\cos(x)^{-1})$$

there is a gamma pole so this solution also fails
we can try using some trig substitution

$$\tan(x) = \sin(x) (\sqrt{\sin(x)^2 + 1})^{-1}$$

but this leads to infinite sum for the product rule and the chain rule that we have proved above it impossible to find with the simple algebra we have is very hard to approximate by itself as we can see there is no simple elegant closed form for $\tan(x)$ in the scientific paper, the reason behind this will be explained later but we can simply say because it has poles, to analytically see it better we need the Maclurin series expansion

$$\tan(x) = x + \frac{x^3}{3!} + \frac{2x^5}{15} + \dots = \sum_{n=0}^{\infty} \frac{B_{2n}(-4)^n(1-4^n)}{(2n)!} x^{2n-1} \quad \text{where } |x| < \frac{\pi}{2}$$

if we look closely we can notice the problem, it has a radius of converges and that by itself is the problem that will be discussed in detail later all what we can

do for now is applying the D^α to the infinite sum as it will be the only analytic closed form way for now
we will get:

$$\begin{aligned} D^\alpha \tan(x) &= \sum_{n=0}^{\infty} \frac{B_{2n}(-4)^n(1-4^n)}{\Gamma(2n)} \left[\frac{\Gamma(2n)}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1} \right] \\ &= \sum_{n=0}^{\infty} \frac{B_{2n}(-4)^n(1-4^n)}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1} \end{aligned}$$

this infinite series shall work for now

the same also works for $\sec(x)$ as it's a quotient so if we tried using the General product rule we will hit a Gamma pole, so the safest answer for now is to go with infinite series

$$\sec(x) = 1 + \frac{x^2}{2!} + \frac{5x^4}{4!} + \dots = \sum_{n=0}^{\infty} \frac{E_{2n}(-1)^n}{(2n)!} x^{2n} \quad \text{where } |x| < \frac{\pi}{2}$$

again we see the same problem with radius of convergence
simply we apply D^α :

$$\begin{aligned} D^\alpha \sec(x) &= \sum_{n=0}^{\infty} \frac{E_{2n}(-1)^n}{\Gamma(2n)} \left[\frac{\Gamma(2n)}{\Gamma(2n-\alpha+1)} x^{2n-\alpha} \right] \\ &= \sum_{n=0}^{\infty} \frac{E_{2n}(-1)^n}{\Gamma(2n-\alpha+1)} x^{2n-\alpha} \end{aligned}$$

Note: these two work for their radius of converges only

Note: they also work for the complex derivative

5.1.3 $\csc(x)$ and $\cot(x)$

now saying $\csc(x)$ and $\cot(x)$ will work the same like the rest of trigonometric functions is a bit of a stretch

since we already know both of their domains aren't $x \in \mathbb{R}$ so they must have some sort of analytical poles and converges radius that isn't \mathbb{R} in their infinite sums

but if we noticed

$$\csc(x) = \frac{1}{\sin(x)}$$

which means that it has a singularity at $x = 0$, in other words simple Taylor series nor simple Maclurin series won't work, we need the General Laurent series for this one The Laurent series is :

$$\csc(x) = \frac{1}{x} + \frac{x}{6} + \frac{7x^3}{360} + \dots = \sum_{n=0}^{\infty} \frac{B_{2n}(-1)^{n+1}(2^{2n}-1)}{(2n)!} x^{2n-1} \quad \text{where } 0 < |x| < \pi$$

but before plugging the D^z operator to the series we can notice a little problem in the beginning, the $\frac{1}{x}$ term simply plugging in the $D^z(x^n)$ will resault a pole, the simple solution is just to the linearity of D^z and differentiate the first term alone and then the rest of the series alone

$$\begin{aligned} D^\alpha \csc(x) &= D^\alpha(x^{-1}) + \sum_{n=1}^{\infty} \frac{B_{2n}(-1)^{n+1}(2^{2n}-1)}{\Gamma(2n)} \left[\frac{\Gamma(2n)}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1} \right] \\ &= \frac{(-1)^\alpha \Gamma(1+\alpha)}{x^{1+\alpha}} + \sum_{n=1}^{\infty} \frac{B_{2n}(-1)^{n+1}(2^{2n}-1)}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1} \end{aligned}$$

of course this is where $0 < |x| < \pi$

The same goes for $\cot(x)$ as it doesn't have any Taylor series but rather Laurent series

The Laurent series for $\cot(x)$ is:

$$\cot(x) = \frac{1}{x} + \sum_{n=1}^{\infty} \frac{(-1)^n 2^{2n} B_{2n}}{(2n)!} x^{2n-1} \quad \text{where } 0 < |x| < \pi$$

Applying D^α to both sides we get

$$D^\alpha \cot(x) = \frac{(-1)^\alpha \Gamma(1+\alpha)}{x^{1+\alpha}} + \sum_{n=1}^{\infty} \frac{(-1)^n 2^{2n} B_{2n}}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1}$$

Note: this works to complex numbers too

5.2 Hyperbolic Functions

5.2.1 $\sinh(x)$, $\cosh(x)$ and $\tanh(x)$

$\sinh(x)$ is pretty straight forward to get from the definition

$$\sinh(x) = \frac{1}{2}(e^x - e^{-x})$$

differentiating both sides to the α

$$D^\alpha \sinh(x) = \frac{1}{2}(D^\alpha e^x - D^\alpha e^{-x}) = \frac{1}{2}(e^x - (-1)^\alpha e^{-x})$$

we can also do the same for $\cosh(x)$

$$D^\alpha \cosh(x) = \frac{1}{2}(D^\alpha e^x + D^\alpha e^{-x}) = \frac{1}{2}(e^x + (-1)^\alpha e^{-x})$$

but if we change the negative sign in $\sinh(x)$ to $+(-1)$ we turn the derivative to

$$D^\alpha \sinh(x) = \frac{1}{2}(e^x + (-1)^{\alpha+1} e^{-x})$$

which is equal to $D^{\alpha+1} \cosh(x)$ and that is because unlike normal $\sin(x)$ and $\cos(x)$ these are the integer integrals and derivatives of their-selves so we can get the negative derivatives to be

$$D^{-1} \sinh(x) = \frac{1}{2}(e^x + e^{-x}) = \cosh(x) \quad D^{-1} \cosh(x) = \frac{1}{2}(e^x - e^{-x}) = \sinh(x)$$

for the complex derivatives we can use the formulas from before
However for $\tanh(x)$ things change, since the definition for it is

$$\tanh(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

we can see, it's a quotient of two functions, which leaves us with nothing but to use its power series
the power series for \tanh is:

$$\tanh(x) = x - \frac{x^3}{3!} + \frac{2x^5}{15} - \dots = \sum_{n=0}^{\infty} \frac{B_{2n} 4^n (1 - 4^n)}{(2n)!} x^{2n-1} \quad \text{where } |x| < \frac{\pi}{2}$$

as predicted there will also be radius of converges here too
but at anyway we get the D^α with this:

$$\begin{aligned} D^\alpha \tanh(x) &= \sum_{n=0}^{\infty} \frac{B_{2n} 4^n (1 - 4^n)}{\Gamma(2n)} \left[\frac{\Gamma(2n)}{\Gamma(2n - \alpha + 1)} x^{2n-\alpha-1} \right] \\ &= \sum_{n=0}^{\infty} \frac{B_{2n} 4^n (1 - 4^n)}{\Gamma(2n - \alpha + 1)} x^{2n-\alpha-1} \end{aligned}$$

5.2.2 $\operatorname{sech}(x)$, $\operatorname{csch}(x)$ and $\operatorname{coth}(x)$

The rest of the hyperbolic functions shall work the same as the trigonometric functions, infact all of the hyperbolic and the trigonometric function's Laurent/Taylor series look identical with little changes, so finding them won't be that difficult

For $\operatorname{sech}(x)$ the Taylor Series is:

$$\operatorname{sech}(x) = 1 - \frac{x^2}{2!} + \frac{5x^4}{4!} - \dots = \sum_{n=0}^{\infty} \frac{E_{2n}}{(2n)!} x^{2n} \quad \text{where } |x| < \frac{\pi}{2}$$

So Applying D^z will be as simple as $\sec(x)$

$$D^\alpha \operatorname{sech}(x) = \sum_{n=0}^{\infty} \frac{E_{2n}}{\Gamma(2n - \alpha + 1)} x^{2n-\alpha} \quad \text{where } |x| < \frac{\pi}{2}$$

the same goes for the Laurent series of $\operatorname{csch}(x)$ and $\operatorname{coth}(x)$

$$\operatorname{csch}(x) = \frac{1}{x} - \frac{x}{6} + \frac{7x^3}{360} - \dots = \frac{1}{x} + \sum_{n=1}^{\infty} \frac{B_{2n} (1 - 2^{2n})}{(2n)!} x^{2n-1} \quad \text{where } 0 < |x| < \pi$$

of course it's similar but not identical , anyway applying D^α gives us

$$\begin{aligned} D^\alpha \operatorname{csch}(x) &= D^\alpha x^{-1} + D^\alpha \sum_{n=0}^{\infty} \frac{B_{2n}(1-2^{2n})}{(2n)!} x^{2n-1} \\ &= \frac{(-1)^\alpha \Gamma(1+\alpha)}{x^{1+\alpha}} + \sum_{n=1}^{\infty} \frac{B_{2n}(1-2^{2n})}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1} \quad \text{where } 0 < |x| < \pi \end{aligned}$$

and for $\operatorname{coth}(x)$:

$$\operatorname{coth}(x) = \frac{1}{x} + \sum_{n=1}^{\infty} \frac{2^{2n} B_{2n}}{(2n)!} x^{2n-1} \quad \text{where } 0 < |x| < \pi$$

and applying D^α operator we get:

$$D^\alpha \operatorname{coth}(x) = \frac{(-1)^\alpha \Gamma(1+\alpha)}{x^{1+\alpha}} + \sum_{n=1}^{\infty} \frac{2^{2n} B_{2n}}{\Gamma(2n-\alpha+1)} x^{2n-\alpha-1}$$

5.3 The Inverse Trigonometric and Hyperbolic Functions

there is a problem with these function that makes them special, if we for example tried to take the derivative for $\sin^{-1}(x)$ and the integral of the same function we get this

$$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}} \quad -1 < x < 1 \quad \int \sin^{-1}(x) dx = x \sin^{-1}(x) + \sqrt{1-x^2} + C$$

as we can see they are a type of **Transformative functions** which will be discussed in later sections

6 Matrix derivative functions

6.1 Whole matrix derivatives

we start with the simple function x^n , and let our matrix be a simple A matrix, we can right the $D^A x^n$ like this:

$$D^A(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-A+1)} x^{n-A}$$

as we can see this shape is hard , and a better simplification for it is to use $e^{\ln(x)}$ so it will be :

$$D^A(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-A+1)} x^n e^{-\ln(x)A}$$

this is better as now we can express it as infinite sum using the Taylor series of function e^x

$$D^A(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-A+1)} x^n \sum_{k=0}^{\infty} \frac{(-1)^k (\ln x)^k}{k!} A^k$$

where $A^k = \underbrace{A \times A \times A \times \dots}_{k \text{ times}}$, and this can be called the simple $D^A x^n$

let's plug in this definition in the Taylor series general formula

$$D^\alpha f(a) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n - \alpha + 1)} (x - a)^{n - \alpha}$$

so now it shall be

$$D^A f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{\Gamma(n - A + 1)} (x - a)^{n - A}$$

and thus the Maclaurin series is

$$\begin{aligned} D^A f(x) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{\Gamma(n - A + 1)} x^{n - A} = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{\Gamma(n - A + 1)} x^n e^{-\ln(x)A} \\ &= \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{\Gamma(n - A + 1)} x^n \sum_{k=0}^{\infty} \frac{(-1)^k (\ln x)^k}{k!} A^k \end{aligned}$$

now we know we can do it to any analytical function, and the simple way to put it in any series is to remove the α from the power of and put the term $\sum_{k=0}^{\infty} \frac{(\ln x)^k}{k!} A^k$

6.2 What is the meaning of matrix derivatives

to understand what even is matrix order derivative we can try and find simple matrix order for matrix $A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ and for the function to be $f(x) = x^2$ so the derivative will be

$$D^A(x^2) = \frac{\Gamma(3)}{\Gamma(3 - A)} x^2 \sum_{k=0}^{\infty} \frac{(-1)^k (\ln x)^k}{k!} A^k$$

and this one is simple, since $A = 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 2I$, we can write the A^k term as $2^k I$, now we can take it out of the summation as it's a constant multiplication forming back to its original form we get $e^{-2 \ln(x)} I$ which is $x^{-2} I$ for the gamma term we can do it also simple subtraction we get $\Gamma(3I - 2I) = \Gamma(I) = I$ so what we now we have this expression

$$D^A(x^2) = \frac{2}{I} x^2 \times x^{-2} I = 2I^2 x^{2-2} = 2I$$

so surprisingly we get the derivative order matrix

6.3 Matrix order properties

Order Matrix: if we take the derivative A of function x^n where A is the identity matrix multiplied by n will return $\Gamma(n+1)I$, in other words any scalar matrix order derivative will return $\Gamma(n+1)$ times identity matrix

Proof:

let A be a matrix order equal to
$$\begin{bmatrix} \alpha & 0 & 0 & \cdots & 0 \\ 0 & \alpha & 0 & \cdots & 0 \\ 0 & 0 & \alpha & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha \end{bmatrix}$$
 and we have function x^α

$$D^A(x^\alpha) = \frac{\Gamma(\alpha+1)}{\Gamma(\alpha-A+1)} x^\alpha \sum_{k=0}^{\infty} \frac{(-1)^k (\ln x)^k}{k!} A^k$$

we know that $A = \alpha \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix} = \alpha I$, the term in the infinite sum

will be $A^n = \alpha^n I$, then any number we can treat like scalar matrix as it works

the same, so the Gamma term we can write as as $\Gamma\left(\begin{bmatrix} \alpha & 0 & 0 & \cdots & 0 \\ 0 & \alpha & 0 & \cdots & 0 \\ 0 & 0 & \alpha & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha \end{bmatrix} - \right.$

$$\left. \begin{bmatrix} \alpha & 0 & 0 & \cdots & 0 \\ 0 & \alpha & 0 & \cdots & 0 \\ 0 & 0 & \alpha & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha \end{bmatrix} + I\right) = \Gamma(I) = I$$

putting everything together we get

$$D^A(x^\alpha) = \Gamma(\alpha+1)I^{-1}x^\alpha \sum_{k=0}^{\infty} \frac{(-1)^k (\ln x)^k}{k!} \alpha^k I^k$$

forming the infinite sum back to its original form we get

$$D^A(x^\alpha) = \Gamma(\alpha+1)Ix^\alpha e^{-\alpha \ln(x)}I = \Gamma(\alpha+1)I^2x^{\alpha-\alpha} = \Gamma(\alpha+1)I$$

Q.E.D

we can consider this as the Matrix version of the **Derivative Order constant** but a prove was needed, another simpler one is treating these scalar matrices as numbers then doing the same proof from section above **Order Addition and Multiplication** Let A, B Be Different matrices with n by n elements

Since $A + B = B + A$ Then $D^A \circ D^B = D^B \circ D^A$ Thus **Matrix order addition is commutative**, that is because the adoptive nature of D^z with it's order field/ring that we have proven in before section

And for the same reason since $AB \neq BA$ then ${}^B D^A \neq {}^A D^B$, Thus Matrix Order Multiplication isn't commutative

7 Functional derivatives

7.1 derivative order as function of itself

The Derivative order being a function is hard, Unless we are talking about function of itself instead of x or y it will be $g(\alpha)$, the formulas for x^n it shall be

$$D^{g(\alpha)} f(x) = \frac{\Gamma(n+1)}{\Gamma(n-g(\alpha)+1)} x^{n-g(\alpha)}$$

this simple formula shall be the way for us to get the Maclurin and Taylor series

$$D^{g(\alpha)} f(x) = \sum_{n=0}^{\infty} \frac{f^n(a)}{\Gamma(n-g(\alpha)+1)} (x-a)^{g(\alpha)}$$

$$D^{g(\alpha)} f(x) = \sum_{n=0}^{\infty} \frac{f^n(0)}{\Gamma(n-g(\alpha)+1)} x^{n-g(\alpha)}$$

7.2 derivative order as function of it's dependent variable

we can now do the same for function that are dependent of the function variable

$$D^{g(x)} f(x) = \frac{\Gamma(n+1)}{\Gamma(n-g(x)+1)} x^{n-g(x)}$$

same formula here ,yet things change a lot, because that is a one time jump, the next time won't go according to the normal formula since we will have x^x term if not more, so the index law nore the multiplication law work here... in fact no normal formula or law will work after the first one, which is a problem but not a big one since we already can jump one time there exists a Maclurin series for it

$$D^{g(x)} f(x) = \sum_{n=0}^{\infty} \frac{f^n(0)}{\Gamma(n-g(x)+1)} x^{n-g(x)}$$

7.3 derivative order as function of another Dimension variable (Dimension changer)

Part II

D^z Analysis

8 The nature of functions under D^z Field

8.1 Explaining what is a fractional derivative

taking an positive Integer derivative gives us the rate of change of a function , talking the derivative of that also gives us the rate of change the derivative function , so on so forth

taking the negative Integer derivative is known as the "Area under the curve" or the function in which the original function is the rate of change of it , or simply the "Anti-derivative"

if we tried to explain D^z operator with the standard definition of integrals and derivatives we can say that

if the first derivative is the rate of change of the function, the first integral is area under the curve then D^z operator is change from being an Area to the function than the rate of the function itself, but this doesn't make sense mostly because these are three different things that work and measure differently how could we explain such things?

The problem becomes bigger and bigger when we say the second derivative which is the rate of change of the rate of change of the function that gives us information about the function like when is it a climax or function convexity, or the second Integral that is the 3D volume of the shape.

then comes the Taylor expansion and and the power series of functions which use infinite derivatives and now we are all out of the meaning of it

even if we though about it physically , for simple cases like position, it's derivative is velocity and it's second derivative is acceleration which at least change the unit powers and doesnt make much sense with D^z

in fact with out standers even if multiple derivatives integer would somehow make sense, the fractional derivative wouldn't

which why is suggest another prespictive for the derivative instead of it being the area under curve or the rate off change , these things are just side effects and the derivative is it's actual object by itself

the fractional derivative can be thought as some change in the being of function itself, as we can see from the x^n deriavtive

$$D^\alpha x^n = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha}$$

as we can see ,what really happen is the function get scalled by a certain amount that is $\frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)}$, and get multiplied by the simple form of it that is $g(x) = x$ raised to the power of the derivative order but from the negative side, if we let $f(x) = x^n$ and $g(x) = x$ and the Scaler S , we can write it like this

$$D^\alpha f(x) = S f(x)g(x)^{-\alpha} = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} \frac{f(x)}{g(x)^\alpha}$$

and if we use the Euler product expansion of the $\Gamma(x)$ we can write the function in it's simplest, algebraic form with no integrals

$$D^\alpha x^n = \frac{\frac{1}{n+1} \prod_{k=1}^{\infty} \left[\frac{1}{1+\frac{n+1}{k}} \left(1 + \frac{1}{k}\right)^{n+1} \right]}{\frac{1}{n-\alpha+1} \prod_{k=1}^{\infty} \left[\frac{1}{1+\frac{n-\alpha+1}{k}} \left(1 + \frac{1}{k}\right)^{n-\alpha+1} \right]} \frac{x^n}{x^\alpha}$$

here it's the derivative in a simpler form that can show us why we can't over derivative the function, because we will get a division by zero, but from what we can see also is that the function works for non-positive integers

From the original function we can see that the more we derivative the more it gets "Discrete" and "Not Smooth enough", and we may give the reason for that is the scaler not the function power itself

because we can go with negative fractional overdrive , the problem is with the not-positive integers

8.2 what it means for a derivative order to be Complex

as we saw how the fractional derivative can say the complex derivative take a weird turn as it actually not a nor rate of change of function...sort of to get an understanding of what i am talking about let's see it on a simple function x^n and see how it works

we know that:

$$D^z(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-z+1)} x^{n-z}$$

substituting $z = i$

$$D^i(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-i+1)} x^{n-i}$$


the nature of this will be explored later but for now if we try to take another imaginary derivative using the **Multiplication Law** we proved earlier we see that

$${}^i D^i(x^n) = D^i(x^n) = \frac{\Gamma(n+1)}{\Gamma(n-(i \times i)+1)} x^{n-(i \times i)} = \frac{\Gamma(n+1)}{\Gamma(n-(-1)+1)} x^{n-(-1)} = \frac{\Gamma(n+1)}{\Gamma(n+2)} x^{n+1}$$


and knowing that $\Gamma(n+1) = n\Gamma(n)$

$${}^i D^i(x^n) = \frac{\Gamma(n+1)}{(n+1)\Gamma(n+1)} x^{n+1} = \frac{x^{n+1}}{n+1}$$

infact this was expected since ${}^iD^i = D^{(i \times i)} = D^{-1}$
the result is kind of weird , clearly it's weird
to see why it's weird (if it'snot already) we can use a geometric interpretation
from the information we have right now we can draw this transformation in D-I
plane




as this map acts as the same as complex plane a we map the transformation
around it like this like this



this


is how the first derivative act when using the complex number i
which means also that means there exist a way (if not multiple) to represent
differentiation and integration geometrically in one same space
we can add more detail to this plane by defining each access as the the order of
the D opreator so we can write it better like this



with the y-axes being the imaginary order part and the x-axis being the normal part, we can of course now express any derivative or integral or any weird result of D^z as points in this space.

that is, of course for one single function as the input is a variable of type of function and the output is of variable function so for now they must stay the same function with the same variables inside of it


so we can now express for example the $D^z(x^n)$ like this



this also allows us to use any root of unity not only i like a normal Re-Im plot plane, of course we can change the perspective and think of them as vectors in a 2D vector plane

for each derivative we can draw them as vectors starting from zero point (the

function itself) to the point point of derivative order



which means that the vector rules also work here fine with the D^z operator properties as we can see adding and subtracting orders will return another vector using the Index Law, and since we know what the plane is for we can express the orders from $D^{\alpha+\beta i}$ as points $(\alpha, \beta i)$ or vectors $\vec{a} = (\alpha, \beta)$, and supposedly polar (r, θ) , of course if we want to make a vector from a point to another we can just use the index law that goes in the path to β which also means that there is infinite starting points for vectors to go to the same point, in another words : infinite vectors for any point and to any point


this can be called the **D(i) Plane**, even when dealing with vectors , at the end this plane isnt really based on the idea of it but rather an extension to it

this may helpful , but i suggest another idea that is based on vectors in such field and I would like to call it the **D(i) Vector Field**

instead of using vectors in the plane as beginning and ending points we will use them as points, arrows and degrees, sort of a mix of the polar form and the vector form of the plane

- 1- the point: will represent it's head, where it's coming out of
- 2- the arrow : will the output of the function derivative rate of change, the bigger it's the higher the rate of change its
- 3- the degree: it will represent how the D^z operator is changing the function itself

for the sake of understanding let's take the function e^{2x} as an example



we can now study how the derivatives of e^{2x} work from this vector field
the degrees between the different derivatives first In the real values we can see
the more we integrate the closer we get to the original function , and the more
we differentiate the more the value of the coefficient grows
the degree of hte vectors represent how quick it grows as we can say it grows by
 2^α , so when we differentiate more the values of outputs of the original function
gets larger and larger for small inputs, and the opposite happens when we inte-
grate ,
we can define the Scaling of both operations like this

$$L_I[D^\alpha(e^{2x})] = e^{2x} \quad L_D[D^\alpha(e^{2x})] = \infty$$

9 Cyclic derivatives

9.1 e^x and Cyclic functions

if we look closely in functions that have **Cyclic-Derivatives** we can see that
there exists a pattren for example

$$D^\alpha e^x = e^x$$

$$D^\alpha \sinh(x) = \frac{1}{2}(e^x - (-1)^\alpha e^{-x}) \quad D^\alpha \cosh(x) = \frac{1}{2}(e^x + (-1)^\alpha e^{-x})$$

$$D^\alpha \sin(x) = \text{Im}(e^{ix}) \quad D^\alpha \cos(x) = \text{Re}(e^{ix})$$

they all have connection with e^x in them
the function e^x by itself is a cyclic derivative function , it always return itself

no matter how many times you differentiate it if we change the function slightly to e^{ax} things now change as it's returns different results

$$\text{when } a > 1 \quad \lim_{\alpha \rightarrow \infty} D^\alpha(e^{ax}) = \infty \qquad \text{when } a < 1 \quad \lim_{\alpha \rightarrow \infty} D^\alpha(e^{ax}) = e^{ax}$$

these are all easy known results of course *when* $a = 1 \quad \lim_{\alpha \rightarrow \infty} D^\alpha(e^{ax}) = e^{ax}$
these are all expected to be the result to repeating multiplication of $a^\alpha e^{ax}$
but there is one case for a where it's not any of the above
let's examine the cyclic derivative functions closely

$$D^\alpha e^x = e^x$$

if we try to write in general e^{ax} formula we get

$$D^\alpha e^{1x} = 1^\alpha e^{1x}$$

which can also explain (with what we already know) why e^x is a repeated derivative of itself

let's look at hyperbolic functions which are cyclic derivatives of order 2

$$\sinh(x) = \frac{e^x - e^{-x}}{2} \qquad \cosh(x) = \frac{e^x + e^{-x}}{2}$$

if we solve for e^{-x} in both functions
we get for $\sinh(x)$

$$2\sinh(x) = e^x - e^{-x} \qquad \therefore 2\sinh(x) - e^x = -e^{-x} \qquad \therefore e^x - 2\sinh(x) = e^{-x}$$

we can do the same for $\cosh(x)$

$$2\cosh(x) = e^x + e^{-x} \qquad \therefore 2\cosh(x) - e^x = e^{-x}$$

let's add both equations together we get

$$2e^{-x} = 2\cosh(x) - e^x + e^x - 2\sinh(x) = 2\cosh(x) - 2\sinh(x)$$

dividing both sides by 2 we get

$$e^{-x} = \cosh(x) - \sinh(x)$$

which is truly an order 2 cyclic derivative function ,if we took the first derivative of the function we get

$$\begin{aligned} D(e^{-x}) &= D(\cosh(x)) - D(\sinh(x)) = \\ &= -e^{-x} = \sinh(x) - \cosh(x) \end{aligned}$$

this is true since if we multiplied by -1 in the first expression we get the same result

of course we can express these in terms of e^x like this

$$e^x = \cosh(x) + \sinh(x)$$

which is true since both sides will return the same result no matter how many times we differentiate them

it looks sort of like Euler formula but for hyperbolics

notice that -1 that is in e^{-x} comes from the equation $a^2 = 1, a = \pm 1$ because if we differentiate any of the e^x or e^{-x} expressions twice we get the the same expressions back

if the idea didn't click in yet let's look at trigonometric

for trigonometric functions but this time we have a formula ready for us

$$e^{ix} = \cos(x) + i \sin(x)$$

differentiating both sides we get

$$ie^{ix} = -\sin(x) + i \cos(x)$$

which again works well ,and if we differentiate again 3 times more we get back to the same expression

notice that here $a = i$ infact not only i but any value that satisfies the expression $a^4 = 1$ works as well , and the returning function will be order 4 cyclic derivative function

the pattern here goes on and on and this is exactly the missing case for a

$$\text{when } a^n = 1, D^n(e^{ax}) = e^{ax} \text{ with } 2 \times n \text{ cyclic order}$$

that is because it holds for both positive and negative values of a, which means that the case $a^1 = 1$ the order is only once cyclic since it has only positive values for case $a^2 = 1$ we get e^{-x} which is 2nd order cyclic derivative and for case $a^4 = 1$ we get e^{ix} which is 4th order cyclic derivative this formula helps us generate any order of cyclic derivatives as for example a 3rd order cyclic derivative will have a as

$$a^3 = 1, a = 1 \text{ or } -\frac{1}{2} + i\frac{\sqrt{3}}{2} \text{ or } -\frac{1}{2} + i\frac{\sqrt{3}}{2}$$

and all of them work exactly as intended, also we can see that some cyclic-derivatives work correctly under the order of another cyclic derivatives like order 2 (sinh and cosh) under order 4 (sin and cos) this is because in order for $a^n = 1$ to be true isn't only n times but it works for n and 2n and 3n so on because every one of them will lead to $1, a^n = a^{2n} = a^{3n} = 1$ in other words the higher cyclic-derivatives that works the same will satisfy the property

$$\text{when } a^n = 1, D^n(e^{ax}) = e^{ax} \text{ with } 2 \times k \text{ cyclic order, where } k \equiv 0 \pmod n$$

getting back to the basics both of case $a^2 = 1$ and case $a^4 = 1$ can be expanded as trigonometric or hyperbolic functions, not only that but both of them satisfy the condition 2^n and both of them can be expanded algebraically, for hyperbolic it's $x^2 - y^2 = 1$ and $x^2 + y^2 = 1$ for trigonometric functions, and since they are

cyclic derivatives they are functions of type FRD, which suggests that they may exist a pattern of function families that are

$$e^{ax} \text{ where } a^n = 1 \text{ that satisfies the condition } 2^n \in \mathbb{Z}^+$$

and these function families may also have algebraic expression that also suggests it will be from high order operator that isn't simple plus or minus also since -1 and i can both be explained geometrically as "rotations" we can also say that *n is the number of unity roots* which also suggests cyclic order derivatives but in 3D or 4D using Quaternions and octonions

9.2 infinity cycles between the cyclic

infact we can do this now

let's start with the first imagery derivative for the cyclic functions

$$D^i e^x = e^x \quad D^i e^{ax} = a^i e^{ax}$$

the first formula is expected since e^x doesn't change by any mean, what is interesting is the second one as it gives some beautiful results let $a = -1$ to get the one for the hyperbolic functions

$$D^i e^{-x} = (-1)^i e^{-x} = (e^{i\pi})^i e^{-x} = e^{-\pi} e^{-x} = e^{-x-\pi} = e^{-(x+\pi)}$$

quite interesting, this means that in between $\cosh(x)$ and $\sinh(x)$ lies exactly $e^{-(x+\pi)}$, and from our e^{-x} formula we can say that

$$e^{-(x+\pi)} = \cosh(x + \pi) - \sinh(x + \pi)$$

and the reason i didn't say that it lies between $\sinh(x)$ becoming $\cosh(x)$ nor the other way around is because it's a mix of both of functions, to find what comes in the middle of any of them we have to differentiate them by themselves so we get

$$D^i \sinh(x) = \frac{e^x - (-1)^i e^{-x}}{2} = \frac{e^x - e^{-(x+\pi)}}{2}$$

which mean that right in the middle of $\sinh(x)$ transforming to $\cosh(x)$ there exists that real function, the same goes for $\cosh(x)$

$$D^i \cosh(x) = \frac{e^x + (-1)^i e^{-x}}{2} = \frac{e^x + e^{-(x+\pi)}}{2}$$

we can do the same for e^{ix} and see what happens

$$D^i e^{ix} = i^i e^{ix} = (e^{\frac{i\pi}{2}})^i e^{ix} = e^{\frac{-\pi}{2}} e^{ix} = e^{ix - \frac{\pi}{2}}$$

we can expand it with euler identity

$$e^{ix - \frac{\pi}{2}} = e^{\frac{-\pi}{2}} \cos(x) + i e^{\frac{-\pi}{2}} \sin(x)$$

now we can expand this to merge the e terms

$$e^{ix-\frac{\pi}{2}} = e^{\frac{-\pi}{2}} \frac{e^{ix} + e^{-ix}}{2i} + ie^{\frac{-\pi}{2}} \frac{e^{ix} - e^{-ix}}{2i} = e^{\frac{-\pi}{2}} \frac{e^{ix-\frac{\pi}{2}} + e^{-ix-\frac{\pi}{2}}}{2i} + i \frac{e^{ix-\frac{\pi}{2}} - e^{-ix-\frac{\pi}{2}}}{2i}$$

then we turn it back to trigonometric we get

$$D^i e^{ix} = \cos(x - \frac{\pi}{2}) + i \sin(x - \frac{\pi}{2})$$


as we can see , the i -th derivative represents a half-rotation between sin and cos changing, which is expected since it's the half way between them
now let's try getting them from their original rules

$$D^i \sin(x) = \sin(\frac{i\pi}{2} + x)$$

this doesn't mean one of the formulas is wrong , since we still can do ${}^i D^i$ for both of them and get the same results , this simply means that one of them operating on real plane with scaling and rotation and the other is on complex plane, same result different prescriptive

9.3 Geometric Interpretation of cyclic derivatives

let's start simple with e^x in the D^z Re-Im Plane



this is expected since e^x is the exact value of itself at any point , even if we tried to represent them as vectors we will get the same results because the it always begins and ends at any value of the order derivative , in other words there is no direction for the vector to point at, but this plane while may show something but isnt so useful, instead we can use the **D(i) vector plane**
of course using it on e^x will give points with no direction as they dont grow or decay not in derivative value or function growth ration

10 The Geometric Differentiel (GDI) theory

10.1 FRD and NFRD

these are two important terms that will be helpful for us

FRD : Full Real Differentiable **NFRD** : Not-Full Real Differentiable

these are two very important terms as they categorize all the functions in D^z since most of the functions are either Integrable and Differentiable or Integrable and not Differentiable and D^z can work on both sides

FRD Functions are functions that have a clean closed form for its α -th derivative that isn't a power series (that is of course if the function itself isn't defined by a power series)

these are functions like x^n with the condition that $\text{Re}(z) \leq n$ and e^x

NFRD functions are functions that have an only closed form being a power series, so functions like $\tan(x)$ and other trigonometric functions are NFRD

this happens of course for a lot of reasons like breaking GDI first principle and

10.2 Transformative Functions

They are the type of functions that transform dependence with derivative, for example

$$\frac{d}{dx}(\cos^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$$

as we can see, when we take the derivative of the function it changes dependence from the value of x to the value of the power of x

10.3 The inverse functions problem

Inverse functions are a problem, at least for being FRD functions as it's not possible for most of them and I have some explanations

Inverse functions are weird compared to the normal functions, we can see some similarities between them and their original functions like x^n and $x^{\frac{1}{n}}$ as they both look identical for odd values because for even numbers we get complex values, but rotated and cut to half, in most of the inverse functions we can find these cuts near always

we can expect this because not every function can be fully inverted

a function is a set of inputs that lead to outputs, one output can be gotten by many inputs but not the way around, so when we try to invert functions like x^2 that has one output for more than one input we get something that doesn't work in the real number systems, and for this example it's the complex values for negative integers

10.4 The Mirror functions

but that doesn't mean we can't have inverse functions of some sort, let's look at the formula for x^n

$$D^\alpha x^n = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} x^{n-\alpha}$$

now let's try to make the numerator go to its limit, since Gamma function doesn't work for non-positive integers we can say that

$$\because n+1 \leq 0 \quad \therefore n \leq -1 \quad \therefore -n \geq 1$$

we can then put it in the x^n derivative

$$D^\alpha x^{-n} = \frac{\Gamma(1-n)}{\Gamma(1-n-\alpha)} x^{-n-\alpha} = \frac{\Gamma(1-n)}{\Gamma(1-(n+\alpha))} x^{-(n+\alpha)}$$

this function works exactly like x^n but for negative values of n $(-(-1), -(-3))$, it behaves the same for derivatives and Integrals as well

Part III

Applications of D^z

11 Bonus Topics

11.0.1 The zeta function $D^z \zeta(n)$ and its nature

the Riemann zeta function is one of the most well known functions in mathematics, its formula is :

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \sum_{n=1}^{\infty} n^{-s}$$

so using the $D^\alpha a^{-x} = (-1)^\alpha a^{-x} \ln(a)^\alpha$ we get

$$D^\alpha \zeta(s) = \sum_{n=1}^{\infty} (-1)^\alpha n^{-s} \ln(n)^\alpha$$

we can put the -1 outside from the series since it's not dependent on n we can now find the first derivative simply

$$D^1 \zeta(s) = (-1) \sum_{n=1}^{\infty} n^{-s} \ln(n) = - \sum_{n=1}^{\infty} \frac{\ln(n)}{n^s}$$

and we can find the half-derivative

$$D^{\frac{1}{2}} \zeta(s) = (-1)^{\frac{1}{2}} \sum_{n=1}^{\infty} n^{-s} \ln(n)^{\frac{1}{2}} = i \sum_{n=1}^{\infty} \frac{\sqrt{\ln(n)}}{n^s}$$

well this is fascinating
 we can also find the i -th derivative

$$D^i \zeta(s) = (-1)^i \sum_{n=1}^{\infty} n^{-s} \ln(n)^i = (e^{i\pi})^i \sum_{n=1}^{\infty} \frac{e^{i \ln(\ln(n))}}{n^s} = e^{-\pi} \sum_{n=1}^{\infty} \frac{e^{i \ln(\ln(n))}}{n^s}$$

12 Fractional and Complex Differential equations

12.1 Matrix equations

We seek all values of n such that for $A = nI, D^A = nI$:

first we need to be reminded with a matrix order propertie we have proven earlier that for any n scaler matrix order derivative will return $\Gamma(n+1)$ times identity matrix

with this we can say $D^A = \Gamma(n+1)I$ thus we can write the equation as this

$$nI = \Gamma(n+1)I$$

then we use the identity of the Gamma $\Gamma(z+1) = z\Gamma(z)$

$$nI = n\Gamma(n)I$$

dividing both sides by nI returns $1 = \Gamma(n)$

substituting $\Gamma(n) = (n-1)!$ we get $(n-1)! = 1$

this only holds true for 2, 1 as $(1-1)! = 0! = 1$ and $(2-1)! = 1! = 1$

if we tried numbers bigger than one we get $(3-1)! = 2! = 2 \neq 1$ then the function grows beyond it's reach

so the only solutions is where $n \in \{1, 2\}$ does the equation satisfy $D^A = \Gamma(n)I$

13 Applications in real world

as sir Issac Newton started it based on motion of everything in the universe , when he didn't find good way to see it he created his way which we shall call today calculus, so of course we are beginning with physics

13.1 Dynamics with D^z operator

dynamics are the beginning of and heart with physics, it's the language of motion in it's simplest form , and so we can begin by stating that $f(x)$ is position and the first derivative with respect to time is Velocity , and the second derivative with respect to time is acceleration , that is for integer derivatives , and for fractional derivatives things tend to get weird

let's take for example the function x^5

$$f(x) = x^5 \quad v(x) = 5x^4 \quad a(x) = 20x^3$$

these are the three main dynamical functions for $f(x)$
now we are going to propose there different functions that works in the middle
so :

$${}_tD^{0.5}f(x) = hf(x) = \frac{120}{\Gamma(5.5)}x^{4.5} \quad {}_tD^{0.5}f(x) = hv(x) = \frac{120}{\Gamma(4.5)}x^{3.5} \quad {}_tD^{0.5}f(x) = ha(x) = \frac{120}{\Gamma(3.5)}x^{2.5}$$

since for every deriavtive we multiply the time scale unit by 1 we can say tha
ttthe units for these are $m/\sqrt{s}, m/\sqrt{s^3}, m/\sqrt{s^5}$
as we can see, these are states in between , hf is between position and veloicity ,
 hv is between velocity and acceleration let's try some other unusual things like
the i -th derivative and $A = 5I$ matrix

$$D^i f(x) = \frac{120}{\Gamma(6-i)}x^{5-i} m/s^i$$

and that is a present , complex time
well it's not fully complex as we can see this time is just for the system itself,
it's for the event , since there is no time squared nor square rooted these are
just for the event itself, yet it's weird to see a complex time of event itself

$$D^A f(x) = \Gamma(6)I = 120I m/s^A$$

and now Matrix Time , and this one is very special since this isn't an Identified
matrix , it can continue for ever , or can simply be written as m/s^5 since both
are technically the same

Contents

I	Algebraic foundation	1
1	The Generalized Operator for $f(x) = x^n$	1
1.1	From Integer to Fractional Order	1
1.2	the x^{-n} problem	3
2	Formulas for Other Algebraic Functions	3
2.1	The General Formula for a^x	3
2.2	The General Formula for e^x	4
2.3	The General Formula for e^{ax}	4
2.4	The problem of $\log_a(x)$	5
3	Proprties of the D^z operator	6
3.1	General power series rule	6
3.2	The linearity of D^z operator	7
3.3	The Index law	8
3.4	The Multiplication Law	9
3.5	Operator inverses and identities	11

4	Deriving the Rules of the Fractional and complex derivatives	13
4.1	General product rule	13
4.2	General chain rule	16
4.2.1	General $f(x^n)$ formula	16
4.2.2	General $f(ax)$ formula	17
4.2.3	General Series form for chain rule	18
4.2.4	General Closed form for chain rule	18
5	Interpretation of fractional and complex derivative in other functions	18
5.1	Trigonometric Functions	18
5.1.1	$\sin(x)$ and $\cos(x)$	18
5.1.2	$\tan(x)$ and $\sec(x)$	20
5.1.3	$\csc(x)$ and $\cot(x)$	21
5.2	Hyperbolic Functions	22
5.2.1	$\sinh(x)$, $\cosh(x)$ and $\tanh(x)$	22
5.2.2	$\operatorname{sech}(x)$, $\operatorname{csch}(x)$ and $\operatorname{coth}(x)$	23
5.3	The Inverse Trigonometric and Hyperbolic Functions	24
6	Matrix derivative functions	24
6.1	Whole matrix derivatives	24
6.2	What is the meaning of matrix derivatives	25
6.3	Matrix order properties	26
7	Functional derivatives	27
7.1	derivative order as function of itself	27
7.2	derivative order as function of it's dependent variable	27
7.3	derivative order as function of another Dimension variable (Dimension changer)	28
II	D^z Analysis	28
8	The nature of functions under D^z Field	28
8.1	Explaining what is a fractional derivative	28
8.2	what it means for a derivative order to be Complex	29
9	Cyclic derivatives	33
9.1	e^x and Cyclic functions	33
9.2	infinity cycles between the cyclic	36
9.3	Geometric Interpretation of cyclic derivatives	37
10	The Geometric DifferentInetgrel (GDI) theory	38
10.1	FRD and NFRD	38
10.2	Transformative Functions	38
10.3	The inverse functions problem	38
10.4	The Mirror functions	39

III	Applications of D^z	39
11	Bonus Topics	39
11.0.1	The zeta function $D^z\zeta(n)$ and it's nature	39
12	Fractional and Complex Differential equations	40
12.1	Matrix equations	40
13	Applications in real world	40
13.1	Dynamics with D^z operator	40