

# cyclic derivatives: Symmetry of Fractional and complex

Fire Fox

December 2025

## 1 Introduction

## 2 The symmetry of fractional cyclics and the complex plane

### 2.1 Overlapping circle in the complex

## 3 Mittag-Leffler and cyclic derivatives

### 3.1 Mittage-Leffler is somewhat a cyclic derivative

In the previous cyclic derivatives research, I pointed out that Cyclic derivatives and Mittag-Leffler are connected, and after some research, I can say that we can express cyclic functions in terms of Mittag-Leffler functions like this  $\cosh_n(x) = E_n(x^n)$

which can be seen in the result of  $E_2(x)$  being  $\cosh(\sqrt{x})$ , when squared we get  $\cosh(x) = E_2(x^2)$

this also allows us to get the integerral definition of  $\cosh_n(x)$  using the  $E_n(x^n)$

$$E_{\alpha,\beta}(x) = \frac{1}{2i\pi} \oint_C \frac{t^{\alpha-\beta} e^t}{t^\alpha - x} dt \quad \text{Re}(\alpha) > 0, \text{Re}(\beta) > 0$$

$$\cosh_n(x) = E_{n,1}(x^n) = \frac{1}{2i\pi} \oint_C \frac{t^{n-1} e^t}{t^n - x^n} dt \quad \text{Re}(n) > 0$$

The same way we can get  $\sinh_n K(x)$  as the  $K$ -th deriavtive of  $E_n(x^n)$ , and if we assume that this function converge and the integral is fixed, we can get the derivative inside of the integral too

$$\sinh_n K(x) = D^K E_n(x^n) = \frac{1}{2i\pi} \oint_C D^K \left( \frac{t^{n-1} e^t}{t^n - x^n} \right) dt \quad \text{Re}(n) > 0, K \in \mathbb{N}$$

using this we can get te first and the second  $\sinh_n K(x)$  functions

$$\sinh_n(x) = \frac{1}{2i\pi} \oint_C t^{n-1} e^t \frac{nx^{n-1}}{(t^n - x^n)^2} dt \quad \text{Re}(n) > 0$$

$$\sinh_n \Pi(x) = \frac{1}{2i\pi} \oint_C t^{n-1} e^{t \frac{n(n-1)x^{n-2}(t^n - x^n) + 2n^n x^{2n-2}}{(t^n - x^n)^2}} dt \quad \text{Re}(n) > 0$$

we can also find  $\cos_{2q|q}(x)$  where  $\cos_{2q|q}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2q}}{(2q)!}$