



Portsmouth Hospitals  
University  
NHS Trust

# Paediatric Drug Calculation Handbook

# Disclaimer

This handbook is intended as a guide for healthcare professionals involved in the care of paediatric patients. While every effort has been made to ensure the accuracy of the information contained herein, the authors and publishers accept no responsibility for any errors, omissions, or outcomes resulting from the use of this material.

Clinical judgment, institutional protocols, and individual patient considerations must always take precedence. It is the responsibility of the healthcare provider to verify drug dosages, contraindications, interactions, and calculations with up-to-date references and official prescribing information before administering any medication.

This handbook is not a substitute for professional training or medical advice and should not be used as the sole resource for patient care decisions.

# Doses

## Energy

4 J/kg rounded to the next increment<sup>(1)</sup>.  
10, 20, 30, 50, 75, 100, 120

## ETT

See chart on page 42

## LMA Ambu AuraGain

As per Ambu™ Data Sheet<sup>(2)</sup>

## Atropine 600 mcg/ml

Minimum dose: 100 mcg<sup>(3)</sup>  
Maximum dose: 600 mcg<sup>(3)</sup> 20 mcg/kg<sup>(4)</sup>

## Glycopyrrolate

Maximum per dose: 200 mcg<sup>(5)</sup> 10 mcg/kg<sup>(5)</sup>

## Sux

2 mg/kg IV  
4 mg/ml IM<sup>(6)</sup>

## Adrenaline

10 mcg/kg<sup>(7)</sup>

## Fluid

10-20 ml/kg<sup>(8)</sup>

## Glucose 10%

2 ml/kg<sup>(9)</sup>

Concentrations in *italics* have reached the upper or lower dosage limit for that drug.

Rocuronium  
Standard - 0.6  
mg/kg  
RSI - 1 mg/kg<sup>(10)</sup>

Ketamine  
RSI - 1-2 mg/kg<sup>(11)</sup>

Fentanyl  
1-3 mcg/kg<sup>(12)</sup>

# 2 kg



**Energy** 4 J/kg

**8 J**

**ETT** (Consider Uncuffed)

**2 - 3 mm**

**LMA** Ambu AuraGain

**Size 1**

**Atropine**

100 mcg = 0.17 ml of  
600mcg/ml. Dilution is  
recommended

**100 mcg**

**Glycopyrrolate**

**0.1 ml**

**Sux** **0.08 ml IV | 0.16 ml IM**

**Adrenaline** 1:10,000  
(Cardiac Arrest)

**0.2 ml**

**Fluid** 10-20ml/kg

**20-40 ml**

**Glucose 10%**

**4 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.1 ml  
RSI - 0.2 ml

**Ketamine**  
RSI - 0.2-0.4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
2-6 mcg

**3 kg**



**NHS**  
Portsmouth Hospitals  
University  
NHS Trust

**Energy** 4 J/kg

**10 J**

**ETT** (Consider Uncuffed)

**2 - 3 mm**

**LMA** Ambu AuraGain

**Size 1**

**Atropine**

100 mcg = 0.17 ml of  
600mcg/ml. Dilution is  
recommended

**100 mcg**

**Glycopyrrolate**

**0.15 ml**

**Sux** 0.12 ml IV | 0.25 ml IM

**Adrenaline** 1:10,000  
(Cardiac Arrest)

**0.3 ml**

**Fluid** 10-20ml/kg

**30-60 ml**

**Glucose 10%**

**6 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.2 ml  
RSI - 0.3 ml

**Ketamine**  
RSI - 0.3-0.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
3-9 mcg

# 4 kg



**Energy** 4 J/kg

**20 J**

**ETT** (Consider Uncuffed)

**2 - 3.5 mm**

**LMA** Ambu AuraGain

**Size 1**

**Atropine**

100 mcg = 0.17 ml of  
600mcg/ml. Dilution is  
recommended

**100 mcg**

**Glycopyrrolate**

**0.2 ml**

**Sux** **0.16 ml IV | 0.3 ml IM**

**Adrenaline** 1:10,000  
(Cardiac Arrest)

**0.4 ml**

**Fluid** 10-20ml/kg

**40-80 ml**

**Glucose 10%**

**8 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.3 ml  
RSI - 0.4 ml

**Ketamine**  
RSI - 0.4-0.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
4-12 mcg

# 5 kg



**Energy** 4 J/kg

**20 J**

**ETT** (Consider Uncuffed)

**2 - 3.5 mm**

**LMA** Ambu AuraGain

**Size 1 or 1½**

**Atropine**

**0.17 ml**

**Glycopyrrolate**

**0.25 ml**

**Sux** **0.2 ml IV | 0.4 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**0.5 ml**

**Fluid** 10-20ml/kg

**50-100 ml**

**Glucose 10%**

**10 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.3 ml  
RSI - 0.5 ml

**Ketamine**  
RSI - 0.5-1 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
5-15 mcg

# 6 kg



**Energy** 4 J/kg **20 J**

**ETT** (Consider Uncuffed) **2 - 3.5 mm**

**LMA** Ambu AuraGain **Size 1 - 1½**

**Atropine** **0.2 ml**

**Glycopyrrolate** **0.3 ml**

**Sux** **0.25 ml IV | 0.5 ml IM**

**Adrenaline** 1:10,000  
(Cardiac Arrest) **0.6 ml**

**Fluid** 10-20ml/kg **60-120 ml**

**Glucose 10%** **12 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.4 ml  
RSI - 0.6 ml

**Ketamine**  
RSI - 0.6-1.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
6-18 mcg

# 7 kg



**Energy** 4 J/kg

**30 J**

**ETT** (Consider Uncuffed)

**2 - 3.5 mm**

**LMA** Ambu AuraGain

**Size 1½**

**Atropine**

**0.23 ml**

**Glycopyrrolate**

**0.35 ml**

**Sux** **0.3 ml IV | 0.6 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**0.7 ml**

**Fluid** 10-20ml/kg

**70-140 ml**

**Glucose 10%**

**14 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.4 ml  
RSI - 0.7 ml

**Ketamine**  
RSI - 0.7-1.4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
7-21 mcg

# 8 kg



**Energy** 4 J/kg **30 J**

**ETT** (Consider Uncuffed) **2 - 3.5 mm**

**LMA** Ambu AuraGain **Size 1½**

**Atropine** **0.27 ml**

**Glycopyrrolate** **0.4 ml**

**Sux** **0.3 ml IV | 0.6 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **0.8 ml**

**Fluid** 10-20ml/kg **80-160 ml**

**Glucose 10%** **16 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.5 ml  
RSI - 0.8 ml

**Ketamine**  
RSI - 0.8-1.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
8-24 mcg

# 9 kg



**Energy** 4 J/kg

**30 J**

**ETT** (Consider Uncuffed)

**2 - 3.5 mm**

**LMA** Ambu AuraGain

**Size 1½**

**Atropine**

**0.3 ml**

**Glycopyrrolate**

**0.45 ml**

**Sux** **0.36 ml IV | 0.7 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**0.9 ml**

**Fluid** 10-20ml/kg

**90-180 ml**

**Glucose 10%**

**18 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.5 ml  
**RSI - 0.9 ml**

**Ketamine**  
RSI - 0.9-1.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
9-27 mcg

# 10 kg



**Energy** 4 J/kg **30 J**

**ETT** (Cuffed) **3.5 - 4 mm**

**LMA** Ambu AuraGain **Size 1½ - 2**

**Atropine** **0.3ml**

**Glycopyrrolate** **0.5ml**

**Sux** **0.4 ml IV | 0.8 ml IM**

**Adrenaline** (Cardiac Arrest) **1:10,000 1ml**

**Fluid** 10-20ml/kg **100-200ml**

**Glucose 10%** **20ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.6 ml  
**RSI - 1 ml**

**Ketamine**  
RSI - 1-2ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
10-30 mcg

# 11 kg



**Energy** 4 J/kg

**50 J**

**ETT** (Cuffed)

**3.5 - 4 mm**

**LMA** Ambu AuraGain

**Size 1½ - 2**

**Atropine**

**0.4 ml**

**Glycopyrrolate**

**0.6 ml**

**Sux** **0.4 ml IV | 0.9 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**1.1 ml**

**Fluid** 10-20ml/kg

**110-220 ml**

**Glucose 10%**

**22 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.7 ml  
**RSI - 1.1 ml**

**Ketamine**  
RSI - 1.1-2.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
11-33 mcg

# 12 kg



**Energy** 4 J/kg **50 J**

**ETT** (Cuffed) **3.5 - 4 mm**

**LMA** Ambu AuraGain **Size 2**

**Atropine** **0.3 ml**

**Glycopyrrolate** **0.6 ml**

**Sux** **0.5 ml IV | 1 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **1.2 ml**

**Fluid** 10-20ml/kg **120 - 240 ml**

**Glucose 10%** **24 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.7 ml  
RSI - 1.2 ml

**Ketamine**  
RSI - 1.2-2.4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
12-36 mcg

# 13 kg



**Energy** 4 J/kg

**50 J**

**ETT** (Cuffed)

**3.5 - 4 mm**

**LMA** Ambu AuraGain

**Size 2**

**Atropine**

**0.43 ml**

**Glycopyrrolate**

**0.65 ml**

**Sux**

**0.5 ml IV | 1 ml IM**

**Adrenaline**

1:10,000  
(Cardiac Arrest)

**1.3 ml**

**Fluid**

10-20ml/kg

**130 - 260 ml**

**Glucose 10%**

**26 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.8 ml  
RSI - 0.13 ml

**Ketamine**  
RSI - 1.3 - 2.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
13 - 39 mcg

# 14 kg



**Energy** 4 J/kg **50 J**

**ETT** (Cuffed) **3.5 - 4 mm**

**LMA** Ambu AuraGain **Size 2**

**Atropine** **0.5 ml**

**Glycopyrrolate** **0.7 ml**

**Sux** **0.7 ml IV | 1.1 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **1.4 ml**

**Fluid** 10-20ml/kg **140 - 280 ml**

**Glucose 10%** **28 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.8 ml  
RSI - 1.4 ml

**Ketamine**  
RSI - 1.4 -2.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
14 - 42 mcg

# 15 kg



NHS  
Portsmouth Hospitals  
University  
NHS Trust

**Energy** 4 J/kg

50 J

**ETT** (Cuffed)

4 - 4.5 mm

**LMA** Ambu AuraGain

Size 2

**Atropine**

0.5 ml

**Glycopyrrolate**

0.8 ml

**Sux** 0.6 ml IV | 1.2 ml IM

**Adrenaline** 1:10,000  
(Cardiac Arrest)

1.5 ml

**Fluid** 10-20ml/kg

150-300 ml

**Glucose 10%**

30 ml

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 0.9 ml  
RSI - 1.5 ml

**Ketamine**  
RSI - 1.5-3ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
15-45 mcg

# 16 kg



**Energy** 4 J/kg

**75 J**

**ETT** (Cuffed) **4 - 4.5 mm**

**LMA** Ambu AuraGain

**Size 2**

**Atropine** **0.5 ml**

**Glycopyrrolate** **0.8 ml**

**Sux** **0.6 ml IV | 1.3 ml IM**

**Adrenaline** 1:10,000  
(Cardiac Arrest) **1.6 ml**

**Fluid** 10-20ml/kg **160 - 320 ml**

**Glucose 10%** **32 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1 ml  
RSI - 1.6 ml

**Ketamine**  
RSI - 1.6-3.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
16-48 mcg

# 17 kg



**Energy** 4 J/kg

**75 J**

**ETT** (Cuffed)

**4 - 4.5 mm**

**LMA** Ambu AuraGain

**Size 2**

**Atropine**

**0.6 ml**

**Glycopyrrolate**

**0.9 ml**

**Sux** **0.7 ml IV | 1.4 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**1.7 ml**

**Fluid** 10-20ml/kg

**170 - 340 ml**

**Glucose 10%**

**34 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1 ml  
**RSI - 1.7 ml**

**Ketamine**  
RSI - 1.7-3.4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
17-51 mcg

# 18 kg



**Energy** 4 J/kg

**75 J**

**ETT** (Cuffed) **4 - 4.5 mm**

**LMA** Ambu AuraGain

**Size 2**

**Atropine** **0.6 ml**

**Glycopyrrolate** **0.9 ml**

**Sux** **0.7 ml IV | 1.4 ml IM**

**Adrenaline** 1:10,000  
(Cardiac Arrest) **1.8 ml**

**Fluid** 10-20ml/kg **180-360 ml**

**Glucose 10%** **36 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.1 ml  
RSI - 1.8 ml

**Ketamine**  
RSI - 1.8-3.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
18-54 mcg

# 19 kg



**Energy** 4 J/kg

**75 J**

**ETT** (Cuffed)

**4.5 - 5 mm**

**LMA** Ambu AuraGain

**Size 2**

**Atropine**

**0.6 ml**

**Glycopyrrolate**

**0.9 ml**

**Sux** **0.8 ml IV | 1.5 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**1.9 ml**

**Fluid** 10-20ml/kg

**190 - 380 ml**

**Glucose 10%**

**38 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.1 ml  
**RSI - 1.9 ml**

**Ketamine**  
RSI - 1.9-3.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
19-57 mcg

# 20 kg



**Energy** 4 J/kg **75 J**

**ETT** (Cuffed) **4.5 - 5 mm**

**LMA** Ambu AuraGain **Size 2 / 2½**

**Atropine** **0.7 ml**

**Glycopyrrolate** **1 ml**

**Sux** **0.8 ml IV | 1.6 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **2 ml**

**Fluid** 10-20ml/kg **200 - 400 ml**

**Glucose 10%** **40 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.2 ml  
RSI - 2 ml

**Ketamine**  
RSI - 2-4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
20-60 mcg

# 21 kg



**Energy** 4 J/kg

**75 J**

**ETT** (Cuffed)

**4.5 - 5 mm**

**LMA** Ambu AuraGain

**Size 2 / 2½**

**Atropine**

**0.7 ml**

**Glycopyrrolate**

**1 ml**

**Sux** **0.8 ml IV | 1.7 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**2.1 ml**

**Fluid** 10-20ml/kg

**210 - 420 ml**

**Glucose 10%**

**42 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.2 ml  
**RSI - 2.1 ml**

**Ketamine**  
RSI - 2.1-4.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
21-63 mcg

# 22 kg



**Energy** 4 J/kg **100 J**

**ETT** (Cuffed) **4.5 - 5 mm**

**LMA** Ambu AuraGain **Size 2½**

**Atropine** **0.7 ml**

**Glycopyrrolate** **1 ml**

**Sux** **0.9 ml IV | 1.8 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **2.2 ml**

**Fluid** 10-20ml/kg **220-240 ml**

**Glucose 10%** **44 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.1 ml  
RSI - 2.2 ml

**Ketamine**  
RSI - 2.2-4.4 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
22-66 mcg

# 23 kg



**Energy** 4 J/kg

**100 J**

**ETT** (Cuffed)

**4.5 - 5 mm**

**LMA** Ambu AuraGain

**Size 2½**

**Atropine**

**0.8 ml**

**Glycopyrrolate**

**1 ml**

**Sux**

**0.9 ml IV | 1.8 ml IM**

**Adrenaline**

1:10,000  
(Cardiac Arrest)

**2.3 ml**

**Fluid**

10-20ml/kg

**230 - 460 ml**

**Glucose 10%**

**46 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**

Standard - 1.4 ml

RSI - 2.3 ml

**Ketamine**

RSI - 2.3-4.6 ml

of 10mg/ml

**Fentanyl**

1-3 mcg/kg

23-69 mcg

# 24 kg



**Energy** 4 J/kg **100 J**

**ETT** (Cuffed) **4.5 - 5 mm**

**LMA** Ambu AuraGain **Size 2½**

**Atropine** **0.8 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1 ml IV | 1.9 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **2.4 ml**

**Fluid** 10-20ml/kg **240 - 480 ml**

**Glucose 10%** **48 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.4 ml  
RSI - 2.4 ml

**Ketamine**  
RSI - 2.4-4.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
24-72 mcg

# 25 kg



**Energy** 4 J/kg

**100 J**

**ETT** (Cuffed)

**4.5 - 5 mm**

**LMA** Ambu AuraGain

**Size 2½**

**Atropine**

**0.8 ml**

**Glycopyrrolate**

**1 ml**

**Sux**

**1 ml IV | 2 ml IM**

**Adrenaline**

1:10,000  
(Cardiac Arrest)

**2.5 ml**

**Fluid**

10-20ml/kg

**250 - 500 ml**

**Glucose 10%**

**50 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**

Standard - 1.5 ml  
RSI - 2.5 ml

**Ketamine**

RSI - 2.5-5 ml  
of 10mg/ml

**Fentanyl**

1-3 mcg/kg  
25-75 mcg

# 26 kg



**Energy** 4 J/kg **100 J**

**ETT** (Cuffed) **4.5 - 5 mm**

**LMA** Ambu AuraGain **Size 2½**

**Atropine** **0.9 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1 ml IV | 2.1 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **2.6 ml**

**Fluid** 10-20ml/kg **260 - 520 ml**

**Glucose 10%** **52 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.6 ml  
RSI - 2.6 ml

**Ketamine**  
RSI - 2.6-5.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
26-78 mcg

# 28 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**4.5 - 5 mm**

**LMA** Ambu AuraGain

**Size 2½**

**Atropine**

**0.9 ml**

**Glycopyrrolate**

**1 ml**

**Sux**

**1.1 ml IV | 2.2 ml IM**

**Adrenaline**

1:10,000  
(Cardiac Arrest)

**2.8 ml**

**Fluid**

10-20ml/kg

**280 - 560 ml**

**Glucose 10%**

**56 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**

Standard - 1.7 ml

**RSI - 2.8 ml**

**Ketamine**

RSI - 2.8-5.6 ml

of 10mg/ml

**Fentanyl**

1-3 mcg/kg

28-84 mcg

# 30 kg



**Energy** 4 J/kg **120 J**

**ETT** (Cuffed) **5 - 5.5 mm**

**LMA** Ambu AuraGain **Size 2½ / 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1.2 ml IV | 2.4 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **3 ml**

**Fluid** 10-20ml/kg **300 - 600 ml**

**Glucose 10%** **60 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 1.8 ml  
RSI - 3 ml

**Ketamine**  
RSI - 3-6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
30-90 mcg

# 32 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**5 - 5.5 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine**

**1 ml**

**Glycopyrrolate**

**1 ml**

**Sux**

**1.3 ml IV | 2.6 ml IM**

**Adrenaline**

1:10,000  
(Cardiac Arrest)

**3.2 ml**

**Fluid**

10-20ml/kg

**320 - 640 ml**

**Glucose 10%**

**64 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**

Standard - 1.9 ml

**RSI - 3.2 ml**

**Ketamine**

RSI - 3.2-6.4 ml

of 10mg/ml

**Fentanyl**

1-3 mcg/kg

32-96 mcg

# 34 kg



**Energy** 4 J/kg **120 J**

**ETT** (Cuffed) **5.5 - 6 mm**

**LMA** Ambu AuraGain **Size 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1.4 ml IV | 2.7 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **3.4 ml**

**Fluid** 10-20ml/kg **340 - 680 ml**

**Glucose 10%** **68 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2 ml  
RSI - 3.4 ml

**Ketamine**  
RSI - 3.4-6.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
34-102 mcg

# 36 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**5.5 - 6 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine**

**1 ml**

**Glycopyrrolate**

**1 ml**

**Sux** **1.4 ml IV | 2.9 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**3.6 ml**

**Fluid** 10-20ml/kg

**360 - 720 ml**

**Glucose 10%**

**72 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.2 ml  
**RSI - 3.6 ml**

**Ketamine**  
RSI - 3.6-7.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
36-108 mcg

# 38 kg



**Energy** 4 J/kg **120 J**

**ETT** (Cuffed) **5.5 - 6 mm**

**LMA** Ambu AuraGain **Size 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1.5 ml IV | 3.0 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **3.8 ml**

**Fluid** 10-20ml/kg **380 - 760 ml**

**Glucose 10%** **76 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.3 ml  
RSI - 3.8 ml

**Ketamine**  
RSI - 3.8-7.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
38-114 mcg

# 40 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**6 - 6.5 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine**

**1 ml**

**Glycopyrrolate**

**1 ml**

**Sux** 1.6 ml IV | 3.2 ml IM

**Adrenaline** 1:10,000 (Cardiac Arrest)

**4 ml**

**Fluid** 10-20ml/kg

**400 - 800 ml**

**Glucose 10%**

**80 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.4 ml  
RSI - 4 ml

**Ketamine**  
RSI - 4-8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
40-120 mcg

# 42 kg



**Energy** 4 J/kg **120 J**

**ETT** (Cuffed) **6 - 6.5 mm**

**LMA** Ambu AuraGain **Size 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1.7 ml IV | 3.4 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **4.2 ml**

**Fluid** 10-20ml/kg **420 - 840 ml**

**Glucose 10%** **84 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.5 ml  
**RSI - 4.2 ml**

**Ketamine**  
RSI - 4.2-8.4ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
42-126 mcg

# 44 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**6 - 6.5 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine**

**1 ml**

**Glycopyrrolate**

**1 ml**

**Sux** **1.8 ml IV | 3.5 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**4.4 ml**

**Fluid** 10-20ml/kg

**440 - 880 ml**

**Glucose 10%**

**88 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.6 ml  
**RSI - 4.4 ml**

**Ketamine**  
RSI - 4.4-8.8 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
44-132 mcg

# 46 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed) **6 - 6.5 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **1.8 ml IV | 3.7 ml IM**

**Adrenaline** **4.6 ml**  
1:10,000 (Cardiac Arrest)

**Fluid** **10-20ml/kg** **460 - 920 ml**

**Glucose 10%** **92 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.8 ml  
**RSI - 4.6 ml**

**Ketamine**  
RSI - 4.6-9.2 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
46-138 mcg

# 48 kg



**Energy** 4 J/kg

**120 J**

**ETT** (Cuffed)

**6 - 6.5 mm**

**LMA** Ambu AuraGain

**Size 3**

**Atropine**

**1 ml**

**Glycopyrrolate**

**1 ml**

**Sux** **1.9 ml IV | 3.8 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest)

**4.8 ml**

**Fluid** 10-20ml/kg

**480 - 960 ml**

**Glucose 10%**

**96 ml**

**Anaphylaxis**

Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 2.9 ml  
**RSI - 4.8 ml**

**Ketamine**  
RSI - 4.8-9.6 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
48-144 mcg

# 50 kg



**Energy** 4 J/kg **120 J**

**ETT** (Cuffed) **6 - 6.5 mm**

**LMA** Ambu AuraGain **Size 3**

**Atropine** **1 ml**

**Glycopyrrolate** **1 ml**

**Sux** **2 ml IV | 4 ml IM**

**Adrenaline** 1:10,000 (Cardiac Arrest) **5 ml**

**Fluid** 10-20ml/kg **500 - 1000 ml**

**Glucose 10%** **100 ml**

**Anaphylaxis** Take arrest dose adrenaline - dilute to 10ml with saline.  
use 0.5 - 1 ml IV boluses to treat hypotension

**Rocuronium**  
Standard - 3 ml  
**RSI - 5 ml**

**Ketamine**  
RSI - 5-10 ml  
of 10mg/ml

**Fentanyl**  
1-3 mcg/kg  
50-150 mcg

Age (Years)	Term - 8 Months	8 months - 2	2 - 4	4 - 6	6 - 8	8 - 10	10 - 12
ETT Size (cuffed) (Stylet)	3.0 (Stylet)	3.5 (Stylet)	4.0	4.5	5.0	5.5	6.0
Length (cm)	11	12	13	14	15	16	17
Bougie	5Fr	10Fr	10Fr	10Fr	10Fr	14Fr	14Fr
Suction Catheter	6	8	10	10	10	14Fr	14Fr

# References

1. Resuscitation Council UK. (2021). Paediatric Advanced Life Support Guidelines. Retrieved from Resuscitation Council UK website: <https://www.resus.org.uk/library/2021-resuscitation-guidelines/paediatric-advanced-life-support-guidelines>
2. Ambu UK. (2025, January 15). AuraGain Data Sheet 2025. Retrieved from Ambu UK website: [https://www.ambu.co.uk/Admin/Public/Download.aspx?file=Files%2FFiles%2FDownloads%2FAmbu+UK%2FAnaesthesia%2FLaryngeal\\_Masks%2FAuraGain+Disposable+Laryngeal+Mask%2FDatasheets%2FDatasheet-AuraGain.pdf](https://www.ambu.co.uk/Admin/Public/Download.aspx?file=Files%2FFiles%2FDownloads%2FAmbu+UK%2FAnaesthesia%2FLaryngeal_Masks%2FAuraGain+Disposable+Laryngeal+Mask%2FDatasheets%2FDatasheet-AuraGain.pdf)
3. Southampton Oxford Retrieval Team. (2024). SORT Drug Calculator. Retrieved from www.sort.nhs.uk website: <https://www.sort.nhs.uk/Guidelines/Drug-calculator.aspx>
4. National Institute for Health and Care Excellence (NICE). (2025). Atropine Sulphate. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/atropine-sulfate/>
5. NICE. (2025). Glycopyrronium Bromide. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/glycopyrronium-bromide/>
6. NICE. (2024). Suxamethonium Chloride. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/suxamethonium-chloride/>
7. NICE. (2025). Adrenaline. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/adrenaline-epinephrine/>
8. Royal College of Paediatrics and Child Health. (2020). Safe Prescription of IV Fluids Workbook. Retrieved from Rcpch.ac.uk website: [https://qicentral.rcpch.ac.uk/wp-content/uploads/sites/9/2020/08/iv\\_fluid\\_workbook\\_0.docx](https://qicentral.rcpch.ac.uk/wp-content/uploads/sites/9/2020/08/iv_fluid_workbook_0.docx)
9. Resuscitation Council UK. (2024). Paediatric emergency algorithms & resources. Retrieved from <https://www.resus.org.uk/sites/default/files/2024-03/RCUK%20Paediatric%20emergency%20algortihms%20and%20resources%20Mar%202024%20V2.2.pdf>
10. NICE. (2025c). Rocuronium Bromide. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/rocuronium-bromide/>
11. NICE. (2025c). Ketamine Hydrochloride. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/ketamine/>
12. NICE. (2022). Fentanyl. Retrieved from NICE website: <https://bnfc.nice.org.uk/drugs/fentanyl/>