

Eidgenössische Technische Hochschule Zürich Swiss Federal Institute of Technology Zurich

Exercise - Even matrices

You are still part of a team to develop a pseudorandom number generator. Your generator has already passed some simple statistical tests, but now it is time to get serious. You arrange a list of bits produced by the generator in an n by n matrix. If

$$M = \begin{pmatrix} x_{1,1} & x_{1,2} & & x_{1,n} \\ x_{2,1} & x_{2,2} & \cdots & x_{2,n} \\ & \vdots & \ddots & \\ x_{n,1} & x_{n,2} & & x_{n,n} \end{pmatrix}$$

was a truly random matrix of bits, then it would have the property that the sum

$$\sum_{i'=i_1}^{i_2} \sum_{j'=j_1}^{j_2} x_{i',j'}$$

is even for about half of the quadruples (i_1,i_2,j_1,j_2) , $1 \leqslant i_1 \leqslant i_2 \leqslant n$ and $1 \leqslant j_1 \leqslant j_2 \leqslant n$.

To check whether this is the case for your generator, you need to be able to count the number of such quadruples.

Input The first line of the input contains the number $t \le 15$ of test cases. Each of the t test cases is described as follows.

- It starts with a line that contains an integer n such that $1 \le n \le 200$.
- This is followed by n lines, where the i-th line contains the n integers $x_{i,1}$... $x_{i,n}$, separated by a space, such that $x_{i,j} \in \{0,1\}$, for all $i \in \{1,\ldots,n\}$ and $j \in \{1,\ldots,n\}$.

Output For each test case output a single line that contains the number of quadruples (i_1, i_2, j_1, j_2) where $1 \le i_1 \le i_2 \le n$ and $1 \le j_1 \le j_2 \le n$ and for which the sum

$$\sum_{i'=i_1}^{i_2} \sum_{j'=j_1}^{j_2} x_{i',j'}$$

is even.

Points There are three groups of test sets, worth 100 points in total.

- 1. For the first group of test sets, worth 30 points, you may assume that $1 \le n \le 10$.
- 2. For the second group of test sets, worth 50 points, you may assume that $1 \le n \le 50$.
- 3. For the third group of test set, worth 20 points, there are no additional assumptions.

Corresponding sample test sets are contained in testi.in/out for $i \in \{1,2,3\}$.

Sample Input

3 2 1 1 1 1 3 1 0 1 0 1 0 0 0 1 4 1 1 0 0 0 0 1 1 1 0 1 0 0 1 0 1

Sample Output

5 15 52