

EXERCISE 6A.3

- 1 **a** quotient is $x + 1$, remainder is $-x - 4$
b quotient is 3, remainder is $-x + 3$
c quotient is $3x$, remainder is $-2x - 1$
d quotient is 0, remainder is $x - 4$
- 2 **a** $1 - \frac{2x}{x^2 + x + 1}$, $x^2 - x + 1 = 1(x^2 + x + 1) - 2x$
b $x - \frac{2x}{x^2 + 2}$, $x^3 = x(x^2 + 2) - 2x$
c $x^2 + x + 3 + \frac{3x - 4}{x^2 - x + 1}$,
 $x^4 + 3x^2 + x - 1 = (x^2 + x + 3)(x^2 - x + 1) + 3x - 4$
d $2x + 4 + \frac{5x + 2}{(x - 1)^2}$,
 $2x^3 - x + 6 = (2x + 4)(x - 1)^2 + 5x + 2$
e $x^2 - 2x + 3 - \frac{4x + 3}{(x + 1)^2}$,
 $x^4 = (x^2 - 2x + 3)(x + 1)^2 - 4x - 3$
f $x^2 - 3x + 5 + \frac{15 - 10x}{(x - 1)(x + 2)}$,
 $x^4 - 2x^3 + x + 5 = (x^2 - 3x + 5)(x - 1)(x + 2) + 15 - 10x$
- 3 quotient is $x^2 + 2x + 3$, remainder is 7
4 quotient is $x^2 - 3x + 5$, remainder is $15 - 10x$

EXERCISE 6B.1

- 1 **a** $4, -\frac{3}{2}$ **b** $-3 \pm \sqrt{10}$ **c** $5 \pm \sqrt{19}$
d $0, \pm 2$ **e** $0, \pm \sqrt{11}$ **f** $\pm 2, \pm \sqrt{2}$
- 2 **a** $1, -\frac{2}{5}$ **b** $-\frac{1}{2}, \pm \sqrt{3}$ **c** $-3, \frac{1}{3}, 2$
d $0, 1 \pm \sqrt{3}$ **e** $0, \pm \sqrt{7}$ **f** $\pm \sqrt{2}, \pm \sqrt{5}$
- 3 **a** $(2x + 3)(x - 5)$ **b** $x(x - 7)(x - 4)$
c $(x - 3 - \sqrt{6})(x - 3 + \sqrt{6})$
d $x(x + 1 + \sqrt{5})(x + 1 - \sqrt{5})$ **e** $x(3x - 2)(2x + 1)$
f $(x + 1)(x - 1)(x + \sqrt{5})(x - \sqrt{5})$
- 4 $P(\alpha) = 0$, $P(\beta) = 0$, $P(\gamma) = 0$
- 5 **a** $P(x) = a(x + 3)(x - 4)(x - 5)$, $a \neq 0$
b $P(x) = a(x + 2)(x - 2)(x - 3)$, $a \neq 0$
c $P(x) = a(x - 3)(x^2 - 2x - 4)$, $a \neq 0$
d $P(x) = a(x + 1)(x^2 + 4x + 2)$, $a \neq 0$
- 6 **a** $P(x) = a(x^2 - 1)(x^2 - 2)$, $a \neq 0$
b $P(x) = a(x - 2)(5x + 1)(x^2 - 3)$, $a \neq 0$
c $P(x) = a(x + 3)(4x - 1)(x^2 - 2x - 1)$, $a \neq 0$
d $P(x) = a(x^2 - 4x - 1)(x^2 + 4x - 3)$, $a \neq 0$

EXERCISE 6B.2

- 1 **a** $a = 2$, $b = 5$, $c = 5$ **b** $a = 3$, $b = 4$, $c = 3$
c $a = 2$, $b = -5$, $c = 4$
- 2 **a** $a = 2$, $b = -2$ or $a = -2$, $b = 2$
b $a = 3$, $b = -1$
- 3 **a** $a = 1$, $b = 6$, $c = -7$ **b** $(x + 3)(x + 7)(x - 1)$
- 4 **a** $p = 2$, $q = 7$, $r = 5$ **b** $x = \frac{1}{2}, -1, -\frac{5}{2}$
- 5 **a** $a = 3$, $b = -2$, $c = 1$
b $3x^3 + 10x^2 - 7x + 4 = (x + 4)(3x^2 - 2x + 1)$
 Δ of $3x^2 - 2x + 1$ is -8 ,
 \therefore the only real zero is -4 .

6 **a** $a = 1$, $b = -2$, $c = -1$, $k = -4$

b $-\frac{2}{3}, 1 \pm \sqrt{2}$

7 **a** $a = -2$, $b = 2$ **b** $-1 \pm \sqrt{3}$

8 $a = -11$, zeros are $\frac{3}{2}, \frac{-3 \pm \sqrt{13}}{2}$

9 **a** $a = -9$, $b = -1$

b $P(x) = 0$ when $x = -1, -\frac{1}{2}, 2, 4$

- 10 **Hint:** Let $x^3 + 3x^2 - 9x + c = (x + a)^2(x + b)$
 When $c = 5$, the cubic is $(x - 1)^2(x + 5)$.
 When $c = -27$, the cubic is $(x + 3)^2(x - 3)$.

EXERCISE 6C

- 1 **a** $P(x) = Q(x)(x - 2) + 7$, $P(x)$ divided by $x - 2$ leaves a remainder of 7.
b $P(-3) = -8$, $P(x)$ divided by $x + 3$ leaves a remainder of -8 .
c $P(5) = 11$, $P(x) = Q(x)(x - 5) + 11$
- 2 **a** 4 **b** -19 **c** 1 **3** 4
- 4 **a** $a = 3$ **b** $a = 2$ **5** $a = -5$, $b = 6$
- 6 $a = -5$, $b = 6$ **7** -7
- 8 **a** $P(x) = Q(x)(2x - 1) + R$
 $P(\frac{1}{2}) = Q(\frac{1}{2})(2 \times \frac{1}{2} - 1) + R$
 $= Q(\frac{1}{2}) \times 0 + R$
 $= R$

b **i** -3 **ii** 7 **iii** -7

9 $a = 3$, $b = 10$ **10** **a** -3 **b** 1

EXERCISE 6D

- 1 **a** factor **b** not a factor **c** factor **d** not a factor
- 2 **a** $c = 2$ **b** $c = -2$ **c** $b = 3$
- 3 $k = -8$, $P(x) = (x + 2)(x - 2)(2x + 1)$
- 4 **a** $k = -8$ **b** $P(x) = (x - 3)(3x^2 + x - 2)$
c $x = -1, \frac{2}{3}, 3$
- 5 $a = 7$, $b = -14$ **6** $a = 3$, $b = 2$
- 7 **a** $a = 7$, $b = -6$ **b** 60
c $P(x) = (x + 3)(2x^2 + 3x - 2)$ **d** $-3, -2, \frac{1}{2}$
- 8 **a** $a = 7$, $b = 2$ **b** $x = -2 \pm \sqrt{6}$
- 9 **a** **i** $P(a) = 0$, $x - a$ is a factor
ii $(x - a)(x^2 + ax + a^2)$
b **i** $P(-a) = 0$, $x + a$ is a factor
ii $(x + a)(x^2 - ax + a^2)$
- 10 $a = 2$

EXERCISE 6E

- 1 **a** $x = 1, 2, 3$ **b** $x = -1, 2$ {2 is a double root}
c $x = 1, -1, -2$ **d** $x = -1, 3, 4$ **e** $x = -5, -4, 4$
f $x = -3, -5$ {-5 is a double root}
- 2 **a** $x = -2, 2, 3$ **b** $x = -3, -2, 6$ **c** $x = -3, 4, 7$

REVIEW SET 6A

- 1 **a** $8x^2 + 6x + 3$ **b** $7x^2 - 9x + 9$
c $15x^4 + 32x^3 + 29x - 4$
- 2 **a** quotient = $2x + 5$, remainder = 3
b quotient = $x^2 - 4x + 2$, remainder = -5
- 3 **a** $\frac{4}{3}, -2$ **b** $-4 \pm \sqrt{5}$

- 4 a** $a = 1$, $b = -2$, $c = 3$
b Δ of $x^2 - 2x + 3$ is -8
 \therefore the only real root is $x = -3$.
5 a 1 **b** -53 **6 a** not a factor **b** factor
7 k = 6 **8 a** = 4, $b = -1$ **9 c** = 3
10 a $a = -19$, $b = -20$ **b** -5 , -1 , 4
11 x = -3 , -1 , 5

REVIEW SET 6B

- 1 a** $12x^4 - 9x^3 + 8x^2 - 26x + 15$
b $4x^4 - 4x^3 + 13x^2 - 6x + 9$
2 a $x^2 - 2x + 4 - \frac{8}{x+2}$ **b** $x - 5 + \frac{19x+30}{(x+2)(x+3)}$
3 $P(x) = a(4x-1)(x^2-2x-4)$, $a \neq 0$
4 For $k = 3$, $b = 27$, $x = 3$ or -3 .
 For $k = -1$, $b = -5$, $x = -1$ or 5 .
5 a -3 **b** -7 **6 a** $a = 5$ **b** -12
7 b $(x-2)(x^2+2x-9)$ **c** $2, -1 \pm \sqrt{10}$
8 a $\frac{8}{7}$, $b = \frac{174}{7}$
9 k = 8, the zeros are $-1, -2$ $\{-2$ is a double root $\}$
10 a $a = -20$, $b = 12$ **b** $f(x) = (2x-1)(x-6)(x+2)$
11 x = $-4, 2, 3$

EXERCISE 7A.1

- 1 a** gradient = 3, y -intercept is 5
b gradient = 4, y -intercept is -2
c gradient = $\frac{1}{5}$, y -intercept is $\frac{3}{5}$
d gradient = -7 , y -intercept is -3
e gradient = $\frac{1}{6}$, y -intercept is $\frac{1}{3}$
f gradient = $-\frac{5}{3}$, y -intercept is $\frac{8}{3}$
2 a $y = x - 2$ **b** $y = -x + 4$ **c** $y = 2x$
d $y = -\frac{1}{2}x + 3$
3 a $y = 4x - 13$ **b** $y = -3x - 5$ **c** $y = -5x + 32$
d $y = \frac{1}{2}x + \frac{7}{2}$ **e** $y = -\frac{1}{3}x + \frac{8}{3}$ **f** $y = 6$
4 a $2x - 3y = -11$ **b** $3x - 5y = -23$ **c** $x + 3y = 5$
d $2x + 7y = -2$ **e** $4x - y = -11$ **f** $2x + y = 7$
g $7x + 2y = 18$ **h** $6x - y = -40$
5 a $y = \frac{5}{2}x - 2$ **b** $y = -2x + 3$ **c** $y = -2$
d $y = -\frac{1}{5}x + \frac{2}{5}$ **e** $y = \frac{1}{6}x - \frac{11}{6}$ **f** $y = -\frac{2}{3}x - \frac{11}{3}$
6 a $x - 3y = -3$ **b** $5x - y = 1$ **c** $x - y = 3$
d $4x - 5y = 10$ **e** $x - 2y = -1$ **f** $2x + 3y = -5$
7 a $\sqrt{45}$ units **b** $(-1, \frac{7}{2})$ **c** $\frac{1}{2}$ **d** $y = \frac{1}{2}x + 4$
8 a $y = \frac{4}{3}x - 1$ **b** $2x - 3y = -13$ **c** $y = x + 1$
d $2x + y = -2$ **e** $y = -\frac{2}{3}x + 2$ **f** $3x + 7y = -9$
9 a $M = \frac{1}{3}p + 2$ **b** $R = -\frac{5}{4}n + 2$ **c** $T = \frac{1}{2}x - 1$
d $F = \frac{1}{10}x + 1$ **e** $H = -\frac{1}{2}z + 2$ **f** $W = -\frac{1}{6}t - 2$
10 a $x + 2y = 13$ **b** $(13, 0)$
11 a $3x + 5y = 10$ **b** $(0, 2)$ **12** 54 units²

EXERCISE 7A.2

- 1 a** $\sqrt{160}$ units **b** $(-1, 1)$ **c** -3 **d** $x - 3y = -4$
2 a $y = x - 4$ **b** $y = 2x + 6$ **c** $y = \frac{6}{5}x + \frac{7}{2}$ **d** $y = 1$
3 15 units²

EXERCISE 7B

- 1 a** $(1, 3)$ **b** $(6, -3)$ **c** $(-5, 3)$ **d** $(-1, -2)$
2 a $3x + 5y = 9$ **b** $(-2, 3)$ **3** $(4, 2)$
4 a $x - 3y = -8$ **b** $y = -3x - 4$ **c** $(-2, 2)$
5 a $(0, -1)$ **b** 25 units²
6 a $(-1, 0)$ **b** 26 units² **7** 30 units²
8 a i $(5, 0)$ **ii** $(7, -4)$ **iii** $(6, -2)$
b Hint: Find the gradients of MN and AC.
c i 15 units² **ii** 20 units²

EXERCISE 7C

- 1** $(-1, -2)$ and $(\frac{11}{5}, -\frac{2}{5})$ **2** $\sqrt{18}$ units
3 $x - 2y = 0$ **4** $(-\frac{4}{3}, -\frac{8}{3})$ and $(2, -1)$
5 $\sqrt{125}$ units **6** $x - 3y = -13$
7 $(3, -\frac{3}{2})$ and $(4, -1)$ **8** $(\frac{7}{3}, \frac{5}{2})$

EXERCISE 7D


- 1 a** $y = \frac{1}{2}x^3 + 2$ **b** $y = 3\sqrt{x} - 1$, $x \geq 0$
c $y = 3 - x^4$ **d** $y = \frac{1}{3} \times 2^x$
e $y = \frac{2}{x} + 1$ **f** $y = -\frac{3}{2} \times 3^x + 11$
2 a i $y = x^2 + 3x$ **ii** $y = 18$
b i $y = -\frac{1}{2}\sqrt{x} + \frac{10}{\sqrt{x}}$, $x > 0$ **ii** $y = \frac{17\sqrt{3}}{6}$
c i $y = \frac{5}{3x} \times 2^x$ **ii** $y = \frac{40}{9}$
d i $y = 2x^3 - 9x$ **ii** $y = 27$
e i $y = \frac{1}{x^2} - \frac{12}{x} + 36$ **ii** $y = 32\frac{1}{9}$
f i $y = (x+2)^2 + 3$ **ii** $y = 28$
3 a $\lg y = 2x - 1$ **b** $y = \frac{1}{10} \times 10^{2x}$
4 $y = 1000 \times 10^{-\frac{3}{2}x}$
5 a $y = \frac{1}{10000} \times 10^x$ **b** $y = 10000 \times (\frac{1}{10})^x$
c $y = 5 \times 4^x$
6 a $y = 10 \times 10^{\frac{1}{3}x}$ **b** $y = 1000$
7 a $\lg y = -\frac{1}{2} \lg x + 2$ **b** $y = \frac{100}{\sqrt{x}}$
8 a $y = x^{\frac{1}{4}}$ **b** $y = \frac{1000}{x}$ **c** $y = x^2 \sqrt{1000}$
9 a $K = 7\sqrt{t}$ **b** $K = 21$ **10 a** 3 **b** $\lg 4$

EXERCISE 7E

- 1 a**


x^2	1	4	9	16
y	2	11	26	47

c $y = 3x^2 - 1$



2 a


x	1	2	3	4
$y\sqrt{x}$	9	14	19	24



c $y = 5\sqrt{x} + \frac{4}{\sqrt{x}}, x > 0$
d $y = 21$

3 a


$\frac{1}{x}$	1	0.5	0.33	0.25
xy	-1	0	0.33	0.48



c $a = 1, b = -2$
d $y = 0.08$

4 a


$x\sqrt{x}$	2.83	8	14.70	22.63
$y\sqrt{x}$	7.41	10	13.35	17.31




c $y = \frac{1}{2}x + \frac{6}{\sqrt{x}}, x > 0$ **d** $y = 6.5$

5 a

t	1	2	3	4	5
$\lg M$	0.600	0.800	1	1.20	1.40



c $M \approx 2.51 \times 1.58^t$
d $\approx 2.51 \text{ g}$



7 Plot xy against \sqrt{x} . $y = \frac{8}{x} - \frac{4}{\sqrt{x}}$ $\{a = 8, b = -4\}$

8 a $a \approx 4.90, b \approx 2.00$ **b** $\approx 44.1 \text{ m}$ **c** $\approx 4.04 \text{ seconds}$

REVIEW SET 7A

1 a $\sqrt{40}$ units **b** $(2, 5)$ **c** $x + 3y = 17$

2 $y = -2x + 6$

3 The gradient of a vertical line is undefined.

4 a $x + 2y = 7$ **b** $(7, 0)$ **5** $(3, -1)$

6 a $(-1, 4)$ **b** $32\frac{1}{2}$ units²


7 $(-\frac{7}{5}, \frac{26}{5})$ and $(2, -5)$ **8** $y = x - 5$

10 a $y = \frac{3}{\sqrt{x}} - \frac{2}{x}, x > 0$ **b** $y = 1$

11 a $\lg y = \frac{1}{2} \lg x + 1$ **b** $y = 10\sqrt{x}$

12 a

x^3	1	8	27	64
xy	8	15	33.99	71



c $y = x^2 + \frac{7}{x}$
d $y = 50$

REVIEW SET 7B

1 $y = -3x + 7$ **2** $\sqrt{80}$ units

3 a $y = 5x - \frac{2}{x}$ **b** $39\frac{3}{4}$


4 a $r = \frac{5}{7}a + 2$ **b** $K = \frac{3}{5}s + 3$

5 $(3, 2)$ **6** $5x - 8y = 31$

7 a

\sqrt{x}	1	1.41	1.73	2
$\frac{y}{\sqrt{x}}$	2.5	3.74	4.69	5.5

c $y = 3x - \frac{\sqrt{x}}{2}$



8 a $bx + ay = ab$

b Hint: $\cos \theta = \frac{a}{\sqrt{a^2 + b^2}}, \sin \theta = \frac{b}{\sqrt{a^2 + b^2}}$

9 a i $(2, 12)$ **ii** $(11, 0)$ **b** 75 units^2

10 $(\frac{7}{8}, \frac{1}{2})$

11 a Plot $\lg y$ against x .

$y = 100 \times (10^{-\frac{1}{3}})^x \quad \{a = 100, b = 10^{-\frac{1}{3}}\}$

b $y \approx 46.4$

EXERCISE 8A

- 1 a** $\frac{\pi}{2}^c$ **b** $\frac{\pi}{3}^c$ **c** $\frac{\pi}{6}^c$ **d** $\frac{\pi}{10}^c$ **e** $\frac{\pi}{20}^c$
f $\frac{3\pi}{4}^c$ **g** $\frac{5\pi}{4}^c$ **h** $\frac{3\pi}{2}^c$ **i** $2\pi^c$ **j** $4\pi^c$
k $\frac{7\pi}{4}^c$ **l** $3\pi^c$ **m** $\frac{\pi}{5}^c$ **n** $\frac{4\pi}{9}^c$ **o** $\frac{23\pi}{18}^c$
2 a 0.641^c **b** 2.39^c **c** 5.55^c **d** 3.83^c **e** 6.92^c
3 a 36° **b** 108° **c** 135° **d** 10° **e** 20°
f 140° **g** 18° **h** 27° **i** 210° **j** 22.5°
4 a 114.59° **b** 87.66° **c** 49.68° **d** 182.14°
e 301.78°

5 a	Degrees	0	45	90	135	180	225	270	315	360
	Radians	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π	$\frac{5\pi}{4}$	$\frac{3\pi}{2}$	$\frac{7\pi}{4}$	2π

b	Deg.	0	30	60	90	120	150	180	210	240	270	300	330	360
	Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	$\frac{11\pi}{6}$	2π

EXERCISE 8B

- 1 a** 49.5 cm , 223 cm^2 **b** 23.0 cm , 56.8 cm^2
2 a 3.14 m **b** 9.30 m^2 **3 a** 5.91 cm **b** 18.9 cm
4 a 0.686^c **b** 0.6^c
5 a $\theta = 0.75^c$, area = 24 cm^2
b $\theta = 1.68^c$, area = 21 cm^2
c $\theta \approx 2.32^c$, area = 126.8 cm^2
6 10 cm , 25 cm^2
8 a 11.7 cm **b** $r \approx 11.7$ **c** 37.7 cm **d** 3.23^c
9 a $\alpha \approx 18.43$ **b** $\theta \approx 143.1$ **c** 387 m^2
10 25.9 cm **11 b** $2 \text{ h } 49 \text{ min}$ **12** 227 m^2

EXERCISE 8C


- 1 a i** $A(\cos 26^\circ, \sin 26^\circ)$, $B(\cos 146^\circ, \sin 146^\circ)$, $C(\cos 199^\circ, \sin 199^\circ)$
ii $A(0.899, 0.438)$, $B(-0.829, 0.559)$, $C(-0.946, -0.326)$
b i $A(\cos 123^\circ, \sin 123^\circ)$, $B(\cos 251^\circ, \sin 251^\circ)$, $C(\cos(-35^\circ), \sin(-35^\circ))$
ii $A(-0.545, 0.839)$, $B(-0.326, -0.946)$, $C(0.819, -0.574)$

2	θ (degrees)	0°	90°	180°	270°	360°	450°
	θ (radians)	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π	$\frac{5\pi}{2}$
	sine	0	1	0	-1	0	1
	cosine	1	0	-1	0	1	0
	tangent	0	undef	0	undef	0	undef


- 3 a i** $\frac{1}{\sqrt{2}} \approx 0.707$ **ii** $\frac{\sqrt{3}}{2} \approx 0.866$

b	θ (degrees)	30°	45°	60°	135°	150°	240°	315°
	θ (radians)	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\frac{4\pi}{3}$	$\frac{7\pi}{4}$
	sine	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{\sqrt{2}}$
	cosine	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	$-\frac{1}{\sqrt{2}}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$\frac{1}{\sqrt{2}}$
	tangent	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	-1	$-\frac{1}{\sqrt{3}}$	$\sqrt{3}$	-1

- 4 a i** 0.985 **ii** 0.985 **iii** 0.866 **iv** 0.866
v 0.5 **vi** 0.5 **vii** 0.707 **viii** 0.707
b $\sin(180^\circ - \theta) = \sin \theta$
c $\sin \theta$ and $\sin(180^\circ - \theta)$ have the same value, as P and Q have the same y-coordinate.



- d i** 135° **ii** 129° **iii** $\frac{2\pi}{3}$ **iv** $\frac{5\pi}{6}$
5 a i 0.342 **ii** -0.342 **iii** 0.5 **iv** -0.5
v 0.906 **vi** -0.906 **vii** 0.174 **viii** -0.174
b $\cos(180^\circ - \theta) = -\cos \theta$
c $\cos(180^\circ - \theta) = -\cos \theta$, as the x-coordinates of P and Q are negatives of each other.



- d i** 140° **ii** 161° **iii** $\frac{4\pi}{5}$ **iv** $\frac{3\pi}{5}$
6 a ≈ 0.6820 **b** ≈ 0.8572 **c** ≈ -0.7986
d ≈ 0.9135 **e** ≈ 0.9063 **f** ≈ -0.6691
7 a

Quadrant	Degree measure	Radian measure	$\cos \theta$	$\sin \theta$	$\tan \theta$
1	$0^\circ < \theta < 90^\circ$	$0 < \theta < \frac{\pi}{2}$	+ve	+ve	+ve
2	$90^\circ < \theta < 180^\circ$	$\frac{\pi}{2} < \theta < \pi$	-ve	+ve	-ve
3	$180^\circ < \theta < 270^\circ$	$\pi < \theta < \frac{3\pi}{2}$	-ve	-ve	+ve
4	$270^\circ < \theta < 360^\circ$	$\frac{3\pi}{2} < \theta < 2\pi$	+ve	-ve	-ve

- b i** 1 and 4 **ii** 2 and 3 **iii** 3 **iv** 2
8 a $\widehat{AOQ} = 180^\circ - \theta$ or $\pi - \theta$ radians
b [OQ] is a reflection of [OP] in the y-axis and so Q has coordinates $(-\cos \theta, \sin \theta)$.
c $\cos(180^\circ - \theta) = -\cos \theta$, $\sin(180^\circ - \theta) = \sin \theta$

9 a	θ°	$\sin \theta$	$\sin(-\theta)$	$\cos \theta$	$\cos(-\theta)$
	0.75	0.682	-0.682	0.732	0.732
	1.772	0.980	-0.980	-0.200	-0.200
	3.414	-0.269	0.269	-0.963	-0.963
	6.25	-0.0332	0.0332	0.999	0.999
	-1.17	-0.921	0.921	0.390	0.390

- b** $\sin(-\theta) = -\sin \theta$, $\cos(-\theta) = \cos \theta$

EXERCISE 8D.1

- 1 a** $\cos \theta = \pm \frac{\sqrt{3}}{2}$ **b** $\cos \theta = \pm \frac{2\sqrt{2}}{3}$ **c** $\cos \theta = \pm 1$
d $\cos \theta = 0$

- 2 **a** $\sin \theta = \pm \frac{3}{5}$ **b** $\sin \theta = \pm \frac{\sqrt{7}}{4}$ **c** $\sin \theta = 0$
d $\sin \theta = \pm 1$
- 3 **a** $\sin \theta = \frac{\sqrt{5}}{3}$ **b** $\cos \theta = -\frac{\sqrt{21}}{5}$ **c** $\cos \theta = \frac{4}{5}$
d $\sin \theta = -\frac{12}{13}$
- 4 **a** $\tan \theta = -\frac{1}{2\sqrt{2}}$ **b** $\tan \theta = -2\sqrt{6}$ **c** $\tan \theta = \frac{1}{\sqrt{2}}$
d $\tan \theta = -\frac{\sqrt{7}}{3}$
- 5 **a** $\sin x = \frac{2}{\sqrt{13}}$, $\cos x = \frac{3}{\sqrt{13}}$
b $\sin x = \frac{4}{5}$, $\cos x = -\frac{3}{5}$
c $\sin x = -\sqrt{\frac{5}{14}}$, $\cos x = -\frac{3}{\sqrt{14}}$
d $\sin x = -\frac{12}{13}$, $\cos x = \frac{5}{13}$
- 6 $\sin \theta = \frac{-k}{\sqrt{k^2+1}}$, $\cos \theta = \frac{-1}{\sqrt{k^2+1}}$

EXERCISE 8D.2

- 1 **a** $\theta \approx 1.33$ or 4.47 **b** $\theta \approx 0.592$ or 5.69
c $\theta \approx 0.644$ or 2.50 **d** $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2}$
e $\theta \approx 0.876$ or 4.02 **f** $\theta \approx 0.674$ or 5.61
g $\theta \approx 0.0910$ or 3.05 **h** $\theta \approx 1.52$ or 4.66
i $\theta \approx 1.35$ or 1.79
- 2 **a** $\theta \approx 1.82$ or 4.46 **b** $\theta = 0, \pi$, or 2π
c $\theta \approx 1.88$ or 5.02 **d** $\theta \approx 3.58$ or 5.85
e $\theta \approx 1.72$ or 4.86 **f** $\theta \approx 1.69$ or 4.59
g $\theta \approx 1.99$ or 5.13 **h** $\theta \approx 2.19$ or 4.10
i $\theta \approx 3.83$ or 5.60

EXERCISE 8E

- 1
- | | a | b | c | d | e |
|---------------|----------------------|----------|-----------------------|----------|-----------------------|
| $\sin \theta$ | $\frac{1}{\sqrt{2}}$ | 1 | $-\frac{1}{\sqrt{2}}$ | 0 | $-\frac{1}{\sqrt{2}}$ |
| $\cos \theta$ | $\frac{1}{\sqrt{2}}$ | 0 | $\frac{1}{\sqrt{2}}$ | -1 | $-\frac{1}{\sqrt{2}}$ |
| $\tan \theta$ | 1 | undef | -1 | 0 | 1 |
- 2
- | | a | b | c | d | e |
|--------------|----------------------|----------------------|-----------------------|-----------------------|-----------------------|
| $\sin \beta$ | $\frac{1}{2}$ | $\frac{\sqrt{3}}{2}$ | $-\frac{1}{2}$ | $-\frac{\sqrt{3}}{2}$ | $-\frac{1}{2}$ |
| $\cos \beta$ | $\frac{\sqrt{3}}{2}$ | $-\frac{1}{2}$ | $-\frac{\sqrt{3}}{2}$ | $\frac{1}{2}$ | $\frac{\sqrt{3}}{2}$ |
| $\tan \beta$ | $\frac{1}{\sqrt{3}}$ | $-\sqrt{3}$ | $\frac{1}{\sqrt{3}}$ | $-\sqrt{3}$ | $-\frac{1}{\sqrt{3}}$ |
- 3 **a** $\cos 120^\circ = -\frac{1}{2}$, $\sin 120^\circ = \frac{\sqrt{3}}{2}$, $\tan 120^\circ = -\sqrt{3}$
b $\cos(-45^\circ) = \frac{1}{\sqrt{2}}$, $\sin(-45^\circ) = -\frac{1}{\sqrt{2}}$, $\tan(-45^\circ) = -1$
- 4 **a** $\cos 270^\circ = 0$, $\sin 270^\circ = -1$
b $\tan 270^\circ$ is undefined
- 5 **a** $\frac{3}{4}$ **b** $\frac{1}{4}$ **c** 3 **d** $\frac{1}{4}$ **e** $-\frac{1}{4}$ **f** 1
g $\sqrt{2}$ **h** $\frac{1}{2}$ **i** $\frac{1}{2}$ **j** 2 **k** -1 **l** $-\sqrt{3}$
- 6 **a** $30^\circ, 150^\circ$ **b** $60^\circ, 120^\circ$ **c** $45^\circ, 315^\circ$
d $120^\circ, 240^\circ$ **e** $135^\circ, 225^\circ$ **f** $240^\circ, 300^\circ$
- 7 **a** $\frac{\pi}{4}, \frac{5\pi}{4}$ **b** $\frac{3\pi}{4}, \frac{7\pi}{4}$ **c** $\frac{\pi}{3}, \frac{4\pi}{3}$
d $0, \pi, 2\pi$ **e** $\frac{\pi}{6}, \frac{7\pi}{6}$ **f** $\frac{2\pi}{3}, \frac{5\pi}{3}$
- 8 **a** $\frac{\pi}{6}, \frac{11\pi}{6}, \frac{13\pi}{6}, \frac{23\pi}{6}$ **b** $\frac{7\pi}{6}, \frac{11\pi}{6}, \frac{19\pi}{6}, \frac{23\pi}{6}$ **c** $\frac{3\pi}{2}, \frac{7\pi}{2}$
- 9 **a** $\theta = \frac{\pi}{3}, \frac{5\pi}{3}$ **b** $\theta = \frac{\pi}{3}, \frac{2\pi}{3}$ **c** $\theta = \pi$
d $\theta = \frac{\pi}{2}$ **e** $\theta = \frac{3\pi}{4}, \frac{5\pi}{4}$ **f** $\theta = \frac{\pi}{2}, \frac{3\pi}{2}$

- g** $\theta = 0, \pi, 2\pi$ **h** $\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$
i $\theta = \frac{5\pi}{6}, \frac{11\pi}{6}$ **j** $\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$
- 10 **a** $\theta = k\pi, k \in \mathbb{Z}$ **b** $\theta = \frac{\pi}{2} + k\pi, k \in \mathbb{Z}$

EXERCISE 8F

- 1 **a** $\frac{2}{\sqrt{3}}$ **b** $-\frac{1}{\sqrt{3}}$ **c** $-\frac{2}{\sqrt{3}}$ **d** undefined
e $-\frac{2}{\sqrt{3}}$ **f** $\sqrt{2}$
- 2 **a** $\operatorname{cosec} x = \frac{5}{3}$, $\sec x = \frac{5}{4}$, $\cot x = \frac{4}{3}$
b $\operatorname{cosec} x = -\frac{3}{\sqrt{5}}$, $\sec x = \frac{3}{2}$, $\cot x = -\frac{2}{\sqrt{5}}$
- 3 **a** $\sin \theta = -\frac{\sqrt{7}}{4}$, $\tan \theta = -\frac{\sqrt{7}}{3}$, $\operatorname{cosec} \theta = -\frac{4}{\sqrt{7}}$,
 $\sec \theta = \frac{4}{3}$, $\cot \theta = -\frac{3}{\sqrt{7}}$
b $\cos x = -\frac{\sqrt{5}}{3}$, $\tan x = \frac{2}{\sqrt{5}}$, $\operatorname{cosec} x = -\frac{3}{2}$,
 $\sec x = -\frac{3}{\sqrt{5}}$, $\cot x = \frac{\sqrt{5}}{2}$
c $\sin x = \frac{\sqrt{21}}{5}$, $\cos x = \frac{2}{5}$, $\tan x = \frac{\sqrt{21}}{2}$,
 $\operatorname{cosec} x = \frac{5}{\sqrt{21}}$, $\cot x = \frac{2}{\sqrt{21}}$
d $\sin \theta = \frac{1}{2}$, $\cos \theta = -\frac{\sqrt{3}}{2}$, $\tan \theta = -\frac{1}{\sqrt{3}}$,
 $\sec \theta = -\frac{2}{\sqrt{3}}$, $\cot \theta = -\sqrt{3}$
e $\sin \beta = -\frac{1}{\sqrt{5}}$, $\cos \beta = -\frac{2}{\sqrt{5}}$, $\operatorname{cosec} \beta = -\sqrt{5}$,
 $\sec \beta = -\frac{\sqrt{5}}{2}$, $\cot \beta = 2$
f $\sin \theta = -\frac{3}{5}$, $\cos \theta = -\frac{4}{5}$, $\tan \theta = \frac{3}{4}$,
 $\operatorname{cosec} \theta = -\frac{5}{3}$, $\sec \theta = -\frac{5}{4}$
- 4 **a** $\theta = k\pi, k \in \mathbb{Z}$ **b** $\theta = \frac{\pi}{2} + k\pi, k \in \mathbb{Z}$
c $\theta = \frac{\pi}{2} + k\pi, k \in \mathbb{Z}$ **d** $\theta = k\pi, k \in \mathbb{Z}$


REVIEW SET 8A

- 1 **a** $\frac{2\pi}{3}$ **b** $\frac{5\pi}{4}$ **c** $\frac{5\pi}{6}$ **d** 3π
- 2 **a** $\frac{\pi}{3}$ **b** 15° **c** 84°
- 3 **a** 0.358 **b** -0.035 **c** 0.259 **d** -0.731
- 4 111 cm² **5** $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$
- 6
- | | a | b | c | d |
|---------------|----------|----------------------|----------|----------------------|
| $\sin \theta$ | 0 | $\frac{\sqrt{3}}{2}$ | 0 | $\frac{\sqrt{3}}{2}$ |
| $\cos \theta$ | 1 | $-\frac{1}{2}$ | -1 | $-\frac{1}{2}$ |
| $\tan \theta$ | 0 | $-\sqrt{3}$ | 0 | $-\sqrt{3}$ |
- 7 $\sin \theta = \pm \frac{\sqrt{7}}{4}$ **8 a** $\frac{\sqrt{3}}{2}$ **b** 0 **c** $\frac{1}{2}$
- 9 **a** $\frac{2}{\sqrt{13}}$ **b** $-\frac{3}{\sqrt{13}}$
- 10 perimeter = 12 units, area = 8 units² **11** $\frac{\sqrt{6}}{\sqrt{11}}$
- 12 **a** $150^\circ, 210^\circ$ **b** $45^\circ, 315^\circ$ **c** $120^\circ, 300^\circ$
- 13 **a** $\theta = \pi$ **b** $\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$
- 14 $\cos x = -\frac{\sqrt{15}}{4}$, $\tan x = \frac{1}{\sqrt{15}}$, $\sec x = -\frac{4}{\sqrt{15}}$,
 $\operatorname{cosec} x = -4$, $\cot x = \sqrt{15}$

REVIEW SET 8B

- 1 a
- 72°
- b
- 225°
- c
- 140°
- d
- 330°


2




- 3 a $\cos(\frac{3\pi}{2}) = 0$, $\sin(\frac{3\pi}{2}) = -1$
 b $\cos(-\frac{\pi}{2}) = 0$, $\sin(-\frac{\pi}{2}) = -1$
 4 a $\sin(\pi - p) = m$ b $\sin(p + 2\pi) = m$
 c $\cos p = \sqrt{1 - m^2}$ d $\tan p = \frac{m}{\sqrt{1 - m^2}}$
 5 a i 60° ii $\frac{\pi}{3}$ b $\frac{\pi}{3}$ units c $\frac{\pi}{6}$ units²
 7 $\sin \theta = \frac{\sqrt{21}}{5}$, $\tan \theta = -\frac{\sqrt{21}}{2}$, $\sec \theta = -\frac{5}{2}$,
 $\operatorname{cosec} \theta = \frac{5}{\sqrt{21}}$, $\cot \theta = -\frac{2}{\sqrt{21}}$
 8 a $2\frac{1}{2}$ b $1\frac{1}{2}$ c $-\frac{1}{2}$
 9 a $\theta \approx 0.841$ or 5.44 b $\theta \approx 3.39$ or 6.03
 c $\theta \approx 1.25$ or 4.39
 10 perimeter ≈ 34.1 cm, area ≈ 66.5 cm²
 11 $r \approx 8.79$ cm, area ≈ 81.0 cm² 12 a 0 b $\sin \theta$
 13 $\sin \alpha = \frac{\sqrt{91}}{10}$, $\cos \alpha = -\frac{3}{10}$, $\tan \alpha = -\frac{\sqrt{91}}{3}$,
 $\operatorname{cosec} \alpha = \frac{10}{\sqrt{91}}$, $\cot \alpha = -\frac{3}{\sqrt{91}}$

EXERCISE 9A


- 1 a periodic b periodic c periodic
 d not periodic e periodic f periodic
 g not periodic h not periodic



- b A curve can be fitted to the data.
 c The data is periodic.
 i $y = 32$ (approx.) ii ≈ 64 cm
 iii ≈ 200 cm iv ≈ 32 cm




Data exhibits periodic behaviour.




Not enough information to say data is periodic.

EXERCISE 9B


1 a




b




c




d



e



f









- 2 a $a = 2$ b $a = 5$ c $a = 11$

- 3 a $b = 3$ b $b = 5$ c $b = 6$ d $b = 4$

- 4 a $c = 3$ b $c = -1$ c $c = 5$

5 a







6 a $a = 3, b = 1, c = 0$ **b** $a = 2, b = 5, c = 6$


c $a = 5, b = 3, c = -2$

7 $m = 2, n = -3$




EXERCISE 9C


1 a




b




c




d




e








f



g






2 a $a = 4, b = 3, c = -1$

b $a = 3, b = 5, c = 3$


3 a $y = 2 \cos 2x$

b $y = \cos\left(\frac{x}{2}\right) + 2$

4 a $a = 5,$
 $b = 1,$
 $c = 1$




5 a $a = \frac{3}{2}, b = 6, c = -\frac{3}{2}$




c $y = -\frac{3}{2} \cos 6x + \frac{3}{2}$

EXERCISE 9D


1 a




b




c




d



e





- 2 a** $b = \frac{3}{2}$, $c = 2$ **b** $b = 2$, $c = -3$
3 $p = \frac{1}{2}$, $q = 1$

EXERCISE 9E.1

- 1 a** $x \approx 0.3, 2.8, 6.6, 9.1, 12.9$ **b** $x \approx 5.9, 9.8, 12.2$
2 a $x \approx 0.9, 5.4, 7.2$ **b** $x \approx 4.4, 8.2, 10.7$
3 a $x \approx 0.4, 1.2, 3.5, 4.3, 6.7, 7.5, 9.8, 10.6, 13.0, 13.7$
b $x \approx 1.7, 3.0, 4.9, 6.1, 8.0, 9.3, 11.1, 12.4, 14.3, 15.6$
4 a i ≈ 1.6 **ii** ≈ -1.1
b i $x \approx 1.1, 4.2, 7.4$ **ii** $x \approx 2.2, 5.3$
5 a $x \approx 0.446, 2.70, 6.73, 8.98$
b $x \approx 2.52, 3.76, 8.80, 10.0$
c $x \approx 0.588, 3.73, 6.87, 10.0$
6 a $x \approx -0.644, 0.644$
b $x \approx -4.56, -1.42, 1.72, 4.87$
c $x \approx -2.76, -0.384, 3.53$
7 a $x \approx 1.57$ **b** $m = -2$ or $m = 1$
c $-1 \leq \sin x \leq 1$, so $m = \sin x = -2$ is not a valid solution.

EXERCISE 9E.2

- 1 a** $x = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \frac{11\pi}{3}$ **b** $x = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{9\pi}{4}, \frac{11\pi}{4}$
c $x = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4}$
2 a $x = -\frac{5\pi}{3}, -\frac{4\pi}{3}, \frac{\pi}{3}, \frac{2\pi}{3}$ **b** $x = -\frac{5\pi}{4}, -\frac{3\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}$
c $x = -\frac{5\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{7\pi}{4}$
3 a $x = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}$ **b** $x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}, \frac{13\pi}{6}, \frac{17\pi}{6}$
4 a $x = \frac{5\pi}{4}, \frac{7\pi}{4}$ **b** $x = \frac{5\pi}{12}, \frac{7\pi}{12}, \frac{13\pi}{12}, \frac{5\pi}{4}, \frac{7\pi}{4}, \frac{23\pi}{12}$
5 a $x = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}, \frac{10\pi}{3}, \frac{14\pi}{3}$
b $x = -330^\circ, -210^\circ, 30^\circ, 150^\circ$
c $x = \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{17\pi}{6}$ **d** $x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}$
e $x = -\frac{8\pi}{9}, -\frac{4\pi}{9}, -\frac{2\pi}{9}, \frac{2\pi}{9}, \frac{4\pi}{9}, \frac{8\pi}{9}$
6 $x = \frac{\pi}{3}, \frac{4\pi}{3}$
a $x = \frac{\pi}{12}, \frac{\pi}{3}, \frac{7\pi}{12}, \frac{5\pi}{6}, \frac{13\pi}{12}, \frac{4\pi}{3}, \frac{19\pi}{12}, \frac{11\pi}{6}$
b $x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$
7 $x = \frac{\pi}{18}, \frac{7\pi}{18}, \frac{13\pi}{18}$
8 a $x = -\frac{2\pi}{3}, \frac{2\pi}{3}$ **b** $x = -\frac{5\pi}{6}, -\frac{2\pi}{3}, \frac{\pi}{6}, \frac{\pi}{3}$
c $x = -\frac{\pi}{2}, \frac{\pi}{2}$
9 a $x = \frac{\pi}{4}, \frac{5\pi}{4}$ **b** $x = \frac{3\pi}{4}, \frac{7\pi}{4}$
c $x = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{3\pi}{4}, \frac{13\pi}{12}, \frac{17\pi}{12}, \frac{7\pi}{4}$
d $x = \frac{\pi}{6}, \frac{2\pi}{3}, \frac{7\pi}{6}, \frac{5\pi}{3}$
10 $x = \frac{\pi}{2}$

EXERCISE 9F.1

- 1 a** $2 \sin \theta$ **b** $3 \cos \theta$ **c** $2 \sin \theta$ **d** $\sin \theta$
e $-2 \tan \theta$ **f** $-3 \cos^2 \theta$
2 a 3 **b** -2 **c** -1 **d** $3 \cos^2 \theta$
e $4 \sin^2 \theta$ **f** $\cos \theta$ **g** $-\sin^2 \theta$ **h** $-\cos^2 \theta$
i $-2 \sin^2 \theta$ **j** 1 **k** $\sin \theta$ **l** $\sin \theta$
3 a $2 \tan x$ **b** $\tan^2 x$ **c** $\sin x$ **d** $\cos x$
e $5 \sin x$ **f** $2 \sec x$ **g** 1 **h** 1
i $\operatorname{cosec} x$ **j** $\cos x$ **k** $\cos x$ **l** $5 \sin x$
4 a $1 + 2 \sin \theta + \sin^2 \theta$ **b** $\sin^2 \alpha - 4 \sin \alpha + 4$
c $\tan^2 \alpha - 2 \tan \alpha + 1$ **d** $1 + 2 \sin \alpha \cos \alpha$
e $1 - 2 \sin \beta \cos \beta$ **f** $-4 + 4 \cos \alpha - \cos^2 \alpha$
5 a $-\tan^2 \beta$ **b** 1 **c** $\sin^2 \alpha$
d $\sin^2 x - \tan^2 x$ **e** 13 **f** $\cos^2 \theta$ **g** 0

EXERCISE 9F.2

- 1 a** $(1 - \sin \theta)(1 + \sin \theta)$
b $(\sin \alpha + \cos \alpha)(\sin \alpha - \cos \alpha)$
c $(\tan \alpha + 1)(\tan \alpha - 1)$ **d** $\sin \beta(2 \sin \beta - 1)$
e $\cos \phi(2 + 3 \cos \phi)$ **f** $3 \sin \theta(\sin \theta - 2)$
g $(\tan \theta + 3)(\tan \theta + 2)$ **h** $(2 \cos \theta + 1)(\cos \theta + 3)$
i $(3 \cos \alpha + 1)(2 \cos \alpha - 1)$ **j** $\tan \alpha(3 \tan \alpha - 2)$
k $(\sec \beta + \operatorname{cosec} \beta)(\sec \beta - \operatorname{cosec} \beta)$
l $(2 \cot x - 1)(\cot x - 1)$
m $(2 \sin x + \cos x)(\sin x + 3 \cos x)$
2 a $1 + \sin \alpha$ **b** $\tan \beta - 1$ **c** $\cos \phi - \sin \phi$
d $\cos \phi + \sin \phi$ **e** $\frac{1}{\sin \alpha - \cos \alpha}$ **f** $\frac{\cos \theta}{2}$
g $\sin \theta$ **h** $\cos \theta$ **i** $\sec \theta + 1$


EXERCISE 9G

- 1 a** $x = 0, \pi, \frac{7\pi}{6}, \frac{11\pi}{6}, 2\pi$ **b** $x = \frac{\pi}{3}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{3}$
c $x = \frac{\pi}{3}, \pi, \frac{5\pi}{3}$ **d** $x = \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6}$
e no solutions **f** $x = 0, 2\pi$
2 a $x = \pi$ **b** $x = \frac{\pi}{6}, \frac{5\pi}{6}$


REVIEW SET 9A


- 1 a** no **b** yes

2 a



b






- 3** **a** minimum = 0, maximum = 2
b minimum = -2, maximum = 2
c minimum = -3, maximum = 3
d minimum = -2, maximum = 0

- 4** **a** 2π **b** $\frac{\pi}{2}$ **c** π **d** $\frac{\pi}{3}$

Function	Period	Amplitude
$y = 3 \sin 2x + 1$	π	3
$y = \tan 2x$	$\frac{\pi}{2}$	undefined
$y = 2 \cos 3x - 3$	$\frac{2\pi}{3}$	2

Function	Domain	Range
$y = 3 \sin 2x + 1$	$x \in \mathbb{R}$	$-2 \leq y \leq 4$
$y = \tan 2x$	$x \neq \pm \frac{\pi}{4}, \pm \frac{3\pi}{4}, \dots$	$y \in \mathbb{R}$
$y = 2 \cos 3x - 3$	$x \in \mathbb{R}$	$-5 \leq y \leq -1$

6 $y = 4 \cos 2x$



8 **a** $x \approx 115^\circ, 245^\circ, 475^\circ, 605^\circ$ **b** $x \approx 25^\circ, 335^\circ, 385^\circ$

9 **a** $x = \frac{7\pi}{6}, \frac{11\pi}{6}, \frac{19\pi}{6}, \frac{23\pi}{6}$ **b** $x = -\frac{7\pi}{4}, -\frac{5\pi}{4}, \frac{\pi}{4}, \frac{3\pi}{4}$

c $x = \frac{4\pi}{9}, \frac{5\pi}{9}, \frac{10\pi}{9}, \frac{11\pi}{9}, \frac{16\pi}{9}, \frac{17\pi}{9}$

d $x = \frac{\pi}{4}, \frac{7\pi}{4}, \frac{9\pi}{4}, \frac{15\pi}{4}$

10 **a** $1 - \cos \theta$ **b** $\frac{1}{\sin \alpha + \cos \alpha}$ **c** $-\frac{\cos \alpha}{2}$
d $\operatorname{cosec} \theta + 1$

12 **a** $x = -\frac{2\pi}{3}, -\frac{\pi}{6}, \frac{\pi}{3}, \frac{5\pi}{6}$ **b** $x = -\frac{2\pi}{3}, -\frac{\pi}{3}, \frac{\pi}{3}, \frac{2\pi}{3}$





REVIEW SET 9B

- 1** **a** The function repeats itself over and over in a horizontal direction, in intervals of length 8 units.


b **i** 8 **ii** 5 **iii** -1

2 **a** $b = 6$ **b** $b = 24$


- 3** **a** minimum = -8, maximum = 2
b minimum = -2, maximum = 4
c minimum = 5, maximum = 13



5 **a** $a = 2, b = 6, c = -2$



6 **a** $y = 2 \tan x + 2$



7 **a** $x \approx -6.1, -3.4$ **b** $x \approx 0.8$

8 $m = 3, n = -1$

9 **a** $x = \frac{3\pi}{2}$ **b** $x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}$

10 a $\cos \theta$ b $-\sin \theta$ c $5 \cos^2 \theta$ d $-\cos \theta$

11 a $4 \sin^2 \alpha - 4 \sin \alpha + 1$ b $1 - 2 \sin \alpha \cos \alpha$

EXERCISE 10A

1 18 **2** a 4 b 8 c 24 **3** 6

4 42 **5** 1680 **6** a 125 b 60

7 17 576 000 **8** a 4 b 9 c 81

EXERCISE 10B

1 a 13 b 20 c 19 d 32 **2** 13

EXERCISE 10C.1

1 1, 1, 2, 6, 24, 120, 720, 5040, 40 320, 362 880, 3 628 800

2 a 6 b 30 c $\frac{1}{7}$ d $\frac{1}{30}$ e 100 f 21

3 a $n, n \geq 1$ b $(n+2)(n+1), n \geq 0$

c $(n+1)n, n \geq 1$

4 a $\frac{7!}{4!}$ b $\frac{10!}{8!}$ c $\frac{11!}{6!}$ d $\frac{13!}{10!3!}$ e $\frac{3!}{6!}$ f $\frac{4!16!}{20!}$

5 a $6 \times 4!$ b $10 \times 10!$ c $73 \times 7!$ d $131 \times 10!$

e $81 \times 7!$ f $62 \times 6!$ g $10 \times 11!$ h $32 \times 8!$

6 a 11! b 9! c 8! d 9

e 34 f $n+1$ g $(n-1)!$ h $(n+1)!$

EXERCISE 10C.2

1 a 3 b 6 c 35 d 210

2 a i 28 ii 28 **3** $k = 3$ or 6

EXERCISE 10D

1 a W, X, Y, Z

b WX, WY, WZ, XW, XY, XZ, YW, YX, YZ, ZW, ZX, ZY

c WXY, WXZ, WYX, WYZ, WZX, WZY, XWY, XWZ, XYW, XYZ, XZW, XZY, YWX, YWZ, YXW, YXZ, YZX, YZW, ZWX, ZWY, ZXW, ZXY, ZYW, ZYX

2 a AB, AC, AD, AE, BA, BC, BD, BE, CA, CB, CD, CE, DA, DB, DC, DE, EA, EB, EC, ED

b ABC, ABD, ABE, ACB, ACD, ACE, ADB, ADC, ADE, AEB, AEC, AED, BAC, BAD, BAE, BCA, BCD, BCE, BDA, BDC, BDE, BEA, BEC, BED, CAB, CAD, CAE, CBA, CBD, CBE, CDA, CDB, CDE, CEA, CEB, CED, DAB, DAC, DAE, DBA, DBC, DBE, DCA, DCB, DCE, DEA, DEB, DEC, EAB, EAC, EAD, EBA, EBC, EBD, ECA, ECB, ECD, EDA, EDB, EDC

3 a 120 b 336 c 5040 **4** 110

5 a 12 b 24 c 36 **6** a 15 120 b 720

7 a 720 b i 24 ii 24 iii 48

8 a 343 b 210 c 120

9 a 648 b 64 c 72 d 136

10 a 6720 b 240 c 4200

11 a 120 b 48 c 72

12 a 3 628 800 b 241 920

13 a 720 b 144 c 72 d 144

14 a 48 b 24 c 15 **15** a 360 b 336 c 288

16 a 3 628 800 b i 151 200 ii 33 600

EXERCISE 10E

1 a permutation b combination

c permutation d combination

2 ABCD, ABCE, ABCF, ABDE, ABDF, ABFE, ACDE, ACDF, ACEF, ADEF, BCDE, BCDF, BCEF, BDEF, CDEF, $\binom{6}{4} = 15$

3 $\binom{17}{11} = 12\,376$ **4** a $\binom{9}{5} = 126$ b $\binom{1}{1} \binom{8}{4} = 70$

5 a $\binom{13}{3} = 286$ b $\binom{1}{1} \binom{12}{2} = 66$

6 a $\binom{12}{5} = 792$

b i $\binom{2}{2} \binom{10}{3} = 120$ ii $\binom{2}{1} \binom{10}{4} = 420$

7 $\binom{3}{3} \binom{1}{0} \binom{11}{6} = 462$

8 a $\binom{1}{1} \binom{9}{3} = 84$ b $\binom{2}{0} \binom{8}{4} = 70$

c $\binom{2}{0} \binom{1}{1} \binom{7}{3} = 35$

9 a $\binom{16}{5} = 4368$ b $\binom{10}{3} \binom{6}{2} = 1800$

c $\binom{10}{5} \binom{6}{0} = 252$

d $\binom{10}{3} \binom{6}{2} + \binom{10}{4} \binom{6}{1} + \binom{10}{5} \binom{6}{0} = 3312$

e $\binom{16}{5} - \binom{10}{5} \binom{6}{0} - \binom{10}{0} \binom{6}{5} = 4110$

10 a 6435 b 2520 c 36 d 4005 **11** 1050

12 a $\binom{6}{2} \binom{3}{1} \binom{7}{2} = 945$ b $\binom{6}{2} \binom{10}{3} = 1800$

c $\binom{16}{5} - \binom{9}{0} \binom{7}{5} = 4347$

13 $\binom{20}{2} - 20 = 170$

14 a i $\binom{12}{2} = 66$ ii $\binom{11}{1} = 11$

b i $\binom{12}{3} = 220$ ii $\binom{11}{2} = 55$

15 $\binom{9}{4} = 126$

16 a Selecting the different committees of 4 from 5 men and 6 women in all possible ways.

b $\binom{m+n}{r}$

17 a $\frac{\binom{12}{6}}{2} = 462$ b $\frac{\binom{12}{4} \binom{8}{4} \binom{4}{4}}{3!} = 5775$

18 a 45, yes b 37 128 c 3 628 800

EXERCISE 10F

1 a $p^3 + 3p^2q + 3pq^2 + q^3$ b $x^3 + 3x^2 + 3x + 1$

c $x^3 - 9x^2 + 27x - 27$ d $8 + 12x + 6x^2 + x^3$

e $27x^3 - 27x^2 + 9x - 1$ f $8x^3 + 60x^2 + 150x + 125$

g $8a^3 - 12a^2b + 6ab^2 - b^3$ h $27x^3 - 9x^2 + x - \frac{1}{27}$

i $8x^3 + 12x + \frac{6}{x} + \frac{1}{x^3}$

2 a $1 + 4x + 6x^2 + 4x^3 + x^4$

b $p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4$

c $x^4 - 8x^3 + 24x^2 - 32x + 16$

d $81 - 108x + 54x^2 - 12x^3 + x^4$

e $1 + 8x + 24x^2 + 32x^3 + 16x^4$

f $16x^4 - 96x^3 + 216x^2 - 216x + 81$

g $16x^4 + 32x^3b + 24x^2b^2 + 8xb^3 + b^4$

h $x^4 + 4x^2 + 6 + \frac{4}{x^2} + \frac{1}{x^4}$

i $16x^4 - 32x^2 + 24 - \frac{8}{x^2} + \frac{1}{x^4}$

3 a $x^5 + 10x^4 + 40x^3 + 80x^2 + 80x + 32$

b $x^5 - 10x^4y + 40x^3y^2 - 80x^2y^3 + 80xy^4 - 32y^5$

c $1 + 10x + 40x^2 + 80x^3 + 80x^4 + 32x^5$

d $x^5 - 5x^3 + 10x - \frac{10}{x} + \frac{5}{x^3} - \frac{1}{x^5}$

4 $64 + 160x^2 + 20x^4$

- 5 a** 1 6 15 20 15 6 1
b i $x^6 + 12x^5 + 60x^4 + 160x^3 + 240x^2 + 192x + 64$
ii $64x^6 - 192x^5 + 240x^4 - 160x^3 + 60x^2 - 12x + 1$
iii $x^6 + 6x^4 + 15x^2 + 20 + \frac{15}{x^2} + \frac{6}{x^4} + \frac{1}{x^6}$
- 6 a** $7 + 5\sqrt{2}$ **b** $161 + 72\sqrt{5}$ **c** $232 - 164\sqrt{2}$
- 7** $\frac{59 + 34\sqrt{3}}{13}$
- 8 a** $64 + 192x + 240x^2 + 160x^3 + 60x^4 + 12x^5 + x^6$
b 65.944 160 601 201
- 9 a** $a = 2$ and $b = e^x$ **b** $T_3 = 6e^{2x}$ and $T_4 = e^{3x}$
- 10** $2x^5 + 11x^4 + 24x^3 + 26x^2 + 14x + 3$
- 11 a** 270 **b** 4320

EXERCISE 10G

- 1 a** $1^{11} + \binom{11}{1}(2x)^1 + \binom{11}{2}(2x)^2 + \dots + \binom{11}{10}(2x)^{10} + (2x)^{11}$
b $(3x)^{15} + \binom{15}{1}(3x)^{14}\left(\frac{2}{x}\right)^1 + \binom{15}{2}(3x)^{13}\left(\frac{2}{x}\right)^2 + \dots$
 $\dots + \binom{15}{14}(3x)^1\left(\frac{2}{x}\right)^{14} + \left(\frac{2}{x}\right)^{15}$
c $(2x)^{20} + \binom{20}{1}(2x)^{19}\left(-\frac{3}{x}\right)^1 + \binom{20}{2}(2x)^{18}\left(-\frac{3}{x}\right)^2 + \dots$
 $\dots + \binom{20}{19}(2x)^1\left(-\frac{3}{x}\right)^{19} + \left(-\frac{3}{x}\right)^{20}$
- 2 a** $T_6 = \binom{15}{5}(2x)^{10}5^5$ **b** $T_4 = \binom{9}{3}(x^2)^6y^3$
c $T_{10} = \binom{17}{9}x^8\left(-\frac{2}{x}\right)^9$ **d** $T_9 = \binom{21}{8}(2x^2)^{13}\left(-\frac{1}{x}\right)^8$
- 3 a** $\binom{12}{4}2^83^4 = 10\,264\,320$ **b** $\binom{12}{7}2^53^7 = 55\,427\,328$
- 4 a** $\binom{10}{3}1^7(-3)^3 = -3240$ **b** $\binom{10}{7}1^3(-3)^7 = -262\,440$
- 5 a** 144 **b** 5376 **c** 2304
- 6 a** $T_{r+1} = \binom{7}{r}x^{7-r}b^r$ **b** $b = -2$
- 7 a** $\binom{15}{5}2^5 = 96\,096$ **b** $\binom{9}{3}(-3)^3 = -2268$
- 8 a** $\binom{10}{5}3^52^5 = 1\,959\,552$ **b** $\binom{6}{3}2^3(-3)^3 = -4320$
c $\binom{6}{3}2^3(-3)^3 = -4320$ **d** $\binom{12}{4}2^8(-1)^4 = 126\,720$
- 9** $k = 5$ **10** $a = 3$ **11** **a** $a = 5$, $b = 2$
- 12** $\binom{8}{6} = 28$ **13** $2\binom{9}{3}3^6x^6 - \binom{9}{4}3^5x^6 = 91\,854x^6$
- 14 a** $\binom{7}{4}3^3(-2)^4 = 15\,120$
b $\binom{7}{4}3^3(-2)^4 + 3\binom{7}{3}3^4(-2)^3 = -52\,920$
- 15 a** $\binom{8}{3}2^5(-5)^3 - 3\binom{8}{1}2^7(-5)^1 = -208\,640$
b $\binom{6}{3}2^3 - \binom{6}{4}2^4 = -80$
- 16** $a = 3$, $b = -2$, $c = 57$ **17** $n = 8$
- 18** $n = 6$ **19** $84x^3$ **20** $k = -2$, $n = 6$

REVIEW SET 10A



- 1 a** $n(n-1)$, $n \geq 2$ **b** $n+2$ **2** 28
- 3 a** 24 **b** 6 **4 a** 900 **b** 180
- 5 a** $a = e^x$ and $b = -e^{-x}$
b $(e^x - e^{-x})^4 = e^{4x} - 4e^{2x} + 6 - 4e^{-2x} + e^{-4x}$
- 6** $362 + 209\sqrt{3}$ **7** It does not have one. **8** $c = 3$
- 9 a** 720 **b** 72 **c** 504 **10** 2500
- 11 a** 252 **b** 246 **12** $\binom{12}{6}2^6(-3)^6 = 43\,110\,144$
- 13** $8\binom{6}{2} - 6\binom{6}{1} = 84$ **14** $a = \pm 4$ **15** $k = 0$ or ± 2

REVIEW SET 10B


- 1 a** $26^2 \times 10^4 = 6\,760\,000$
b $5 \times 26 \times 10^4 = 1\,300\,000$
c $26 \times 25 \times 10 \times 9 \times 8 \times 7 = 3\,276\,000$
- 2 a** 3003 **b** 980 **c** 2982
- 3 a** $x^3 - 6x^2y + 12xy^2 - 8y^3$
b $81x^4 + 216x^3 + 216x^2 + 96x + 16$
- 4** 20 000 **5** 60 **6** $-103 + 74\sqrt{2}$ **7** 4200
- 8** $\binom{5}{2}2^3 - 3\binom{5}{1}2^4 = -160$
- 9 a** 3024 **b** 840 **c** 42
- 10** $q = 0$ or $\pm\sqrt{\frac{3}{35}}$ **11** 4320 **12** $k = 180$
- 13 a** 43 758 teams **b** 11 550 teams **c** 41 283 teams
d 3861 teams
- 14** $n = 7$ **15** $k = -\frac{1}{4}$, $n = 16$

EXERCISE 11A


- 1 a** $\begin{pmatrix} 7 \\ 3 \end{pmatrix}$, $7\mathbf{i} + 3\mathbf{j}$ **b** $\begin{pmatrix} -6 \\ 0 \end{pmatrix}$, $-6\mathbf{i}$
c $\begin{pmatrix} 2 \\ -5 \end{pmatrix}$, $2\mathbf{i} - 5\mathbf{j}$ **d** $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$, $6\mathbf{j}$
e $\begin{pmatrix} -6 \\ 3 \end{pmatrix}$, $-6\mathbf{i} + 3\mathbf{j}$ **f** $\begin{pmatrix} -5 \\ -5 \end{pmatrix}$, $-5\mathbf{i} - 5\mathbf{j}$
- 2 a** $3\mathbf{i} + 4\mathbf{j}$ **b** $2\mathbf{i}$



- c** $2\mathbf{i} - 5\mathbf{j}$




- d** $-\mathbf{i} - 3\mathbf{j}$




- 3 a i** $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$, $4\mathbf{i} + \mathbf{j}$ **ii** $\begin{pmatrix} -4 \\ -1 \end{pmatrix}$, $-4\mathbf{i} - \mathbf{j}$
iii $\begin{pmatrix} -1 \\ -5 \end{pmatrix}$, $-\mathbf{i} - 5\mathbf{j}$ **iv** $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$, $2\mathbf{i}$
v $\begin{pmatrix} 3 \\ -4 \end{pmatrix}$, $3\mathbf{i} - 4\mathbf{j}$ **vi** $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$, $4\mathbf{i} + \mathbf{j}$

- b** \overrightarrow{AB} and \overrightarrow{DE} . They have the same magnitude and direction.
c \overrightarrow{BA} is the negative of both \overrightarrow{AB} and \overrightarrow{DE} . They have the same magnitude but opposite direction.


- 4 a** $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$




- b** $\begin{pmatrix} -1 \\ 3 \end{pmatrix}$



c $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$



d $\begin{pmatrix} 4 \\ -2 \end{pmatrix}$



5 a $\begin{pmatrix} -1 \\ -4 \end{pmatrix}$ b $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$ c $\begin{pmatrix} -5 \\ 2 \end{pmatrix}$ d $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$

EXERCISE 11B

- 1 a 5 units b 5 units c 2 units
 d $\sqrt{8}$ units e 3 units
 2 a $\sqrt{2}$ units b 13 units c $\sqrt{17}$ units
 d 3 units e $|k|$ units
 3 a unit vector b unit vector c not a unit vector
 d unit vector e not a unit vector
 4 a $k = \pm 1$ b $k = \pm 1$ c $k = 0$
 d $k = \pm \frac{1}{\sqrt{2}}$ e $k = \pm \frac{\sqrt{3}}{2}$
 5 $p = \pm 3$

EXERCISE 11C


- 1 a $\begin{pmatrix} -2 \\ 6 \end{pmatrix}$ b $\begin{pmatrix} -2 \\ 6 \end{pmatrix}$ c $\begin{pmatrix} -1 \\ -1 \end{pmatrix}$ d $\begin{pmatrix} -1 \\ -1 \end{pmatrix}$
 e $\begin{pmatrix} -5 \\ -3 \end{pmatrix}$ f $\begin{pmatrix} -5 \\ -3 \end{pmatrix}$ g $\begin{pmatrix} -6 \\ 4 \end{pmatrix}$ h $\begin{pmatrix} -4 \\ 1 \end{pmatrix}$
 2 a $\begin{pmatrix} -3 \\ 7 \end{pmatrix}$ b $\begin{pmatrix} -4 \\ -3 \end{pmatrix}$ c $\begin{pmatrix} -8 \\ -1 \end{pmatrix}$ d $\begin{pmatrix} -6 \\ 9 \end{pmatrix}$
 e $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$ f $\begin{pmatrix} 6 \\ -9 \end{pmatrix}$
 3 a $\mathbf{a} + \mathbf{0} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} a_1 + 0 \\ a_2 + 0 \end{pmatrix} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \mathbf{a}$
 b $\mathbf{a} - \mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} - \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \begin{pmatrix} a_1 - a_1 \\ a_2 - a_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \mathbf{0}$
 4 a $\begin{pmatrix} -3 \\ -15 \end{pmatrix}$ b $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$ c $\begin{pmatrix} 0 \\ 14 \end{pmatrix}$ d $\begin{pmatrix} 5 \\ -3 \end{pmatrix}$
 e $\begin{pmatrix} \frac{5}{2} \\ \frac{11}{2} \end{pmatrix}$ f $\begin{pmatrix} -7 \\ 7 \end{pmatrix}$ g $\begin{pmatrix} 5 \\ 11 \end{pmatrix}$ h $\begin{pmatrix} 3 \\ \frac{17}{3} \end{pmatrix}$
 5 a $\begin{pmatrix} 8 \\ -1 \end{pmatrix}$ b $\begin{pmatrix} 8 \\ -1 \end{pmatrix}$ c $\begin{pmatrix} 8 \\ -1 \end{pmatrix}$

In each case, the result is $2\mathbf{p} + 3\mathbf{q} = \begin{pmatrix} 8 \\ -1 \end{pmatrix}$.

- 6 a $\sqrt{13}$ units b $\sqrt{17}$ units c $5\sqrt{2}$ units
 d $\sqrt{10}$ units e $\sqrt{29}$ units
 7 a $\sqrt{10}$ units b $2\sqrt{10}$ units c $2\sqrt{10}$ units
 d $3\sqrt{10}$ units e $3\sqrt{10}$ units f $2\sqrt{5}$ units
 g $8\sqrt{5}$ units h $8\sqrt{5}$ units i $\sqrt{5}$ units
 j $\sqrt{5}$ units
 8 a $3\mathbf{i} + 2\mathbf{j}$ b $-\mathbf{i} + 9\mathbf{j}$ c $6\mathbf{i} - \mathbf{j}$ d $7\mathbf{j}$
 e 2 units f $2\sqrt{10}$ units

EXERCISE 11D

- 1 a $\begin{pmatrix} 2 \\ 4 \end{pmatrix}$ b $\begin{pmatrix} -2 \\ 5 \end{pmatrix}$ c $\begin{pmatrix} 3 \\ -3 \end{pmatrix}$ d $\begin{pmatrix} 1 \\ -5 \end{pmatrix}$
 e $\begin{pmatrix} 6 \\ -5 \end{pmatrix}$ f $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$
 2 a B(4, 2) b C(2, 2) 3 a $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ b Q(3, 3)
 4 a $\begin{pmatrix} 5 \\ 1 \end{pmatrix}$ b $\begin{pmatrix} -5 \\ -1 \end{pmatrix}$ c D(-1, -2)
 5 a $\vec{AB} = \begin{pmatrix} 4 \\ k-3 \end{pmatrix}$, $|\vec{AB}| = \sqrt{16 + (k-3)^2} = 5$ units
 b $k = 0$ or 6 c



- 6 a $\vec{AB} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$, $\vec{AC} = \begin{pmatrix} 3 \\ -3 \end{pmatrix}$
 b $\vec{BC} = \vec{BA} + \vec{AC} = -\vec{AB} + \vec{AC}$ c $\vec{BC} = \begin{pmatrix} 1 \\ -6 \end{pmatrix}$
 7 a $\begin{pmatrix} -5 \\ 4 \end{pmatrix}$ b $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ c $\begin{pmatrix} 6 \\ -5 \end{pmatrix}$
 8 a M(1, 4) b $\vec{CA} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$, $\vec{CM} = \begin{pmatrix} 5 \\ 3 \end{pmatrix}$, $\vec{CB} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$

EXERCISE 11E

- 1 $r = 3$ 2 $a = -6$
 3 a \vec{AB} is parallel and in the same direction as \vec{CD} , and 3 times its length.
 b \vec{RS} is parallel and in the opposite direction to \vec{KL} , and half its length.
 c A, B, and C are collinear. \vec{AB} is parallel and in the same direction as \vec{BC} , and twice its length.
 4 a $\begin{pmatrix} 4 \\ 8 \end{pmatrix}$ b $\begin{pmatrix} -1 \\ -2 \end{pmatrix}$
 5 a $\frac{1}{\sqrt{5}}\mathbf{i} + \frac{2}{\sqrt{5}}\mathbf{j}$ b $\frac{1}{\sqrt{10}}\mathbf{i} - \frac{3}{\sqrt{10}}\mathbf{j}$ c $\frac{2}{\sqrt{5}}\mathbf{i} - \frac{1}{\sqrt{5}}\mathbf{j}$
 6 a $\mathbf{v} = \frac{3}{\sqrt{5}}\begin{pmatrix} 2 \\ -1 \end{pmatrix}$ b $\mathbf{v} = \frac{2}{\sqrt{17}}\begin{pmatrix} 1 \\ 4 \end{pmatrix}$
 7 a $\vec{AB} = \begin{pmatrix} 2\sqrt{2} \\ -2\sqrt{2} \end{pmatrix}$ b $\vec{OB} = \begin{pmatrix} 3 + 2\sqrt{2} \\ 2 - 2\sqrt{2} \end{pmatrix}$
 c B($3 + 2\sqrt{2}$, $2 - 2\sqrt{2}$)

EXERCISE 11F

- 1 a $\begin{array}{c} 6 \text{ m s}^{-1} \quad 1 \text{ m s}^{-1} \\ \hline 7 \text{ m s}^{-1} \end{array} \quad \therefore 7 \text{ m s}^{-1}$
 b $\begin{array}{c} 6 \text{ m s}^{-1} \\ \hline 5 \text{ m s}^{-1} \quad 1 \text{ m s}^{-1} \end{array} \quad \therefore 5 \text{ m s}^{-1}$
 2 a 1.34 m s^{-1} in the direction 26.6° to the right of intended line
 b i 30° to the left of Q ii 1.04 m s^{-1}

- 3 a** 24.6 km h^{-1} **b** $\approx 9.93^\circ$ east of south
4 a 82.5 m **b** 23.3° to the left of directly across **c** 48.4 s
5 a The plane's speed in still air would be $\approx 437 \text{ km h}^{-1}$.
 The wind slows the plane down to 400 km h^{-1} .
b 4.64° north of due east


EXERCISE 11G


- 1 a i** $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ -4 \end{pmatrix} + t \begin{pmatrix} 1 \\ 4 \end{pmatrix}, t \in \mathbb{R}$
ii $x = 3 + t, y = -4 + 4t, t \in \mathbb{R}$ **iii** $4x - y = 16$
b i $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -6 \\ 0 \end{pmatrix} + t \begin{pmatrix} 3 \\ 7 \end{pmatrix}, t \in \mathbb{R}$
ii $x = -6 + 3t, y = 7t, t \in \mathbb{R}$ **iii** $7x - 3y = -42$
c i $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 \\ 11 \end{pmatrix} + t \begin{pmatrix} -2 \\ 1 \end{pmatrix}, t \in \mathbb{R}$
ii $x = -1 - 2t, y = 11 + t, t \in \mathbb{R}$ **iii** $x + 2y = 21$
2 a $x = -1 + 2t, y = 4 - t, t \in \mathbb{R}$
b

t	0	1	3	-1	-4
Point	$(-1, 4)$	$(1, 3)$	$(5, 1)$	$(-3, 5)$	$(-9, 8)$

3 a When $t = 1, x = 3, y = -2, \therefore$ yes **b** $k = -5$
4 a $(0, 8)$ **b** It is a non-zero scalar multiple of $\begin{pmatrix} -1 \\ 3 \end{pmatrix}$.
c $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \end{pmatrix} + s \begin{pmatrix} 1 \\ -3 \end{pmatrix}, s \in \mathbb{R}$

EXERCISE 11H


- 1 a** $(1, 2)$ **b**



c $\begin{pmatrix} 2 \\ -5 \end{pmatrix}$
d $\sqrt{29} \text{ cm s}^{-1}$
2 a $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} + t \begin{pmatrix} 4 \\ -5 \end{pmatrix}, t \geq 0$ **b** $(8, -4.5)$
c 45 minutes
3 a $\begin{pmatrix} -3 + 2t \\ -2 + 4t \end{pmatrix}$ **d**


b $\begin{pmatrix} 2 \\ 8 \end{pmatrix}$
c i $t = 1.5 \text{ s}$
ii $t = 0.5 \text{ s}$
4 a i $(-4, 3)$ **ii** $\begin{pmatrix} 12 \\ 5 \end{pmatrix}$ **iii** 13 m s^{-1}
b i $(3, 0)$ **ii** $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$ **iii** $\sqrt{5} \text{ m s}^{-1}$
5 a $\begin{pmatrix} 120 \\ -90 \end{pmatrix}$ **b** $\begin{pmatrix} 20\sqrt{5} \\ 10\sqrt{5} \end{pmatrix}$ **c** $\begin{pmatrix} 35 \\ -84 \end{pmatrix}$
7 a A is at $(4, 5)$, B is at $(1, -8)$
b For A it is $\begin{pmatrix} 1 \\ -2 \end{pmatrix}$. For B it is $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$.
c For A, speed is $\sqrt{5} \text{ km h}^{-1}$. For B, speed is $\sqrt{5} \text{ km h}^{-1}$.
d Yacht A: $2x + y = 13$, Yacht B: $x - 2y = 17$

- e** Yacht A moves with gradient -2 ; Yacht B with gradient $\frac{1}{2}$.
 So, their paths are perpendicular.
f no
8 a $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} -5 \\ 4 \end{pmatrix} + t \begin{pmatrix} 3 \\ -1 \end{pmatrix}, t \geq 0$
 $\therefore x_1(t) = -5 + 3t, y_1(t) = 4 - t, t \geq 0$
b speed $= \sqrt{10} \text{ km min}^{-1}$
c a minutes later, $(t - a) \text{ min}$ have elapsed.
 $\therefore \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} 15 \\ 7 \end{pmatrix} + (t - a) \begin{pmatrix} -4 \\ -3 \end{pmatrix}, t \geq 0$
 $\therefore x_2(t) = 15 - 4(t - a), y_2(t) = 7 - 3(t - a), t \geq 0$
d Torpedo is fired at 1:35:28 pm and the explosion occurs at 1:37:42 pm.

REVIEW SET 11A

- 1 a** $\mathbf{x} = \begin{pmatrix} 5 \\ 1 \end{pmatrix} = 5\mathbf{i} + \mathbf{j}, \mathbf{y} = \begin{pmatrix} 1 \\ -2 \end{pmatrix} = \mathbf{i} - 2\mathbf{j}$
b i $6\mathbf{i} - \mathbf{j}$ **ii** $-9\mathbf{i} - 4\mathbf{j}$
2 a $\begin{pmatrix} 3 \\ -1 \end{pmatrix}$ **b**


c $\begin{pmatrix} -3 \\ 1 \end{pmatrix}$
d $\sqrt{10}$ units
3 a $k = \pm \frac{1}{\sqrt{2}}$ **b** $\begin{pmatrix} -2\sqrt{5} \\ \sqrt{5} \end{pmatrix}$
4 a $\begin{pmatrix} 3 \\ -3 \end{pmatrix}$ **b** $\begin{pmatrix} 7 \\ -3 \end{pmatrix}$ **c** 5 units
5 a $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$ **b** $\begin{pmatrix} -3 \\ -6 \end{pmatrix}$ **c** $\sqrt{34}$ units
6 a $\overrightarrow{BC} = \begin{pmatrix} k+3 \\ 3 \end{pmatrix}$ **c**


 $|\overrightarrow{BC}| = 5$
b $k = -7$ or 1
7 a 11.5° east of due north **b** $\approx 343 \text{ km h}^{-1}$
8 a $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -6 \\ 3 \end{pmatrix} + t \begin{pmatrix} 4 \\ -3 \end{pmatrix}, t \in \mathbb{R}$
b $x = -6 + 4t, y = 3 - 3t, t \in \mathbb{R}$ **c** $3x + 4y = -6$
9 $m = 10$ **10** $\begin{pmatrix} 6\sqrt{10} \\ -2\sqrt{10} \end{pmatrix}$
11 a $(5, 2)$ **b** $\begin{pmatrix} 4 \\ 10 \end{pmatrix}$ is a non-zero scalar multiple of $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$
c $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 2 \end{pmatrix} + s \begin{pmatrix} 4 \\ 10 \end{pmatrix}, s \in \mathbb{R}$
12 a $(-4, 3)$ **b** $(28, 27)$ **c** $\begin{pmatrix} 8 \\ 6 \end{pmatrix}$ **d** 10 m s^{-1}

REVIEW SET 11B

- 1 a i** $\overrightarrow{AB} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} = 4\mathbf{i}$ **ii** $\overrightarrow{BC} = \begin{pmatrix} -2 \\ -4 \end{pmatrix} = -2\mathbf{i} - 4\mathbf{j}$
iii $\overrightarrow{CA} = \begin{pmatrix} -2 \\ 4 \end{pmatrix} = -2\mathbf{i} + 4\mathbf{j}$

b \vec{BC} and \vec{CA} . Each vector has components of the same magnitude, but differing signs (which do not affect the length of \vec{BC} or \vec{CA}).


c \vec{AC} , $\begin{pmatrix} 2 \\ -4 \end{pmatrix}$, $2\mathbf{i} - 4\mathbf{j}$

2 a $\sqrt{13}$ units **b** $\sqrt{10}$ units **c** $\sqrt{109}$ units

3 a $k = \pm \frac{12}{13}$ **b** $k = \pm \frac{1}{\sqrt{2}}$ **4** $\begin{pmatrix} 1 \\ 4 \end{pmatrix}$

5 a $M(-2, 4)$ **b** 5 units **6** $m = 5$

7 a



b i isosceles triangle \therefore 2 remaining angles $= 89^\circ$ each, breeze makes angle of $180 - 89 = 91^\circ$ to intended direction of the arrow.

ii bisect angle 2° and use $\sin 1^\circ = \frac{\frac{1}{2} \text{ speed}}{|\mathbf{v}|}$
 \therefore speed $= 2|\mathbf{v}| \sin 1^\circ$

8 $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \end{pmatrix} + t \begin{pmatrix} 5 \\ 4 \end{pmatrix}$, $t \in \mathbb{R}$

9 a i $-6\mathbf{i} + 10\mathbf{j}$ **ii** $-5\mathbf{i} - 15\mathbf{j}$

iii $(-6 - 5t)\mathbf{i} + (10 - 15t)\mathbf{j}$, $t \geq 0$

b $t = 40 \text{ min } (\frac{2}{3} \text{ h})$, yacht is $\approx 9.33 \text{ km}$ away from the beacon.

10 a i $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ -3 \end{pmatrix} + t \begin{pmatrix} 4 \\ -1 \end{pmatrix}$, $t \in \mathbb{R}$

ii $x = 2 + 4t$, $y = -3 - t$, $t \in \mathbb{R}$

b i $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 \\ 6 \end{pmatrix} + t \begin{pmatrix} 3 \\ -4 \end{pmatrix}$, $t \in \mathbb{R}$

ii $x = -1 + 3t$, $y = 6 - 4t$, $t \in \mathbb{R}$

11 a $x_1(t) = 2 + t$, $y_1(t) = 4 - 3t$, $t \geq 0$

b $x_2(t) = 13 - t$, $y_2(t) = [3 - 2a] + at$, $t \geq 2$

c interception occurred at 2:22:30 pm

d bearing $\approx 12.7^\circ$ west of due south, at $\approx 4.54 \text{ km min}^{-1}$

EXERCISE 12A

1 a 1×4 **b** 2×1 **c** 2×2 **d** 3×3

2 a $\begin{pmatrix} 2 & 1 & 6 & 1 \end{pmatrix}$ **b** $\begin{pmatrix} 1.95 \\ 2.35 \\ 0.45 \\ 2.95 \end{pmatrix}$

c total cost of groceries

3 $\begin{pmatrix} 1000 & 1500 & 1250 \\ 1500 & 1000 & 1000 \\ 800 & 2300 & 1300 \\ 1200 & 1200 & 1200 \end{pmatrix}$ **4** $\begin{pmatrix} 40 & 50 & 55 & 40 \\ 25 & 65 & 44 & 30 \\ 35 & 40 & 40 & 35 \\ 35 & 40 & 35 & 50 \end{pmatrix}$

EXERCISE 12B.1

1 a $\begin{pmatrix} 9 & 1 \\ 3 & 3 \end{pmatrix}$ **b** $\begin{pmatrix} 6 & 8 \\ -1 & 1 \end{pmatrix}$
c $\begin{pmatrix} 3 & 4 \\ -6 & -1 \end{pmatrix}$ **d** $\begin{pmatrix} 0 & 0 \\ -11 & -3 \end{pmatrix}$

2 a $\begin{pmatrix} 20 & 1 & -8 \\ 8 & 10 & -2 \\ 1 & -5 & 18 \end{pmatrix}$ **b** $\begin{pmatrix} -14 & 9 & -14 \\ 12 & -6 & 14 \\ -5 & 3 & -4 \end{pmatrix}$

c $\begin{pmatrix} 14 & -9 & 14 \\ -12 & 6 & -14 \\ 5 & -3 & 4 \end{pmatrix}$

3 a Friday Saturday
 $\begin{pmatrix} 85 \\ 92 \\ 52 \end{pmatrix}$ $\begin{pmatrix} 102 \\ 137 \\ 49 \end{pmatrix}$ **b** $\begin{pmatrix} 187 \\ 229 \\ 101 \end{pmatrix}$

4 a i $\begin{pmatrix} 1.72 \\ 27.85 \\ 0.92 \\ 2.53 \\ 3.56 \end{pmatrix}$ **ii** $\begin{pmatrix} 1.79 \\ 28.75 \\ 1.33 \\ 2.25 \\ 3.51 \end{pmatrix}$

b subtract cost price from selling price **c** $\begin{pmatrix} 0.07 \\ 0.90 \\ 0.41 \\ -0.28 \\ -0.05 \end{pmatrix}$

5 a L R
 $\begin{pmatrix} 23 & 19 \\ 17 & 29 \\ 31 & 24 \end{pmatrix}$ fr
st
mi

b L R
 $\begin{pmatrix} 18 & 25 \\ 7 & 13 \\ 36 & 19 \end{pmatrix}$ fr
st
mi

c L R
 $\begin{pmatrix} 41 & 44 \\ 24 & 42 \\ 67 & 43 \end{pmatrix}$ fr
st
mi

6 a $x = -2$, $y = -2$

b $x = 0$, $y = 0$

7 a $\mathbf{A} + \mathbf{B} = \begin{pmatrix} 1 & 3 \\ 5 & 2 \end{pmatrix}$, $\mathbf{B} + \mathbf{A} = \begin{pmatrix} 1 & 3 \\ 5 & 2 \end{pmatrix}$

8 a $(\mathbf{A} + \mathbf{B}) + \mathbf{C} = \begin{pmatrix} 6 & 3 \\ -1 & 6 \end{pmatrix}$, $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = \begin{pmatrix} 6 & 3 \\ -1 & 6 \end{pmatrix}$

EXERCISE 12B.2

1 a $\begin{pmatrix} 12 & 24 \\ 48 & 12 \end{pmatrix}$ **b** $\begin{pmatrix} 2 & 4 \\ 8 & 2 \end{pmatrix}$

c $\begin{pmatrix} \frac{1}{2} & 1 \\ 2 & \frac{1}{2} \end{pmatrix}$ **d** $\begin{pmatrix} -3 & -6 \\ -12 & -3 \end{pmatrix}$

2 a $\begin{pmatrix} 3 & 5 & 6 \\ 2 & 8 & 7 \end{pmatrix}$ **b** $\begin{pmatrix} 1 & 1 & 4 \\ 0 & 4 & 1 \end{pmatrix}$

c $\begin{pmatrix} 5 & 8 & 11 \\ 3 & 14 & 11 \end{pmatrix}$ **d** $\begin{pmatrix} 5 & 7 & 14 \\ 2 & 16 & 9 \end{pmatrix}$

3 $12\mathbf{F} = \begin{pmatrix} 12 \\ 48 \\ 24 \\ 12 \end{pmatrix}$

4 a $\mathbf{A} = \begin{pmatrix} 75 \\ 27 \\ 102 \end{pmatrix}$ $\begin{matrix} \leftarrow \text{DVD} \\ \leftarrow \text{Blu-ray} \\ \leftarrow \text{games} \end{matrix}$ $\mathbf{B} = \begin{pmatrix} 136 \\ 43 \\ 129 \end{pmatrix}$ $\begin{matrix} \leftarrow \text{DVD} \\ \leftarrow \text{Blu-ray} \\ \leftarrow \text{games} \end{matrix}$

b $5\mathbf{A} + 2\mathbf{B} = \begin{pmatrix} 647 \\ 221 \\ 768 \end{pmatrix}$ $\begin{matrix} \leftarrow \text{DVD} \\ \leftarrow \text{Blu-ray} \\ \leftarrow \text{games} \end{matrix}$

c total weekly average hirings

5 a A B C D **b** A B C D
 $\begin{pmatrix} 35 & 46 & 46 & 69 \\ 58 & 46 & 35 & 86 \\ 46 & 46 & 58 & 58 \\ 12 & 23 & 23 & 17 \end{pmatrix}$ $\begin{pmatrix} 26 & 34 & 34 & 51 \\ 43 & 34 & 26 & 64 \\ 34 & 34 & 43 & 43 \\ 9 & 17 & 17 & 13 \end{pmatrix}$

EXERCISE 12B.3

- 1 a** $3A$ **b** O **c** $-C$ **d** O **e** $2A + 2B$
f $-A - B$ **g** $-2A + C$ **h** $4A - B$ **i** $3B$
2 a $X = A - B$ **b** $X = C - B$ **c** $X = 2C - 4B$
d $X = \frac{1}{2}A$ **e** $X = \frac{1}{3}B$ **f** $X = A - B$
g $X = 2C$ **h** $X = \frac{1}{2}B - A$ **i** $X = \frac{1}{4}(A - C)$

3 a $X = \begin{pmatrix} 3 & 6 \\ 9 & 18 \end{pmatrix}$ **b** $X = \begin{pmatrix} \frac{1}{2} & -\frac{1}{4} \\ \frac{3}{4} & \frac{5}{4} \end{pmatrix}$
c $X = \begin{pmatrix} -1 & -6 \\ 1 & -\frac{1}{2} \end{pmatrix}$

EXERCISE 12C.1

- 1 a** (11) **b** (22) **c** (16) **2 b** $\begin{pmatrix} w & x & y & z \end{pmatrix} \begin{pmatrix} \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{pmatrix}$
3 a $P = \begin{pmatrix} 27 & 35 & 39 \end{pmatrix}$, $Q = \begin{pmatrix} 4 \\ 3 \\ 2 \end{pmatrix}$
b total cost = $\begin{pmatrix} 27 & 35 & 39 \end{pmatrix} \begin{pmatrix} 4 \\ 3 \\ 2 \end{pmatrix} = \291
4 a $P = \begin{pmatrix} 10 & 6 & 3 & 1 \end{pmatrix}$, $N = \begin{pmatrix} 3 \\ 2 \\ 4 \\ 2 \end{pmatrix}$
b total points = $\begin{pmatrix} 10 & 6 & 3 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ 4 \\ 2 \end{pmatrix} = 56 \text{ points}$

EXERCISE 12C.2

- 1** Number of columns in A does not equal number of rows in B .
2 a $m = n$ **b** 2×3 **c** B has 3 columns, A has 2 rows
3 a does not exist **b** (28 29)
4 a (8) **b** $\begin{pmatrix} 2 & 0 & 3 \\ 8 & 0 & 12 \\ 4 & 0 & 6 \end{pmatrix}$
5 a (3 5 3) **b** $\begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}$
6 a $Q = \begin{pmatrix} 32 & 24 \\ 25 & 16 \\ 13 & 9 \end{pmatrix}$ **b** $P = \begin{pmatrix} 1.19 \\ 1.55 \end{pmatrix}$
c $QP = \begin{pmatrix} 32 & 24 \\ 25 & 16 \\ 13 & 9 \end{pmatrix} \begin{pmatrix} 1.19 \\ 1.55 \end{pmatrix} = \begin{pmatrix} 75.28 \\ 54.55 \\ 29.42 \end{pmatrix}$
 It represents the total value of sales for each pen colour.
d $\$75.28 + \$54.55 + \$29.42 = \159.25
7 a $C = \begin{pmatrix} 12.5 \\ 9.5 \end{pmatrix}$ $N = \begin{pmatrix} 2375 & 5156 \\ 2502 & 3612 \end{pmatrix}$
b $\begin{pmatrix} 78\,669.5 \\ 65\,589 \end{pmatrix}$ income from day 1 **c** $\$144\,258.50$ income from day 2
8 a $R = \begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 2 & 3 \end{pmatrix}$ **b** $P = \begin{pmatrix} 7 & 3 & 19 \\ 6 & 2 & 22 \end{pmatrix}$

c $\begin{pmatrix} 48 & 70 \\ 52 & 76 \end{pmatrix}$ **d** $\text{\$}48$ **ii** $\text{\$}76$

- e** The elements of PR tell us that, if all the items are to be bought at one store, it is cheapest to do so at store A for both you and your friend. However, the cheapest way is to buy paint from store A, and hammers and screwdrivers from store B.

EXERCISE 12C.3

- 1 a** $A^2 + A$ **b** $B^2 + 2B$ **c** $A^3 - 2A^2 + A$
d $A^3 + A^2 - 2A$ **e** $AC + AD + BC + BD$
f $A^2 + AB + BA + B^2$ **g** $A^2 - AB + BA - B^2$
h $A^2 + 2A + I$ **i** $9I - 6B + B^2$
2 a $A^3 = 3A - 2I$, $A^4 = 4A - 3I$
b $B^3 = 3B - 2I$, $B^4 = 6I - 5B$, $B^5 = 11B - 10I$
c $C^3 = 13C - 12I$, $C^5 = 121C - 120I$
3 a **i** $I + 2A$ **ii** $2I - 2A$ **iii** $10A + 6I$
b $A^2 + A + 2I$ **c** **i** $-3A$ **ii** $-2A$ **iii** A
4 a $A^2 = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$
b false as $A(A - I) = O$ does not imply that $A = O$ or $A - I = O$
c $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} a & b \\ \frac{a-a^2}{b} & 1-a \end{pmatrix}$, $b \neq 0$
5 For example, $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ gives $A^2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$.
6 a $a = 3$, $b = -4$ **b** $a = 1$, $b = 8$
7 a $p = -2$, $q = 1$ **b** $A^3 = 5A - 2I$
c $A^4 = -12A + 5I$

EXERCISE 12D.1

- 1 a** $\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} = 3I$, $\begin{pmatrix} 1 & -2 \\ -\frac{2}{3} & \frac{5}{3} \end{pmatrix}$
b $\begin{pmatrix} 10 & 0 \\ 0 & 10 \end{pmatrix} = 10I$, $\begin{pmatrix} 0.2 & 0.4 \\ -0.1 & 0.3 \end{pmatrix}$
2 a -2 **b** -1 **c** 0 **d** 1
3 a 26 **b** 6 **c** -1 **d** $a^2 + a$
4 a -3 **b** -3 **c** -12 **5 Hint:** Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$
6 a **i** $\det A = ad - bc$ **ii** $\det B = wz - xy$
iii $AB = \begin{pmatrix} aw + by & ax + bz \\ cw + dy & cx + dz \end{pmatrix}$
iv $\det AB = (ad - bc)(wz - xy)$
7 a $\det A = -2$, $\det B = -1$
b **i** $\det(2A) = -8$ **ii** $\det(-A) = -2$
iii $\det(-3B) = -9$ **iv** $\det(AB) = 2$
8 a $\frac{1}{14} \begin{pmatrix} 5 & -4 \\ 1 & 2 \end{pmatrix}$ **b** $\begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}$ **c** does not exist
d $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ **e** $\frac{1}{10} \begin{pmatrix} 2 & 0 \\ 1 & 5 \end{pmatrix}$ **f** does not exist
g $-\frac{1}{15} \begin{pmatrix} 7 & -2 \\ -4 & -1 \end{pmatrix}$ **h** $\frac{1}{10} \begin{pmatrix} 2 & -4 \\ 1 & 3 \end{pmatrix}$ **i** $\begin{pmatrix} -3 & -1 \\ 2 & 1 \end{pmatrix}$
9 a $\frac{1}{2k+6} \begin{pmatrix} 2 & -1 \\ 6 & k \end{pmatrix}$, $k \neq -3$ **b** $\frac{1}{3k} \begin{pmatrix} k & 1 \\ 0 & 3 \end{pmatrix}$, $k \neq 0$

$$\begin{aligned} \text{c } & \frac{1}{(k+2)(k-1)} \begin{pmatrix} k & -2 \\ -1 & k+1 \end{pmatrix}, \quad k \neq -2 \text{ or } 1 \\ \text{d } & \frac{1}{k(k+1)} \begin{pmatrix} k & -k \\ 3 & k-2 \end{pmatrix}, \quad k \neq 0 \text{ or } -1 \\ \text{e } & \frac{1}{k(2-k)} \begin{pmatrix} 1 & 1-k \\ -2k & k^2 \end{pmatrix}, \quad k \neq 0 \text{ or } 2 \\ \text{f } & \frac{1}{(k+4)(k-1)} \begin{pmatrix} 3k & -2 \\ -k^2-2 & k+1 \end{pmatrix}, \quad k \neq -4 \text{ or } 1 \end{aligned}$$

EXERCISE 12D.2

$$1 \quad \mathbf{X} = \begin{pmatrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 0 \end{pmatrix} \quad 2 \quad \mathbf{a} \quad \mathbf{X} = \mathbf{ABZ} \quad \mathbf{b} \quad \mathbf{Z} = \mathbf{B}^{-1}\mathbf{A}^{-1}\mathbf{X}$$

$$3 \quad \mathbf{A}^2 = 2\mathbf{A} - \mathbf{I}, \quad \mathbf{A}^{-1} = 2\mathbf{I} - \mathbf{A}$$

$$4 \quad \mathbf{a} \quad \mathbf{A}^{-1} = 4\mathbf{I} - \mathbf{A} \quad \mathbf{b} \quad \mathbf{A}^{-1} = 5\mathbf{I} + \mathbf{A}$$

$$\text{c } \mathbf{A}^{-1} = \frac{3}{2}\mathbf{A} - 2\mathbf{I}$$

$$6 \quad \text{If } \mathbf{A}^{-1} \text{ exists, that is, } \det \mathbf{A} \neq 0.$$

EXERCISE 12E

$$1 \quad \mathbf{a} \quad \begin{pmatrix} 3 & -1 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 8 \\ 6 \end{pmatrix}$$

$$\mathbf{b} \quad \begin{pmatrix} 4 & -3 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 11 \\ -5 \end{pmatrix}$$

$$\mathbf{c} \quad \begin{pmatrix} 3 & -1 \\ 2 & 7 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 6 \\ -4 \end{pmatrix}$$

$$2 \quad \mathbf{a} \quad x = \frac{32}{7}, \quad y = \frac{22}{7} \quad \mathbf{b} \quad x = -\frac{37}{23}, \quad y = -\frac{75}{23}$$

$$\mathbf{c} \quad x = \frac{17}{13}, \quad y = -\frac{37}{13} \quad \mathbf{d} \quad x = \frac{59}{13}, \quad y = -\frac{25}{13}$$

$$\mathbf{e} \quad x = -40, \quad y = -24 \quad \mathbf{f} \quad x = \frac{1}{34}, \quad y = \frac{55}{34}$$

$$3 \quad \mathbf{b} \quad \mathbf{i} \quad \mathbf{X} = \begin{pmatrix} -\frac{4}{3} & \frac{13}{9} \\ -1 & \frac{4}{3} \end{pmatrix} \quad \mathbf{ii} \quad \mathbf{X} = \begin{pmatrix} -1 & 3 \\ 2 & 4 \end{pmatrix}$$

$$\mathbf{iii} \quad \mathbf{X} = \begin{pmatrix} \frac{13}{7} & \frac{3}{7} \\ -\frac{2}{7} & -\frac{8}{7} \end{pmatrix} \quad \mathbf{iv} \quad \mathbf{X} = \begin{pmatrix} \frac{19}{7} & \frac{6}{7} \\ \frac{20}{7} & -\frac{25}{7} \end{pmatrix}$$

$$4 \quad \mathbf{a} \quad \mathbf{i} \quad \begin{pmatrix} 2 & -3 \\ 4 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 8 \\ 11 \end{pmatrix}, \quad \det \mathbf{A} = 10$$

$$\mathbf{ii} \quad \text{Yes, } x = 2.5, \quad y = -1$$

$$\mathbf{b} \quad \mathbf{i} \quad \begin{pmatrix} 2 & k \\ 4 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 8 \\ 11 \end{pmatrix}, \quad \det \mathbf{A} = -2 - 4k$$

$$\mathbf{ii} \quad k \neq -\frac{1}{2}, \quad x = \frac{8 + 11k}{2 + 4k}, \quad y = \frac{5}{1 + 2k}$$

$$\mathbf{iii} \quad k = -\frac{1}{2}, \quad \text{no solutions}$$

REVIEW SET 12A

$$1 \quad \mathbf{a} \quad \begin{pmatrix} 4 & 2 \\ -2 & 3 \end{pmatrix} \quad \mathbf{b} \quad \begin{pmatrix} 9 & 6 \\ 0 & -3 \end{pmatrix} \quad \mathbf{c} \quad \begin{pmatrix} -2 & 0 \\ 4 & -8 \end{pmatrix}$$

$$\mathbf{d} \quad \begin{pmatrix} 2 & 2 \\ 2 & -5 \end{pmatrix} \quad \mathbf{e} \quad \begin{pmatrix} -5 & -4 \\ -2 & 6 \end{pmatrix} \quad \mathbf{f} \quad \begin{pmatrix} 7 & 6 \\ 4 & -11 \end{pmatrix}$$

$$\mathbf{g} \quad \begin{pmatrix} -1 & 8 \\ 2 & -4 \end{pmatrix} \quad \mathbf{h} \quad \begin{pmatrix} 3 & 2 \\ -6 & -8 \end{pmatrix} \quad \mathbf{i} \quad \begin{pmatrix} \frac{1}{3} & \frac{2}{3} \\ 0 & -1 \end{pmatrix}$$

$$\mathbf{j} \quad \begin{pmatrix} 9 & 4 \\ 0 & 1 \end{pmatrix} \quad \mathbf{k} \quad \begin{pmatrix} -3 & -10 \\ 6 & 8 \end{pmatrix} \quad \mathbf{l} \quad \begin{pmatrix} \frac{1}{3} & \frac{2}{3} \\ \frac{1}{6} & \frac{1}{12} \end{pmatrix}$$

$$2 \quad \mathbf{a} \quad a = 0, \quad b = 5, \quad c = 1, \quad d = -4$$

$$\mathbf{b} \quad a = 2, \quad b = -1, \quad c = 3, \quad d = 8$$

$$3 \quad \mathbf{a} \quad \mathbf{Y} = \mathbf{B} - \mathbf{A} \quad \mathbf{b} \quad \mathbf{Y} = \frac{1}{2}(\mathbf{A} - \mathbf{C}) \quad \mathbf{c} \quad \mathbf{Y} = \mathbf{A}^{-1}\mathbf{B}$$

$$\mathbf{d} \quad \mathbf{Y} = \mathbf{CB}^{-1} \quad \mathbf{e} \quad \mathbf{Y} = \mathbf{A}^{-1}(\mathbf{C} - \mathbf{B}) \quad \mathbf{f} \quad \mathbf{Y} = \mathbf{B}^{-1}\mathbf{A}$$

$$4 \quad \mathbf{a} \quad 4\mathbf{L} \quad \mathbf{b} \quad -2\mathbf{L}$$

$$5 \quad \mathbf{a} \quad \begin{pmatrix} 10 & -12 \\ -10 & 4 \end{pmatrix} \quad \mathbf{b} \quad \begin{pmatrix} 2 & 6 & -3 \\ -4 & -2 & 11 \end{pmatrix} \quad \mathbf{c} \quad \text{not possible}$$

$$6 \quad \mathbf{a} \quad \mathbf{A} - \mathbf{A}^2 \quad \mathbf{b} \quad \mathbf{AB} + \mathbf{A}^2 - \mathbf{B}^2 - \mathbf{BA}$$

$$\mathbf{c} \quad 4\mathbf{A}^2 - 4\mathbf{A} + \mathbf{I}$$

$$7 \quad \mathbf{A}^3 = 27\mathbf{A} + 10\mathbf{I}, \quad \mathbf{A}^4 = 145\mathbf{A} + 54\mathbf{I}$$

$$8 \quad a = 4, \quad b = -7$$

$$9 \quad \mathbf{a} \quad \begin{pmatrix} \frac{7}{2} & -4 \\ -\frac{5}{2} & 3 \end{pmatrix} \quad \mathbf{b} \quad \text{does not exist} \quad \mathbf{c} \quad \begin{pmatrix} 1 & \frac{5}{3} \\ -2 & -\frac{11}{3} \end{pmatrix}$$

$$10 \quad \text{Unique solution if } k \neq \frac{3}{4}.$$

$$11 \quad \mathbf{a} \quad x = 0, \quad y = -\frac{1}{2} \quad \mathbf{b} \quad x = \frac{12}{7}, \quad y = \frac{13}{7}$$

$$12 \quad \mathbf{b} \quad (\mathbf{A} - \mathbf{I})(\mathbf{A} + 3\mathbf{I}) = 2\mathbf{A} - \mathbf{I}$$

REVIEW SET 12B

$$1 \quad \mathbf{a} \quad \begin{pmatrix} 4 & 2 \\ 2 & 4 \\ 3 & 4 \end{pmatrix} \quad \mathbf{b} \quad \begin{pmatrix} 2 & -2 \\ 0 & 4 \\ -1 & -2 \end{pmatrix} \quad \mathbf{c} \quad \begin{pmatrix} -\frac{3}{2} & 3 \\ \frac{1}{2} & -4 \\ 2 & \frac{7}{2} \end{pmatrix}$$

$$2 \quad \mathbf{a} \quad \mathbf{A} - \mathbf{B} = \begin{pmatrix} 2 & 2 \\ 4 & 1 \\ 3 & 1 \end{pmatrix} \quad \mathbf{b} \quad \mathbf{i} \quad \text{Book 2 (hard cover)} \\ \mathbf{ii} \quad \$101$$

$$4 \quad \mathbf{a} \quad \mathbf{X} = \frac{1}{2}(\mathbf{B} - \mathbf{A}) \quad \mathbf{b} \quad \mathbf{X} = \frac{1}{3}(2\mathbf{B} - 3\mathbf{A})$$

$$\mathbf{c} \quad \mathbf{X} = \frac{1}{4}(\mathbf{B} - \mathbf{A})$$

$$5 \quad \mathbf{X} = \begin{pmatrix} -\frac{1}{2} & -\frac{3}{2} \\ 0 & \frac{3}{2} \end{pmatrix}$$

$$6 \quad \mathbf{a} \quad \begin{pmatrix} 4 & 8 \\ 0 & 2 \\ 6 & 4 \end{pmatrix} \quad \mathbf{b} \quad \begin{pmatrix} 1 & 2 \\ 0 & \frac{1}{2} \\ \frac{3}{2} & 1 \end{pmatrix} \quad \mathbf{c} \quad \begin{pmatrix} 11 & 12 \end{pmatrix}$$

$$\mathbf{d} \quad \mathbf{BA} \text{ does not exist}$$

$$7 \quad \mathbf{a} \quad \det \mathbf{B} \neq 0 \quad \mathbf{b} \quad \mathbf{AB} = \mathbf{BA}$$

$$8 \quad \mathbf{a} \quad \det \mathbf{A} = 5 \quad \mathbf{b} \quad \det(-2\mathbf{A}) = 20 \quad \mathbf{c} \quad \det(\mathbf{A}^2) = 25$$

$$9 \quad \mathbf{a} \quad x = \frac{14}{3}, \quad y = \frac{1}{3} \quad \mathbf{b} \quad x = -1, \quad y = 3$$

$$10 \quad k \in \mathbb{R}, \quad k \neq 3, -2, 2$$

$$11 \quad \text{Unique solution for } k \neq -3 \text{ or } 1. \quad x = \frac{-6}{k-1}, \quad y = \frac{2}{k-1}$$

$$12 \quad \mathbf{A}(\frac{5}{3}\mathbf{A} - 2\mathbf{I}) = \mathbf{I}, \quad \mathbf{A}^{-1} = \frac{5}{3}\mathbf{A} - 2\mathbf{I}$$

EXERCISE 13A

$$1 \quad \mathbf{a} \quad 7 \quad \mathbf{b} \quad 7 \quad \mathbf{c} \quad 11 \quad \mathbf{d} \quad 16 \quad \mathbf{e} \quad 0 \quad \mathbf{f} \quad 5$$

$$2 \quad \mathbf{a} \quad 5 \quad \mathbf{b} \quad 7 \quad \mathbf{c} \quad c$$


$$3 \quad \mathbf{a} \quad -2 \quad \mathbf{b} \quad 7 \quad \mathbf{c} \quad -1 \quad \mathbf{d} \quad 1$$

$$4 \quad \mathbf{a} \quad -3 \quad \mathbf{b} \quad 5 \quad \mathbf{c} \quad -1 \quad \mathbf{d} \quad 6 \quad \mathbf{e} \quad -4 \quad \mathbf{f} \quad -8$$

$$\mathbf{g} \quad 1 \quad \mathbf{h} \quad 2 \quad \mathbf{i} \quad 5$$

EXERCISE 13B

$$1 \quad \mathbf{a} \quad \begin{array}{l} \text{no} \\ \mathbf{b} \quad \text{no} \\ \mathbf{c} \quad \mathbf{i} \quad 0 \text{ ms}^{-1} \\ \mathbf{ii} \quad 9.6 \text{ ms}^{-1} \\ \mathbf{iii} \quad 19.2 \text{ ms}^{-1} \\ \mathbf{iv} \quad 28.8 \text{ ms}^{-1} \end{array}$$



2 a	x	Point B	Gradient of AB
	0	(0, 0)	2
	1	(1, 1)	3
	1.5	(1.5, 2.25)	3.5
	1.9	(1.9, 3.61)	3.9
	1.99	(1.99, 3.9601)	3.99
	1.999	(1.999, 3.996 001)	3.999

x	Point B	Gradient of AB
5	(5, 25)	7
3	(3, 9)	5
2.5	(2.5, 6.25)	4.5
2.1	(2.1, 4.41)	4.1
2.01	(2.01, 4.0401)	4.01
2.001	(2.001, 4.004 001)	4.001

b $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = 4$

The gradient of the tangent to $y = x^2$ at the point (2, 4) is 4.

EXERCISE 13C

- 1 a** $f(2) = 3$ **b** $f'(2) = 0$
2 a $f(0) = 4$ **b** $f'(0) = -1$ **3** $f(2) = 3, f'(2) = 1$

EXERCISE 13D

- 1 a** $f'(x) = 1$ **b** $f'(x) = 0$ **c** $f'(x) = 2$
2 a $\frac{dy}{dx} = -1$ **b** $\frac{dy}{dx} = 2x - 3$ **c** $\frac{dy}{dx} = 4x + 1$
3 a 3 **b** -12 **c** 9 **d** 10

EXERCISE 13E

- 1 a** $f'(x) = 3x^2$ **b** $f'(x) = 6x^2$
c $f'(x) = 14x$ **d** $f'(x) = \frac{3}{\sqrt{x}}$
e $f'(x) = \frac{1}{3\sqrt{x^2}}$ **f** $f'(x) = 2x + 1$
g $f'(x) = -4x$ **h** $f'(x) = 2x + 3$
i $f'(x) = 2x^3 - 12x$ **j** $f'(x) = \frac{6}{x^2}$
k $f'(x) = -\frac{2}{x^2} + \frac{6}{x^3}$ **l** $f'(x) = 2x - \frac{5}{x^2}$
m $f'(x) = 2x + \frac{3}{x^2}$ **n** $f'(x) = -\frac{1}{2x\sqrt{x}}$
o $f'(x) = 8x - 4$ **p** $f'(x) = 3x^2 + 12x + 12$
2 a $\frac{dy}{dx} = 7.5x^2 - 2.8x$ **b** $\frac{dy}{dx} = 2\pi x$
c $\frac{dy}{dx} = -\frac{2}{5x^3}$ **d** $\frac{dy}{dx} = 100$
e $\frac{dy}{dx} = 10$ **f** $\frac{dy}{dx} = 12\pi x^2$
3 a 6 **b** $\frac{3\sqrt{x}}{2}$ **c** $2x - 10$ **d** $2 - 9x^2$ **e** $2x - 1$
f $-\frac{2}{x^3} + \frac{3}{\sqrt{x}}$ **g** $4 + \frac{1}{4x^2}$ **h** $6x^2 - 6x - 5$
4 a 4 **b** $-\frac{16}{729}$ **c** -7 **d** $\frac{13}{4}$ **e** $\frac{1}{8}$ **f** -11
5 $b = 3, c = -4$

- 6 a** $f'(x) = \frac{2}{\sqrt{x}} + 1$ **b** $f'(x) = \frac{1}{3\sqrt[3]{x^2}}$
c $f'(x) = \frac{1}{x\sqrt{x}}$ **d** $f'(x) = 2 - \frac{1}{2\sqrt{x}}$
e $f'(x) = -\frac{2}{x\sqrt{x}}$ **f** $f'(x) = 6x - \frac{3}{2}\sqrt{x}$
g $f'(x) = \frac{-25}{2x^3\sqrt{x}}$ **h** $f'(x) = 2 + \frac{9}{2x^2\sqrt{x}}$

- 7 a** $\frac{dy}{dx} = 4 + \frac{3}{x^2}$, $\frac{dy}{dx}$ is the gradient function of $y = 4x - \frac{3}{x}$ from which the gradient at any point can be found.
b $\frac{dS}{dt} = 4t + 4$ ms⁻¹, $\frac{dS}{dt}$ is the instantaneous rate of change in position at the time t , or the velocity function.
c $\frac{dC}{dx} = 3 + 0.004x$ \$ per toaster, $\frac{dC}{dx}$ is the instantaneous rate of change in cost as the number of toasters changes.

EXERCISE 13F.1

- 1 a** $gf(x) = (2x + 7)^2$ **b** $gf(x) = 2x^2 + 7$
c $gf(x) = \sqrt{3 - 4x}$ **d** $gf(x) = 3 - 4\sqrt{x}$
e $gf(x) = \frac{2}{x^2 + 3}$ **f** $gf(x) = \frac{4}{x^2} + 3$
2 Note: There may be other answers.
a $g(x) = x^3, f(x) = 3x + 10$
b $g(x) = \frac{1}{x}, f(x) = 2x + 4$
c $g(x) = \sqrt{x}, f(x) = x^2 - 3x$
d $g(x) = \frac{10}{x^3}, f(x) = 3x - x^2$

EXERCISE 13F.2

- 1 a** $u^{-2}, u = 2x - 1$ **b** $u^{\frac{1}{2}}, u = x^2 - 3x$
c $2u^{-\frac{1}{2}}, u = 2 - x^2$ **d** $u^{\frac{1}{3}}, u = x^3 - x^2$
e $4u^{-3}, u = 3 - x$ **f** $10u^{-1}, u = x^2 - 3$
2 a $\frac{dy}{dx} = 8(4x - 5)$ **b** $\frac{dy}{dx} = 2(5 - 2x)^{-2}$
c $\frac{dy}{dx} = \frac{1}{2}(3x - x^2)^{-\frac{1}{2}} \times (3 - 2x)$
d $\frac{dy}{dx} = -12(1 - 3x)^3$ **e** $\frac{dy}{dx} = -18(5 - x)^2$
f $\frac{dy}{dx} = \frac{1}{3}(2x^3 - x^2)^{-\frac{2}{3}} \times (6x^2 - 2x)$
g $\frac{dy}{dx} = -60(5x - 4)^{-3}$
h $\frac{dy}{dx} = -4(3x - x^2)^{-2} \times (3 - 2x)$
i $\frac{dy}{dx} = 6 \left(x^2 - \frac{2}{x} \right)^2 \times \left(2x + \frac{2}{x^2} \right)$
3 a $-\frac{1}{\sqrt{3}}$ **b** -18 **c** -8 **d** -4 **e** $-\frac{3}{32}$ **f** 0
4 a 3, $b = 1$ **5 a** 2, $b = 1$
6 a $\frac{dy}{dx} = 3x^2, \frac{dx}{dy} = \frac{1}{3}y^{-\frac{2}{3}}$ **Hint:** Substitute $y = x^3$
b $\frac{dy}{dx} \times \frac{dx}{dy} = \frac{dy}{dy}$ {chain rule} = 1

EXERCISE 13G

- 1 a $f'(x) = 2x - 1$ b $f'(x) = 4x + 2$
 c $f'(x) = 2x(x+1)^{\frac{1}{2}} + \frac{1}{2}x^2(x+1)^{-\frac{1}{2}}$
 2 a $\frac{dy}{dx} = 2x(2x-1) + 2x^2$
 b $\frac{dy}{dx} = 4(2x+1)^3 + 24x(2x+1)^2$
 c $\frac{dy}{dx} = 2x(3-x)^{\frac{1}{2}} - \frac{1}{2}x^2(3-x)^{-\frac{1}{2}}$
 d $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}}(x-3)^2 + 2\sqrt{x}(x-3)$
 e $\frac{dy}{dx} = 10x(3x^2-1)^2 + 60x^3(3x^2-1)$
 f $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}}(x-x^2)^3 + 3\sqrt{x}(x-x^2)^2(1-2x)$
 3 a -48 b $406\frac{1}{4}$ c $\frac{13}{3}$ d $\frac{11}{2}$
 4 b $x = 3$ or $\frac{3}{5}$ c $x \leq 0$ 5 $x = -1$ and $x = -\frac{5}{3}$

EXERCISE 13H

- 1 a $\frac{dy}{dx} = \frac{7}{(2-x)^2}$ b $\frac{dy}{dx} = \frac{2x(2x+1) - 2x^2}{(2x+1)^2}$
 c $\frac{dy}{dx} = \frac{(x^2-3) - 2x^2}{(x^2-3)^2}$
 d $\frac{dy}{dx} = \frac{\frac{1}{2}x^{-\frac{1}{2}}(1-2x) + 2\sqrt{x}}{(1-2x)^2}$
 e $\frac{dy}{dx} = \frac{2x(3x-x^2) - (x^2-3)(3-2x)}{(3x-x^2)^2}$
 f $\frac{dy}{dx} = \frac{(1-3x)^{\frac{1}{2}} + \frac{3}{2}x(1-3x)^{-\frac{1}{2}}}{1-3x}$
 2 a 1 b 1 c $-\frac{7}{324}$ d $-\frac{28}{27}$
 3 b i never { $\frac{dy}{dx}$ is undefined at $x = -1$ }
 ii $x \leq 0$ and $x = 1$
 4 b i $x = -2 \pm \sqrt{11}$ ii $x = -2$

EXERCISE 13I

- 1 a $f'(x) = 4e^{4x}$ b $f'(x) = e^x$
 c $f'(x) = -2e^{-2x}$ d $f'(x) = \frac{1}{2}e^{\frac{x}{2}}$
 e $f'(x) = -e^{-\frac{x}{2}}$ f $f'(x) = 2e^{-x}$
 g $f'(x) = 2e^{\frac{x}{2}} + 3e^{-x}$ h $f'(x) = \frac{e^x - e^{-x}}{2}$
 i $f'(x) = -2xe^{-x^2}$ j $f'(x) = e^{\frac{1}{x}} \times \frac{-1}{x^2}$
 k $f'(x) = 20e^{2x}$ l $f'(x) = 40e^{-2x}$
 m $f'(x) = 2e^{2x+1}$ n $f'(x) = \frac{1}{4}e^{\frac{x}{4}}$
 o $f'(x) = -4xe^{1-2x^2}$ p $f'(x) = -0.02e^{-0.02x}$
 2 a $e^x + xe^x$ b $3x^2e^{-x} - x^3e^{-x}$
 c $\frac{xe^x - e^x}{x^2}$ d $\frac{1-x}{e^x}$
 e $2xe^{3x} + 3x^2e^{3x}$ f $\frac{xe^x - \frac{1}{2}e^x}{x\sqrt{x}}$
 g $\frac{1}{2}x^{-\frac{1}{2}}e^{-x} - x^{\frac{1}{2}}e^{-x}$ h $\frac{e^x + 2 + 2e^{-x}}{(e^{-x} + 1)^2}$

- 3 a 108 b -1 c $\frac{9}{\sqrt{19}}$ 4 $k = -9$

- 5 a $\frac{dy}{dx} = 2^x \ln 2$ 6 P = (0, 0) or $(2, \frac{4}{e^2})$

EXERCISE 13J

- 1 a $\frac{dy}{dx} = \frac{1}{x}$ b $\frac{dy}{dx} = \frac{2}{2x+1}$ c $\frac{dy}{dx} = \frac{1-2x}{x-x^2}$
 d $\frac{dy}{dx} = -\frac{2}{x}$ e $\frac{dy}{dx} = 2x \ln x + x$
 f $\frac{dy}{dx} = \frac{1-\ln x}{2x^2}$ g $\frac{dy}{dx} = e^x \ln x + \frac{e^x}{x}$
 h $\frac{dy}{dx} = \frac{2 \ln x}{x}$ i $\frac{dy}{dx} = \frac{1}{2x\sqrt{\ln x}}$
 j $\frac{dy}{dx} = \frac{e^{-x}}{x} - e^{-x} \ln x$ k $\frac{dy}{dx} = \frac{\ln(2x)}{2\sqrt{x}} + \frac{1}{\sqrt{x}}$
 l $\frac{dy}{dx} = \frac{\ln x - 2}{\sqrt{x}(\ln x)^2}$ m $\frac{dy}{dx} = \frac{4}{1-x}$
 n $\frac{dy}{dx} = \ln(x^2+1) + \frac{2x^2}{x^2+1}$
 2 a $\frac{dy}{dx} = \ln 5$ b $\frac{dy}{dx} = \frac{3}{x}$ c $\frac{dy}{dx} = \frac{4x^3+1}{x^4+x}$
 d $\frac{dy}{dx} = \frac{1}{x-2}$ e $\frac{dy}{dx} = \frac{6}{2x+1} [\ln(2x+1)]^2$
 f $\frac{dy}{dx} = \frac{1-\ln(4x)}{x^2}$ g $\frac{dy}{dx} = -\frac{1}{x}$
 h $\frac{dy}{dx} = \frac{1}{x \ln x}$ i $\frac{dy}{dx} = \frac{-1}{x(\ln x)^2}$
 3 a $\frac{dy}{dx} = \frac{-1}{1-2x}$ b $\frac{dy}{dx} = \frac{-2}{2x+3}$ c $\frac{dy}{dx} = 1 + \frac{1}{2x}$
 d $\frac{dy}{dx} = \frac{1}{x} - \frac{1}{2(2-x)}$ e $\frac{dy}{dx} = \frac{1}{x+3} - \frac{1}{x-1}$
 f $\frac{dy}{dx} = \frac{2}{x} + \frac{1}{3-x}$ g $f'(x) = \frac{9}{3x-4}$
 h $f'(x) = \frac{1}{x} + \frac{2x}{x^2+1}$ i $f'(x) = \frac{2x+2}{x^2+2x} - \frac{1}{x-5}$
 4 a 2 b $-\frac{5}{3}$ 5 a = 3, b = -e

EXERCISE 13K

- 1 a $\frac{dy}{dx} = 2 \cos(2x)$ b $\frac{dy}{dx} = \cos x - \sin x$
 c $\frac{dy}{dx} = -3 \sin(3x) - \cos x$ d $\frac{dy}{dx} = \cos(x+1)$
 e $\frac{dy}{dx} = 2 \sin(3-2x)$ f $\frac{dy}{dx} = \frac{5}{\cos^2(5x)}$
 g $\frac{dy}{dx} = \frac{1}{2} \cos\left(\frac{x}{2}\right) + 3 \sin x$ h $\frac{dy}{dx} = \frac{3\pi}{\cos^2(\pi x)}$
 i $\frac{dy}{dx} = 4 \cos x + 2 \sin(2x)$
 2 a $2x - \sin x$ b $\frac{1}{\cos^2 x} - 3 \cos x$
 c $e^x \cos x - e^x \sin x$ d $-e^{-x} \sin x + e^{-x} \cos x$
 e $\frac{\cos x}{\sin x}$ f $2e^{2x} \tan x + \frac{e^{2x}}{\cos^2 x}$ g $3 \cos(3x)$
 h $-\frac{1}{2} \sin\left(\frac{x}{2}\right)$ i $\frac{6}{\cos^2(2x)}$ j $\cos x - x \sin x$
 k $\frac{x \cos x - \sin x}{x^2}$ l $\tan x + \frac{x}{\cos^2 x}$

- 3 a** $2x \cos(x^2)$ **b** $-\frac{1}{2\sqrt{x}} \sin(\sqrt{x})$ **c** $-\frac{\sin x}{2\sqrt{\cos x}}$
d $2 \sin x \cos x$ **e** $-3 \sin x \cos^2 x$
f $-\sin x \sin(2x) + 2 \cos x \cos(2x)$
g $\sin x \sin(\cos x)$ **h** $-12 \sin(4x) \cos^2(4x)$
i $-\frac{\cos x}{\sin^2 x}$ **j** $\frac{2 \sin(2x)}{\cos^2(2x)}$
k $-\frac{8 \cos(2x)}{\sin^3(2x)}$ **l** $-\frac{12}{\cos^2(\frac{\pi}{2}) \tan^4(\frac{\pi}{2})}$
- 4 a** $-\frac{9}{8}$ **b** 0

EXERCISE 13L

- 1 a** $f''(x) = 6$ **b** $f''(x) = \frac{3}{2x^{\frac{5}{2}}}$
c $f''(x) = 12x - 6$ **d** $f''(x) = \frac{12 - 6x}{x^4}$
e $f''(x) = 24 - 48x$ **f** $f''(x) = \frac{20}{(2x - 1)^3}$
- 2 a** $\frac{d^2y}{dx^2} = -6x$ **b** $\frac{d^2y}{dx^2} = 2 - \frac{30}{x^4}$
c $\frac{d^2y}{dx^2} = -\frac{9}{4}x^{-\frac{5}{2}}$ **d** $\frac{d^2y}{dx^2} = \frac{8}{x^3}$
e $\frac{d^2y}{dx^2} = 6(x^2 - 3x)(5x^2 - 15x + 9)$
f $\frac{d^2y}{dx^2} = 2 + \frac{2}{(1 - x)^3}$
- 3 a** $f(2) = 9$ **b** $f'(2) = 10$ **c** $f''(2) = 12$
- 5 a** $x = 1$ **6**

x	-1	0	1
$f(x)$	-	0	+
$f'(x)$	+	-	+
$f''(x)$	-	0	+
- b** $x = 0, \pm\sqrt{6}$

- 7 b** $f''(x) = 3 \sin x \cos 2x + 6 \cos x \sin 2x$
8 a $\frac{d^2y}{dx^2} = \frac{1}{x^2}$ **b** $\frac{d^2y}{dx^2} = \frac{1}{x}$
c $\frac{d^2y}{dx^2} = \frac{2}{x^2}(1 - \ln x)$
9 a $f(1) = 0$ **b** $f'(1) = 3$ **c** $f''(1) = 0$

10 Hint: Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ and substitute into the equation.

REVIEW SET 13A

- 1 a** -1 **b** -1 **c** 8
2 a $f'(x) = 2x + 2$ **b** $\frac{dy}{dx} = -6x$
3 a $f'(t) = -9.6t \text{ ms}^{-1}$
b $f'(2) = -19.2 \text{ ms}^{-1}$
 (the negative sign indicates travelling downwards)
4 a $f(3) = -17$ **b** $f'(3) = -17$ **c** $f''(3) = -6$
5 a $\frac{dy}{dx} = 6x - 4x^3$ **b** $\frac{dy}{dx} = 1 + \frac{1}{x^2}$
6 (0, 0) **7 a** $\frac{dy}{dx} = 3x^2e^{x^3+2}$ **b** $\frac{dy}{dx} = \frac{1}{x+3} - \frac{2}{x}$
9 a $5 + 3x^{-2}$ **b** $4(3x^2 + x)^3(6x + 1)$
c $2x(1 - x^2)^3 - 6x(x^2 + 1)(1 - x^2)^2$

- 10** (-2, 19) and (1, -2)

11 a $\frac{dy}{dx} = -2(5 - 4x)^{-\frac{1}{2}}$ **b** $\frac{d^2y}{dx^2} = -4(5 - 4x)^{-\frac{3}{2}}$

12 a $5 \cos(5x) \ln x + \frac{\sin(5x)}{x}$
b $\cos x \cos(2x) - 2 \sin x \sin(2x)$
c $-2e^{-2x} \tan x + \frac{e^{-2x}}{\cos^2 x}$

13 $\frac{\sqrt{3}}{2}$

14 a $f'(x) = 8x(x^2 + 3)^3$
b $g'(x) = \frac{\frac{1}{2}x(x+5)^{-\frac{1}{2}} - 2(x+5)^{\frac{1}{2}}}{x^3}$

15 a $f''(2) = \frac{23}{4}$ **b** $f''(2) = -\frac{1}{8\sqrt{2}}$

16 a $10 - 10 \cos(10x)$ **b** $\tan x$
c $5 \cos(5x) \ln(2x) + \frac{\sin(5x)}{x}$

REVIEW SET 13B

- 1 a** -3 **b** 3 **c** -1 **2** $f'(1) = 3$
3 a $\frac{dy}{dx} = 4x$ **b** when $x = 4$, gradient = 16
c when gradient = -12, $x = -3$
4 a $\frac{dy}{dx} = 3x^2(1 - x^2)^{\frac{1}{2}} - x^4(1 - x^2)^{-\frac{1}{2}}$
b $\frac{dy}{dx} = \frac{(2x - 3)(x + 1)^{\frac{1}{2}} - \frac{1}{2}(x^2 - 3x)(x + 1)^{-\frac{1}{2}}}{x + 1}$
5 a $\frac{d^2y}{dx^2} = 36x^2 - \frac{4}{x^3}$ **b** $\frac{d^2y}{dx^2} = 6x + \frac{3}{4}x^{-\frac{5}{2}}$
6 (1, e) **7 a** $f'(x) = \frac{e^x}{e^x + 3}$ **b** $f'(x) = \frac{3}{x + 2} - \frac{1}{x}$
8 When $x = 1$, $\frac{dy}{dx} = 0$.

9 a $\frac{dy}{dx} = \frac{3x^2 - 3}{x^3 - 3x}$ **b** $\frac{dy}{dx} = \frac{e^x(x - 2)}{x^3}$

10 $x = -\frac{1}{2}, \frac{3}{2}$

11 a $f(\pi) = \pi + 1$ **b** $f'(\frac{\pi}{2}) = 2$ **c** $f''(\frac{3\pi}{4}) = -\frac{\sqrt{2}}{2}$

12 a $f'(x) = \frac{1}{2}x^{-\frac{1}{2}} \cos(4x) - 4x^{\frac{1}{2}} \sin(4x)$,
 $f''(x) = -\frac{1}{4}x^{-\frac{3}{2}} \cos(4x) - 4x^{-\frac{1}{2}} \sin(4x)$
 $- 16x^{\frac{1}{2}} \cos(4x)$
b $f'(\frac{\pi}{16}) \approx -0.455$, $f''(\frac{\pi}{8}) \approx -6.38$

14 a $x = -6 \pm \sqrt{33}$ **b** $x = \pm\sqrt{3}$ **c** $x = 0, \pm 3$

15 a $f(x) = -5 \sin 4x$
b $f'(x) = 0$ when $x = \frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}$, $0 \leq x \leq \pi$

16 $\frac{dy}{dx} = 3b \cos(bx) + 2a \sin(2x)$, $a = 2$, $b = \pm 1$













EXERCISE 14A

- 1 a** $y = -7x + 11$ **b** $x - 4y = -8$ **c** $y = -2x - 2$
d $y = -2x + 6$ **e** $y = -5x - 9$ **f** $y = -5x - 1$
2 a $x + 6y = 57$ **b** $x + 7y = 26$ **c** $x - 3y = -11$
d $x + 6y = 43$

- 3 $y = 21$ and $y = -6$
 5 $k = -5$
 7 $a = -4$, $b = 7$
 10 **a** $x - 3y = -5$
c $x - 16y = 3$
 11 **a** $y = 2x - \frac{7}{4}$
c $4x + 57y = 1042$
 12 $a = 4$, $b = 3$
 13 **a** $x + ey = 2$
c $2x + e^2y = \frac{2}{e^2} - e^2$
 15 **a** $y = x$ **b** $y = x$ **c** $2x - y = \frac{\pi}{3} - \frac{\sqrt{3}}{2}$ **d** $x = \frac{\pi}{4}$
 16 **a** $(-4, -64)$ **b** $(4, -31)$
 17 **a** $f'(x) = 2x - \frac{8}{x^3}$ **b** $x = \pm\sqrt{2}$ **c** tangent is $y = 4$
 18 A is $(\frac{2}{3}, 0)$, B is $(0, -2e)$
 19 **a** $y = (2a - 1)x - a^2 + 9$
b $y = 5x$, contact at $(3, 15)$, $y = -7x$, contact at $(-3, 21)$
 20 $y = 0$, $y = 27x + 54$ **21** $y = -\sqrt{14}x + 4\sqrt{14}$
 22 $y = e^ax + e^a(1 - a)$ so $y = ex$ is the tangent to $y = e^x$ from the origin.
 23 **a** **Hint:** They must have the same y -coordinate at $x = b$ and the same gradient.
c $a = \frac{1}{2e}$ **d** $y = e^{-\frac{1}{2}}x - \frac{1}{2}$
 24 $\approx 63.43^\circ$
 25 **a** **Hint:** $y = f(a) + f'(a)(x - a)$
b **Hint:** Expand $f(x) = 4 - 8(x + 1) - (x + 1)^2 + 2(x + 1)^3$
c Notice the first 2 terms in **b** are the same as the tangent line found in part **a**.

EXERCISE 14B

- 1 **a** A - local max, B - stationary inflection, C - local min.



- 3 $x = -\frac{b}{2a}$, local min if $a > 0$, local max if $a < 0$

4 $a = 9$

5 **a** $a = -12$, $b = -13$

b $(-2, 3)$ local max., $(2, -29)$ local min.

6 **a** local maximum at $(1, e^{-1})$

b local maximum at $(-2, 4e^{-2})$, local minimum at $(0, 0)$

c local minimum at $(1, e)$





d local maximum at $(-1, e)$


7 **a** $x > 0$

8 **a** Greatest value is 63 when $x = 5$, least value is -18 when $x = 2$.

b Greatest value is 4 when $x = 3$ and $x = 0$, least value is -16 when $x = -2$.

9 $P(x) = -9x^3 - 9x^2 + 9x + 2$





- 11 Hint:** Find $\frac{dy}{dx}$, then determine the nature of the stationary points.
- 12 Hint:** Show that as $x \rightarrow 0$, $f(x) \rightarrow -\infty$, and as $x \rightarrow \infty$, $f(x) \rightarrow 0$.
- 13 a Hint:** Find $f'(x)$, then determine the nature of the stationary points.
- b Hint:** Show that $f(x) \geq 1$ for all $x > 0$.

EXERCISE 14C.1

- 1 a** 7 m s^{-1} **b** $(h + 5) \text{ m s}^{-1}$
- c** $5 \text{ m s}^{-1} = s'(1)$ is the instantaneous velocity at $t = 1$ s
- d** average velocity $= (2t + h + 3) \text{ m s}^{-1}$,
 $\lim_{h \rightarrow 0} (2t + h + 3) = 2t + 3 \text{ m s}^{-1}$ is the instantaneous velocity at time t seconds.
- 2 a** -14 cm s^{-1} **b** $(-8 - 2h) \text{ cm s}^{-1}$
- c** $-8 \text{ cm s}^{-1} = s'(2)$
 \therefore instantaneous velocity $= -8 \text{ cm s}^{-1}$ at $t = 2$
- d** $-4t = s'(t) = v(t)$ is the instantaneous velocity at time t seconds.
- 3 a** $\frac{2}{3} \text{ cm s}^{-2}$ **b** $\frac{2\sqrt{1+h}-2}{h} \text{ cm s}^{-2}$
- c** $1 \text{ cm s}^{-2} = v'(1)$ is the instantaneous accn. at $t = 1$ s
- d** $\frac{1}{\sqrt{t}} \text{ cm s}^{-2} = v'(t)$, the instantaneous accn. at time t
- 4 a** velocity at $t = 4$ **b** acceleration at $t = 4$

EXERCISE 14C.2

- 1 a** $v(t) = 2t - 4 \text{ cm s}^{-1}$, $a(t) = 2 \text{ cm s}^{-2}$
-
- b** $s(0) = 3 \text{ cm}$, $v(0) = -4 \text{ cm s}^{-1}$, $a(0) = 2 \text{ cm s}^{-2}$
 The object is initially 3 cm to the right of the origin and is moving to the left at 4 cm s^{-1} . It is accelerating at 2 cm s^{-2} to the right.
- c** $s(2) = -1 \text{ cm}$, $v(2) = 0 \text{ cm s}^{-1}$, $a(2) = 2 \text{ cm s}^{-2}$
 The object is instantaneously stationary, 1 cm to the left of the origin and is accelerating to the right at 2 cm s^{-2} .
- d** At $t = 2$, $s(2) = 1 \text{ cm}$ to the left of the origin.
- e**
-
- f** $0 \leq t \leq 2$
- 2 a** $v(t) = 98 - 9.8t \text{ m s}^{-1}$, $a(t) = -9.8 \text{ m s}^{-2}$
-
- b** $s(0) = 0 \text{ m}$ above the ground, $v(0) = 98 \text{ m s}^{-1}$ skyward
- c** $t = 5 \text{ s}$ Stone is 367.5 m above the ground and moving skyward at 49 m s^{-1} . Its speed is decreasing.
 $t = 12 \text{ s}$ Stone is 470.4 m above the ground and moving groundward at 19.6 m s^{-1} . Its speed is increasing.


- d** 490 m **e** 20 seconds

- 3 a** 1.2 m
- b** $s'(t) = 28.1 - 9.8t$ represents the instantaneous velocity of the ball.
- c** $t = 2.87 \text{ s}$. The ball has reached its maximum height and is instantaneously at rest.
- d** 41.5 m
- e** **i** 28.1 m s^{-1} **ii** 8.5 m s^{-1} **iii** 20.9 m s^{-1}
 $s'(t) \geq 0$ when the ball is travelling upwards.
 $s'(t) \leq 0$ when the ball is travelling downwards.
- f** 5.78 s
- g** $s''(t)$ is the rate of change of $s'(t)$, or the instantaneous acceleration.

- 4 a** $v(t) = 3t^2 - 18t + 24 \text{ m s}^{-1}$ $a(t) = 6t - 18 \text{ m s}^{-2}$




- b** $x(2) = 20$, $x(4) = 16$



- c** **i** $0 \leq t \leq 2$ and $3 \leq t \leq 4$ **ii** $0 \leq t \leq 3$
- d** 28 m

- 5 a** $v(t) = 100 - 40e^{-\frac{t}{5}} \text{ cm s}^{-1}$, $a(t) = 8e^{-\frac{t}{5}} \text{ cm s}^{-2}$
- b** $s(0) = 200 \text{ cm}$ on positive side of origin
 $v(0) = 60 \text{ cm s}^{-1}$, $a(0) = 8 \text{ cm s}^{-2}$

- c**



- 6 a** $x(0) = -1 \text{ cm}$, $v(0) = 0 \text{ cm s}^{-1}$, $a(0) = 2 \text{ cm s}^{-2}$
- b** At $t = \frac{\pi}{4}$ seconds, the particle is $(\sqrt{2} - 1) \text{ cm}$ left of the origin, moving right at $\sqrt{2} \text{ cm s}^{-1}$, with increasing speed.
- c** changes direction when $t = \pi$, $x(\pi) = 3 \text{ cm}$
- d** $0 \leq t \leq \frac{\pi}{2}$ and $\pi \leq t \leq \frac{3\pi}{2}$


- 7 Hint:** Assume that $s(t) = at^2 + bt + c$
 $s'(t) = v(t)$ and $s''(t) = a(t) = g$
 Show that $a = \frac{1}{2}g$, $b = v(0)$, $c = 0$.

- 8 a** 0.675 s
- b** **i** $S'(t) = u + at \text{ m s}^{-1}$ **ii** $t = -\frac{u}{a} \text{ s}$
- iii** $a = -\frac{640}{99} \approx -6.46 \text{ m s}^{-2}$
- iv Hint:** Substitute $t = -\frac{u}{a}$ into $S(t)$.

- v** If the speed u is doubled, then the braking distance is quadrupled ($2^2 = 4$ times).



EXERCISE 14D

- 1 a** \$118 000 **b** $\frac{dP}{dt} = 4t - 12$, \$1000s per year
- c** $\frac{dP}{dt}$ is the rate of change in profit with time
- d** **i** $0 \leq t \leq 3$ years **ii** $t > 3$ years
- e** minimum profit is \$100 000 when $t = 3$

- f** $\left. \frac{dP}{dt} \right|_{t=4} = 4$ Profit is increasing at \$4000 per year after 4 years.
 $\left. \frac{dP}{dt} \right|_{t=10} = 28$ Profit is increasing at \$28 000 per year after 10 years.
 $\left. \frac{dP}{dt} \right|_{t=25} = 88$ Profit is increasing at \$88 000 per year after 25 years.
- 2 a** **i** $Q(0) = 100$ **ii** $Q(25) = 50$ **iii** $Q(100) = 0$
b **i** decr. 1 unit per year **ii** decr. $\frac{1}{\sqrt{2}}$ units per year
c $Q'(t) = -\frac{5}{\sqrt{t}} < 0$
- 3 a** 0.5 m
b $t = 4$: 9.17 m, $t = 8$: 12.5 m, $t = 12$: 14.3 m
c $t = 0$: 3.9 m year⁻¹, $t = 5$: 0.975 m year⁻¹, $t = 10$: 0.433 m year⁻¹
d As $\frac{dH}{dt} = \frac{97.5}{(t+5)^2} > 0$ for all $t \geq 0$, the tree is always growing.
- 4 a** $C'(x) = 0.0009x^2 + 0.04x + 4$ dollars per pair
b $C'(220) = \$56.36$ per pair. This estimates the additional cost of making one more pair of jeans if 220 pairs are currently being made.
c \$56.58 This is the actual increase in cost to make an extra pair of jeans (221 rather than 220).
d $C''(x) = 0.0018x + 0.04$
 $C''(x) = 0$ when $x = -22.2$. This is where the rate of change is a minimum, however it is out of the bounds of the model (you cannot make < 0 jeans!).
- 5 a** **i** €4500 **ii** €4000
b **i** decrease of €210.22 per km h⁻¹
ii increase of €11.31 per km h⁻¹
c $\frac{dC}{dv} = 0$ at $v = \sqrt[3]{500\,000} \approx 79.4$ km h⁻¹
- 6 a** $\frac{dV}{dt} = -1250 \left(1 - \frac{t}{80}\right)$ L min⁻¹
b at $t = 0$ when the tap was first opened
c $\frac{d^2V}{dt^2} = \frac{125}{8}$ L min⁻²
- 
- This shows that the rate of change of V is constantly increasing, so the outflow is decreasing at a constant rate.
- 7 a** The near part of the lake is 2 km from the sea, the furthest part is 3 km.
b $\frac{dy}{dx} = \frac{3}{10}x^2 - x + \frac{3}{5}$
 $\left. \frac{dy}{dx} \right|_{x=\frac{1}{2}} = 0.175$, height of hill is increasing as gradient is positive.
 $\left. \frac{dy}{dx} \right|_{x=1\frac{1}{2}} = -0.225$, height of hill is decreasing as gradient is negative.
 \therefore top of the hill is between $x = \frac{1}{2}$ and $x = 1\frac{1}{2}$.
c 2.55 km from the sea, 63.1 m deep
- 8 a** $k = \frac{1}{50} \ln 2 \approx 0.0139$
b **i** 20 grams **ii** 14.3 grams **iii** 1.95 grams
c 9 days and 6 minutes (216 hours)
d **i** -0.0693 g h⁻¹ **ii** -2.64×10^{-7} g h⁻¹
e Hint: You should find $\frac{dW}{dt} = -\frac{1}{50} \ln 2 \times 20e^{-\frac{1}{50} \ln 2t}$

- 9 a** $k = \frac{1}{15} \ln \left(\frac{19}{3}\right) \approx 0.123$ **b** 100°C
c $c = -k \approx -0.123$
d **i** decreasing at 11.7°C min⁻¹
ii decreasing at 3.42°C min⁻¹
iii decreasing at 0.998°C min⁻¹
- 10 a** 43.9 cm **b** 10.4 years
c **i** growing at 5.45 cm per year
ii growing at 1.88 cm per year
- 11 a** $A(0) = 0$
b **i** $k = \frac{\ln 2}{3}$ (≈ 0.231)
ii 0.728 litres of alcohol produced per hour
- 12** $\frac{21}{\sqrt{2}}$ cm² per radian
- 13 a** rising at 2.73 m per hour **b** rising
- 14 a** **i** 0 **ii** 1 **iii** ≈ 1.11

EXERCISE 14E

- 1** 250 items
- 2 b** $L_{\min} \approx 28.3$ m, $x \approx 7.07$ m **c**
- 
- 3** 10 blankets **4** 14.8 km h⁻¹ **5** at 4.41 months old
- 6 a** Hint: $V = 200 = 2x \times x \times h$
b Hint: Show $h = \frac{100}{x^2}$ and substitute into the surface area equation.
c $SA_{\min} \approx 213$ cm², $x \approx 4.22$ cm **d**
- 
- 7** 20 kettles **8** $C\left(\frac{1}{\sqrt{2}}, e^{-\frac{1}{2}}\right)$
- 9 a** Recall that $V_{\text{cylinder}} = \pi r^2 h$ and that 1 L = 1000 cm³.
b Recall that $SA_{\text{cylinder}} = 2\pi r^2 + 2\pi r h$.
c radius ≈ 5.42 cm, height ≈ 10.8 cm
- 10 b** $\theta \approx 1.91$, $A \approx 237$ cm² **11 b** 6 cm \times 6 cm
- 12 a** $0 \leq x \leq 63.7$
b $l = 100$ m, $x = \frac{100}{\pi} \approx 31.83$ m, $A = \frac{20\,000}{\pi} \approx 6366$ m²
- 13** after 13.8 weeks **14** after 40 minutes
- 15 c** $\theta = 30^\circ$, $A \approx 130$ cm²
- 16 a** Hint: Show that $AC = \frac{\theta}{360} \times 2\pi \times 10$
b Hint: Show that $2\pi r = AC$
c Hint: Use the result from **b** and Pythagoras' theorem.
d $V = \frac{1}{3}\pi \left(\frac{\theta}{36}\right)^2 \sqrt{100 - \left(\frac{\theta}{36}\right)^2}$ **e** $\theta \approx 294^\circ$
- 17** 1 hour 34 min 53 s when $\theta \approx 36.9^\circ$ **18** 9.87 m

EXERCISE 14F

- 1** a is decreasing at 7.5 units per second
2 increasing at 1 cm per minute
3 a 4 π m² per second **b** 8 π m² per second
4 increasing at 6 π m² per minute

5 decreasing at 0.16 m^3 per minute 6 $\frac{20}{3} \text{ cm}$ per minute

7 $\frac{25\sqrt{3}}{6} \approx 7.22 \text{ cm}$ per minute

8 decreasing at $\frac{250}{13} \approx 19.2 \text{ m s}^{-1}$

9 a 0.2 m s^{-1} b $\frac{4}{45} \text{ m s}^{-1}$

10 decreasing at $\frac{\sqrt{2}}{100}$ radians per second

11 increasing at 0.12 radians per minute

REVIEW SET 14A


1 a $y = 4x + 2$ b $y = 4x + 4 \ln 2 - 4$ c $y = e^2$

2 $a = \frac{5}{2}$, $b = -\frac{3}{2}$

3 a $a = -6$

b local max. $(-\sqrt{2}, 4\sqrt{2})$, local min. $(\sqrt{2}, -4\sqrt{2})$

c




4 a $y = \frac{1}{5}x - \frac{11}{5}$ (or $x - 5y = 11$) b $y = -4x + 14$

5 $\frac{3267}{152} \text{ units}^2$ 6 $a = 64$ 7 P(0, 7.5), Q(3, 0)

9 $3x - 4y = -5$

10 a $x > 0$


b Sign diagram of $f'(x)$




$f(x)$ is increasing for all $x > 0$.

d normal is $x + 2y = 3$

11 a



c





b $y = -\frac{4}{k^2}x + \frac{8}{k}$

c A(2k, 0), B(0, $\frac{8}{k}$)

d Area = 8 units²

e $k = 2$

12 a $v(t) = (6t^2 - 18t + 12) \text{ cm s}^{-1}$, $a(t) = (12t - 18) \text{ cm s}^{-2}$



b $s(0) = -5 \text{ cm}$ (5 cm to the left of origin)


$v(0) = 12 \text{ cm s}^{-1}$ towards origin

$a(0) = -18 \text{ cm s}^{-2}$ (reducing speed)

c At $t = 2$, particle is 1 cm to the left of the origin, is stationary and is accelerating towards the origin.

d $t = 1$, $s = 0$ and $t = 2$, $s = -1$

e



f $1 \leq t \leq 1\frac{1}{2}$ and $t \geq 2$

13 b $k = 9$

14 a $x(0) = 3 \text{ cm}$, $x'(0) = 2 \text{ cm s}^{-1}$, $x''(0) = 0 \text{ cm s}^{-2}$

b $t = \frac{\pi}{4} \text{ s}$ and $\frac{3\pi}{4} \text{ s}$ c 4 cm

15 6 cm from each end

16 a $y = \frac{1}{x^2}$, $x > 0$

c base is 1.26 m square, height 0.630 m

17 a $v(t) = 15 + \frac{120}{(t+1)^3} \text{ cm s}^{-1}$, $a(t) = \frac{-360}{(t+1)^4} \text{ cm s}^{-2}$

b At $t = 3$, particle is 41.25 cm to the right of the origin, moving to the right at 16.88 cm s^{-1} and decelerating at 1.41 cm s^{-2} .

c speed is never increasing

18 A $(\frac{1}{2}, \frac{1}{e})$ 19 $\frac{20\sqrt{10}}{3} \approx 21.1 \text{ m per minute}$

20 a $V(r) = \frac{8}{9}\pi r^3 \text{ m}^3$

b $\frac{dr}{dt} = -\frac{8}{375\pi} \approx -0.00679 \text{ m min}^{-1}$

REVIEW SET 14B

1 a $x = 1$ b $ex - 2y = e - \frac{2}{e}$ c $y = 16x - \frac{127}{2}$

2 $a = -14$, $b = 21$

3 a $f(3) = 2$, $f'(3) = -1$ b $f(x) = x^2 - 7x + 14$

4 a $2x + 3y = \frac{2\pi}{3} + 2\sqrt{3}$ b $\sqrt{2}y - 4x = 1 - 2\pi$

5 $p = 1$, $q = -8$ 6 $(-2, -25)$ 7 $a = \frac{1}{2}$


8 a local minimum at (0, 1) c $f''(x) = e^x$

9 (0, $\ln 4 - 1$)

10 a y-intercept = -35 b $x = 1, \frac{7}{2}, 5$

c local maximum at (2, 9), local minimum at $(\frac{13}{3}, -\frac{100}{27})$

d



11 BC = $\frac{8\sqrt{10}}{3} \text{ units}$

12 a 60 cm b i 4.24 years ii 201 years

c i 16 cm per year ii 1.95 cm per year

13 a $v(t) = -8e^{-\frac{t}{10}} - 40 \text{ m s}^{-1}$


$a(t) = \frac{4}{5}e^{-\frac{t}{10}} \text{ m s}^{-2} \{t \geq 0\}$

b $s(0) = 80 \text{ m}$

$v(0) = -48 \text{ m s}^{-1}$

$a(0) = 0.8 \text{ m s}^{-2}$

d $t = 10 \ln 2$ seconds



14 a i \$535 ii \$1385.79

b i $-\$0.267 \text{ per km h}^{-1}$ ii $\$2.33 \text{ per km h}^{-1}$

c 51.3 km h^{-1}

15 a $v(t) = 3 - \frac{1}{2\sqrt{t+1}}$ $a(t) = \frac{1}{4(t+1)^{\frac{3}{2}}}$




b $x(0) = -1$, $v(0) = 2.5$, $a(0) = 0.25$

Particle is 1 cm to the left of the origin, is travelling to the right at 2.5 cm s^{-1} , and accelerating at 0.25 cm s^{-2} .

c Particle is 21 cm to the right of the origin, is travelling to the right at 2.83 cm s^{-1} , and accelerating at $0.00926 \text{ cm s}^{-2}$.

d never changes direction **e** never decreasing

16 b $A = 200x - 2x^2 - \frac{1}{2}\pi x^2$



17 a $v(0) = 0 \text{ cm s}^{-1}$, $v(\frac{1}{2}) = -\pi \text{ cm s}^{-1}$, $v(1) = 0 \text{ cm s}^{-1}$,
 $v(\frac{3}{2}) = \pi \text{ cm s}^{-1}$, $v(2) = 0 \text{ cm s}^{-1}$

b $0 \leq t \leq 1$, $2 \leq t \leq 3$, $4 \leq t \leq 5$, etc.

So, for $2n \leq t \leq 2n+1$, $n \in \{0, 1, 2, 3, \dots\}$

18 $x = \frac{k}{2} \left(1 - \frac{1}{\sqrt{3}}\right)$ **19** 3.60 m s^{-1}

20 increasing at 0.128 radians per second

21 a $\frac{\sqrt{3}}{2}\pi \text{ cm s}^{-1}$ **b** 0 cm s^{-1}

22 a i $y = -\frac{a^2}{4b} + \frac{a}{2b}x$ **ii** when $y = 0$, $x = \frac{a}{2}$

b i $y = -\frac{2b}{a}x + b$ **ii** when $x = 0$, $y = b$

iii Hint: Let P'' be the point on the line $y = -b$ where the distance to P is shortest. Show that $FP = P''P$.

c i Hint: Show that $\triangle FPP' \cong \triangle P''PP'$.

ii Hint: Show that the tangents meet at $\left(\frac{a+c}{2}, \frac{ac}{4b}\right)$.

EXERCISE 15A.1

1 a i 0.6 units² **ii** 0.4 units² **b** 0.5 units²

2 a 0.737 units² **b** 0.653 units²

3

n	A_L	A_U
10	2.1850	2.4850
25	2.2736	2.3936
50	2.3034	2.3634
100	2.3184	2.3484
500	2.3303	2.3363

converges to $\frac{7}{3}$

4 a i

n	A_L	A_U
5	0.16000	0.36000
10	0.20250	0.30250
50	0.24010	0.26010
100	0.24503	0.25503
500	0.24900	0.25100
1000	0.24950	0.25050
10000	0.24995	0.25005

ii

n	A_L	A_U
5	0.40000	0.60000
10	0.45000	0.55000
50	0.49000	0.51000
100	0.49500	0.50500
500	0.49900	0.50100
1000	0.49950	0.50050
10000	0.49995	0.50005

iii

n	A_L	A_U
5	0.54974	0.74974
10	0.61051	0.71051
50	0.65610	0.67610
100	0.66146	0.67146
500	0.66565	0.66765
1000	0.66616	0.66716
10000	0.66662	0.66672

iv

n	A_L	A_U
5	0.61867	0.81867
10	0.68740	0.78740
50	0.73851	0.75851
100	0.74441	0.75441
500	0.74893	0.75093
1000	0.74947	0.75047
10000	0.74995	0.75005

b i $\frac{1}{4}$ **ii** $\frac{1}{2}$ **iii** $\frac{2}{3}$ **iv** $\frac{3}{4}$ **c** area = $\frac{1}{a+1}$


5 a

n	Rational bounds for π
10	$2.9045 < \pi < 3.3045$
50	$3.0983 < \pi < 3.1783$
100	$3.1204 < \pi < 3.1604$
200	$3.1312 < \pi < 3.1512$
1000	$3.1396 < \pi < 3.1436$
10000	$3.1414 < \pi < 3.1418$

b $n = 10000$

EXERCISE 15A.2

1 a



b

n	A_L	A_U
5	0.5497	0.7497
10	0.6105	0.7105
50	0.6561	0.6761
100	0.6615	0.6715
500	0.6656	0.6676

c $\int_0^1 \sqrt{x} dx \approx 0.67$


2 a $A_L = \frac{2}{n} \sum_{i=0}^{n-1} \sqrt{1+x_i^3}$, $A_U = \frac{2}{n} \sum_{i=1}^n \sqrt{1+x_i^3}$

b

n	A_L	A_U
50	3.2016	3.2816
100	3.2214	3.2614
500	3.2373	3.2453

c $\int_0^2 \sqrt{1+x^3} dx \approx 3.24$

3 a


 b upper ≈ 1.2506 , lower ≈ 1.2493

 c upper ≈ 1.2506 , lower ≈ 1.2493

 d $\int_{-3}^3 e^{-\frac{x^2}{2}} dx \approx 2.4999$ compared to $\sqrt{2\pi} \approx 2.5066$

4 a 18

b 4.5

 c 2π

EXERCISE 15B

 1 a i $\frac{x^2}{2}$ ii $\frac{x^3}{3}$ iii $\frac{x^6}{6}$ iv $-\frac{1}{x}$
 v $-\frac{1}{3x^3}$ vi $\frac{3}{4}x^{\frac{4}{3}}$ vii $2\sqrt{x}$

 b The antiderivative of x^n is $\frac{x^{n+1}}{n+1}$ ($n \neq -1$).

 2 a i $\frac{1}{2}e^{2x}$ ii $\frac{1}{5}e^{5x}$ iii $2e^{\frac{1}{2}x}$ iv $100e^{0.01x}$
 v $\frac{1}{\pi}e^{\pi x}$ vi $3e^{\frac{\pi}{3}}$

 b The antiderivative of e^{kx} is $\frac{1}{k}e^{kx}$.

 3 a $\frac{d}{dx}(x^3 + x^2) = 3x^2 + 2x$
 \therefore the antiderivative of $6x^2 + 4x = 2x^3 + 2x^2$

 b $\frac{d}{dx}(e^{3x+1}) = 3e^{3x+1}$
 \therefore the antiderivative of $e^{3x+1} = \frac{1}{3}e^{3x+1}$

 c $\frac{d}{dx}(x\sqrt{x}) = \frac{3}{2}\sqrt{x}$
 \therefore the antiderivative of $\sqrt{x} = \frac{2}{3}x\sqrt{x}$

 d $\frac{d}{dx}(2x+1)^4 = 8(2x+1)^3$
 \therefore the antiderivative of $(2x+1)^3 = \frac{1}{8}(2x+1)^4$

EXERCISE 15C

 1 a $\frac{1}{4}$ units² b $2\frac{1}{3}$ units² c $\frac{2}{3}$ units²
 3 a $3\frac{3}{4}$ units² b $24\frac{2}{3}$ units² c $\frac{-2+4\sqrt{2}}{3}$ units²
 d ≈ 3.48 units² e 2 units²

 4 c i $\int_0^1 (-x^2) dx = -\frac{1}{3}$, the area between $y = -x^2$ and the x -axis from $x = 0$ to $x = 1$ is $\frac{1}{3}$ units².
 ii $\int_0^1 (x^2 - x) dx = -\frac{1}{6}$, the area between $y = x^2 - x$ and the x -axis from $x = 0$ to $x = 1$ is $\frac{1}{6}$ units².
 iii $\int_{-2}^0 3x dx = -6$, the area between $y = 3x$ and the x -axis from $x = -2$ to $x = 0$ is 6 units²
 d $-\pi$

EXERCISE 15D

 1 $\frac{dy}{dx} = 7x^6$, $\int x^6 dx = \frac{1}{7}x^7 + c$
 2 $\frac{dy}{dx} = 3x^2 + 2x$, $\int (3x^2 + 2x) dx = x^3 + x^2 + c$

 3 $\frac{dy}{dx} = 2e^{2x+1}$, $\int e^{2x+1} dx = \frac{1}{2}e^{2x+1} + c$

 4 $\frac{dy}{dx} = 8(2x+1)^3$, $\int (2x+1)^3 dx = \frac{1}{8}(2x+1)^4 + c$

 5 $\frac{dy}{dx} = \frac{3}{2}\sqrt{x}$, $\int \sqrt{x} dx = \frac{2}{3}x\sqrt{x} + c$

 6 $\frac{dy}{dx} = -\frac{1}{2x\sqrt{x}}$, $\int \frac{1}{x\sqrt{x}} dx = -\frac{2}{\sqrt{x}} + c$

 7 $\frac{dy}{dx} = -2\sin 2x$, $\int \sin 2x dx = -\frac{1}{2}\cos 2x + c$

 8 $\frac{dy}{dx} = -5\cos(1-5x)$,
 $\int \cos(1-5x) dx = -\frac{1}{5}\sin(1-5x) + c$

 9 $\int (2x-1)(x^2-x)^2 dx = \frac{1}{3}(x^2-x)^3 + c$

 11 $\frac{dy}{dx} = \frac{-2}{\sqrt{1-4x}}$, $\int \frac{1}{\sqrt{1-4x}} dx = -\frac{1}{2}\sqrt{1-4x} + c$

EXERCISE 15E.1

 1 a $\frac{x^5}{5} - \frac{x^3}{3} - \frac{x^2}{2} + 2x + c$ b $x^5 - x^4 - 2x^3 - 7x + c$

 c $\frac{2}{3}x^{\frac{3}{2}} + e^x + c$ d $3e^x + \frac{1}{3}x^3 + c$

 e $\frac{2}{5}x^{\frac{5}{2}} - 2x + c$ f $-2x^{-\frac{1}{2}} + 2x^2 + c$

 g $\frac{1}{8}x^4 - \frac{1}{5}x^5 + \frac{3}{4}x^{\frac{4}{3}} + c$ h $\frac{x^2}{4} + \frac{1}{3}x^3 - e^x + c$

 i $5e^x + \frac{1}{12}x^4 - \frac{2}{3}x^{\frac{3}{2}} + c$

 2 a $-3\cos x - 2x + c$ b $2x^2 - 2\sin x + c$

 c $-\cos x - 2\sin x + e^x + c$ d $\frac{2}{7}x^3\sqrt{x} + 10\cos x + c$

 e $\frac{1}{9}x^3 - \frac{1}{6}x^2 + \sin x + c$ f $\cos x + \frac{4}{3}x\sqrt{x} + c$

 3 a $\frac{1}{3}x^3 + \frac{3}{2}x^2 - 2x + c$ b $\frac{2}{3}x^{\frac{3}{2}} - 2x^{\frac{1}{2}} + c$

 c $2e^x + \frac{1}{x} + c$ d $-2x^{-\frac{1}{2}} - 8x^{\frac{1}{2}} + c$

 e $\frac{4}{3}x^3 + 2x^2 + x + c$ f $\frac{1}{3}x^3 + 2x - \frac{1}{x} + c$

 g $\frac{4}{3}x^{\frac{3}{2}} - 2x^{\frac{1}{2}} + c$ h $2x^{\frac{1}{2}} + 8x^{-\frac{1}{2}} - \frac{20}{3}x^{-\frac{3}{2}} + c$

 i $\frac{1}{4}x^4 + x^3 + \frac{3}{2}x^2 + x + c$

 4 a $\frac{2}{3}x^{\frac{3}{2}} + \frac{1}{2}\sin x + c$ b $2e^t + 4\cos t + c$

 c $3\sin t + \cos t + c$

 5 a $y = 6x + c$ b $y = \frac{4}{3}x^3 + c$

 c $y = \frac{10}{3}x\sqrt{x} - \frac{1}{3}x^3 + c$ d $y = -\frac{1}{x} + c$

 e $y = 2e^x - 5x + c$ f $y = x^4 + x^3 + c$

 6 a $f(x) = x - 2x^2 + \frac{4}{3}x^3 + c$

 b $f(x) = \frac{2}{3}x^{\frac{3}{2}} - 4\sqrt{x} + c$ c $f(x) = x + \frac{5}{x} + c$

EXERCISE 15E.2

 1 a $f(x) = x^2 - x + 3$ b $f(x) = x^3 + x^2 - 7$

 c $f(x) = e^x + 2\sqrt{x} - 1 - e$ d $f(x) = \frac{1}{2}x^2 - 4\sqrt{x} + \frac{11}{2}$

 2 a $f(x) = \frac{x^3}{3} - 4\sin x + 3$

 b $f(x) = 2\sin x + 3\cos x - 2\sqrt{2}$

- 3 a $f(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + \frac{1}{3}$
 b $f(x) = 4x^{\frac{5}{2}} + 4x^{\frac{3}{2}} - 4x + 5$
 c $f(x) = -\cos x - x + 4$ d $f(x) = \frac{1}{3}x^3 - \frac{16}{3}x + 5$

EXERCISE 15F




- 1 a $\frac{1}{8}(2x+5)^4 + c$ b $\frac{1}{2(3-2x)} + c$
 c $\frac{-2}{3(2x-1)^3} + c$ d $\frac{1}{32}(4x-3)^8 + c$
 e $\frac{2}{9}(3x-4)^{\frac{3}{2}} + c$ f $-4\sqrt{1-5x} + c$
 g $-\frac{3}{5}(1-x)^5 + c$ h $-2\sqrt{3-4x} + c$
 2 a $-\frac{1}{3}\cos(3x) + c$ b $-\frac{1}{2}\sin(-4x) + x + c$
 c $6\sin\left(\frac{x}{2}\right) + c$ d $-\frac{3}{2}\cos(2x) + e^{-x} + c$
 e $-\cos\left(2x + \frac{\pi}{6}\right) + c$ f $3\sin\left(\frac{\pi}{4} - x\right) + c$
 g $\frac{1}{2}\sin(2x) - \frac{1}{2}\cos(2x) + c$
 h $-\frac{2}{3}\cos(3x) + \frac{5}{4}\sin(4x) + c$
 i $\frac{1}{16}\sin(8x) + 3\cos x + c$
 3 $y = \frac{1}{3}(2x-7)^{\frac{3}{2}} + 2$ 4 $(-8, -19)$
 5 a $\frac{1}{2}(2x-1)^3 + c$ b $\frac{1}{5}x^5 - \frac{1}{2}x^4 + \frac{1}{3}x^3 + c$
 c $-\frac{1}{12}(1-3x)^4 + c$ d $x - \frac{2}{3}x^3 + \frac{1}{5}x^5 + c$
 e $-\frac{8}{3}(5-x)^{\frac{3}{2}} + c$ f $\frac{1}{7}x^7 + \frac{3}{5}x^5 + x^3 + x + c$
 6 a $2e^x + \frac{5}{2}e^{2x} + c$ b $\frac{3}{5}e^{5x-2} + c$
 c $-\frac{1}{3}e^{7-3x} + c$ d $\frac{1}{2}e^{2x} + 2x - \frac{1}{2}e^{-2x} + c$
 e $-\frac{1}{2}e^{-2x} - 4e^{-x} + 4x + c$ f $\frac{1}{2}x^2 + 5(1-x)^{-1} + c$
 7 $y = x - 2e^x + \frac{1}{2}e^{2x} + \frac{11}{2}$
 8 $p = -\frac{1}{4}$, $f(x) = \frac{1}{2}\cos\left(\frac{1}{2}x\right) + \frac{1}{2}$
 10 $f(x) = -e^{-2x} + 4$
 11 $f(x) = \frac{2}{3}x^{\frac{3}{2}} - \frac{1}{8}e^{-4x} + \frac{1}{8}e^{-4} - \frac{2}{3}$

EXERCISE 15G

- 1 a $\int_1^4 \sqrt{x} dx = \frac{14}{3}$, $\int_1^4 (-\sqrt{x}) dx = -\frac{14}{3}$
 b $\int_0^1 x^7 dx = \frac{1}{8}$, $\int_0^1 (-x^7) dx = -\frac{1}{8}$
 2 a $\frac{1}{3}$ b $\frac{7}{3}$ c $\frac{8}{3}$ d 1
 3 a -4 b 6.25 c 2.25 4 a $\frac{1}{3}$ b $\frac{2}{3}$ c 1
 5 a $\frac{1}{4}$ b $\frac{2}{3}$ c $e-1$ (≈ 1.72) d $\frac{1}{2}$
 e $1\frac{1}{2}$ f $6\frac{2}{3}$ g $\ln 3$ (≈ 1.10) h $\frac{1}{2}$
 i ≈ 1.52 j 2 k $e-1$ (≈ 1.72) l $\frac{1}{3}$
 6 $m = -1$ or $\frac{4}{3}$ 7 a $\frac{\pi}{8} + \frac{1}{4}$ b $\frac{\pi}{4}$
 8 a 6.5 b -9 c 0 d -2.5
 9 a 2π b -4 c $\frac{\pi}{2}$ d $\frac{5\pi}{2} - 4$
 10 a $\int_2^7 f(x) dx$ b $\int_1^9 g(x) dx$
 11 a -5 b 4
 12 a 4 b 0 c -8 d $k = -\frac{7}{4}$ 13 0

REVIEW SET 15A

1 a



n	A_L	A_U
5	2.9349	3.3349
50	3.1215	3.1615
100	3.1316	3.1516
500	3.1396	3.1436

c $\int_0^1 \frac{4}{1+x^2} dx \approx 3.1416$

2 a 2π

b 4

3 a $8\sqrt{x} + c$ b $-\frac{1}{4}\cos(4x-5) + c$ c $-\frac{1}{3}e^{4-3x} + c$ 4 a $12\frac{4}{9}$ b $\sqrt{2}$

5 $\frac{dy}{dx} = \frac{x}{\sqrt{x^2-4}}$, $\int \frac{x}{\sqrt{x^2-4}} dx = \sqrt{x^2-4} + c$

6 $b = \frac{\pi}{4}, \frac{3\pi}{4}$

7 a $y = \frac{1}{5}x^5 - \frac{2}{3}x^3 + x + c$ b $y = 400x + 40e^{-\frac{x}{2}} + c$

8 $f(x) = 3x^3 + 5x^2 + 6x - 1$ 9 $a = \ln \sqrt{2}$

10 a $f(x) = \frac{1}{4}x^4 + \frac{1}{3}x^3 - \frac{10}{3}x + 3$ b $3x + 26y = 84$

11 a $e^{3x} + 6e^{2x} + 12e^x + 8$ b $\frac{1}{3}e^3 + 3e^2 + 12e - 7\frac{1}{3}$

REVIEW SET 15B

1 a $A = \frac{17}{4}$, $B = \frac{25}{4}$ b $\int_0^2 (4-x^2) dx \approx \frac{21}{4}$

2 a $-2e^{-x} + 3x + c$ b $\frac{2}{3}x\sqrt{x} - 2\sqrt{x} + c$
 c $9x + 3e^{2x-1} + \frac{1}{4}e^{4x-2} + c$

3 $f(x) = \frac{1}{3}x^3 - \frac{3}{2}x^2 + 2x + 2\frac{1}{6}$ 4 $\frac{2}{3}(\sqrt{5} - \sqrt{2})$

5 $\frac{d}{dx}(3x^2+x)^3 = 3(3x^2+x)^2(6x+1)$
 $\int (3x^2+x)^2(6x+1) dx = \frac{1}{3}(3x^2+x)^3 + c$

6 a 6 b 3 7 $f\left(\frac{\pi}{2}\right) = 3 - \frac{\pi}{2}$ 8 $e^{-\pi}$

9 $\frac{1}{2(n+1)}(2x+3)^{n+1} + c$, $n \neq -1$

10 $a = \frac{1}{3}$, $f'(x) = 2\sqrt{x} + \frac{1}{3\sqrt{x}}$ is never 0 as $\sqrt{x} \geq 0$ for all x
 $\therefore f'(x) > 0$ for all x


11 $a = 0$ or ± 3

EXERCISE 16A

- 1 a 30 units² b $\frac{9}{2}$ units² c $\frac{27}{2}$ units² d 2 units²
 2 a $\frac{1}{3}$ units² b 2 units² c $63\frac{3}{4}$ units²
 d $(e-1)$ units² e $20\frac{5}{6}$ units² f 18 units²
 g $\frac{1}{2}$ units² h $4\frac{1}{2}$ units² i $(2e - \frac{2}{e})$ units²
 3 $\frac{2}{3}$ units²


EXERCISE 16B

- 1 a $4\frac{1}{2}$ units² b $(1 + e^{-2})$ units² c $1\frac{5}{27}$ units²
 d 2 units² e $2\frac{1}{4}$ units² f $(\frac{\pi}{2} - 1)$ units²
 2 $10\frac{2}{3}$ units²
 3 a




- b (1, -2) and (3, 0)
 c $1\frac{1}{3}$ units²

- 4 $\frac{1}{3}$ units²




- b (0, 0) and (ln 2, 1)
 c enclosed area = $3 \ln 2 - 2$ (≈ 0.0794) units²

- 6 $\frac{1}{2}$ units²



enclosed area = $\frac{1}{12}$ units²



- a Rearranging $x^2 + y^2 = 9$ gives $y = \pm\sqrt{9 - x^2}$. The upper half has $y \geq 0$, so $y = \sqrt{9 - x^2}$.
 b $\frac{9\pi}{4}$

- 9 a $40\frac{1}{2}$ units² b 8 units² c 8 units²

- 10 a C_1 is $y = \sin x$, C_2 is $y = 3 \sin x$ b 4 units²

- 11 a $\int_3^5 f(x) dx = -$ (area between $x = 3$ and $x = 5$)

$$b \int_1^3 f(x) dx - \int_3^5 f(x) dx + \int_5^7 f(x) dx$$

- 12 a C_1 is $y = \frac{1}{2} + \frac{1}{2} \cos(2x)$, C_2 is $y = \cos(2x)$

- b A(0, 1), B($\frac{\pi}{4}$, 0), C($\frac{\pi}{2}$, 0), D($\frac{3\pi}{4}$, 0), E(π , 1)

$$c \text{ Area} = \int_0^\pi (\frac{1}{2} + \frac{1}{2} \cos(2x) - \cos(2x)) dx$$

- 13 If $h(x) \geq 0$ on $a \leq x \leq b$, the area between $y = h(x)$ and the x -axis is $\int_a^b h(x) dx$. If $h(x) < 0$ on $a \leq x \leq b$, the area between $y = h(x)$ and the x -axis is $\int_a^b -h(x) dx$.
 \therefore the area between $y = h(x)$ and the x -axis on $a \leq x \leq b$ is $\int_a^b |h(x)| dx$.

Letting $h(x) = f(x) - g(x)$, the area between $y = f(x) - g(x)$ and the x -axis $y = 0$ on $a \leq x \leq b$ is $\int_a^b |f(x) - g(x)| dx$.


Equivalently, the area between $y = f(x)$ and $y = g(x)$ on $a \leq x \leq b$ is $\int_a^b |f(x) - g(x)| dx$.

- 14 $b \approx 1.3104$

- 15 $a = \sqrt{3}$

EXERCISE 16C.1

- 1 110 m
 2 a i travelling forwards
 ii travelling backwards (opposite direction)
 b 16 km c 8 km from starting point (on positive side)



EXERCISE 16C.2

- 1 a $s(t) = t - t^2 + 2$ cm b $\frac{1}{2}$ cm c 0 cm
 2 a $s(t) = \frac{1}{3}t^3 - \frac{1}{2}t^2 - 2t$ cm b $5\frac{1}{6}$ cm
 c $1\frac{1}{2}$ cm left of its starting point

- 3 $\frac{\sqrt{3}+2}{4}$ m

- 4 a $s(t) = 32t + 2t^2 + 16$ m

- b no change of direction

$$\text{so displacement} = s(t_1) - s(0) = \int_0^{t_1} (32 + 4t) dt$$

- c acceleration = 4 m s^{-2}


- 5 a 41 units b 34 units 6 b 2 m

- 7 a 40 m s^{-1} b 47.8 m s^{-1} c 1.39 seconds

- d as $t \rightarrow \infty$, $v(t) \rightarrow 50$ from below

- e $a(t) = 5e^{-0.5t}$ and as $e^x > 0$ for all x , $a(t) > 0$ for all t .

- f $g \approx 134.5$ m



- 8 a $v(t) = -\frac{1}{(t+1)^2} + 1 \text{ m s}^{-1}$

- b $s(t) = \frac{1}{t+1} + t - 1$ m

- c** The particle is $\frac{4}{3}$ m to the right of the origin, moving to the right at $\frac{8}{9} \text{ m s}^{-1}$, and accelerating at $\frac{2}{27} \text{ m s}^{-2}$.
- 9 a** $v(t) = \frac{t^2}{20} - 3t + 45 \text{ m s}^{-1}$
- b** $\int_0^{60} v(t) dt = 900$. The train travels a total of 900 m in the first 60 seconds.
- 10 a** Show that $v(t) = 100 - 80e^{-\frac{1}{20}t} \text{ m s}^{-1}$ and as $t \rightarrow \infty$, $v(t) \rightarrow 100 \text{ m s}^{-1}$.
- b** 370.4 m

REVIEW SET 16A

- 1** $A = \int_a^b [f(x) - g(x)] dx + \int_b^c [g(x) - f(x)] dx$
 $+ \int_c^d [f(x) - g(x)] dx$
- 2 a** $2 + \pi$ **b** -2 **c** π
- 3** No, total area shaded $= \int_{-1}^1 f(x) dx - \int_1^3 f(x) dx$.
- 4** $k = \sqrt[3]{16}$ **5** 4.5 units^2


- 6 a**
-

- b** The particle moves in the positive direction initially, then at $t = 2$, $6\frac{2}{3}$ m from its starting point, it changes direction. It changes direction again at $t = 4$, $5\frac{1}{3}$ m from its starting point, and at $t = 5$, it is $6\frac{2}{3}$ m from its starting point again.


- c** $6\frac{2}{3} \text{ m}$ **d** $9\frac{1}{3} \text{ m}$
- 7** $(3 - \ln 4) \text{ units}^2$ **8** 2.35 m

REVIEW SET 16B

- 1 a** $v(t) = 3t^2 - 30t + 27 \text{ cm s}^{-1}$
- b** -162 cm (162 cm to the left of the origin)



- c** $(1 - \frac{\pi}{4}) \text{ units}^2$
- 3** $a = \ln 3$, $b = \ln 5$
- 4 a** $a(t) = 2 - 6t \text{ m s}^{-2}$ **b** $s(t) = t^2 - t^3 + c \text{ m}$
- c** -4 m (4 m to the left)
- 5** $k = \frac{4}{3}$ **6** $m = \frac{\pi}{3}$ **7** $(\frac{4}{\pi} - 1) \text{ units}^2$
- 8 a** $v(0) = 25 \text{ m s}^{-1}$, $v(3) = 4 \text{ m s}^{-1}$
- b** as $t \rightarrow \infty$, $v(t) \rightarrow 0$ from above



- d** 3 seconds **e** $a(t) = \frac{-200}{(t+2)^3} \text{ m s}^{-2}$, $t \geq 0$
- f** $k = \frac{1}{5}$

INDEX

absolute value	46	degree	202	gradient function	340
acceleration	446	degree of polynomial	156	gradient of tangent	339
addition rule	345, 424	DeMorgan's law	25	gradient of normal	370
algebraic test	37	derivative	342	gradient-intercept form	177
amplitude	228	derivative function	340	horizontal asymptote	118, 149
antiderivative	416	determinant	324	horizontal line test	39
antidifferentiation	416	differential equation	399	idempotent law	25
arc	205	differentiation	344	identity function	55
arc length	202, 205	discriminant	73, 85	identity law	25
area of sector	205	disjoint sets	14, 22	identity matrix	320
area under curve	410, 438	displacement	446	image	41
associativity law	25	displacement function	380	indefinite integration	422
average acceleration	381	distance between points	176	index	107
average speed	337	distance travelled	444	index laws	108
average velocity	381	distinct real root	73	inequality	72
axis of symmetry	76	distributive law	25	infinite set	13
base	107	division algorithm	159	initial conditions	383
base unit vector	281	divisor	158	instantaneous acceleration	381
binomial	270	domain	36, 43	instantaneous velocity	381
binomial coefficient	261, 273	double root	73	integer	12
binomial expansion	270	element	12	integral	422
binomial theorem	273	empty set	12	intersection	13, 22
Cartesian equation	296	equal matrices	309	interval	16
Cartesian plane	17	equal vectors	282	interval notation	43
chain rule	350	equation	37, 72	inverse function	54
change of base rule	147	equation of a line	177	inverse operation	54
chord	205, 338	equation of normal	370	invertible matrix	324
closed interval	16	exponent	107	laws of logarithms	135
coincident lines	328	exponential equation	116	leading coefficient	156
column matrix	307	exponential function	118	length of vector	280, 284
column vector	307	factor	162	limit	335
combination	267	factor theorem	169	linear factor	162
commutative law	25	factorial numbers	259	linear function	41, 65
complement	18, 25	finite set	13	local maximum	375
completing the square	67, 81	first derivative	363	local minimum	375
composite function	49	first principles formula	343	lower rectangles	411
concave downwards	76	function	37	magnitude of vector	280, 284
concave upwards	76	function value	41	major	205
conic sections	75	fundamental theorem		mapping	37
constant of integration	425	of calculus	419	matrix	307
constant term	156	general cosine function	236	matrix algebra	314, 320
cosine	208	general form	177	matrix multiplication	315
cosine function	236	general sine function	232	maximum point	228
counting numbers	12	general tangent function	239	maximum turning point	76
cubic function	65	general term	273	maximum value	95
cycloid	228	geometric test	37	mean line	228
definite integral	413, 431	global maximum	375	member	12
		global minimum	375	midpoint	176, 182
		golden ratio	94	minimum point	228
		gradient	177	minimum turning point	76

minimum value	95	quadratic formula	69	unit circle	208
minor	205	quadratic function	65, 75	unit vector	281, 284, 292
modulus	46	quadratic graph	85	universal set	18
motion graph	380	quartic function	65	upper rectangles	410
multiplicative inverse	323	quotient	158	variable	37, 156
mutually exclusive sets	14, 22	quotient rule	353	vector	280
natural domain	43	radian	202	vector equation of line	296
natural exponential	123	radical	102	velocity	446
natural logarithm	142	radical conjugates	105	velocity function	381
natural numbers	12	radius	202	Venn diagram	21
negative definite	85	range	36, 43	vertex	76
negative integer	12	rate	336	vertical asymptote	149
negative matrix	314	rational number	12	vertical line test	37
negative reciprocals	177	real number	12	vertical translation	232
negative vector	283	real polynomial	156	x -intercept	76
non-singular matrix	324	relation	17, 37	y -intercept	76
normal to a curve	370	remainder	158	zero	51, 162
number line	15	remainder theorem	167	zero matrix	313
one-one function	39	repeated root	73	zero vector	282
open interval	16	resultant vector	294		
optimisation	95, 393	Riemann integral	434		
optimum solution	393	root	51, 65, 162		
order of matrix	307	row matrix	307		
parabola	75	row vector	307		
parallel lines	177, 328	scalar	280, 287		
parallel vectors	292	scalar multiplication	287		
parametric equations	216, 296	second derivative	363, 377		
Pascal's triangle	271	sector	205		
period	228	segment	205		
periodic function	228	self-inverse function	57		
periodic phenomena	226	set	12		
permutation	262	sign diagram	51, 376, 383		
perpendicular bisector	182	sign test	384		
perpendicular lines	177	sine	208		
point of discontinuity	335	sine curve	230		
point of inflection	369	sine waves	227		
polynomial	65, 156	singular matrix	324		
polynomial function	156	solution	162		
position vector	280, 289	speed	337, 384		
positive definite	85	square matrix	307		
positive integer	12	standard basis	281		
power of a binomial	270	stationary inflection	375		
power rule	424	stationary point	375		
principal axis	228	subset	13, 21		
principal domain	241	summation notation	156		
product principle	256	surd	102		
product rule	352	tangent	338		
proper subset	13	tangent function	209, 238		
quadrant	210	tangent to a curve	369		
quadratic equation	65	union	13, 22		