

What is BPH Caused by?

As men age, the prostate enlarges

Hormonal changes



Increased risk of developing BPH for:

- Men with a family history of it
- Men with certain health conditions (diabetes, heart disease, overweight)
- Men who do not stay active
- Being on certain medications

If any of the above sound familiar, and bother you or interfere with your sleep, its usually time to intervene and past the “watchful waiting” period.



A Small Change for your Prostate, A Big Change for your Day and Night.
Talk to Us about your options for BPH: Restore your Flow with Prostate Centers of America.



How can I find out if I have BPH?

- 1 Through a symptom questionnaire and discussion with your provider
- 2 A blood test
- 3 And an MRI



What are Treatment Options for BPH?

Certain lifestyle changes

Medications

Minimally Invasive Treatments

Surgeries



What is Benign Prostatic Hyperplasia (BPH)?

A very common, non-cancerous condition characterized by enlargement of the prostate gland. The growth of the prostate gland squeezes the urethra (the tube connecting the bladder to the body's exterior), preventing urine from fully leaving the body. Over time the bladder tries to compensate for the blocked exit and it thickens and becomes less functional/elastic.

Symptoms BPH can cause:

- Frequency:** needing to pee (urinate) multiple times during the day
- Nocturia:** needing to urinate multiple times during sleeping hours
- Urgency:** needing to urinate right away
- Incomplete emptying or retention:** the sensation of needing to urinate shortly after urinating
- Weak stream:** weak urination stream or dribbling
- Hesitancy:** difficulty starting urination
- Straining:** needing to strain to start or continue urinating
- Intermittency:** starting and stopping during urination
- Urine leakage**

Call 800 - 000 - 0000

