Proofs From Zero to Velleman

A Conversational Reader for Building Mathematical Maturity

Engineering Math to Pure Math

May 2026

Contents

Preface: What This Reader Is **2**

1 The Change in Game **3**

1.1 From calculation to justification . . . 3

1.2 The big shift . . . 3

2 Statements, Predicates, and the First Filter **4**

2.1 Predicates . . . 4

2.2 Why this matters . . . 4

2.3 Exercises . . . 4

3 Variables Are Not All the Same **6**

3.1 Fixed but unknown . . . 6

3.2 Arbitrary . . . 6

3.3 Witness . . . 6

3.4 Bound and free variables . . . 6

3.5 Exercises . . . 7

4 Quantifiers: The Tiny Symbols That Run the Machine **8**

4.1 Universal statements . . . 8

4.2 Existential statements . . . 8

4.3 The order matters . . . 8

4.4 A game interpretation . . . 8

5 Negation: Turning Claims Inside Out **10**

5.1 The basic laws . . . 10
5.2 Negating implication . . . 10
5.3 Nested negations . . . 10
5.4 Exercises . . . 11

6 If-Then: The Spine of Theorems 12
6.1 Direct proof . . . 12
6.2 What happened? . . . 12
6.3 Converse . . . 12

7 Definitions Are Machines 13
7.1 Opening and closing definitions . . . 13
7.2 Worked proof: even plus even . . . 13
7.3 Worked proof: divisibility is transitive . . . 13

8 The Three First Proof Moves 14
8.1 Move 1: Direct proof . . . 14
8.2 Move 2: Contrapositive . . . 14
8.3 Move 3: Contradiction . . . 14

9 Examples and Counterexamples 16
9.1 Example . . . 16
9.2 Non-example . . . 16
9.3 Counterexample . . . 16
9.4 The useful cruelty of counterexamples . . . 16
9.5 Exercises . . . 16

10 Sets: The Grammar of Modern Mathematics 17
10.1 Basic operations . . . 17
10.2 Subset . . . 17
10.3 Worked proof . . . 17
10.4 Historical note . . . 18

11 Set Proofs: Element Chasing 19
11.1 What to notice . . . 19
11.2 Exercises . . . 19

12 Functions: More Than Formulas 20

12.1 Why codomain matters . . . 20
12.2 Injective . . . 20
12.3 Surjective . . . 20
12.4 Bijective . . . 20

13 Function Proofs **21**
13.1 Inverse images . . . 21
13.2 Why inverse images matter later . . . 21

14 Relations: When Objects Count as Connected **22**
14.1 Equivalence relations . . . 22
14.2 Congruence modulo n . . . 22
14.3 Why relations matter later . . . 22

15 Induction: Infinite Proof With Finite Labor **23**
15.1 Worked proof . . . 23
15.2 Common induction bug . . . 23

16 Strong Induction and Factorization **24**
16.1 Prime factorization existence . . . 24
16.2 Why this matters . . . 24

17 Equality, Identity, and Isomorphism **25**
17.1 Equality . . . 25
17.2 Isomorphism . . . 25
17.3 Why this matters . . . 25

18 Infinity: Countable and Uncountable **26**
18.1 The integers are countable . . . 26
18.2 The rationals are countable . . . 26
18.3 The reals are uncountable . . . 26
18.4 Why this matters later . . . 26

19 First Epsilon Proofs **27**
19.1 Worked proof . . . 27
19.2 The secret . . . 27

20 Continuity: Local Control **28**

20.1 Worked proof . . . 28
20.2 Connection to topology . . . 28

21 Algebra Begins With Symmetry 29
21.1 Worked proof: identity is unique . . . 29
21.2 What to notice . . . 29

22 Topology Begins With Nearness Without Rulers 30
22.1 Physical intuition . . . 30
22.2 Warning . . . 30

23 Differential Geometry: Calculus on Curved Spaces 31
23.1 Connection to previous material . . . 31
23.2 Physical connection . . . 31

24 Algebraic Geometry: Equations Become Shapes 32
24.1 Affine varieties . . . 32
24.2 Ideals . . . 32
24.3 The bridge . . . 32

25 How to Read a Definition 33
25.1 The five-part definition routine . . . 33
25.2 Example: injective . . . 33

26 How to Read a Theorem 34
26.1 The theorem-reading template . . . 34

27 How to Write a Proof 35
27.1 Proof skeleton . . . 35
27.2 Sentences to avoid . . . 35
27.3 A better replacement . . . 35

28 Proof Debugging 36
28.1 The first broken line . . . 36
28.2 Gap types . . . 36
28.3 Bad proof, repaired . . . 36

29 Exercises for Walking 37

30 A Four-Week Beginner-to-Velleman Launch **38**
30.1 Daily rhythm . . . 38
30.2 When to start Velleman . . . 38

31 Milestone Proof Portfolio **39**

32 Closing: The Shape of the Road **40**

A Appendix A: Notation You Should Be Able to Read **41**
A.1 Notation drills . . . 41

B Appendix B: A Short History of Why Rigor Arrived **42**
B.1 Euclid and the axiomatic dream . . . 42
B.2 Calculus and the problem of motion . . . 42
B.3 Cauchy and Weierstrass . . . 42
B.4 Cantor and infinity . . . 42
B.5 Noether and structure . . . 42
B.6 Hilbert and formal clarity . . . 42
B.7 Grothendieck and the expansion of geometry . . . 43

C Appendix C: Extra Proof Drills **44**
C.1 Drill block 1: Definitions only . . . 44
C.2 Drill block 2: Counterexamples . . . 44
C.3 Drill block 3: Proof strategy choice . . . 44

D Appendix D: Hints Without Full Solutions **45**
D.1 Definitions only . . . 45
D.2 Counterexamples . . . 45
D.3 Strategy choice . . . 45

E Appendix E: Reading Velleman Without Getting Lost **46**
E.1 Before each section . . . 46
E.2 During each section . . . 46
E.3 After each section . . . 46
E.4 What “understanding” should feel like . . . 46

F Appendix F: The One-Page Proof Debugger **47**

G Appendix G: Final Readiness Exam **48**

G.1 How to score yourself 48

Preface: What This Reader Is

This is the full self-contained reader rather than a syllabus or summary checklist: it works through the definitions, examples, and proof techniques directly.

The goal is simple to say and serious to do: take someone who knows useful engineering mathematics and help them become comfortable with proof-based pure mathematics. That means learning to read slowly, learning to use definitions as tools, learning to make examples and counterexamples, learning to write proofs that survive inspection, and learning why these little habits eventually open the doors to real analysis, abstract algebra, topology, differential geometry, and algebraic geometry.

The first major book is Velleman's *How to Prove It*. This reader is meant to make the opening of that book feel familiar instead of foggy. We will start before Velleman starts: with mathematical sentences themselves.

The tone here is friendly, but the destination is rigor. A proof is not a vibe. It is not a pile of plausible sentences. It is a sequence of justified moves from assumptions to conclusion. But rigor does not have to arrive wearing a cold face. Mathematics was built by human beings trying to understand number, shape, motion, symmetry, infinity, and structure. The formal language came later because the ideas were powerful enough to need protection.

Focus

Read with a pencil. Do the small exercises. Say definitions aloud. If a sentence feels slippery, do not push through it. Stop and ask: what are the objects, what is being assumed, and what must be shown?

1 The Change in Game

Engineering mathematics often trains excellent reflexes. You see a differential equation; you try separation, integrating factors, Laplace transforms, eigenvalues. You see an integral; you look for substitution, parts, symmetry, a known transform. You see a matrix; you think rank, determinant, eigenvectors, conditioning.

That is real skill. Do not throw it away.

But pure mathematics asks a different question first. Instead of "What method solves this?", it asks:

> What exactly is being claimed, and why must it follow from the assumptions?

That one question changes the whole room.

1.1 From calculation to justification

Calculation is still present. Pure mathematicians compute all the time. But computation is no longer the whole story. A computation can suggest a pattern; a proof explains why the pattern cannot fail.

For example:

$$1+3=4, \qquad 1+3+5=9, \qquad 1+3+5+7=16.$$

You might guess:

$$1+3+5+\cdots+(2n-1)=n^2.$$

A calculator can check the first thousand cases. A proof covers all cases at once.

1.2 The big shift

The shift is from answer-getting to structure-seeing. You are learning to ask:

- What kind of object is this?
- What is the exact definition?
- What examples satisfy it?
- What examples almost satisfy it but fail?
- What is the theorem really promising?
- What would break it?
- Which assumptions are doing work?

Walking question

On a walk, take one theorem you know from calculus, like "continuous functions on closed intervals attain maximum values." Ask: which words are essential? Continuous? Closed? Bounded? Interval? What breaks if one word disappears?

2 Statements, Predicates, and the First Filter

Mathematics begins with sentences that can be judged true or false.

Definition 1. A *statement* is a sentence that is either true or false.

Example 1. "5 is prime" is a statement. It is true.

Example 2. "9 is prime" is a statement. It is false.

Example 3. "Find the derivative of x^2" is not a statement. It is a command.

2.1 Predicates

Definition 2. A *predicate* is a sentence with one or more variables that becomes a statement when the variables are assigned values from a domain.

The sentence

$$x^2 > 10$$

is not true or false until x is known. If $x = 4$, true. If $x = 2$, false. So $x^2 > 10$ is a predicate.

In software terms, a predicate is like a Boolean-valued function:

$$P(x) = \text{“}x^2 > 10\text{”}.$$

But mathematics demands something software often hides: the domain. Are we allowing $x \in \mathbb{N}$, $x \in \mathbb{Z}$, $x \in \mathbb{R}$, or $x \in \mathbb{C}$? The sentence may change character when the domain changes.

2.2 Why this matters

A beginner often reads a theorem too quickly. The theorem is a sentence with machinery inside it. Variables have domains. Quantifiers control choice. Hypotheses give permissions. Conclusions give obligations.

Before proving anything, learn to parse.

2.3 Exercises

For each item, decide whether it is a statement, predicate, command, or expression.

1. $2 + 2 = 4$.
2. $x + 2 = 4$.
3. Solve $x + 2 = 4$.
4. There are infinitely many prime numbers.
5. n is divisible by 3.
6. $\int_0^1 x^2\,dx$.
7. Every polynomial of odd degree with real coefficients has a real root.

8. Is every continuous function differentiable?

Focus

Training target: never begin a proof until you know what the sentence is saying.

3 Variables Are Not All the Same

One source of mathematical fog is that every variable looks like a letter, but not every variable has the same role.

Some variables are fixed. Some are arbitrary. Some are chosen as witnesses. Some are temporary. Some are bound by quantifiers. Some are free.

3.1 Fixed but unknown

When a proof says:

> Let $n \in \mathbb{Z}$.

It means n is now fixed for the discussion. We may not know which integer it is, but we are treating it as one particular integer.

3.2 Arbitrary

When a proof says:

> Let $n \in \mathbb{Z}$ be arbitrary.

It means we are not using any special property of n except that it is an integer. If the argument works, it works for every integer.

This is how universal proofs begin.

3.3 Witness

When proving an existential statement, a variable may be chosen to prove existence.

Claim:

$$\exists n \in \mathbb{Z} \text{ such that } n^2 = 81.$$

Proof:

> Take $n = 9$. Then $n \in \mathbb{Z}$ and $n^2 = 81$.

Here 9 is the witness.

3.4 Bound and free variables

In

$$\forall x \in \mathbb{R},\ x^2 \geq 0,$$

the variable x is bound by the quantifier $\forall$. It is local to that sentence.

In

$$x^2 \geq 0,$$

the variable x is free unless context has already assigned it a meaning.

This is close to scope in programming. A loop variable belongs to the loop. A local variable belongs to the function. A mathematical variable belongs to the sentence or proof context that introduced it.

3.5 Exercises

In each sentence, identify the role of the variable.

1. Let $x \in \mathbb{R}$ be arbitrary.
2. Choose $n = 12$.
3. For every $a \in A$, there exists $b \in B$ such that $f(a) = b$.
4. Suppose p is prime.
5. Let $\varepsilon > 0$.

4 Quantifiers: The Tiny Symbols That Run the Machine

The two most important symbols at the beginning are:

$$\forall \quad \text{and} \quad \exists.$$

$$\forall = \text{for every}, \qquad \exists = \text{there exists}.$$

Small symbols. Large consequences.

4.1 Universal statements

A universal statement says every object in a domain has some property:

$$\forall x \in D,\ P(x).$$

To prove it, take an arbitrary $x \in D$ and prove $P(x)$.

To disprove it, find one counterexample.

4.2 Existential statements

An existential statement says at least one object in a domain has some property:

$$\exists x \in D \text{ such that } P(x).$$

To prove it, provide a witness or prove one must exist.

To disprove it, prove every object in the domain fails the property.

4.3 The order matters

Compare:

$$\forall x \in \mathbb{R},\ \exists y \in \mathbb{R} \text{ such that } x + y = 0.$$

$$\exists y \in \mathbb{R} \text{ such that } \forall x \in \mathbb{R},\ x + y = 0.$$

The first is true. Each x gets its own $y = -x$.

The second is false. It asks for one y that cancels every x.

This is the sort of thing that makes proofs feel mysterious if you read too fast. The symbols are not decoration. They describe a game of choice.

4.4 A game interpretation

Think of $\forall$ as an adversary choosing an object. Think of $\exists$ as you responding with an object.

For

$$\forall x \in \mathbb{R},\ \exists y \in \mathbb{R},\ x + y = 0,$$

the adversary hands you x. You respond with $y = -x$. You win every time.

For

$$\exists y \in \mathbb{R},\ \forall x \in \mathbb{R},\ x + y = 0,$$

you must choose y first. Then the adversary gets to choose x. Whatever y you choose, the adversary can choose $x = 1 - y$, and then $x + y = 1$, not 0. You lose.

Walking question

Ask yourself: in this theorem, who chooses first? Do I get to choose an object after seeing the input, or must I produce one object that works for all inputs?

5 Negation: Turning Claims Inside Out

Negation is not just putting "not" at the front. Good negation preserves the exact structure while reversing the truth.

5.1 The basic laws

$$\neg(\forall x \in D,\ P(x)) \iff \exists x \in D \text{ such that } \neg P(x).$$

$$\neg(\exists x \in D,\ P(x)) \iff \forall x \in D,\ \neg P(x).$$

If someone says, "Every bridge is open," one closed bridge disproves it. If someone says, "There is an open bridge," you must show every bridge is closed to disprove it.

5.2 Negating implication

The statement

$$P \Rightarrow Q$$

is false exactly when P is true and Q is false.

So:

$$\neg(P \Rightarrow Q) \iff P \wedge \neg Q.$$

This matters when you test the boundary of a theorem.

Claim:

> If a function is continuous, then it is differentiable.

To disprove it, you need a function that is continuous and not differentiable. The absolute value function $f(x) = |x|$ at 0 is the classic first example.

5.3 Nested negations

Negate:

$$\forall \varepsilon > 0,\ \exists N \in \mathbb{N} \text{ such that } \forall n \geq N,\ |a_n - L| < \varepsilon.$$

Step by step:

$$\exists \varepsilon > 0 \text{ such that } \forall N \in \mathbb{N},\ \exists n \geq N \text{ such that } |a_n - L| \geq \varepsilon.$$

This is the negation of sequence convergence. In words: there is some fixed tolerance ε such that no matter how far out you go, the sequence still eventually escapes that tolerance.

That sentence is not just logic practice. It is analysis beginning to breathe.

5.4 Exercises

Negate the following.

1. Every integer is even.
2. There exists a real number x such that $x^2 < 0$.
3. For every $x \in A$, there exists $y \in B$ such that $R(x, y)$.
4. If n^2 is even, then n is even.
5. Every continuous function on $\mathbb{R}$ is bounded.

6 If-Then: The Spine of Theorems

Many theorems have the form:

$$P \Rightarrow Q.$$

The hypothesis P is what you may assume. The conclusion Q is what you must prove.

6.1 Direct proof

The direct proof is the plain road:

1. Assume P.
2. Use definitions and known facts.
3. Derive Q.

Theorem 1. *If n is divisible by 6, then n is divisible by 3.*

Proof. Suppose n is divisible by 6. Then there exists $k \in \mathbb{Z}$ such that $n = 6k$. Since $6k = 3(2k)$ and $2k \in \mathbb{Z}$, there exists an integer $m = 2k$ such that $n = 3m$. Therefore n is divisible by 3. □

6.2 What happened?

The proof opened the definition of divisibility by 6 and rearranged it into the definition of divisibility by 3.

That is a large fraction of beginner proof writing.

Focus

When stuck, ask: what does the hypothesis mean by definition, and what does the conclusion require by definition?

6.3 Converse

The converse of $P \Rightarrow Q$ is $Q \Rightarrow P$.

The theorem "if n is divisible by 6, then n is divisible by 3" is true. Its converse says "if n is divisible by 3, then n is divisible by 6." False. Example: $n = 3$.

Many false proofs accidentally prove or assume the converse.

7 Definitions Are Machines

Definitions are where rigor begins. A definition tells you exactly what must be shown and exactly what may be used.

Definition 3. An integer n is even if there exists $k \in \mathbb{Z}$ such that $n = 2k$.

Definition 4. An integer n is odd if there exists $k \in \mathbb{Z}$ such that $n = 2k + 1$.

Definition 5. For integers a, b, we say a divides b, written $a \mid b$, if there exists $k \in \mathbb{Z}$ such that $b = ak$.

7.1 Opening and closing definitions

If you know n is even, you may open the definition:

$$n = 2k \text{ for some } k \in \mathbb{Z}.$$

If you want to prove n is even, you must close the definition: find an integer k such that $n = 2k$.

7.2 Worked proof: even plus even

Theorem 2. *The sum of two even integers is even.*

Proof. Let $a, b \in \mathbb{Z}$ be even. Since a is even, there exists $m \in \mathbb{Z}$ such that $a = 2m$. Since b is even, there exists $n \in \mathbb{Z}$ such that $b = 2n$. Then

$$a + b = 2m + 2n = 2(m + n).$$

Since $m + n \in \mathbb{Z}$, the number $a + b$ has the form $2k$ for some integer k. Therefore $a + b$ is even. □

7.3 Worked proof: divisibility is transitive

Theorem 3. *If $a, b, c \in \mathbb{Z}$, $a \mid b$, and $b \mid c$, then $a \mid c$.*

Proof. Suppose $a \mid b$ and $b \mid c$. Since $a \mid b$, there exists $m \in \mathbb{Z}$ such that $b = am$. Since $b \mid c$, there exists $n \in \mathbb{Z}$ such that $c = bn$. Substituting $b = am$ into $c = bn$, we get

$$c = (am)n = a(mn).$$

Since $mn \in \mathbb{Z}$, this shows $a \mid c$. □

Walking question

Pick a definition you know. Ask: if I assume it, what algebraic or logical object do I get? If I need to prove it, what object must I produce?

8 The Three First Proof Moves

At the start, you do not need a giant toolbox. You need a few reliable moves.

8.1 Move 1: Direct proof

Use it when the hypothesis naturally gives the conclusion.

$$P \Rightarrow Q.$$

Assume P. Prove Q.

8.2 Move 2: Contrapositive

The contrapositive of

$$P \Rightarrow Q$$

is

$$\neg Q \Rightarrow \neg P.$$

These are logically equivalent.

Use contrapositive when $\neg Q$ gives more concrete information than P.

Theorem 4. *If n^2 is even, then n is even.*

Proof. We prove the contrapositive: if n is not even, then n^2 is not even. For integers, not even means odd. Suppose n is odd. Then $n = 2k + 1$ for some $k \in \mathbb{Z}$. Then

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1.$$

Since $2k^2 + 2k \in \mathbb{Z}$, n^2 is odd. Therefore n^2 is not even. By contrapositive, if n^2 is even, then n is even. □

8.3 Move 3: Contradiction

To prove P, assume $\neg P$ and derive an impossibility.

Contradiction is powerful, but beginners sometimes use it when direct proof would be cleaner. It is a sharp knife. Use it with care.

Theorem 5. *$\sqrt{2}$ is irrational.*

Proof. Suppose, for contradiction, that $\sqrt{2}$ is rational. Then $\sqrt{2} = a/b$ for integers a, b with $b \neq 0$, and we may choose the fraction in lowest terms. Squaring gives

$$2 = \frac{a^2}{b^2},$$

so

$$a^2 = 2b^2.$$

Thus a^2 is even, so a is even. Write $a = 2k$. Then

$$(2k)^2 = 2b^2,$$

so

$$4k^2 = 2b^2,$$

and hence

$$b^2 = 2k^2.$$

Thus b^2 is even, so b is even. Therefore both a and b are even, contradicting the assumption that a/b was in lowest terms. Hence $\sqrt{2}$ is irrational. □

9 Examples and Counterexamples

Examples are not decorations. They are how you test whether your understanding has touched the ground.

9.1 Example

A function $f : \mathbb{R} \to \mathbb{R}$ that is injective:

$$f(x) = 2x + 1.$$

If $f(a) = f(b)$, then $2a + 1 = 2b + 1$, so $a = b$.

9.2 Non-example

A function $f : \mathbb{R} \to \mathbb{R}$ that is not injective:

$$f(x) = x^2.$$

Because $f(1) = 1 = f(-1)$ but $1 \neq -1$.

9.3 Counterexample

A counterexample is an example that disproves a universal statement.

Claim:

$$\forall x, y \in \mathbb{R},\ x^2 = y^2 \Rightarrow x = y.$$

Counterexample: $x = 1$, $y = -1$. Then $x^2 = y^2$, but $x \neq y$.

9.4 The useful cruelty of counterexamples

Counterexamples are not mean. They are honest. A theorem with no boundary is usually not a theorem; it is a guess.

When a theorem is false, a counterexample shows where it breaks. When a theorem is true, failed counterexamples teach you which assumptions are secretly holding the roof up.

9.5 Exercises

Find counterexamples.

1. Every function $f : \mathbb{R} \to \mathbb{R}$ is surjective.
2. If $ab = 0$, then $a = 0$ and $b = 0$.
3. If $A \cup B = A \cup C$, then $B = C$.
4. Every nonempty subset of $\mathbb{R}$ has a smallest element.
5. Every continuous function is differentiable.

10 Sets: The Grammar of Modern Mathematics

A set is a collection of objects. The objects are called elements.

$$x \in A$$

means x is an element of A.

$$x \notin A$$

means x is not an element of A.

Sets look innocent, but they became the common language of modern mathematics because they let us talk about collections of numbers, functions, points, symmetries, shapes, solutions, and open neighborhoods with one grammar.

10.1 Basic operations

$$A \cup B = \{x : x \in A \text{ or } x \in B\}.$$

$$A \cap B = \{x : x \in A \text{ and } x \in B\}.$$

$$A \setminus B = \{x : x \in A \text{ and } x \notin B\}.$$

10.2 Subset

Definition 6. $A \subseteq B$ means every element of A is an element of B.

To prove $A \subseteq B$:

> Let $x \in A$ be arbitrary. Show $x \in B$.

To prove $A = B$:

> Prove $A \subseteq B$ and $B \subseteq A$.

10.3 Worked proof

Theorem 6. *For any sets A and B,*

$$A \setminus B = A \cap B^c$$

where B^c is the complement of B in the ambient universe.

Proof. Let x be arbitrary. Then

$$x \in A \setminus B$$

if and only if $x \in A$ and $x \notin B$. But $x \notin B$ means $x \in B^c$. Therefore $x \in A$ and $x \in B^c$, which means $x \in A \cap B^c$. Thus $A \setminus B = A \cap B^c$. □

10.4 Historical note

Set theory became central in the nineteenth century, especially through Cantor's work on infinity. Before Cantor, infinity often behaved like smoke in the room: useful, present, but hard to hold. Set theory gave mathematicians a way to compare infinite collections with precision.

11 Set Proofs: Element Chasing

Set proofs often reduce to following one element through definitions.

Theorem 7. *For all sets A, B, C,*

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$$

Proof. We prove both inclusions.

First suppose $x \in A \cap (B \cup C)$. Then $x \in A$ and $x \in B \cup C$. Since $x \in B \cup C$, either $x \in B$ or $x \in C$.

If $x \in B$, then $x \in A \cap B$, so $x \in (A \cap B) \cup (A \cap C)$.

If $x \in C$, then $x \in A \cap C$, so $x \in (A \cap B) \cup (A \cap C)$.

So $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$.

Now suppose $x \in (A \cap B) \cup (A \cap C)$. Then either $x \in A \cap B$ or $x \in A \cap C$.

If $x \in A \cap B$, then $x \in A$ and $x \in B$. Hence $x \in B \cup C$, so $x \in A \cap (B \cup C)$.

If $x \in A \cap C$, then $x \in A$ and $x \in C$. Hence $x \in B \cup C$, so $x \in A \cap (B \cup C)$.

So $(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C)$. Therefore the sets are equal. □

11.1 What to notice

The proof did not draw a Venn diagram. Venn diagrams are useful for intuition, but they are not proof enough for arbitrary sets. The rigorous proof follows membership.

Focus

Set identities are where proof writing becomes grammar practice. Let x live on one side. Translate definitions until it lives on the other.

11.2 Exercises

Prove:

1. $A \cap B \subseteq A$.
2. $A \subseteq A \cup B$.
3. $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$.
4. $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$.

12 Functions: More Than Formulas

A function is often introduced as a formula, but a function is not just a formula.

Definition 7. A function $f : A \to B$ assigns to every element of A exactly one element of B.

The set A is the domain. The set B is the codomain.

The image is:

$$\mathrm{im}(f) = \{f(a) : a \in A\}.$$

12.1 Why codomain matters

The formula $f(x) = x^2$ can define many different functions:

$$f : \mathbb{R} \to \mathbb{R}, \qquad f : \mathbb{R} \to [0, \infty), \qquad f : \mathbb{N} \to \mathbb{N}.$$

Same formula. Different function, because the domain and codomain are part of the data.

12.2 Injective

Definition 8. $f : A \to B$ is injective if for all $a_1, a_2 \in A$,

$$f(a_1) = f(a_2) \Rightarrow a_1 = a_2.$$

Injective means no merging. If two outputs are equal, the inputs must have been equal.

12.3 Surjective

Definition 9. $f : A \to B$ is surjective if for every $b \in B$, there exists $a \in A$ such that $f(a) = b$.

Surjective means every target in the codomain is hit.

12.4 Bijective

Definition 10. f is bijective if it is both injective and surjective.

Bijections are mathematical pairings. They say two sets have the same structure of elements, at least from the viewpoint of counting.

13 Function Proofs

Theorem 8. *If $f : A \to B$ and $g : B \to C$ are injective, then $g \circ f : A \to C$ is injective.*

Proof. Let $a_1, a_2 \in A$ and suppose $(g \circ f)(a_1) = (g \circ f)(a_2)$. Then

$$g(f(a_1)) = g(f(a_2)).$$

Since g is injective, $f(a_1) = f(a_2)$. Since f is injective, $a_1 = a_2$. Therefore $g \circ f$ is injective. □

13.1 Inverse images

If $f : A \to B$ and $S \subseteq B$, the inverse image of S is:

$$f^{-1}(S) = \{a \in A : f(a) \in S\}.$$

This notation does not require f to have an inverse function. It is a set operation.

Theorem 9. *For $f : A \to B$ and $S, T \subseteq B$,*

$$f^{-1}(S \cap T) = f^{-1}(S) \cap f^{-1}(T).$$

Proof. Let $a \in A$. Then

$$a \in f^{-1}(S \cap T)$$

if and only if

$$f(a) \in S \cap T.$$

This holds if and only if $f(a) \in S$ and $f(a) \in T$. That holds if and only if $a \in f^{-1}(S)$ and $a \in f^{-1}(T)$. Therefore $a \in f^{-1}(S) \cap f^{-1}(T)$. Hence the sets are equal. □

13.2 Why inverse images matter later

Topology defines continuity by inverse images:

$$f \text{ is continuous if } f^{-1}(U) \text{ is open whenever } U \text{ is open.}$$

That sentence is not beginner fluff. It is a bridge to topology. The habit of understanding $f^{-1}(S)$ now pays off later.

14 Relations: When Objects Count as Connected

A relation on a set A is a rule that tells whether two elements of A are related.

Examples:

- $a = b$ on $\mathbb{Z}$.
- $a \leq b$ on $\mathbb{R}$.
- $a \mid b$ on $\mathbb{Z}$.
- $a \equiv b \pmod{n}$ on $\mathbb{Z}$.

14.1 Equivalence relations

Definition 11. A relation $\sim$ on A is an equivalence relation if it is reflexive, symmetric, and transitive.

Reflexive:

$$a \sim a.$$

Symmetric:

$$a \sim b \Rightarrow b \sim a.$$

Transitive:

$$a \sim b \text{ and } b \sim c \Rightarrow a \sim c.$$

14.2 Congruence modulo n

Define $a \equiv b \pmod{n}$ if $n \mid (a - b)$.

Theorem 10. *Congruence modulo n is an equivalence relation on $\mathbb{Z}$.*

Proof. Reflexive: For any $a \in \mathbb{Z}$, $a - a = 0$, and $n \mid 0$, so $a \equiv a \pmod{n}$.

Symmetric: Suppose $a \equiv b \pmod{n}$. Then $n \mid (a - b)$, so $a - b = nk$ for some $k \in \mathbb{Z}$. Then $b - a = -nk = n(-k)$, so $n \mid (b - a)$. Hence $b \equiv a \pmod{n}$.

Transitive: Suppose $a \equiv b \pmod{n}$ and $b \equiv c \pmod{n}$. Then $n \mid (a - b)$ and $n \mid (b - c)$. So $a - b = nm$ and $b - c = n\ell$ for integers m, ℓ. Adding,

$$a - c = (a - b) + (b - c) = nm + n\ell = n(m + \ell).$$

Thus $n \mid (a - c)$, so $a \equiv c \pmod{n}$. □

14.3 Why relations matter later

Equivalence relations create quotient structures. Quotients are everywhere: modular arithmetic, quotient groups, quotient rings, projective spaces, manifolds built by gluing, and more. A quotient says: we will treat these different-looking things as the same under a chosen rule.

15 Induction: Infinite Proof With Finite Labor

Induction is one of the first places mathematics feels like a magic trick until you see the mechanism.

Theorem 11 (Principle of induction)**.** *Suppose $P(n)$ is a statement for each integer $n \geq 1$. If $P(1)$ is true, and for every $k \geq 1$, $P(k) \Rightarrow P(k+1)$, then $P(n)$ is true for all $n \geq 1$.*

The image is dominoes. Knock down the first. Show each knocks down the next. Then all fall.

15.1 Worked proof

Theorem 12. *For every $n \geq 1$,*

$$1 + 2 + \cdots + n = \frac{n(n+1)}{2}.$$

Proof. Base case: For $n = 1$, the left side is 1 and the right side is $\frac{1(2)}{2} = 1$. So the statement holds for $n = 1$.

Induction step: Assume for some $k \geq 1$ that

$$1 + 2 + \cdots + k = \frac{k(k+1)}{2}.$$

We must prove

$$1 + 2 + \cdots + k + (k+1) = \frac{(k+1)(k+2)}{2}.$$

Using the induction hypothesis,

$$1 + 2 + \cdots + k + (k+1) = \frac{k(k+1)}{2} + (k+1).$$

Then

$$\frac{k(k+1)}{2} + (k+1) = \frac{k(k+1) + 2(k+1)}{2} = \frac{(k+1)(k+2)}{2}.$$

So the statement holds for $k+1$. By induction, it holds for all $n \geq 1$. □

15.2 Common induction bug

Do not assume what you are trying to prove for $k+1$. The induction hypothesis gives you $P(k)$, not $P(k+1)$.

Focus

In induction, always label the base case, induction hypothesis, and induction step.

16 Strong Induction and Factorization

Strong induction lets you assume all earlier cases, not just the immediately previous one.

Theorem 13 (Strong induction)**.** *Suppose $P(n)$ is a statement for integers $n \geq n_0$. If $P(n_0)$ is true, and for every $k \geq n_0$, the truth of $P(n_0), P(n_0+1), \ldots, P(k)$ implies $P(k+1)$, then $P(n)$ is true for all $n \geq n_0$.*

16.1 Prime factorization existence

Theorem 14. *Every integer $n \geq 2$ is a product of primes.*

Proof. We use strong induction on n.

Base case: 2 is prime, so it is a product of primes.

Induction step: Suppose every integer m with $2 \leq m \leq k$ is a product of primes. We prove $k+1$ is a product of primes.

If $k+1$ is prime, then it is already a product of primes.

If $k+1$ is not prime, then $k+1 = ab$ for integers a, b with $2 \leq a \leq k$ and $2 \leq b \leq k$. By the induction hypothesis, both a and b are products of primes. Therefore $ab = k+1$ is a product of primes.

Thus by strong induction, every integer $n \geq 2$ is a product of primes. □

16.2 Why this matters

This proof is a doorway to number theory. Later, unique factorization becomes a central theme. In ordinary integers, factorization behaves well. In other number systems, it can fail. That failure forced mathematicians to invent ideals, which later became essential in algebraic geometry.

Walking question

Think about software dependencies. A package can depend on smaller packages, which depend on smaller packages, until you reach primitives. Strong induction is the proof version of trusting all smaller dependencies.

17 Equality, Identity, and Isomorphism

Beginners often think mathematics only cares about equality. Later mathematics cares just as much about sameness of structure.

17.1 Equality

Equality means two expressions name the same object.

$$2 + 3 = 5.$$

17.2 Isomorphism

An isomorphism means two structures may have different labels but the same pattern.

For example, the even integers under addition behave like the integers under addition:

$$\mathbb{Z} \to 2\mathbb{Z}, \qquad n \mapsto 2n.$$

This map preserves addition:

$$2(a + b) = 2a + 2b.$$

The labels changed. The additive structure did not.

17.3 Why this matters

Abstract algebra is full of this idea. Topology has homeomorphism. Differential geometry has diffeomorphism. Category theory makes structure-preserving maps the main characters.

The deep habit: do not ask only whether two things are equal. Ask what structure is preserved by the map between them.

18 Infinity: Countable and Uncountable

Infinity is where mathematics learned humility.

The natural numbers are infinite:

$$\mathbb{N} = \{1, 2, 3, \ldots\}.$$

The even positive integers are also infinite:

$$\{2, 4, 6, \ldots\}.$$

There are "half as many" evens in an everyday sense, but the map

$$n \mapsto 2n$$

pairs every natural number with exactly one even positive integer. So the two sets have the same cardinality.

Definition 12. A set is countably infinite if its elements can be listed in a sequence indexed by $\mathbb{N}$.

18.1 The integers are countable

List them:

$$0, 1, -1, 2, -2, 3, -3, \ldots$$

Every integer appears eventually.

18.2 The rationals are countable

Fractions seem denser. Between any two rationals, there is another rational. Still, $\mathbb{Q}$ is countable.

Imagine a grid of fractions p/q with $p \in \mathbb{Z}$ and $q \in \mathbb{N}$. Sweep through the grid by increasing $|p| + q$, skipping duplicates like $2/4 = 1/2$. Every rational appears eventually.

18.3 The reals are uncountable

Cantor's diagonal argument shows no list of real numbers in $(0, 1)$ can be complete. Given any proposed list of decimals, build a new decimal that differs from the first number in the first digit, the second number in the second digit, and so on. The new number differs from every listed number, so it was missing.

Infinity has sizes.

18.4 Why this matters later

Analysis lives among uncountable sets. Topology studies spaces where infinity can behave subtly. Algebraic geometry often studies solution sets with infinitely many points controlled by finite equations.

19 First Epsilon Proofs

Engineering calculus often treats limits as motion. A variable approaches. A graph settles. A value tends toward another value.

Real analysis asks for the contract.

Definition 13. A sequence (a_n) converges to L if for every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$,

$$|a_n - L| < \varepsilon.$$

Translation: no matter how small a tolerance band you choose around L, the sequence eventually stays inside it forever.

19.1 Worked proof

Theorem 15.

$$\lim_{n\to\infty} \frac{1}{n} = 0.$$

Proof. Let $\varepsilon > 0$ be arbitrary. We need to find $N \in \mathbb{N}$ such that for all $n \geq N$,

$$\left|\frac{1}{n} - 0\right| < \varepsilon.$$

Choose $N \in \mathbb{N}$ such that $N > 1/\varepsilon$. Then if $n \geq N$,

$$0 < \frac{1}{n} \leq \frac{1}{N} < \varepsilon.$$

Thus $\left|\frac{1}{n} - 0\right| < \varepsilon$. Therefore $1/n \to 0$. □

19.2 The secret

You do not choose ε. The challenger chooses ε. You choose N in response.

Walking question

Think of ε as a tolerance in a sensor. Someone sets the tolerance. Your job is to say how far into the sequence you must go before the readings stay within tolerance forever.

20 Continuity: Local Control

The epsilon-delta definition of continuity looks intimidating because it has nested quantifiers. But the idea is physical:

> Small changes in input force small changes in output.

Definition 14. A function $f : \mathbb{R} \to \mathbb{R}$ is continuous at $c \in \mathbb{R}$ if for every $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in \mathbb{R}$, if $|x - c| < \delta$, then $|f(x) - f(c)| < \varepsilon$.

20.1 Worked proof

Theorem 16. *The function $f(x) = 3x + 2$ is continuous at every $c \in \mathbb{R}$.*

Proof. Let $c \in \mathbb{R}$ and let $\varepsilon > 0$ be arbitrary. We need $\delta > 0$ such that $|x-c| < \delta$ implies $|f(x)-f(c)| < \varepsilon$.

Now

$$|f(x) - f(c)| = |(3x + 2) - (3c + 2)| = |3x - 3c| = 3|x - c|.$$

Choose $\delta = \varepsilon/3$. If $|x - c| < \delta$, then

$$|f(x) - f(c)| = 3|x - c| < 3\delta = 3(\varepsilon/3) = \varepsilon.$$

Therefore f is continuous at c. □

20.2 Connection to topology

Topology later replaces epsilon-delta with open sets. Instead of measuring distances directly, it says a function is continuous when inverse images of open sets are open. That is a huge abstraction, but it is not alien. It is the same idea: local neighborhoods are respected.

21 Algebra Begins With Symmetry

Abstract algebra can look like someone removed all the numbers and left only laws. That is partly right. Algebra asks what can be known when the objects change but the operation rules remain.

Definition 15. A group is a set G with an operation $\cdot$ such that:

1. closure: if $a, b \in G$, then $a \cdot b \in G$,
2. associativity: $(a \cdot b) \cdot c = a \cdot (b \cdot c)$,
3. identity: there exists $e \in G$ such that $e \cdot a = a \cdot e = a$ for all $a \in G$,
4. inverses: for each $a \in G$, there exists $a^{-1} \in G$ such that $a \cdot a^{-1} = a^{-1} \cdot a = e$.

Examples:

- $\mathbb{Z}$ under addition.
- Nonzero real numbers under multiplication.
- Symmetries of a square under composition.
- Invertible matrices under multiplication.

21.1 Worked proof: identity is unique

Theorem 17. *The identity element of a group is unique.*

Proof. Suppose e and e' are both identity elements in a group G. Since e is an identity, $e \cdot e' = e'$. Since e' is an identity, $e \cdot e' = e$. Therefore $e = e'$. Hence the identity element is unique. □

21.2 What to notice

The proof did not compute. It used only the definition. That is abstract algebra in miniature.

22 Topology Begins With Nearness Without Rulers

Topology studies continuity and shape in a way that does not always require measuring distance.

Definition 16. A topology on a set X is a collection $\mathcal{T}$ of subsets of X, called open sets, such that:

1. $\emptyset$ and X are in $\mathcal{T}$,
2. arbitrary unions of sets in $\mathcal{T}$ are in $\mathcal{T}$,
3. finite intersections of sets in $\mathcal{T}$ are in $\mathcal{T}$.

This definition looks sparse. But it captures enough structure to talk about continuity.

Definition 17. If X and Y are topological spaces, a function $f : X \to Y$ is continuous if for every open set $U \subseteq Y$, the inverse image $f^{-1}(U)$ is open in X.

Now the earlier work on inverse images returns.

22.1 Physical intuition

Topology cares about neighborhoods. If a point lies in an open set, it has some room around it. The exact distance may not matter. What matters is the pattern of which neighborhoods exist.

This is why a coffee mug and a donut are famously similar in topology: each has one hole. Topology ignores bending and stretching but notices tearing and gluing.

22.2 Warning

The coffee mug story is memorable, but do not let it replace definitions. The definitions are the engine. The story is the spark plug.

23 Differential Geometry: Calculus on Curved Spaces

Differential geometry begins when calculus leaves flat Euclidean space.

A curve can be studied with derivatives. A surface can be studied with tangent planes. A manifold is the general idea: a space that may be globally complicated but locally resembles $\mathbb{R}^n$.

Definition 18. Informally, an n-dimensional manifold is a space that near each point looks like $\mathbb{R}^n$.

The phrase "looks like" becomes precise using charts.

Definition 19. A chart is a coordinate map from part of a manifold to an open subset of $\mathbb{R}^n$.

23.1 Connection to previous material

To understand manifolds, you need:

- functions and inverse images,
- open sets and continuity,
- linear algebra,
- multivariable calculus,
- proof comfort.

The tangent space at a point is a linear approximation to a curved object. The derivative becomes a map between tangent spaces.

23.2 Physical connection

General relativity uses differential geometry because spacetime is modeled as a manifold with curvature. The mathematics of curved spaces becomes physics when curvature tells matter how to move and matter tells curvature how to change.

24 Algebraic Geometry: Equations Become Shapes

Algebraic geometry begins with a simple observation:

> Polynomial equations define geometric sets.

For example:

$$x^2 + y^2 = 1$$

defines a circle in the plane.

$$y^2 = x^3 - x$$

defines an elliptic curve.

24.1 Affine varieties

Definition 20. Given polynomials $f_1, \ldots, f_m$ in variables $x_1, \ldots, x_n$ over a field k, their common zero set is

$$V(f_1, \ldots, f_m) = \{a \in k^n : f_i(a) = 0 \text{ for all } i\}.$$

This is a first version of an affine variety.

24.2 Ideals

Instead of studying one polynomial at a time, algebra studies collections of polynomials closed under addition and multiplication by arbitrary polynomials. These are ideals.

Definition 21. An ideal I in a ring R is a subset such that:

1. if $a, b \in I$, then $a + b \in I$,
2. if $r \in R$ and $a \in I$, then $ra \in I$.

Ideals are algebraic machines for encoding equations.

24.3 The bridge

Geometry asks about shapes. Algebra asks about rings and ideals. Algebraic geometry learns to translate between them.

That is why the path here matters. Sets, functions, relations, quotients, topology, rings, ideals: these are not random hurdles. They are the grammar of the later language.

25 How to Read a Definition

When you meet a definition, do not only underline it. Work it.

25.1 The five-part definition routine

For every important definition, write:

1. the formal definition,
2. a plain-English version,
3. two examples,
4. one near miss,
5. one theorem where the definition is used.

25.2 Example: injective

Formal:

$$f : A \to B \text{ is injective if } f(a_1) = f(a_2) \Rightarrow a_1 = a_2.$$

Plain English: the function does not collapse two different inputs to one output.

Examples:

$$f : \mathbb{R} \to \mathbb{R}, \quad f(x) = 2x + 1.$$

$$g : \mathbb{N} \to \mathbb{N}, \quad g(n) = n + 10.$$

Near miss:

$$h : \mathbb{R} \to \mathbb{R}, \quad h(x) = x^2.$$

Theorem:

If f and g are injective, then $g \circ f$ is injective.

Focus

Definitions become memory when you can produce examples and non-examples without looking.

26 How to Read a Theorem

A theorem should be interrogated before it is proved.

Take:

> If $f : A \to B$ and $g : B \to C$ are injective, then $g \circ f : A \to C$ is injective.

Ask:

1. What are the objects? Sets A, B, C and functions f, g.
2. What are the assumptions? f and g are injective.
3. What must be shown? $g \circ f$ is injective.
4. What does the conclusion mean by definition? If $(g \circ f)(a_1) = (g \circ f)(a_2)$, then $a_1 = a_2$.
5. Which assumptions can help? Injectivity of g first, then injectivity of f.

26.1 The theorem-reading template

```
Objects:
Hypotheses:
Conclusion:
Definitions involved:
Likely proof strategy:
Possible counterexample if an assumption is removed:
```

This template slows you down in the right place.

27 How to Write a Proof

A proof is a piece of communication. It should show why the claim follows.

27.1 Proof skeleton

Before writing full sentences, write the skeleton.

For a universal implication:

> Let [object] be arbitrary. Suppose [hypothesis]. We must show [conclusion]. By definition, [unpack]. Therefore [derive]. Hence [conclusion].

For set equality:

> We prove both inclusions. First let $x \in$ left side. ... Thus left $\subseteq$ right. Now let $x \in$ right side. ... Thus right $\subseteq$ left.

For existence:

> Choose [witness]. Verify [property]. Therefore such an object exists.

27.2 Sentences to avoid

Avoid:

- “Clearly”
- “Obviously”
- “It is easy to see”
- “By math”

These are sometimes harmless in expert writing, but early on they often hide the one step that needs attention.

27.3 A better replacement

Instead of:

> Clearly, $a + b$ is even.

Write:

> Since $a + b = 2(m + n)$ and $m + n \in \mathbb{Z}$, the number $a + b$ is even by definition.

28 Proof Debugging

Proof debugging is the habit that turns confusion into progress.

28.1 The first broken line

When a proof is wrong, do not rewrite the whole thing immediately. Find the first line that does not follow.

28.2 Gap types

Gap	What it looks like	Repair
Definition gap	A term is used but not unpacked	Write the definition
Quantifier gap	A variable appears without scope	Introduce it with domain and role
Existence gap	An object is claimed without witness	Construct or cite existence
Case gap	One possibility is handled	List all cases
Implication gap	A conclusion jumps too far	Insert a lemma or algebra step
Converse error	$P \Rightarrow Q$ treated as $Q \Rightarrow P$	Check direction

28.3 Bad proof, repaired

Claim: If n^2 is even, then n is even.

Bad proof:

> Since n^2 is even, $n^2 = 2k$. Taking square roots, $n = \sqrt{2k}$, so n is even.

Problem: being $\sqrt{2k}$ does not mean being twice an integer. The proof has an implication gap.

Repair: use contrapositive, as shown earlier.

29 Exercises for Walking

These are not calculation exercises. They are thinking exercises for when you are away from paper.

1. What does it mean to prove a statement for an arbitrary object?
2. Why does one counterexample disprove a universal claim?
3. What is the difference between a function's codomain and image?
4. Why does injective mean "no merging"?
5. Why does surjective mean "full coverage"?
6. What is a quotient trying to forget?
7. Why is induction not circular?
8. In an epsilon proof, who chooses ε?
9. Why does topology care about inverse images?
10. What does algebra preserve when it stops caring about particular numbers?

Try answering in full sentences. If you cannot explain an idea while walking, you probably know its symbol better than its meaning.

30 A Four-Week Beginner-to-Velleman Launch

Week	Theme	Work	Output
1	Sentences and quantifiers	Statements, predicates, domains, $\forall$, $\exists$, negation	30 translations and 20 negations
2	Definitions and direct proof	Even, odd, divides, subset, direct proof	10 short proofs from definitions
3	Sets and functions	Subsets, equality, injective, surjective, inverse image	6 set proofs and 4 function proofs
4	Relations and induction	Equivalence relations, modular arithmetic, induction	3 relation proofs and 3 induction proofs

30.1 Daily rhythm

1. Read 3 to 5 pages.
2. Copy one definition by hand.
3. Make two examples and one non-example.
4. Prove one small claim.
5. Debug yesterday's proof.
6. Review five spaced repetition prompts.

30.2 When to start Velleman

Start Velleman when you can:

- negate nested quantifiers without panic,
- prove simple parity and divisibility facts,
- prove set equality by two inclusions,
- prove injectivity statements from the definition,
- write an induction proof with all parts labeled.

31 Milestone Proof Portfolio

Before moving deeply into Velleman, write these ten proofs cleanly.

1. The sum of two even integers is even.
2. The product of two odd integers is odd.
3. If $a \mid b$ and $b \mid c$, then $a \mid c$.
4. If n^2 is even, then n is even.
5. $\sqrt{2}$ is irrational.
6. $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$.
7. $f^{-1}(S \cap T) = f^{-1}(S) \cap f^{-1}(T)$.
8. The composition of injective functions is injective.
9. Congruence modulo n is an equivalence relation.
10. $1 + 2 + \cdots + n = n(n+1)/2$ for all $n \geq 1$.

For each proof, include:

- theorem statement,
- definitions used,
- proof strategy,
- final proof,
- one sentence explaining the key idea.

Focus

This portfolio is the bridge. If you can write these proofs cleanly, Velleman's first chapters will feel like organized terrain rather than fog.

32 Closing: The Shape of the Road

The road from engineering mathematics to pure mathematics is not a rejection of computation. It is a widening of what counts as understanding.

At first, proof feels slow because every little word matters. Then the slowness becomes sight. You begin to notice why a theorem needs each hypothesis. You begin to hear the difference between "there exists" and "for every." You begin to see definitions as tools, examples as tests, counterexamples as boundary stones, and proofs as honest explanations.

From here, Velleman teaches the craft more systematically. Real analysis will sharpen your relationship with infinity and approximation. Abstract algebra will train you to recognize structure. Topology will separate continuity from measurement. Differential geometry will bring calculus onto curved spaces. Algebraic geometry will let equations and shapes speak to each other.

The beginning is humble: parse the sentence. Find the objects. Open the definitions. Move one justified step at a time.

That is not a small beginning. It is the beginning of doing mathematics.

A Appendix A: Notation You Should Be Able to Read

Notation	Meaning	What to ask yourself
$x \in A$	x is an element of A	What kind of objects live in A?
$A \subseteq B$	Every element of A is in B	How would I prove this with an arbitrary x?
$A \cup B$	Union	Is membership "or"?
$A \cap B$	Intersection	Is membership "and"?
$A \setminus B$	Difference	Is x in A but not in B?
$\mathcal{P}(A)$	Power set	What are the subsets of A?
$f : A \to B$	Function from A to B	What are domain and codomain?
$f(S)$	Image of S	Which outputs come from inputs in S?
$f^{-1}(T)$	Inverse image of T	Which inputs land in T?
$\forall$	For every	Who chooses the object?
$\exists$	There exists	What witness proves it?
$P \Rightarrow Q$	If P, then Q	What is assumed and what is owed?
$P \Leftrightarrow Q$	If and only if	Can I prove both directions?
$a \mid b$	a divides b	Can I write $b = ak$?
$a \equiv b \pmod{n}$	Same remainder mod n	Is $n \mid (a - b)$?
ϵ, δ	small positive tolerances	Which one is chosen first?

A.1 Notation drills

Translate each into English.

1. $\forall x \in A,\ x \in B$.
2. $\exists x \in A$ such that $x \notin B$.
3. $f^{-1}(S \cup T) = f^{-1}(S) \cup f^{-1}(T)$.
4. $\forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n \geq N,\ |a_n - L| < \varepsilon$.
5. $\forall a, b \in G,\ \phi(ab) = \phi(a)\phi(b)$.

B Appendix B: A Short History of Why Rigor Arrived

Mathematics did not begin as a perfectly formal machine. It began with counting, measuring, astronomy, land, trade, architecture, music, and motion. Rigor arrived in waves whenever intuition became too powerful to trust without guardrails.

B.1 Euclid and the axiomatic dream

Euclid's *Elements* organized geometry around definitions, postulates, and propositions. It was not modern formalism, but it established a model: start from agreed assumptions and reason carefully. A proof became a public object, something another mind could inspect.

B.2 Calculus and the problem of motion

Newton and Leibniz gave mathematics a language for change. Calculus worked spectacularly. It explained falling bodies, planetary motion, tangents, areas, and rates. But early calculus leaned on infinitesimals and limiting arguments that were powerful but not always clean.

The results were too useful to abandon, so mathematicians sharpened the foundations.

B.3 Cauchy and Weierstrass

The nineteenth century rebuilt calculus into analysis. Limits became epsilon-delta statements. Continuity became a quantified contract. Infinite series were no longer trusted just because they looked well-behaved.

That is why your work with quantifiers matters. The formal definition of a limit is not a hazing ritual. It is the historical repair of a brilliant machine.

B.4 Cantor and infinity

Cantor showed that infinity has sizes. The natural numbers, integers, and rationals are countable. The real numbers are not. This was shocking because it made infinity into something that could be compared, classified, and proved about.

B.5 Noether and structure

Emmy Noether helped push algebra away from individual equations and toward structures: groups, rings, fields, ideals. This shift is one reason modern mathematics feels abstract. Used well, abstraction compresses many concrete calculations into a single theorem that covers them all at once.

B.6 Hilbert and formal clarity

Hilbert championed axiomatic thinking. The idea was not to remove meaning from mathematics, but to clarify exactly what assumptions were being used. This habit still shapes modern proof writing.

B.7 Grothendieck and the expansion of geometry

Grothendieck transformed algebraic geometry by changing the level of abstraction. Instead of studying only solution sets of polynomials over familiar fields, he built a language where geometry could be studied through rings, schemes, and maps. That world is far away from this reader, but the early tools are already here: sets, functions, equivalence, quotients, topology, and algebra.

Focus

Rigor usually appears after intuition succeeds. First people find a powerful idea. Then they discover the idea needs a stronger language.

C Appendix C: Extra Proof Drills

C.1 Drill block 1: Definitions only

Prove each using only definitions and basic algebra.

1. If n is even, then n^2 is even.
2. If m is even and n is odd, then mn is even.
3. If $a \mid b$ and $a \mid c$, then $a \mid (b + c)$.
4. If $a \mid b$, then $a \mid bc$ for every $c \in \mathbb{Z}$.
5. If a and b are rational, then $a + b$ is rational.
6. If a and b are rational, then ab is rational.

C.2 Drill block 2: Counterexamples

Disprove each false statement.

1. If $a \mid bc$, then $a \mid b$ or $a \mid c$.
2. If $x < y$, then $x^2 < y^2$.
3. Every injective function is surjective.
4. Every surjective function is injective.
5. If $A \subseteq B \cup C$, then $A \subseteq B$ or $A \subseteq C$.
6. If $A \cap B = A \cap C$, then $B = C$.

C.3 Drill block 3: Proof strategy choice

For each theorem, choose a likely strategy before proving it.

1. If n^2 is odd, then n is odd.
2. There is no smallest positive real number.
3. The intersection of two subgroups of a group is a subgroup.
4. If $A \subseteq B$, then $\mathcal{P}(A) \subseteq \mathcal{P}(B)$.
5. If $f : A \to B$ is injective and $S, T \subseteq A$, then $f(S \cap T) = f(S) \cap f(T)$.

D Appendix D: Hints Without Full Solutions

D.1 Definitions only

1. For n^2 even: write $n = 2k$, then square.
2. For even times odd: write $m = 2a$ and $n = 2b + 1$.
3. For divisibility over sums: write $b = ak$ and $c = a\ell$.
4. For $a \mid bc$: start from $b = ak$.
5. For rational sums: write $a = p/q$ and $b = r/s$ with nonzero denominators.
6. For rational products: multiply numerators and denominators.

D.2 Counterexamples

1. Try $a = 6$, $b = 2$, $c = 3$.
2. Try negative numbers.
3. Think of $f : \mathbb{N} \to \mathbb{N}$ given by $f(n) = n + 1$.
4. Think of a map from a two-element set to a one-element set.
5. Let A contain two points, one in B and one in C.
6. Make A empty.

D.3 Strategy choice

1. Contrapositive is clean.
2. Contradiction is natural: assume a smallest positive real exists and halve it.
3. Direct proof from subgroup definition.
4. Let $S \in \mathcal{P}(A)$ and prove $S \in \mathcal{P}(B)$.
5. Prove both inclusions; injectivity is needed for the harder direction.

E Appendix E: Reading Velleman Without Getting Lost

When you open Velleman, do not read it like a novel and do not read it like a formula sheet. Read it like a manual for a new instrument.

E.1 Before each section

1. Skim the section headings.
2. Write down the symbols you expect to see.
3. Predict what kind of proof method may appear.

E.2 During each section

1. Copy every definition that seems central.
2. After each example proof, write the strategy in one line.
3. Mark any sentence where a variable is introduced.
4. Pause at every "therefore" and ask what justifies it.

E.3 After each section

1. Rewrite one proof from memory.
2. Make one example and one non-example for each definition.
3. Do at least one exercise badly before looking for help.
4. Debug the bad attempt before writing a polished version.

E.4 What "understanding" should feel like

Understanding does not mean every proof feels obvious. It means you can locate the difficulty. You can say: "I understand the definitions, but I do not see why this witness works," or "I see the strategy, but the induction step has an algebra gap."

That is progress. Fog becomes a map when you can name where you are.

F Appendix F: The One-Page Proof Debugger

Use this every time a proof feels shaky.

1. State the claim exactly.
2. Circle all quantifiers.
3. List the objects and their domains.
4. Separate hypotheses from conclusion.
5. Expand every definition in the conclusion.
6. Expand useful definitions in the hypotheses.
7. Choose a strategy.
8. Write a skeleton.
9. Fill in one line at a time.
10. For each line, ask: why does this follow?
11. Check that arbitrary objects were really arbitrary.
12. Check that witnesses were actually constructed.
13. Check all cases.
14. Remove “clearly” unless the next sentence explains why.
15. Read the final proof aloud.

Focus

The proof debugger is not remedial. Professionals debug proofs too. The difference is that they have better names for the failure modes.

G Appendix G: Final Readiness Exam

Try this without notes. It is not a speed test.

1. Define statement, predicate, theorem, proof, counterexample.
2. Negate: $\forall x \in A, \exists y \in B,\ R(x, y)$.
3. Prove that if n is odd, then n^2 is odd.
4. Prove that if $a \mid b$ and $a \mid c$, then $a \mid (b - c)$.
5. Prove $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$.
6. Give a counterexample to: if $f(S \cap T) = f(S) \cap f(T)$, then f is injective.
7. Prove that if f and g are surjective, then $g \circ f$ is surjective.
8. Prove congruence modulo n is transitive.
9. Prove by induction that $2^n \geq n + 1$ for all $n \geq 1$.
10. Explain, in words, why $\mathbb{Q}$ is countable but $\mathbb{R}$ is not.

G.1 How to score yourself

- If you cannot start: review definitions.
- If you start but drift: review quantifiers and proof skeletons.
- If your proof has gaps: use the debugger.
- If your proof is correct but ugly: good. Polish comes after correctness.