

Notes on “Finite-Dimensional Vector Spaces”

by Paul R. Halmos

October 10, 2022

Each \section corresponds to the scope of one member’s assignment, and each \subsection corresponds to one theorem or exercise in the textbook, specified in the format $m.n$ where m is the section number and n is the theorem/exercise number. If n is not given, we use $n = 1$ instead.

1 Toga (2022/09/19)

1.1 Exercise 1.1

- (a) Since addition is commutative, $0 + \alpha = \alpha + 0$ holds. We also have $\alpha + 0 = \alpha$ by definition, hence $0 + \alpha = \alpha$.

2 Mohehe (2022/09/27)

2.1 Exercise 1.1

- (b) If $\alpha + \beta = \alpha + \gamma$, we have $\beta = \beta + 0 = 0 + \beta = (\alpha + (-\alpha)) + \beta = ((-\alpha) + \alpha) + \beta = (-\alpha) + (\alpha + \beta) = (-\alpha) + (\alpha + \gamma) = ((-\alpha) + \alpha) + \gamma = (\alpha + (-\alpha)) + \gamma = 0 + \gamma = \gamma + 0 = \gamma$ by definition. Therefore, $\beta = \gamma$ holds.
- (c) We have $\alpha + (\beta - \alpha) = \alpha + (\beta + (-\alpha)) = \alpha + ((-\alpha) + \beta) = (\alpha + (-\alpha)) + \beta = 0 + \beta = \beta + 0 = \beta$ by definition. Therefore, $\alpha + (\beta - \alpha) = \beta$ holds.
- (d) We have $\alpha 0 + \alpha 0 = \alpha(0 + 0) = \alpha 0 = \alpha + 0$ by definition, hence $\alpha 0 = 0$ by Exercise 1(b). We also have $\alpha \cdot 0 = 0 \cdot \alpha$ by definition. Therefore, $\alpha \cdot 0 = 0 \cdot \alpha = 0$.
- (e) We have $\alpha + (-1)\alpha = 1\alpha + (-1)\alpha = (1 + (-1))\alpha = 0\alpha = 0$ by definition and Exercise 1(d). Since the additive inverse is unique, we obtain $(-1)\alpha = -\alpha$.
- (f) We have $(-\alpha)(-\beta) = ((-1)\alpha)((-1)\beta) = (\alpha(-1))((-1)\beta) = \alpha((-1)((-1)\beta)) = \alpha((-1)(-1)\beta)$ by Exercise 1(e) and definition. We also have $(-1)(-1) = 0 + (-1)(-1) = (1 + (-1)) + (-1)(-1) = 1 + (-1) + (-1)(-1) = 1 +$

$(-1)((-1) + 1) = 1 + (-1)(1 + (-1)) = 1 + (-1)0 = 1 + 0 = 1$ by definition. By it and definition, $\alpha((-1)(-1)\beta) = \alpha(1\beta) = \alpha(\beta 1) = \alpha\beta$ holds. Therefore, $(-\alpha)(-\beta) = \alpha\beta$ holds.

- (g) If $\alpha\beta = 0$, suppose $\alpha \neq 0$ and $\beta \neq 0$ hold. By supposition and definition, we have $0 = \alpha^{-1}0 = \alpha^{-1}(\alpha\beta) = (\alpha^{-1}\alpha)\beta = (\alpha\alpha^{-1})\beta = 1\beta = \beta 1 = \beta$, hence $\beta = 0$. However, this result contradicts supposition, " $\alpha \neq 0$ and $\beta \neq 0$ ". Therefore, if $\alpha\beta = 0$, then either $\alpha = 0$ or $\beta = 0$ (or both).

3 Joh (2022/09/19)

3.1 Exercise 1.2

- (a) The set of positive integers is not a field since there is no additive inverse for 1.
- (b) The set of integers is not a field since there is no multiplicative inverse for 2.
- (c) There exists a bijective map φ from \mathbb{N} (or \mathbb{Z}) to \mathbb{Q} [1], where \mathbb{Q} is a field [2]. We can make \mathbb{N} a field by re-defining (i) addition by $a \oplus b = \varphi^{-1}(\varphi(a) + \varphi(b))$ and (ii) multiplication by $a \otimes b = \varphi^{-1}(\varphi(a)\varphi(b))$ for each $a, b \in \mathbb{N}$. Note that the additive and multiplicative identities become $\varphi^{-1}(0)$ and $\varphi^{-1}(1)$, respectively. For each $\alpha \in \mathbb{N}$, the additive inverse becomes $\varphi^{-1}(-\varphi(\alpha))$, and the multiplicative inverse becomes $\varphi^{-1}(1/\varphi(\alpha))$ if $\alpha \neq \varphi^{-1}(0)$.

Let $\alpha, \beta, \gamma, \alpha', \beta' \in \mathbb{N}$. Note that

- 1) $\alpha \oplus \beta = \varphi^{-1}(\varphi(\alpha) + \varphi(\beta)) = \varphi^{-1}(\varphi(\beta) + \varphi(\alpha)) = \beta \oplus \alpha$ holds. (addition is commutative)
(from here, mohehe)
- 2) $\alpha \oplus (\beta \oplus \gamma) = \alpha \oplus (\varphi^{-1}(\varphi(\beta) + \varphi(\gamma))) = \varphi^{-1}(\varphi(\alpha) + \varphi(\varphi^{-1}(\varphi(\beta) + \varphi(\gamma)))) = \varphi^{-1}(\varphi(\alpha) + (\varphi(\beta) + \varphi(\gamma))) = \varphi^{-1}((\varphi(\alpha) + \varphi(\beta)) + \varphi(\gamma)) = \varphi^{-1}(\varphi(\varphi^{-1}(\varphi(\alpha) + \varphi(\beta))) + \varphi(\gamma)) = \varphi^{-1}(\varphi(\alpha) + \varphi(\beta)) \oplus \gamma = (\alpha \oplus \beta) \oplus \gamma$ holds. (addition is associative)
- 3) $\alpha \oplus \varphi^{-1}(0) = \varphi^{-1}(\varphi(\alpha) + \varphi(\varphi^{-1}(0))) = \varphi^{-1}(\varphi(\alpha) + 0) = \varphi^{-1}(\varphi(\alpha)) = \alpha$ holds. (there exists additive identity, $\varphi^{-1}(0)$) If α' and β' are additive identity, we have $\alpha' = \alpha' \oplus \beta' = \beta' \oplus \alpha' = \beta'$ by 1) and the definition of additive identity. (additive identity is unique)
- 4) $-\varphi(\alpha) \in \mathbb{Q}$ holds by definition, so $\varphi^{-1}(-\varphi(\alpha)) \in \mathbb{N}$ holds. Therefore, $\alpha \oplus \varphi^{-1}(-\varphi(\alpha)) = \varphi^{-1}(\varphi(\alpha) + \varphi(\varphi^{-1}(-\varphi(\alpha)))) = \varphi^{-1}(\varphi(\alpha) + (-\varphi(\alpha))) = \varphi^{-1}(0)$ holds. (for each α ($\alpha \in \mathbb{N}$), there exists additive inverse) For each α , if α' and β' are additive inverse, we have $\alpha' = \alpha' \oplus \varphi^{-1}(0) = \alpha' \oplus (\alpha \oplus \beta') = (\alpha' \oplus \alpha) \oplus \beta' = (\alpha \oplus \alpha') \oplus \beta' = \varphi^{-1}(0) \oplus \beta' = \beta \oplus \varphi^{-1}(0) = \beta'$ by 1), 2), 3) and the definition of additive inverse. (additive inverse is unique)

- 5) $\alpha \otimes \beta = \varphi^{-1}(\varphi(\alpha)\varphi(\beta)) = \varphi^{-1}(\varphi(\beta)\varphi(\alpha)) = \beta \otimes \alpha$ holds.(multiplication is commutative)
- 6) $\alpha \otimes (\beta \otimes \gamma) = \alpha \otimes (\varphi^{-1}(\varphi(\beta)\varphi(\gamma))) = \varphi^{-1}(\varphi(\alpha)\varphi(\varphi^{-1}(\varphi(\beta)\varphi(\gamma)))) = \varphi^{-1}(\varphi(\alpha)\varphi(\beta)\varphi(\gamma)) = \varphi^{-1}(\varphi(\varphi^{-1}(\varphi(\alpha)\varphi(\beta)))\varphi(\gamma)) = \varphi^{-1}(\varphi(\alpha)\varphi(\beta)) \otimes \gamma = (\alpha \otimes \beta) \otimes \gamma$ holds.(multiplication is associative)
- 7) $\alpha \otimes \varphi^{-1}(1) = \varphi^{-1}(\varphi(\alpha)\varphi(\varphi^{-1}(1))) = \varphi^{-1}(\varphi(\alpha) \cdot 1) = \alpha$ holds.(there exists additive identity, $\varphi^{-1}(1)$) If α' and β' are additive identity, we have $\alpha' = \alpha' \otimes \beta' = \beta' \otimes \alpha' = \beta'$ by 5) and definition of multiplicative identity.(multiplicative identity is unique)
- 8) For each α ($\alpha \neq \varphi^{-1}(0)$), $(1/\varphi(\alpha)) \in \mathbb{Q}$ holds by definition, so $\varphi^{-1}(1/\varphi(\alpha)) \in \mathbb{N}$ holds. Therefore, $\alpha \otimes \varphi^{-1}(1/\varphi(\alpha)) = \varphi^{-1}(\varphi(\alpha)\varphi(\varphi^{-1}(1/\varphi(\alpha)))) = \varphi^{-1}(\varphi(\alpha)(1/\varphi(\alpha))) = \varphi^{-1}(1)$ holds.(for each α ($\alpha \in \mathbb{N}$), there exists multiplicative inverse) For each α ($\alpha \neq \varphi^{-1}(0)$), if α' and β' are multiplicative inverse, we have $\alpha' = \alpha' \otimes \varphi^{-1}(1) = \alpha' \otimes (\alpha \otimes \beta') = (\alpha' \otimes \alpha) \otimes \beta' = (\alpha \otimes \alpha') \otimes \beta' = \varphi^{-1}(1) \otimes \beta' = \beta' \otimes \varphi^{-1}(1) = \beta'$ by 5), 6), 7) and the definition of multiplicative inverse.(multiplicative inverse is unique)
- 9) $\alpha \otimes (\beta \oplus \gamma) = \alpha \otimes (\varphi^{-1}(\varphi(\beta) + \varphi(\gamma))) = \varphi^{-1}(\varphi(\alpha)\varphi(\varphi^{-1}(\varphi(\beta) + \varphi(\gamma)))) = \varphi^{-1}(\varphi(\alpha)(\varphi(\beta) + \varphi(\gamma))) = \varphi^{-1}(\varphi(\alpha)\varphi(\beta) + \varphi(\alpha)\varphi(\gamma)) = \varphi^{-1}(\varphi(\varphi^{-1}(\varphi(\alpha)\varphi(\beta))) + \varphi(\varphi^{-1}(\varphi(\alpha)\varphi(\gamma)))) = \varphi^{-1}(\varphi(\alpha \otimes \beta) + \varphi(\alpha \otimes \gamma)) = \alpha \otimes \beta \oplus \alpha \otimes \gamma$ holds.(distributive law stands)

4 Mohehe (2022/10/8)

4.1 Exercise 1.3

For two integers a and b , we denote by $a \% b$ the remainder after dividing a by b , and write $b \mid a$ if and only if $a \% b = 0$. For clarity, we denote the ordinary sum and product of two integers a and b by $a +_{\mathbb{Z}} b$ and $a \cdot_{\mathbb{Z}} b$, respectively. Note that $\alpha + \beta = (\alpha +_{\mathbb{Z}} \beta) \% m$ and $\alpha\beta = (\alpha \cdot_{\mathbb{Z}} \beta) \% m$ for $\alpha, \beta \in \mathbb{Z}_m$.

(a) Let $\alpha, \beta, \gamma \in \mathbb{Z}_m$, $k \in \mathbb{Z}$

1' Proof : if m is a prime, \mathbb{Z}_m is a field.

Suppose m is a prime,

- 1) $\alpha + \beta = (\alpha +_{\mathbb{Z}} \beta) \% m = (\beta +_{\mathbb{Z}} \alpha) \% m = \beta + \alpha$ (addition is commutative)
- 2) Since $\alpha +_{\mathbb{Z}} (\beta + \gamma) = \alpha +_{\mathbb{Z}} (\beta +_{\mathbb{Z}} \gamma) \% m \equiv \alpha +_{\mathbb{Z}} (\beta +_{\mathbb{Z}} \gamma) = (\alpha +_{\mathbb{Z}} \beta) +_{\mathbb{Z}} \gamma \equiv (\alpha +_{\mathbb{Z}} \beta) \% m +_{\mathbb{Z}} \gamma = (\alpha + \beta) +_{\mathbb{Z}} \gamma \pmod{m}$ holds, $\alpha + (\beta + \gamma) = (\alpha +_{\mathbb{Z}} (\beta + \gamma)) \% m = ((\alpha + \beta) +_{\mathbb{Z}} \gamma) \% m = (\alpha + \beta) + \gamma$ holds.(addition is associative)
- 3) $\alpha + 0 = (\alpha +_{\mathbb{Z}} 0) \% m = \alpha \% m = \alpha$ (there exists additive identity)
By it and 1), if β and γ are additive identity, $\beta = \beta + \gamma = \gamma + \beta = \gamma$ (additive identity is unique)

- 4) If $\alpha +_{\mathbb{Z}} \beta = m$, $\alpha + \beta = (\alpha +_{\mathbb{Z}} \beta) \% m = m \% m = 0$ (there exists additive inverse)
- 5) $\alpha\beta = (\alpha \cdot_{\mathbb{Z}} \beta) \% m = (\beta \cdot_{\mathbb{Z}} \alpha) \% m = \beta\alpha$ (multiplication is commutative)
- 6) Since $\alpha \cdot_{\mathbb{Z}} (\beta\gamma) = \alpha \cdot_{\mathbb{Z}} ((\beta \cdot_{\mathbb{Z}} \gamma) \% m) \equiv \alpha \cdot_{\mathbb{Z}} (\beta \cdot_{\mathbb{Z}} \gamma) = (\alpha \cdot_{\mathbb{Z}} \beta) \cdot_{\mathbb{Z}} \gamma \equiv ((\alpha \cdot_{\mathbb{Z}} \beta) \% m) \cdot_{\mathbb{Z}} \gamma = (\alpha\beta) \cdot_{\mathbb{Z}} \gamma \pmod{m}$ holds, $\alpha(\beta\gamma) = (\alpha \cdot_{\mathbb{Z}} (\beta\gamma)) \% m = ((\alpha\beta) \cdot_{\mathbb{Z}} \gamma) \% m = (\alpha\beta)\gamma$ holds. (multiplication is associative)
- 7) $\alpha 1 = (\alpha \cdot_{\mathbb{Z}} 1) \% m = \alpha \% m = \alpha$ (there exists multiplicative identity)
By it and 5), if β and γ are multiplicative identity, $\beta = \beta\gamma = \gamma\beta = \gamma$ (multiplicative identity is unique)
- 8) For all $\alpha (\alpha \neq 0)$, suppose there doesn't exist β that makes $\alpha\beta = 1$.
There exist $\beta, \gamma \in \mathbb{Z}_m$ with $\beta \neq \gamma$ and $\alpha\beta = \alpha\gamma$, because β is any one from 0 to $m-1$ and $\alpha\beta$ is any one from 0 to $m-1$ except 1.
Therefore, $(\alpha \cdot_{\mathbb{Z}} \beta +_{\mathbb{Z}} (-\alpha \cdot_{\mathbb{Z}} \gamma)) = \alpha \cdot_{\mathbb{Z}} (\beta +_{\mathbb{Z}} (-\gamma)) = km$ holds.
The right side has divisor m , but it contradicts that the left side doesn't have divisor of m except 1, because $0 < \alpha < (m-1)$ and $((-m) < (\beta +_{\mathbb{Z}} (-\gamma)) < 0$ or $0 < (\beta +_{\mathbb{Z}} (-\gamma)) < m$) holds.
Thus, there exists β that makes $\alpha\beta = 1$. (there exists multiplicative inverse)
- 9) $\alpha(\beta + \gamma) = (\alpha \cdot_{\mathbb{Z}} (\beta + \gamma)) \% m = (\alpha \cdot_{\mathbb{Z}} ((\beta +_{\mathbb{Z}} \gamma) \% m)) \% m \equiv (\alpha \cdot_{\mathbb{Z}} (\beta +_{\mathbb{Z}} \gamma)) \% m = (\alpha \cdot_{\mathbb{Z}} \beta + \alpha \cdot_{\mathbb{Z}} \gamma) \% m \equiv ((\alpha \cdot_{\mathbb{Z}} \beta) \% m +_{\mathbb{Z}} (\alpha \cdot_{\mathbb{Z}} \gamma) \% m) \% m \equiv (\alpha \cdot_{\mathbb{Z}} \beta) \% m + (\alpha \cdot_{\mathbb{Z}} \gamma) \% m \equiv \alpha\beta + \alpha\gamma$ holds. (distributive law stands)

In conclusion, if m is a prime, \mathbb{Z}_m is a field.

2' Proof : If \mathbb{Z}_m is a field, m is a prime.

By contraposition, it is equivalent to prove "If m is not a prime, \mathbb{Z}_m is not a field." We can show 1) to 7) and 9) in the same way as 1'. For each m , suppose there exist α and β that make $\alpha\beta = 1$. m is not a prime, so let p be one of prime factors of m and then we have $m = p \cdot_{\mathbb{Z}} p' (p' \in \mathbb{Z} \text{ and } 1 < p' < m)$

If $\alpha = p$, by $\alpha\beta = 1$ and $m = p \cdot_{\mathbb{Z}} p'$, we have $\alpha \cdot_{\mathbb{Z}} \beta = k \cdot_{\mathbb{Z}} m + 1 (k \in \mathbb{Z}) \Leftrightarrow p \cdot_{\mathbb{Z}} \beta = k \cdot_{\mathbb{Z}} p \cdot_{\mathbb{Z}} p' +_{\mathbb{Z}} 1 \Leftrightarrow (\beta +_{\mathbb{Z}} (-k \cdot_{\mathbb{Z}} p')) \cdot_{\mathbb{Z}} p = 1$. The right side is 1 but the left one is not 1 because of $1 < p$ and $\beta +_{\mathbb{Z}} (-k \cdot_{\mathbb{Z}} p') \in \mathbb{Z}$. Therefore It is contradicted. For each m , there doesn't exist α and β that make $\alpha\beta = 1$. In conclusion, "If m is not a prime, \mathbb{Z}_m is not a field." and "If \mathbb{Z}_m is a field, m is a prime."

Because of 1' and 2', \mathbb{Z}_m is a field if and only if m is a prime.

(b) 4

(c) 5

4.2 Exercise 1.4

Define $\alpha_0, \alpha_1, \dots, \in \mathfrak{F}$ as below:

$$\alpha_0 = 0, \tag{1}$$

$$\alpha_n = \alpha_{n-1} + 1 \quad (n > 0). \tag{2}$$

We have $\alpha_m \alpha_n = \alpha_{mn}$ for all m and n .

It is proved by the following.

$$\begin{aligned} \bullet \text{ We have } \alpha_n &= 0 + \overbrace{1 + 1 + \dots + 1}^{n \text{ terms}} = \overbrace{1 + 1 + \dots + 1}^{n \text{ terms}}. \text{ Therefore, } \alpha_m \alpha_n = \\ &\alpha_m (\overbrace{1 + 1 + \dots + 1}^{n \text{ terms}}) = \overbrace{\alpha_m + \alpha_m + \dots + \alpha_m}^{n \text{ terms}} = \\ &\underbrace{(1 + 1 + \dots + 1) + (1 + 1 + \dots + 1) + \dots + (1 + 1 + \dots + 1)}_{\alpha_{mn}} = \overbrace{1 + 1 + \dots + 1}^{mn \text{ terms}} = \end{aligned}$$

Assume there exists an n with $\alpha_n = 0$ but $\alpha_k \neq 0$ for any $k < n$. It suffices to prove that n is a prime.

References

[1] https://proofwiki.org/wiki/Rational_Numbers_are_Countably_Infinite

[2] https://proofwiki.org/wiki/Rational_Numbers_form_Field