

Fraction	Dates (Ma)				Composition					Isotopic Ratios										Fraction
	²⁰⁶ Pb/ ²³⁸ U ^a	±σ abs	²⁰⁷ Pb/ ²³⁵ U ^a	±σ abs	²⁰⁷ Pb/ ²⁰⁶ Pb ^g	±σ abs	Corr. coef.	% disc ^b	Th/ U ^c	Pb* (pg) ^d	Pbc (pg) ^e	Pb*/ Pbc ^f	²⁰⁶ Pb/ ²⁰⁴ Pb ^g	²⁰⁶ Pb/ ²³⁸ U ^h	±σ	²⁰⁷ Pb/ ²³⁵ U ^h	±σ	²⁰⁷ Pb/ ²⁰⁶ Pb ^g	±σ	
zircon																				
z1	70.568	0.038	70.42	0.20	65.4	6.5	0.525	-7.95	0.08	20.1	0.49	41	2793	0.0110070	0.054	0.07181	0.30	0.04732	0.27	z1
z2	70.556	0.036	70.74	0.14	76.9	4.4	0.577	8.30	0.31	42.0	0.66	64	4045	0.0110051	0.051	0.07215	0.21	0.047550	0.19	z2
z3	70.544	0.034	70.54	0.11	70.5	3.3	0.565	-0.04	0.61	58.7	0.55	107	6238	0.0110033	0.048	0.07194	0.16	0.047422	0.14	z3
z4	70.562	0.036	70.53	0.13	69.6	4.0	0.594	-1.35	0.28	34.4	0.44	78	4950	0.0110060	0.051	0.07194	0.19	0.047404	0.17	z4
z5	70.566	0.038	70.48	0.23	67.7	7.2	0.547	-4.31	0.34	26.9	0.72	38	2363	0.0110067	0.054	0.07188	0.33	0.04736	0.30	z5
z6	70.524	0.039	70.60	0.24	73.3	7.8	0.491	3.78	0.47	19.3	0.50	39	2354	0.0110001	0.056	0.07201	0.35	0.04748	0.33	z6
z7	70.557	0.044	70.51	0.27	68.9	8.8	0.536	-2.40	0.36	13.4	0.40	34	2106	0.0110053	0.062	0.07191	0.40	0.04739	0.37	z7
z8	70.573	0.035	70.62	0.12	72.2	3.7	0.588	2.20	0.36	49.1	0.59	83	5171	0.0110077	0.049	0.07202	0.18	0.047454	0.16	z8
z9	70.556	0.035	70.64	0.13	73.6	3.8	0.663	4.11	0.83	44.7	0.42	105	5808	0.0110051	0.050	0.07205	0.19	0.047483	0.16	z9
z10	70.570	0.035	70.66	0.13	73.7	4.1	0.548	4.26	0.80	41.6	0.44	95	5257	0.0110073	0.049	0.07207	0.19	0.047485	0.17	z10
z11	70.559	0.036	70.61	0.14	72.2	4.4	0.485	2.34	0.78	39.3	0.44	90	5020	0.0110056	0.051	0.07201	0.20	0.047456	0.18	z11
z12	70.530	0.035	70.55	0.12	71.2	3.6	0.587	0.92	0.85	52.4	0.47	111	6075	0.0110011	0.049	0.07195	0.17	0.047435	0.15	z12
z13	70.538	0.034	70.58	0.11	72.0	3.4	0.548	2.06	0.59	68.5	0.68	100	5890	0.0110023	0.048	0.07198	0.16	0.047452	0.14	z13
z14	70.533	0.037	70.62	0.18	73.6	5.7	0.580	4.15	0.80	41.1	0.73	56	3133	0.0110016	0.052	0.07203	0.27	0.04748	0.24	z14

a Isotopic dates calculated using the decay constants ²³⁸ = 1.55125E-10 and ²³⁵ = 9.8485E-10 (Jaffey et al. 1971).

b % discordance = 100 - (100 * (²⁰⁶Pb/²³⁸U date) / (²⁰⁷Pb/²⁰⁶Pb date))

c Th contents calculated from radiogenic ²⁰⁸Pb and the ²⁰⁷Pb/²⁰⁶Pb date of the sample, assuming concordance between U-Th and Pb systems.

d Total mass of radiogenic Pb.

e Total mass of common Pb.

f Ratio of radiogenic Pb (including ²⁰⁸Pb) to common Pb.

g Measured ratio corrected for fractionation and spike contribution only.

h Measured ratios corrected for fractionation, tracer and blank.