

Galaxies and Extragalactic Astronomy

Active Galactic Nuclei

Mar Mezcua
mezcua@ice.csic.es
Office D3-18

Question


What are AGN?

Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies


Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies



Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies




TEST: WHO IS WHO?

- A) Star within the Milky Way
- B) Quasar:
9 billion light-years away
- C) Elliptical galaxy:
7 billion light-years away

Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies




TEST: WHO IS WHO?

- A) Star within the Milky Way
- B) Quasar:
9 billion light-years away
- C) Elliptical galaxy:
7 billion light-years away

Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies




TEST: WHO IS WHO?

- A) Star within the Milky Way
- B) Quasar:
9 billion light-years away
- C) Elliptical galaxy:
7 billion light-years away

Introduction

- Active Galactic Nuclei (AGN) are powerful sources of radiation residing at the centre of 1-10% of all galaxies
- Galaxies hosting an AGN are known as active galaxies



TEST: WHO IS WHO?

- A) Star within the Milky Way
- B) Quasar:
9 billion light-years away
- C) Elliptical galaxy:
7 billion light-years away

Introduction

- AGN are the most luminous long-lived objects in the Universe
- Huge span of observed AGN luminosities: $L \sim 10^{40} - 10^{47}$ erg/s
- The more luminous AGN outshine their host galaxies by factors > 1000

Introduction

- AGN are the most luminous long-lived objects in the Universe
- Huge span of observed AGN luminosities: $L \sim 10^{40} - 10^{47}$ erg/s
- The more luminous AGN outshine their host galaxies by factors > 1000

Quasar 3C273:

Magnitude = 12.8

Virgo constellation

Distance: ~2.5 billion light years




Historical discovery

INITIAL AIM: clarify the nature of “spiral nebulae”

- Fath (1908), Lick Observatory:
spectrum of NGC1068 ➔ emission lines as in
planetary nebulae
- Confirmed by Slipher (1917) at Lowell
- Hubble (1926) studied more systems
- Seyfert (1941,1943): galaxies whose
spectra show high-ionization emission lines,
forbidden emission lines, broad emission
lines from small, bright nucleus




Seyfert
galaxies



SPIRAL NEBULA *Messier 51 Canum Venaticorum*
Photographed with 60-Inch Reflector on Sead 23 plate, February 7 and 8, 1910
Exposure ab 25m. Enlargement from negative 6.4 diameter.


MS1 spiral nebula
Ritchey (1910)

Seyfert spectrum: high-ionization, forbidden, and broad emission lines




Caccianiga et al. (2014)

Seyfert spectrum: high-ionization, forbidden, and broad emission lines




Caccianiga et al. (2014)

Seyfert spectrum: high-ionization, forbidden, and broad emission lines



Caccianiga et al. (2014)

Seyfert spectrum: high-ionization, forbidden, and broad emission lines



Caccianiga et al. (2014)

Questions

Why are forbidden lines
called forbidden?


Questions

Why are forbidden lines
called forbidden?

Why are broad lines broad?

Historical discovery

- After II World War with the development of radioastronomy, radiogalaxies were discovered
- Quasars were discovered in 1963 when Maarten Schmidt interpreted the optical emission lines of the known radio source 2C273 as redshifted hydrogen Balmer lines.



Bill Keel, UAlabama

Question

What powers AGN?

Supermassive black holes: the AGN central engine


- Current paradigm: the fundamental power source of all AGN is accretion onto a supermassive black hole
- Presence of a supermassive black hole at the center of most galaxies predicted by galaxy formation models...

Supermassive black holes: the AGN central engine

- Current paradigm: the fundamental power source of all AGN is accretion onto a supermassive black hole
- Presence of a supermassive black hole at the center of most galaxies predicted by galaxy formation models...
...and observations!


Supermassive black holes: the AGN central engine

- Current paradigm: the fundamental power source of all AGN is accretion onto a supermassive black hole
 - Presence of a supermassive black hole at the center of most galaxies predicted by galaxy formation models...
...and observations!



Supermassive black holes: the AGN central engine

- Current paradigm: the fundamental power source of all AGN is accretion onto a supermassive black hole
- Presence of a supermassive black hole at the center of most galaxies predicted by galaxy formation models...
...and observations!



Supermassive black holes: the AGN central engine

Accretion onto the supermassive black hole

 significant quantities of gas are deposited onto the central parts of a galaxy and enter the region where the black hole mass dominates the gravitational potential

Supermassive black holes: the AGN central engine

Accretion onto the supermassive black hole



significant quantities of gas are deposited onto the central parts of a galaxy and enter the region where the black hole mass dominates the gravitational potential

Stochastic processes

- interstellar medium (ISM) of the host galaxy
- stellar mass loss associated with a nuclear star cluster
- non-axisymmetric potentials resulting from galactic bar formation

Supermassive black holes: the AGN central engine


Accretion onto the supermassive black hole



significant quantities of gas are deposited onto the central parts of a galaxy and enter the region where the black hole mass dominates the gravitational potential

Stochastic processes

- interstellar medium (ISM) of the host galaxy
- stellar mass loss associated with a nuclear star cluster
- non-axisymmetric potentials resulting from galactic bar formation



Supermassive black holes: the AGN central engine


Accretion onto the supermassive black hole



significant quantities of gas are deposited onto the central parts of a galaxy and enter the region where the black hole mass dominates the gravitational potential

Stochastic processes → Dominate late-type AGN host galaxies

- interstellar medium (ISM) of the host galaxy
- stellar mass loss associated with a nuclear star cluster
- non-axisymmetric potentials resulting from galactic bar formation



Supermassive black holes: the AGN central engine

Merger-driven processes

- non-axisymmetric potentials resulting from a nearby companion galaxy can further increase the mass flow rate onto the central regions

Supermassive black holes: the AGN central engine


Merger-driven processes  Dominate early-type AGN host galaxies

- non-axisymmetric potentials resulting from a nearby companion galaxy can further increase the mass flow rate onto the central regions

Supermassive black holes: the AGN central engine

Merger-driven processes  Dominate early-type AGN host galaxies


- non-axisymmetric potentials resulting from a nearby companion galaxy can further increase the mass flow rate onto the central regions



Supermassive black holes: the AGN central engine

Merger-driven processes ➡ Dominate early-type AGN host galaxies

- non-axisymmetric potentials resulting from a nearby companion galaxy can further increase the mass flow rate onto the central regions



Supermassive black holes: accretion disk

- Gas inevitably posses a non-zero angular momentum when it enters the region dominated by the potential of the black hole
 - the inflowing gas will form a flattened, rotationally-supported structure:

Supermassive black holes: accretion disk

- Gas inevitably possesses a non-zero angular momentum when it enters the region dominated by the potential of the black hole
 - the inflowing gas will form a flattened, rotationally-supported structure:


ACCRETION
DISK

Supermassive black holes: accretion disk

- Gas inevitably possesses a non-zero angular momentum when it enters the region dominated by the potential of the black hole
 - the inflowing gas will form a flattened, rotationally-supported structure:

ACCRETION
DISK

Standard model




advection-dominated cooling,
radiatively inefficient

high radiative
cooling efficiency

Supermassive black holes: accretion disk

- Accretion disks “viscosity” transports angular momentum outwards and allows matter to flow inwards
- The gravitational potential energy of the infalling material is transformed into observable radiation and kinetic energy of jets/outflows
- A self-sustaining magnetic dynamo probably provides the required angular momentum transport



Roen Kelly

Supermassive black holes: jet-accretion disk




Hotaka Shiokawa

Supermassive black holes: jet-accretion disk



Hotaka Shiokawa

AGN components according to Unified Model



AGN components according to Unified Model

- Black hole anatomy

Characteristics:

AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

Schwarzschild BH:

AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

Schwarzschild BH: non-rotating, no charge

AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

Schwarzschild BH: non-rotating, no charge

Kerr BH: rotating


AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

Schwarzschild BH: non-rotating, no charge

Kerr BH: rotating




AGN components according to Unified Model

- Black hole anatomy

Characteristics: mass, spin, charge

Schwarzschild BH: non-rotating, no charge

Kerr BH: rotating



AGN components according to Unified Model

Black hole anatomy


Characteristics: mass, spin, charge

Schwarzschild BH: non-rotating, no charge

Kerr BH: rotating

Event horizon
Boundary region
within nothing
can escape

Schwarzschild radius
Radius of event horizon
 $R_{\text{Sch}} = 2GM/c^2$



Photon sphere

Photons moving on tangential trajectories are trapped in circular orbits around the BH
Radius = $1.5 \times R_{\text{Sch}}$

Singularity
BH density & gravity = ∞

AGN components according to Unified Model

Unified model of AGN:


All AGN contain the same components,
simply viewed at different angles

AGN components according to Unified Model

Unified model of AGN:

All AGN contain the same components,
simply viewed at different angles

- Supermassive black hole of $10^{6-10} M_{\odot}$
- Accretion disk and corona:



AGN components according to Unified Model


Unified model of AGN:

All AGN contain the same components, simply viewed at different angles

- Supermassive black hole of $10^{6-10} M_{\odot}$

- Accretion disk and corona:

- Heated by magnetic and/or viscous processes so that it radiates at optical through soft X-ray energies
- Size: a few AU




AGN components according to Unified Model

Unified model of AGN:

All AGN contain the same components, simply viewed at different angles

- Supermassive black hole of $10^{6-10} M_{\odot}$



- Accretion disk and corona:

- Heated by magnetic and/or viscous processes so that it radiates at optical through soft X-ray energies
- Size: a few AU


- Broad line region:

- Dense gas clouds orbiting central engine at distances 0.1-1 pc
- High velocity (~ 2000 km/s)
- Ionized by continuum emission coming from accretion disk: produces broad emission lines

AGN components according to Unified Model

- Narrow line region:

- Low-density gas clouds at distances 100pc - 4kpc from central engine
- Lower velocity (\sim 500 km/s)
- Ionized by accretion disk and massive stars, produces intense narrow forbidden emission lines




AGN components according to Unified Model

- Narrow line region:

- Low-density gas clouds at distances 100pc - 4kpc from central engine
- Lower velocity (\sim 500 km/s)
- Ionized by accretion disk and massive stars, produces intense narrow forbidden emission lines

- Obscuring torus:




- Cold gas and dust that surrounds the central engine at a distance of up to 10 pc
- Constitutes the source of material used by the accretion disk to feed the SMBH
- Its dust grains absorb the optical radiation coming from the nucleus and re-radiate it at infrared (IR) wavelengths
- Has a non-uniform clumpy structure

AGN components according to Unified Model

- Narrow line region:

- Low-density gas clouds at distances 100pc - 4kpc from central engine
- Lower velocity (\sim 500 km/s)
- Ionized by accretion disk and massive stars, produces intense narrow forbidden emission lines

- Obscuring torus:



- Cold gas and dust that surrounds the central engine at a distance of up to 10 pc
- Constitutes the source of material used by the accretion disk to feed the SMBH
- Its dust grains absorb the optical radiation coming from the nucleus and re-radiate it at infrared (IR) wavelengths
- Has a non-uniform clumpy structure


- Jets:

- Pair of highly collimated relativistic plasma outflows emerging in opposite directions from within $10-100R_{\text{Sch}}$ from the black hole
- Can extend beyond the host galaxy, several kpc or Mpc
- Radiate in all wavebands from radio to gamma-rays via synchrotron and inverse-Compton emission processes

AGN components according to Unified Model

A new view of AGN heated dust:

- Thanks to mid-infrared interferometry & imaging
- Polar dust component in addition to torus



Ramos Almeida & Ricci (2017)

Question

How do we know
there is a SMBH?

Data supporting the presence of a SMBH

How do we know there is a supermassive black hole?

Virial theorem: $M \sim v^2 R / G$

R: distance of clouds/star to black hole
v: velocity of gas/stars

Data supporting the presence of a SMBH

How do we know there is a supermassive black hole?


Virial theorem: $M \sim v^2 R / G$

R: distance of clouds/star to black hole
v: velocity of gas/stars

- Kinetic/rotation curves:

Velocity dispersion of stars in many galaxies increases at the center (e.g. within 1 arcsec)

mass concentration at the galactic nucleus: M_{BH}



Data supporting the presence of a SMBH

How do we know there is a supermassive black hole?

Virial theorem: $M \sim v^2 R / G$

R: distance of clouds/star to black hole
v: velocity of gas/stars


- Kinetic/rotation curves:

Velocity dispersion of stars in many galaxies increases at the center (e.g. within 1 arcsec)

→ mass concentration at the galactic nucleus: M_{BH}

- High-resolution imaging/spectroscopy reveal rotating disks in small radii

→ indicates large masses: SMBH



Data supporting the presence of a SMBH

How do we know there is a supermassive black hole?

Virial theorem: $M \sim v^2 R / G$

R: distance of clouds/star to black hole
v: velocity of gas/stars

- Kinetic/rotation curves:

Velocity dispersion of stars in many galaxies increases at the center (e.g. within 1 arcsec)



➤ mass concentration at the galactic nucleus: M_{BH}

- High-resolution imaging/spectroscopy reveal rotating disks in small radii

➤ indicates large masses: SMBH

ALMA dust continuum, 694 GHz
García-Burillo et al. (2016)


Kormendy & Richstone (1995)



Data supporting the presence of a SMBH


- Dynamics of stars around black hole

Sgr A* in the IR



ESO

1 Light-year




$$M_{\text{BH}} = 3 \times 10^6 M_{\odot}$$

Data supporting the presence of a SMBH


- Dynamics of stars around black hole

Sgr A* in the IR



ESO

1 light-year



1995.5





$$M_{\text{BH}} = 3 \times 10^6 M_{\odot}$$

Data supporting the presence of a SMBH

- Dynamics of stars around black hole

Sgr A* in the IR



$$M_{\text{BH}} = 3 \times 10^6 M_{\odot}$$





Data supporting the presence of a SMBH

- Many AGN produce H₂O and OH masers thanks to intrinsic narrow width of maser emission lines, velocity maps can be derived in very small radius via line frequency measurement with VLBI



Keplerian motion at ~ 0.5 pc from the source
Potential dominated by point-like mass of $\sim 10^7 M_\odot$

VLBI, 22 GHz
Miyoshi et al. (1995)



NGC 4258

0.5 ly

10,000 ly


Data supporting the presence of a SMBH

- Reverberation mapping


- measure time difference between variations in continuum emission & in broad line emission:



distance between BLR
and black hole



UC Davis



Data supporting the presence of a SMBH

- Reverberation mapping


- measure time difference between variations in continuum emission & in broad line emission:
- width of broad emission lines:




distance between BLR
and black hole



gas velocity



UC Davis



Data supporting the presence of a SMBH

- Reverberation mapping

- measure time difference between variations in continuum emission & in broad line emission:
- width of broad emission lines:
- $M \sim v^2 R / G$




distance between BLR
and black hole




gas velocity



black hole mass



UC Davis




Data supporting the presence of a SMBH

- Accretion produces radiation pressure

$$\text{Flux: } F = L/4\pi R^2$$

$$\text{Radiation pressure: } P_{\text{rad}} = F/c = L/4\pi R^2 c$$

$$\text{Radiation force: } F_{\text{rad}} = P_{\text{rad}} \kappa(m_p + m_e)$$



Data supporting the presence of a SMBH

- Accretion produces radiation pressure


$$\text{Flux: } F = L/4\pi R^2$$

$$\text{Radiation pressure: } P_{\text{rad}} = F/c = L/4\pi R^2 c$$

$$\text{Radiation force: } F_{\text{rad}} = P_{\text{rad}} \kappa (m_p + m_e)$$

if accreting material is mostly ionized hydrogen and opacity is provided by Thomson scattering:
cross-section from radiation pressure on electrons, but mass dominated by protons

$$\kappa = \sigma_T / m_p$$



Data supporting the presence of a SMBH

- Accretion produces radiation pressure

$$\text{Flux: } F = L/4\pi R^2$$

$$\text{Radiation pressure: } P_{\text{rad}} = F/c = L/4\pi R^2 c$$


$$\text{Radiation force: } F_{\text{rad}} = P_{\text{rad}} \kappa (m_p + m_e)$$

if accreting material is mostly ionized hydrogen and opacity is provided by Thomson scattering:
cross-section from radiation pressure on electrons, but mass dominated by protons

$$\kappa = \sigma_T / m_p$$



$$F_{\text{rad}} = P_{\text{rad}} \kappa (m_p + m_e) \sim P_{\text{rad}} \kappa m_p = P_{\text{rad}} \sigma_T$$



Data supporting the presence of a SMBH

- Accretion produces radiation pressure

$$\text{Flux: } F = L/4\pi R^2$$

$$\text{Radiation pressure: } P_{\text{rad}} = F/c = L/4\pi R^2 c$$


$$\text{Radiation force: } F_{\text{rad}} = P_{\text{rad}} \kappa (m_p + m_e)$$

if accreting material is mostly ionized hydrogen and opacity is provided by Thomson scattering:
cross-section from radiation pressure on electrons, but mass dominated by protons

$$\kappa = \sigma_T / m_p$$



$$\begin{aligned} F_{\text{rad}} &= P_{\text{rad}} \kappa (m_p + m_e) \sim P_{\text{rad}} \kappa m_p = P_{\text{rad}} \sigma_T \\ &= L \sigma_T / 4\pi R^2 c \end{aligned}$$



Data supporting the presence of a SMBH

- Accretion produces radiation pressure

$$\text{Flux: } F = L/4\pi R^2$$

$$\text{Radiation pressure: } P_{\text{rad}} = F/c = L/4\pi R^2 c$$

$$\text{Radiation force: } F_{\text{rad}} = P_{\text{rad}} \kappa (m_p + m_e)$$

if accreting material is mostly ionized hydrogen and opacity is provided by Thomson scattering:


cross-section from radiation pressure on electrons, but mass dominated by protons

$$\kappa = \sigma_T / m_p$$



$$\begin{aligned} F_{\text{rad}} &= P_{\text{rad}} \kappa (m_p + m_e) \sim P_{\text{rad}} \kappa m_p = P_{\text{rad}} \sigma_T \\ &= L \sigma_T / 4\pi R^2 c \end{aligned}$$

$$\text{Gravitational force: } F_{\text{grav}} = GM(m_p + m_e)/R^2 \sim GMm_p/R^2$$



Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$



$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$


If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$



$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$

$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

 The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$

$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

 The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Data supporting the presence of a SMBH

Eddington limit or luminosity:

maximum luminosity at which

$$F_{\text{grav}} = F_{\text{rad}}$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Energy budget: $E = \epsilon Mc^2$

Radiated energy: $L = dE/dt = \epsilon c^2 dM/dt = \epsilon c^2 \dot{M}$

Data supporting the presence of a SMBH

Eddington limit or luminosity:

maximum luminosity at which

$$F_{\text{grav}} = F_{\text{rad}}$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T/4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Energy budget: $E = \epsilon Mc^2$ \dot{M} = accretion rate

Radiated energy: $L = dE/dt = \epsilon c^2 dM/dt = \epsilon c^2 \dot{M}$

Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$

$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Energy budget: $E = \epsilon Mc^2$ \dot{M} = accretion rate

Radiated energy: $L = dE/dt = \epsilon c^2 dM/dt = \epsilon c^2 \dot{M}$

Eddington accretion rate:
limiting accretion rate above which infalling matter will be blown away by radiation pressure

Data supporting the presence of a SMBH

Eddington limit or luminosity:
maximum luminosity at which
 $F_{\text{grav}} = F_{\text{rad}}$

$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T / 4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Energy budget: $E = \epsilon Mc^2$ \dot{M} = accretion rate

Radiated energy: $L = dE/dt = \epsilon c^2 dM/dt = \epsilon c^2 \dot{M}$

Eddington accretion rate:
limiting accretion rate above which infalling matter will be blown away by radiation pressure

$$\epsilon c^2 \dot{M}_{\text{Edd}} = L_{\text{Edd}}$$

$$\dot{M}_{\text{Edd}} = L_{\text{Edd}} / \epsilon c^2$$

$$\dot{M}_{\text{Edd}} (M_{\odot}/\text{yr}) = 2.2 \times M_{\text{BH}} (M_{\odot}) / 10^8$$

Data supporting the presence of a SMBH

Eddington limit or luminosity:

maximum luminosity at which

$$F_{\text{grav}} = F_{\text{rad}}$$

If AGN luminosity $\gtrsim 10^{44}$ erg/s
(typical Seyfert)



$$GMm_p/R^2 = L_{\text{Edd}} \sigma_T/4\pi R^2 c$$

$$L_{\text{Edd}} = 4\pi c G M m_p / \sigma_T$$

$$L_{\text{Edd}} (\text{erg/s}) = 1.3 \times 10^{38} M_{\text{BH}} (M_{\odot})$$

$M_{\text{BH}} \gtrsim 10^6 M_{\odot}$ so that accretion does not halt due to radiation pressure

The only known object with $10^6 M_{\odot}$ in a $100 R_{\text{Sch}}$ radius is a black hole

- Energy comes from mass-energy conversion

Energy budget: $E = \epsilon Mc^2$

\dot{M} = accretion rate

Radiated energy: $L = dE/dt = \epsilon c^2 dM/dt = \epsilon c^2 \dot{M}$

ϵ = efficiency

Eddington accretion rate:

limiting accretion rate above which infalling matter will be blown away by radiation pressure



$$\epsilon c^2 \dot{M}_{\text{Edd}} = L_{\text{Edd}}$$

$$\dot{M}_{\text{Edd}} = L_{\text{Edd}} / \epsilon c^2$$

$$\dot{M}_{\text{Edd}} (M_{\odot}/\text{yr}) = 2.2 \times M_{\text{BH}} (M_{\odot}) / 10^8$$

Data supporting the presence of a SMBH

- Efficiency of matter conversion into energy, ϵ

Potential energy: $U = GMm/R$

Schwarzchild radius: $R_{\text{Sch}} = 2GMc^2$

Most of optical/UV emission from $5R_{\text{Sch}}$

Data supporting the presence of a SMBH

- Efficiency of matter conversion into energy, ϵ

Potential energy: $U = GMm/R$

Schwarzchild radius: $R_{\text{Sch}} = 2GMc^2$

Most of optical/UV emission from $5R_{\text{Sch}}$



$$U = GMm/5R_{\text{Sch}} = 0.1mc^2$$

Data supporting the presence of a SMBH

- Efficiency of matter conversion into energy, ϵ

Potential energy: $U = GMm/R$

Schwarzchild radius: $R_{\text{Sch}} = 2GMc^2$

Most of optical/UV emission from $5R_{\text{Sch}}$



$$U = GMm/5R_{\text{Sch}} = 0.1mc^2$$

$$\epsilon \sim 0.1$$

Higher than that from
thermonuclear processes!
(H burning into He has $\epsilon = 0.007$)

Data supporting the presence of a SMBH

- Efficiency of matter conversion into energy, ϵ

Potential energy: $U = GMm/R$

Schwarzchild radius: $R_{\text{Sch}} = 2GMc^2$

Most of optical/UV emission from $5R_{\text{Sch}}$



$$U = GMm/5R_{\text{Sch}} = 0.1mc^2$$

$$\epsilon \sim 0.1$$

Higher than that from
thermonuclear processes!
(H burning into He has $\epsilon = 0.007$)

The only process so efficient (apart from particle annihilation)
is accretion of matter onto massive compact object

Data supporting the presence of a SMBH

- Variability

- Interval of variations $\Delta t \geq R/c$ (light crossing time)  upper bound on R
e.g.: $\Delta t \sim 10^5$ s $\Rightarrow R < 10^{-3}$ pc


Data supporting the presence of a SMBH

• Variability

- Interval of variations $\Delta t \geq R/c$ (light crossing time) upper bound on R
e.g.: $\Delta t \sim 10^5$ s $\Rightarrow R < 10^{-3}$ pc

record (BL Lac):

$$\Delta t \sim 11\text{min} \Rightarrow R < 2 \times 10^{13} \text{ cm}$$



BL Lac SS 0716+714

Gupta et al. (2012)

Data supporting the presence of a SMBH

Variability


- Interval of variations $\Delta t \geq R/c$ (light crossing time) upper bound on R
e.g.: $\Delta t \sim 10^5$ s $\Rightarrow R < 10^{-3}$ pc

record (BL Lac):

$$\Delta t \sim 11\text{min} \Rightarrow R < 2 \times 10^{13} \text{ cm}$$

- Mass from $M = L\Delta t/\epsilon$
 $\epsilon = 10\%$


$$M \sim 10^{8-10} M_\odot$$




BL Lac SS 0716+714

Gupta et al. (2012)

Data supporting the presence of a SMBH


- Broadened X-ray fluorescent Fe K α emission lines




Claudio Ricci

Data supporting the presence of a SMBH

- Broadened X-ray fluorescent Fe K α emission lines




Claudio Ricci




Binding energy of the most internal electrons of the Fe atom = 7.1 keV. X-ray photons with energies higher than this threshold can be absorbed and produce photoelectric effect.

Data supporting the presence of a SMBH

- Broadened X-ray fluorescent Fe K α emission lines



Claudio Ricci




Binding energy of the most internal electrons of the Fe atom = 7.1 keV. X-ray photons with energies higher than this threshold can be absorbed and produce photoelectric effect.

Emission line Fe K α at 6.4 keV: produced by an X-ray photon (e.g. of 50 keV) that removes one electron from the K shell of Fe. Subsequent recombination of electrons in the higher levels => cascade of photons => fluorescence emission

Data supporting the presence of a SMBH

Emission line Fe K α at 6.4 keV: thought to originate from the innermost regions of the accretion disk




Data supporting the presence of a SMBH

Emission line Fe K α at 6.4 keV: thought to originate from the innermost regions of the accretion disk

=> mildly-relativistic orbital velocities

=> strong gravitational field: gravitational redshift




Data supporting the presence of a SMBH


Emission line Fe K α at 6.4 keV: thought to originate from the innermost regions of the accretion disk

=> mildly-relativistic orbital velocities

=> strong gravitational field: gravitational redshift



Accretion disk




Fabian et al. (2000)

Data supporting the presence of a SMBH

Emission line Fe K α at 6.4 keV: thought to originate from the innermost regions of the accretion disk


=> mildly-relativistic orbital velocities

=> strong gravitational field: gravitational redshift




Skewed profile:

- extended red wing due to gravitational redshift
- prominent blue wing relativistically boosted due to high orbital velocities of the disk



Fabian et al. (2000)


Data supporting the presence of a SMBH



Fe line profile in agreement with that expected from material in a thin accretion disk near a black hole


Data supporting the presence of a SMBH

- The first image of a black hole



Data supporting the presence of a SMBH

- The first image of a black hole




Question

What are we
exactly seeing?


Data supporting the presence of a SMBH

Space-time curvature




Data supporting the presence of a SMBH

Space-time curvature




Data supporting the presence of a SMBH

Space-time curvature




On galactic scales




Data supporting the presence of a SMBH

Space-time curvature




On galactic scales




Data supporting the presence of a SMBH

The sketch...




Data supporting the presence of a SMBH


The sketch...




...the reality!



Data supporting the presence of a SMBH



Data supporting the presence of a SMBH





EHT collaboration

Data supporting the presence of a SMBH

SIMULATIONS




420 combinations of rotation and fluid composition



Data supporting the presence of a SMBH


SIMULATIONS

420 combinations of rotation and fluid composition




Plasma in clockwise rotation around the black hole
 $M_{\text{BH}} = 6.5 \times 10^9 M_{\odot}$

Data supporting the presence of a SMBH




Data supporting the presence of a SMBH



$M_{\text{BH}} = 4 \times 10^6 M_{\odot}$
Rotation period = 30 min




Data supporting the presence of a SMBH



$M_{\text{BH}} = 4 \times 10^6 M_{\odot}$
Rotation period = 30 min

Frequency of the
different images

Data supporting the presence of a SMBH



M87

Distance: 53 million light-year

Sgr A*

Distance: 27000 light-year

Question

How many AGN
types do you know?

AGN Zoology


- Radio galaxies
- Radio Quasars
- BL Lac Objects
- Optically Violent Variables (OVV's)
- Radio Quiet Quasars (QSOs)
- Seyfert I galaxies (SyI)
- Seyfert II galaxies (SyII)
- Low Ionization Nuclear Emission-Line Regions (LINERs)

AGN Classification

- There is a large diversity in the nature of the observed radiation from different AGN
- The classification scheme has varied with time
- A fundamental division in the classification scheme is based on the radio properties of AGN:

radio-loud or radio-quiet

AGN Classification





AGN Classification

Radio-loud



Radio-quiet



QSO Host Galaxies

HST images



10''



AGN Classification

Radio-quiet AGN

- Do not show large scale collimated jets (extent <0.5 kpc)
- Within radio-quiet AGN: classification on the basis of optical/UV spectral properties

Radio-quiet AGN

- Do not show large scale collimated jets (extent <0.5 kpc)
- Within radio-quiet AGN: classification on the basis of optical/UV spectral properties

Seyfert I galaxies/radio-quiet quasars

- Spectrum is complex and multicomponent
- Presence of broad (FWHM~2000-20000 km/s) permitted optical/UV emission lines
- Possible presence of low-ionization broad lines (MgII λ 2798 and hydrogen lines) and high-ionization broad lines (CIV λ 1549, OVI λ 1035 and NeVIII λ 774)


AGN Classification

Radio-quiet AGN

- Do not show large scale collimated jets (extent <0.5 kpc)
- Within radio-quiet AGN: classification on the basis of optical/UV spectral properties

Seyfert I galaxies/radio-quiet quasars

- Spectrum is complex and multicomponent
- Presence of broad (FWHM~2000-20000 km/s) permitted optical/UV emission lines
- Possible presence of low-ionization broad lines (MgII λ 2798 and hydrogen lines) and high-ionization broad lines (CIV λ 1549, OVI λ 1035 and NeVIII λ 774)



Broad line region:
dense ($n_e \sim 10^9 \text{ cm}^{-3}$)
photoionized clouds
with small volume
filling factor

AGN Classification


Seyfert I galaxies/radio-quiet quasars

- Line widths due to cloud motions
- Collisional de-excitation dominates over forbidden line emission due to high density
- Optical/UV narrow (~ 500 km/s) permitted and forbidden emission lines also present in spectra
 - e.g.: hydrogen lines, [OII] $\lambda 3727$, [OIII] $\lambda 5007$, [NII] $\lambda 6583$, [SII] $\lambda 6716/6731$
- The NLR can be spatially resolved in nearby AGN and is found to have typical sizes of 0.1 kpc

AGN Classification

Seyfert I galaxies/radio-quiet quasars

- Line widths due to cloud motions
- Collisional de-excitation dominates over forbidden line emission due to high density
- Optical/UV narrow (~ 500 km/s) permitted and forbidden emission lines also present in spectra
- e.g.: hydrogen lines, [OII] $\lambda 3727$, [OIII] $\lambda 5007$, [NII] $\lambda 6583$, [SII] $\lambda 6716/6731$
- The NLR can be spatially resolved in nearby AGN and is found to have typical sizes of 0.1 kpc



Narrow line region:
material more
tenuous than BLR,
at larger distances
from black hole

AGN Classification


Seyfert I galaxies vs radio-quiet quasars

- Distinction between Seyfert I and radio-quiet quasar (RQQ or quasi-stellar object QSO) based on whether the galaxy or AGN is discovered first
- SyI are generally lower-luminosity than RQQ:
Dividing luminosity $L_{\text{bol}} \sim 10^{45} \text{ erg/s}$, $L_x \sim 10^{42} \text{ erg/s}$
- In practice, SyI and RQQs form a continuous sequence in luminosity
- Both display strong continuum radiation from IR through gamma-ray wavelengths
- The higher-energy emission is often observed to vary rapidly

AGN Classification

Seyfert II galaxies

- Their spectra display narrow permitted and forbidden emission lines
- Unlike SyI galaxies, no broad lines are observable and the optical/UV radiation is much weaker
- Some display broad optical/UV lines in polarimetry or NIR observations



Antonucci (1993), Urry & Padovani (1995)


AGN Classification

Seyfert II galaxies

- Their spectra display narrow permitted and forbidden emission lines
- Unlike SyI galaxies, no broad lines are observable and the optical/UV radiation is much weaker
- Some display broad optical/UV lines in polarimetry or NIR observations



some SyII are reddened SyI in which the BLR lies behind the obscuring material



Antonucci (1993), Urry & Padovani (1995)

AGN Classification


Seyfert II galaxies

- Their spectra display narrow permitted and forbidden emission lines
- Unlike SyI galaxies, no broad lines are observable and the optical/UV radiation is much weaker
- Some display broad optical/UV lines in polarimetry or NIR observations



some SyII are reddened SyI in which the BLR lies behind the obscuring material


- X-ray observations reveal photoelectric absorption from cold gas presumably associated with the dust



Antonucci (1993), Urry & Padovani (1995)

AGN Classification

Narrow line Seyfert I galaxies (NLS1)




Masetti et al. (2009)

AGN Classification

Narrow line Seyfert I galaxies (NLS1)

- Permitted optical/UV lines significantly narrower than SyI (but still broader than lines from NLR): FWHM ($H\beta$) ≤ 2000 km/s
- Ratio $[OIII]\lambda 5007/H\beta < 3$, a criterion used to differentiate SyI from SyII




Masetti et al. (2009)

AGN Classification

Narrow line Seyfert I galaxies (NLS1)

- Permitted optical/UV lines significantly narrower than SyI (but still broader than lines from NLR): FWHM ($H\beta$) ≤ 2000 km/s
- Ratio $[OIII]\lambda 5007/H\beta < 3$, a criterion used to differentiate SyI from SyII
- Display strong optical line complexes due to FeII emission: $FeII/H\beta > 0.5$
- X-ray continuum softer and more variable than normal SyI



Masetti et al. (2009)

AGN Classification

Narrow Emission Line galaxies (NELGs)

- Show narrow (FWHM<1000 km/s) emission lines, typically of H α , H β , [OIII] $\lambda 5007$
- X-ray luminosities $L_x \sim 10^{42}$ erg/s (two orders of magnitude brighter than normal galaxies)
- Often classified as AGN but part of the emission may be due to star formation
- Good candidates for producing the X-ray background due to number of sources and hard spectra

AGN Classification

Radio-loud AGN

- Emit collimated jets of plasma that feed energy and high-energy particles into extended halo or lobe. Large-scale jet (extent > 0.5 kpc). The halo can be $> 1\text{Mpc}$
- Jets and radio halo: sources of continuum radio emission (synchrotron emission from relativistic electrons in jet/halo plasma)

Perseus cluster
Minihalo



Radio, optical
Gendron-Marsolais et al. (2017)
NRAO, NASA, SDSS

AGN Classification

Radio-loud AGN


- Emit collimated jets of plasma that feed energy and high-energy particles into extended halo or lobe. Large-scale jet (extent > 0.5 kpc). The halo can be $> 1\text{Mpc}$
- Jets and radio halo: sources of continuum radio emission (synchrotron emission from relativistic electrons in jet/halo plasma)
- Well-collimated jets linking AGN and halo are observed

AGN Classification

Radio-loud AGN

- Emit collimated jets of plasma that feed energy and high-energy particles into extended halo or lobe. Large-scale jet (extent > 0.5 kpc). The halo can be $> 1\text{Mpc}$
- Jets and radio halo: sources of continuum radio emission (synchrotron emission from relativistic electrons in jet/halo plasma)
- Well-collimated jets linking AGN and halo are observed

Hercules A




X-ray, radio, optical
NASA, NRAO

AGN Classification

Radio-loud AGN

- Emit collimated jets of plasma that feed energy and high-energy particles into extended halo or lobe. Large-scale jet (extent > 0.5 kpc). The halo can be $> 1\text{Mpc}$
- Jets and radio halo: sources of continuum radio emission (synchrotron emission from relativistic electrons in jet/halo plasma)
- Well-collimated jets linking AGN and halo are observed

Hercules A




X-ray, radio, optical
NASA, NRAO

AGN Classification

Radio-loud AGN

- Some present only one observable jet




Antonucci (1993), Urry & Padovani (1995)

AGN Classification

Radio-loud AGN

- Some present only one observable jet




Antonucci (1993), Urry & Padovani (1995)

AGN Classification

Radio-loud AGN

- Some present only one observable jet





- Outflowing blobs within the jets

AGN Classification


Radio-loud AGN

- Some present only one observable jet



Antonucci (1993), Urry & Padovani (1995)

Superluminal Motion in the M87 Jet




Biretta (1999)

- Outflowing blobs within the jets

Blobs superluminal motion:

apparent proper motion can exceed c .
This results from relativistic motion in
a direction \sim towards the observer



Radio-loud AGN

- Within radio-loud AGN: classification on the basis of both radio morphology and optical/UV spectral properties


AGN Classification

Radio-loud AGN

- Within radio-loud AGN: classification on the basis of both radio morphology and optical/UV spectral properties

FR-I (Fanaroff-Riley-I) radio galaxies

- Edge-darkened morphology: radio surface brightness profile falls continuously from nucleus to the edge of radio lobes
- Low luminosity ($L_{\text{radio}} < 10^{42} \text{ erg/s}$)
- Display narrow emission lines from NLR, but no broad emission lines



AGN Classification

FR-II (Fanaroff-Riley-II) radio galaxies


- Edge-brightened morphology: nucleus apart, radio surface brightness low at the centre and dramatically brightens towards edge of halo due to jet(s) terminating at strong shocks
- High luminosity ($L_{\text{radio}} > 10^{42} \text{ erg/s}$)
- Optical properties either SyI-like or SyII-like: broad line radio galaxies (BLRG) and narrow line radio galaxies (NLRG), respectively



AGN Classification

Radio-loud quasars (RLQ)

- Blurred distinction between BLRG and RLQ (or simply quasar)
- RLQ = BLRG in which AGN continuum emission dominates over radio emission
- RLQ often posses one-sided jets with superluminal motion



AGN Classification


Blazars

- Radio-loud objects displaying very strong and variable continuum at all wavelengths
- High optical polarization ($\geq 3\%$)
- Optical/UV emission lines very weak or absent

AGN Classification

Blazars

- Radio-loud objects displaying very strong and variable continuum at all wavelengths
- High optical polarization ($\geq 3\%$)
- Optical/UV emission lines very weak or absent



Sub-classification:

AGN Classification

Blazars

- Radio-loud objects displaying very strong and variable continuum at all wavelengths
- High optical polarization ($\geq 3\%$)
- Optical/UV emission lines very weak or absent



Sub-classification: optically-violent
variable (OVV)

AGN Classification

Blazars

- Radio-loud objects displaying very strong and variable continuum at all wavelengths
- High optical polarization ($\geq 3\%$)
- Optical/UV emission lines very weak or absent
 - Sub-classification:
 - optically-violent variable (OVV)
 - BL Lac object

AGN Classification

Blazars

- Radio-loud objects displaying very strong and variable continuum at all wavelengths
- High optical polarization ($\geq 3\%$)
- Optical/UV emission lines very weak or absent
 - Sub-classification:
 - optically-violent variable (OVV)
 - BL Lac object
- Flat radio spectrum: flat-spectrum radio quasars (FSRQ)

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES				SOME	SOME
Radio-quiet quasars	YES	YES				WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES				WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO				NO	NO
OVV quasars	YES	YES				YES	YES
BL Lac objects	YES	YES				YES	YES
Seyferts type 1	YES	YES				SOME	WEAK
Seyferts type 2	NO	YES				NO	SOME
LINERs	NO	NO				NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES				WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES				WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO				NO	NO
OVV quasars	YES	YES				YES	YES
BL Lac objects	YES	YES				YES	YES
Seyferts type 1	YES	YES				SOME	WEAK
Seyferts type 2	NO	YES				NO	SOME
LINERs	NO	NO				NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES				WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO				NO	NO
OVV quasars	YES	YES				YES	YES
BL Lac objects	YES	YES				YES	YES
Seyferts type 1	YES	YES				SOME	WEAK
Seyferts type 2	NO	YES				NO	SOME
LINERs	NO	NO				NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized	
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME	
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK	
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK	
Narrow line radio galaxies (FR1 and FR2)	NO	NO					NO	NO
OVV quasars	YES	YES					YES	YES
BL Lac objects	YES	YES					YES	YES
Seyferts type 1	YES	YES					SOME	WEAK
Seyferts type 2	NO	YES					NO	SOME
LINERs	NO	NO					NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized	
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME	
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK	
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK	
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO	
OVV quasars	YES	YES					YES	YES
BL Lac objects	YES	YES					YES	YES
Seyferts type 1	YES	YES					SOME	WEAK
Seyferts type 2	NO	YES					NO	SOME
LINERs	NO	NO					NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized	
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME	
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK	
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK	
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO	
OVV quasars	YES	YES	YES	YES	YES	YES	YES	
BL Lac objects	YES	YES					YES	YES
Seyferts type 1	YES	YES					SOME	WEAK
Seyferts type 2	NO	YES					NO	SOME
LINERs	NO	NO					NO	NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO
OVV quasars	YES	YES	YES	YES	YES	YES	YES
BL Lac objects	YES	YES	NO	NO	YES	YES	YES
Seyferts type 1	YES	YES					SOME
Seyferts type 2	NO	YES					WEAK
LINERs	NO	NO					NO

AGN Classification

AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO
OVV quasars	YES	YES	YES	YES	YES	YES	YES
BL Lac objects	YES	YES	NO	NO	YES	YES	YES
Seyferts type 1	YES	YES	YES	YES	WEAK	SOME	WEAK
Seyferts type 2	NO	YES				NO	SOME
LINERs	NO	NO				NO	NO

AGN Classification

AGN bestiary


Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO
OVV quasars	YES	YES	YES	YES	YES	YES	YES
BL Lac objects	YES	YES	NO	NO	YES	YES	YES
Seyferts type 1	YES	YES	YES	YES	WEAK	SOME	WEAK
Seyferts type 2	NO	YES	NO	YES	WEAK	NO	SOME
LINERs	NO	NO				NO	NO

AGN Classification


AGN bestiary

Beast	Point like	Broad-band	Broad lines	Narrow lines	Radio	Variable	Polarized
Radio-loud quasars	YES	YES	YES	YES	YES	SOME	SOME
Radio-quiet quasars	YES	YES	YES	YES	WEAK	WEAK	WEAK
Broad line radio galaxies (FR 2 only)	YES	YES	YES	YES	YES	WEAK	WEAK
Narrow line radio galaxies (FR1 and FR2)	NO	NO	NO	YES	YES	NO	NO
OVV quasars	YES	YES	YES	YES	YES	YES	YES
BL Lac objects	YES	YES	NO	NO	YES	YES	YES
Seyferts type 1	YES	YES	YES	YES	WEAK	SOME	WEAK
Seyferts type 2	NO	YES	NO	YES	WEAK	NO	SOME
LINERs	NO	NO	NO	YES	NO	NO	NO


AGN Classification: find the Seyfert 1 & BLRG




AGN Classification: find the Seyfert 1 & BLRG




AGN Classification: find the Seyfert 2 & NLRG




AGN Classification: find the Seyfert 2 & NLRG




AGN Classification: find the BL Lac




AGN Classification: find the BL Lac




AGN Classification: find the normal galaxy



AGN Classification: find the normal galaxy




AGN Classification: find the normal galaxy



AGN Unification

Radio-Quiet AGN unification

- Strongest unification for SyI and SyII galaxies
- Prototypical SyII galaxy NGC1068: broad optical emission lines when observed in polarized light. If polarization due to scattering, there are directions from which NGC1068 displays a SyI spectrum




Miller & Antonucci (1983), Antonucci & Miller (1985), Miller, Goodrich & Mathews (1991)

AGN Unification

Radio-Quiet AGN unification

- Strongest unification for SyI and SyII galaxies
- Prototypical SyII galaxy NGC1068: broad optical emission lines when observed in polarized light. If polarization due to scattering, there are directions from which NGC1068 displays a SyI spectrum

SyII appear to be SyI
with BLR and
continuum radiation
heavily obscured




Miller & Antonucci (1983), Antonucci & Miller (1985), Miller, Goodrich & Mathews (1991)

AGN Unification

Radio-Quiet AGN unification model


Antonucci & Miller (1985): all Seyfert galaxies possess a dusty torus in between the BLR and NLR that intercepts the sight of the BLR




AGN Unification

Radio-Quiet AGN unification model

Antonucci & Miller (1985): all Seyfert galaxies possess a dusty torus in between the BLR and NLR that intercepts the sight of the BLR



Type 2



Antonucci (1993), Urry & Padovani (1995)

Radio-Loud AGN unification

- Blazars, radio-loud quasars and FR-II galaxies unified under a similar scheme to the Seyfert one
- The only difference is the inclusion of relativistic jets of synchrotron emitting plasma
 - Relativistic beaming and Doppler shifts: source of anisotropy in addition to torus obscuration

Radio-Loud AGN unification

- Blazars, radio-loud quasars and FR-II galaxies unified under a similar scheme to the Seyfert one
- The only difference is the inclusion of relativistic jets of synchrotron emitting plasma
 - Relativistic beaming and Doppler shifts: source of anisotropy in addition to torus obscuration
- Blazars: line-of-sight within jet cone. Strong relativistic aberration and jet Doppler shift produces highly variable continuum-dominated emission

Radio-Loud AGN unification

- Blazars, radio-loud quasars and FR-II galaxies unified under a similar scheme to the Seyfert one
- The only difference is the inclusion of relativistic jets of synchrotron emitting plasma
 - Relativistic beaming and Doppler shifts: source of anisotropy in addition to torus obscuration
- Blazars: line-of-sight within jet cone. Strong relativistic aberration and jet Doppler shift produces highly variable continuum-dominated emission
- RLQ: line-of-sight closet (but not within) jet cone. Continuum strongly beamed but does not swamp emission from BLR

Radio-Loud AGN unification

- Blazars, radio-loud quasars and FR-II galaxies unified under a similar scheme to the Seyfert one
- The only difference is the inclusion of relativistic jets of synchrotron emitting plasma
 - Relativistic beaming and Doppler shifts: source of anisotropy in addition to torus obscuration
- Blazars: line-of-sight within jet cone. Strong relativistic aberration and jet Doppler shift produces highly variable continuum-dominated emission
- RLQ: line-of-sight closet (but not within) jet cone. Continuum strongly beamed but does not swamp emission from BLR
- FR-II BLRG: larger orientation angle, central continuum flux falls


Radio-Loud AGN unification

- Blazars, radio-loud quasars and FR-II galaxies unified under a similar scheme to the Seyfert one
- The only difference is the inclusion of relativistic jets of synchrotron emitting plasma
 - Relativistic beaming and Doppler shifts: source of anisotropy in addition to torus obscuration
- Blazars: line-of-sight within jet cone. Strong relativistic aberration and jet Doppler shift produces highly variable continuum-dominated emission
- RLQ: line-of-sight closet (but not within) jet cone. Continuum strongly beamed but does not swamp emission from BLR
- FR-II BLRG: larger orientation angle, central continuum flux falls
- FR-II NLRG: yet larger angle, torus obscures central continuum source and BLR

AGN Unification

The observed properties, and thus the classification, of a given AGN depend upon its **orientation**

AGN Unification




AGN Unification: caveats

Caveats

- Unclear how FR-I radio galaxies fit into the unification scheme
- Source power rather than orientation produces different radio morphologies for FR-I and FR-II
- E.g. blazars can be beamed FR-I, but there is a lack of BL FR-I torus has small opening angle? absence of BLR in FR-I sources?
- Dust lanes from galactic scales can explain nuclear obscuration in some low-luminosity type 2 AGN


AGN Unification: caveats



Prieto, Mezcua et al. (2014)

Nucleus behind dust lane:

- Obscures it ($A_v=3$ mag)
- Optical peak is not the nucleus!
Shift of 17 pc!




AGN Unification: caveats

Caveats

- Unclear how FR-I radio galaxies fit into the unification scheme
- Source power rather than orientation produces different radio morphologies for FR-I and FR-II
- E.g. blazars can be beamed FR-I, but there is a lack of BL FR-I
 - torus has small opening angle? absence of BLR in FR-I sources?
- Dust lanes from galactic scales can explain nuclear obscuration in some low-luminosity type 2 AGN
 - no torus needed to obscure BLR?



AGN Spectral Energy Distribution

X-ray Spectral Energy Distribution (SED)



AGN Spectral Energy Distribution


X-ray Spectral Energy Distribution (SED)




Primary X-ray radiation:
power-law, $\Gamma=1.8-2$ ($\Gamma=1-\alpha$)

Reflected X-ray radiation:
- reflection continuum, peak at 30-40 keV
- Fe K α line at 6.4 keV
- soft excess at ~ 2 keV

AGN Spectral Energy Distribution




AGN Spectral Energy Distribution




Largely unobservable due to absorption in the IGM

AGN Spectral Energy Distribution



Prieto et al. (2010)

AGN Spectral Energy Distribution




Prieto et al. (2010)

Question

How do SMBHs form?

How do stellar-mass black holes form?


Stellar-black hole formation



NASA

How do stellar-mass black holes form?

Stellar-black hole formation




Crab nebula
(year 1054)



ESA/HST

How do stellar-mass black holes form?

Stellar-black hole formation




Crab nebula
(year 1054)



ESA/HST

How do stellar-mass black holes form?

Stellar-black hole formation



Crab nebula
(year 1054)




ESA/HST

How do SMBHs form?

SMBH formation

- Slow accretion onto stellar-mass black hole



NASA

- Mergers of stellar-mass BHs in stellar cluster + slow accretion



NASA

- SMBH merger in galaxy merger



How do SMBHs form?

Galaxy merger: simulation vs observations


How do SMBHs form?

Galaxy merger: simulation vs observations

How do SMBHs form?

The challenge of the first quasars


- Quasars of $10^{10} M_{\odot}$ already existed when Universe was < 1 Gyr



How do SMBHs form?


The challenge of the first quasars

- Quasars of $10^{10} M_{\odot}$ already existed when Universe was < 1 Gyr



Valiante et al. (2017)

No time to grow from stellar-mass BHs!




Had to start as **seed BHs** of $100 - 10^5 M_{\odot}$

Grow fast via accretion and mergers

How do SMBHs form?

Seed black hole formation




How do SMBHs form?

Where can we find seed IMBHs?


How do SMBHs form?

Where can we find seed IMBHs?



How do SMBHs form?


Where can we find seed IMBHs?



How do SMBHs form?


Where can we find seed IMBHs?

In dwarf galaxies!





How do SMBHs form?

Seed black hole formation



How do SMBHs form?

Seed black hole formation



How do SMBHs form?

Dwarf galaxies

- Have low mass and low metallicity
- Do not (presumably) significantly grow through mergers and accretion
- Regulated by supernova feedback: stellar outflows - evacuate gas from nucleus
 - hamper BH growth



Resemble the first galaxies formed in the early Universe

How do SMBHs form?

Dwarf galaxies

- Have low mass and low metallicity
- Do not (presumably) significantly grow through mergers and accretion
- Regulated by supernova feedback: stellar outflows - evacuate gas from nucleus
 - hamper BH growth




Resemble the first galaxies formed in the early Universe

IMBHs as AGN in dwarf galaxies

- First two discoveries serendipitous:
 - Broad emission lines $M_{\text{BH}} \sim 3 \times 10^5 M_{\odot}$
 - X-ray emission signature of accretion



Mt. Lemmon Sky Center



Thornton et al. (2008)

How do SMBHs form?

IMBHs as AGN in dwarf galaxies

- Searches using optical/IR emission lines:
 - ~Hundreds of candidates, tens confirmed by X-ray emission
 - Limited to $z < 1$, $M_{\text{BH}} \sim 10^4 - 10^6 M_{\odot}$ from broad lines

Baldassare et al. (2015)




How do SMBHs form?

IMBHs as AGN in dwarf galaxies


- Searches using optical/IR emission lines:
 - ~Hundreds of candidates, tens confirmed by X-ray emission
 - Limited to $z < 1$, $M_{\text{BH}} \sim 10^4 - 10^6 M_{\odot}$ from broad lines

Baldassare et al. (2015)



- Searches in X-ray and radio surveys:
 - Out to $z \sim 3.4$
 - M_{BH} from BH-galaxy scaling relations

Mezcua et al. (2018)



How do SMBHs form?

IMBHs as AGN in dwarf galaxies


- Searches using optical/IR emission lines:
 - ~Hundreds of candidates, tens confirmed by X-ray emission
 - Limited to $z < 1$, $M_{\text{BH}} \sim 10^4 - 10^6 M_{\odot}$ from broad lines

Baldassare et al. (2015)



- Searches in X-ray and radio surveys:
 - Out to $z \sim 3.4$
 - M_{BH} from BH-galaxy scaling relations
- At $z=10$?



Mezcua et al. (2018)



How do SMBHs form?

James Webb Space Telescope


SMBHs in low-mass galaxies at $z \sim 4-10!!$



How do SMBHs form?

Gravitational waves

- Waves in the space-time due to massive objects in motion (e.g. two BHs orbiting each other)
- Gravitational waves propagate at the speed of light
- They were predicted by Einstein's Theory of Relativity!!




NASA, Caltech

How do SMBHs form?

Gravitational waves

- Waves in the space-time due to massive objects in motion (e.g. two BHs orbiting each other)
- Gravitational waves propagate at the speed of light
- They were predicted by Einstein's Theory of Relativity!!




NASA, Caltech

Coalescence of two stellar-mass black holes yield an IMBH




Black hole mergers



How do SMBHs form?

The Nobel Prize in Physics
2017



Gravitational waves


- Waves in the space-time due to massive objects in motion (e.g. two BHs orbiting each other)
- Gravitational waves propagate at the speed of light
- They were predicted by Einstein's Theory of Relativity!!

Coalescence of two stellar-mass black holes yield an IMBH



2020


Black hole mergers




How do SMBHs form?

Wait for future observatories to detect seed BH at high-z


LISA (2034?)



Athena X-ray observatory (?)



Lynx X-ray observatory
(concept study)



Question

What have we
learned?

AGN: summary


- AGN are powered by SMBHs (of $>10^6 M_\odot$)
- Found in galaxies of all masses and types
- Observed AGN properties and classification depend on

AGN: summary

- AGN are powered by SMBHs (of $>10^6 M_\odot$)
- Found in galaxies of all masses and types
- Observed AGN properties and classification depend on **orientation**

AGN: summary

- AGN are powered by SMBHs (of $>10^6 M_\odot$)
- Found in galaxies of all masses and types
- Observed AGN properties and classification depend on **orientation**



Beckman et al. (2012)

AGN: summary

- SMBH presence and BH mass inferred from:

AGN: summary

- SMBH presence and BH mass inferred from:

Rotation curves

Dynamics of stars

Water masers

Reverberation mapping

Accretion/radiation arguments

Broadened X-ray K α emission line

Direct imaging (M87)

AGN: summary

- SMBH presence and BH mass inferred from:

Rotation curves

Dynamics of stars

Water masers

Reverberation mapping 

Accretion/radiation arguments

Broadened X-ray K α emission line

Direct imaging (M87)

AGN: summary

- SMBH presence and BH mass inferred from:

Rotation curves

Dynamics of stars

Water masers


Reverberation mapping



Accretion/radiation arguments

Broadened X-ray K α emission line

Direct imaging (M87)



Kaspi et
al. (2000)

AGN: summary

- SMBH presence and BH mass inferred from:

Rotation curves

Dynamics of stars


Water masers

Reverberation mapping 

Accretion/radiation arguments

Broadened X-ray K α emission line

Direct imaging (M87)



Kaspi et al. (2000)

- Quasars detect at z~7 (Universe ~0.8 Gyr old)



SMBHs formed from seed IMBHs of $100 - 10^5 M_{\odot}$

Relics of seed IMBHs possibly found in local dwarf galaxies