

Partial Differential Equations

TA Homework 4

Francisco Jose Castillo Carrasco

August 24, 2018

Problem 2

Verify that the function \langle, \rangle defined in Example 0.3 is an inner product.

Solution: Given the inner product on C^2 defined by

$$\langle v, w \rangle = (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix},$$

it is easy to check the properties.

- Positivity:
- Conjugate symmetry:

$$\begin{aligned} \langle v, w \rangle &= (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \\ &= (2\bar{w}_1 + i\bar{w}_2 \quad 3\bar{w}_2 - i\bar{w}_1) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \\ &= 2v_1\bar{w}_1 + iv_1\bar{w}_2 + 3v_2\bar{w}_2 - iv_2\bar{w}_1 \\ &= \overline{2\bar{v}_1w_1 - i\bar{v}_1w_2 + 3\bar{v}_2w_2 + i\bar{v}_2w_1} \\ &= \overline{(2\bar{v}_1 + i\bar{v}_2 \quad 3\bar{v}_2 - i\bar{v}_1) \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}} \\ &= (\bar{v}_1 \quad \bar{v}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} \\ &= \overline{\langle w, v \rangle} \end{aligned}$$

- Homogeneity:

$$\begin{aligned} \langle cv, w \rangle &= (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} cv_1 \\ cv_2 \end{pmatrix} \\ &= (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} c \\ &= \langle v, w \rangle c \\ &= c \langle v, w \rangle, \end{aligned}$$

where we have taken the complex scalar c out of the vector v since it is common in all its components.

- Linearity:

$$\begin{aligned}
\langle u + v, w \rangle &= (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} u_1 + v_1 \\ u_2 + v_2 \end{pmatrix} \\
&= (\bar{w}_1 \quad \bar{w}_2) (2(u_1 + v_1) - i(u_2 + v_2) \quad i(u_1 + v_1) + 3(u_2 + v_2)) \\
&= (\bar{w}_1 \quad \bar{w}_2) (2u_1 - iu_2 + 2v_1 - iv_2 \quad iu_1 + 3u_2 + iv_1 + 3v_2) \\
&= (\bar{w}_1 \quad \bar{w}_2) [(2u_1 - iu_2 \quad iu_1 + 3u_2) + (2v_1 - iv_2 \quad iv_1 + 3v_2)] \\
&= (\bar{w}_1 \quad \bar{w}_2) \left[\begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \right] \\
&= (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + (\bar{w}_1 \quad \bar{w}_2) \begin{pmatrix} 2 & -i \\ i & 3 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \\
&= \langle u, w \rangle + \langle v, w \rangle
\end{aligned}$$

Problem 7

For $n > 0$, let

$$f_n(t) = \begin{cases} \sqrt{n}, & 0 \leq t \leq 1/n^2 \\ 0, & \text{otherwise} \end{cases}$$

Show that $f_n \rightarrow 0$ in $L^2[0, 1]$ but that $f_n(0)$ does not converge to zero.

Solution: For the first proof we need to prove that

$$\|f_n(t) - 0\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Then,

$$\begin{aligned} \|f_n(t) - 0\| &= \sqrt{\langle f_n - 0, f_n - 0 \rangle_{L^2}} \\ &= \sqrt{\int_0^1 [f_n(t) - 0]^2 dt} \\ &= \sqrt{\int_0^1 [f_n(t)]^2 dt} \\ &= \sqrt{\int_0^{1/n^2} n dt + \int_{1/n^2}^1 0 dt} \\ &= \sqrt{\frac{1}{2} n \frac{1}{n^2}} \\ &= \sqrt{\frac{1}{2n}}. \end{aligned}$$

Hence,

$$\lim_{n \rightarrow \infty} \|f_n(t) - 0\| = \lim_{n \rightarrow \infty} \sqrt{\frac{1}{2n}} = 0.$$

However,

$$\lim_{n \rightarrow \infty} f_n(0) = \lim_{n \rightarrow \infty} \sqrt{n} = \infty \neq 0.$$

Problem 11

Show that if a differentiable function , f , is orthogonal to $\cos(t)$ on $L^2[0, \pi]$ then f' is orthogonal to $\sin(t)$ on $L^2[0, \pi]$.

Solution: Since f is orthogonal to $\cos(t)$ we know that

$$\langle f, \cos(t) \rangle = \int_0^\pi f(t) \cos(t) dt = 0.$$

Then,

$$\begin{aligned} \langle f', \sin(t) \rangle &= \int_0^\pi f'(t) \sin(t) dt \\ &= \int_0^\pi \left(\frac{d}{dt} [f(t) \sin(t)] - f(t) \cos(t) \right) dt \\ &= \int_0^\pi d[f(t) \sin(t)] - \int_0^\pi f(t) \cos(t) dt \\ &= \cancel{f(t) \sin(t)} \Big|_0^\pi - \int_0^\pi f(t) \cos(t) dt \\ &= - \int_0^\pi f(t) \cos(t) dt = 0, \end{aligned}$$

where we have used that $\sin(\pi) = \sin(0) = 0$ and the fact that f is orthogonal to $\cos(t)$. Hence, it has been proved that f' is orthogonal to $\sin(t)$ provided that f is orthogonal to $\cos(t)$.

Problem 14

Find the $L^2[-\pi, \pi]$ projection of the function $f(x) = x^2$ onto the space $V_n \in L^2[-\pi, \pi]$ spanned by for $n=2$. Plot these projections along with f using a computer algebra system. Repeat for $g(x) = x^3$.

Solution:

Problem 23

Show that a set of orthonormal vectors is linearly independent.

Solution: