

# Advanced Numerical Methods for PDEs

## Homework 1

Francisco Castillo

February 9, 2021

Consider the initial - boundary value problem for the scalar advection diffusion equation

$$\partial_t u(x, t) + a \partial_x u(x, t) - b \partial_x^2 u(x, t) = 0, \quad u(x, t = 0) = u^I(x), \quad (1)$$

on the interval  $x \in [-1, 1]$  with periodic boundary conditions  $u(x + 2, t) = u(x, t), \forall x, t$ . Consider the explicit difference method

$$U(x, t + \Delta t) = U(x, t) - \frac{a \Delta t}{2 \Delta x} (T - T^{-1}) U(x, t) + \frac{b}{\Delta x^2} (T - 2 + T^{-1}) U(x, t) \quad (2)$$

for the problem (1).

### Problem 1

1. Derive an analytic expression for the solution  $u(x, t)$  of problem (1) for a general initial function  $u^I(x)$  and general stepsizes  $\Delta x, \Delta t$ , using Fourier transforms.

**Solution:** We will use the Fourier transform

$$\hat{u}(w, t) = \frac{1}{\sqrt{2\pi}} \int_{-1}^1 u(x, t) e^{-i\omega x} dx, \quad (3)$$

to turn our 1 -  $D$  PDE into an ODE. Taking the Fourier transform of the PDE (1) we obtain

$$\partial_t \hat{u}(w, t) + iaw \hat{u}(w, t) + bw^2 \hat{u}(w, t) = 0,$$

which can be manipulated into

$$\partial_t \hat{u}(w, t) = -(bw^2 + iaw) \hat{u}(w, t).$$

The previous equation has a simple analytical solution,

$$\hat{u}(w, t) = e^{-(bw^2 + iaw)t} \hat{u}(w, 0),$$

which we can conveniently rewrite as

$$\hat{u}(w, t) = e^{-b\omega^2 t} \hat{u}^I(w) e^{-ia\omega t} \quad (4)$$

To obtain the solution, we will use the following property of Fourier transforms,

$$\mathcal{F}[f * g] = \mathcal{F}[f] \mathcal{F}[g], \quad (5)$$

where  $*$  represents convolution. In our case, we have

$$\begin{aligned} \mathcal{F}[f](w, t) &= e^{-b\omega^2 t}, \\ \mathcal{F}[g](w, t) &= \hat{u}^I(w) e^{-ia\omega t}. \end{aligned}$$

By using the inverse Fourier transform on the previous equations we obtain

$$\begin{aligned} f(x, t) &= \mathcal{F}^{-1}[\mathcal{F}[f](w, t)](x, t) = \frac{e^{-x^2/4bt}}{\sqrt{2bt}}, \\ g(x, t) &= \mathcal{F}^{-1}[\mathcal{F}[g](w, t)](x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{u}^I(w) e^{-i\omega(x-at)} d\omega \\ &= u^I(x - at). \end{aligned} \quad (6)$$

Hence, we have proved so far that

$$\hat{u}(w, t) = e^{-b\omega^2 t} \hat{u}^I(w) e^{-ia\omega t} = \mathcal{F}[f](w, t) \mathcal{F}[g](w, t) = \mathcal{F}[f * g](w, t), \quad (7)$$

Then, we can finally obtain the solution to the problem,

$$\begin{aligned} u(x, t) &= \mathcal{F}^{-1}[\hat{u}(w, t)](x, t) \\ &= \mathcal{F}^{-1}[\mathcal{F}[f * g](w, t)](x, t) \\ &= [f * g](x, t). \end{aligned}$$

To conclude,

$$u(x, t) = [f * u^I(x - at)](x, t), \quad (8)$$

with  $f$  given by (6).

2. Derive an analytic expression for the solution  $U(x, t)$  of problem (2) for a general initial function  $u^I(x)$  and general stepsizes  $\Delta x, \Delta t$ , using discrete Fourier transforms.

**Solution:** CACA

## Problem 2

## Problem 3