

# Numerical Methods for PDEs

## Homework 4

Francisco Jose Castillo Carrasco

October 15, 2018

### Problem 1

- a. Show that the Jacobi spectral radius  $\mu = \cos(\pi h)$  for Laplace's equation on the unit square with second-order accurate central differences. Write-up only the 1D version. *Hint:* In 1D, set  $A = \text{tridiag}[-1 \ 2 \ -1]$ . Then the iteration matrix  $B = \frac{1}{2}\text{tridiag}[1 \ 0 \ 1]$ . Then show that  $Bv = \cos(\pi h)v$  where the 1D eigenvector

$$v = [\sin(\pi h), \sin(2\pi h), \dots, \sin(n\pi h)].$$

Note that here  $h = 1/(n+1)$ .

**Solution:** Since  $A = L + D + U = \text{diag}([-1 \ 2 \ -1])$  and we are using the Jacobi iteration,  $M = D = \text{diag}(2)$ . Hence,

$$B = M^{-1}(M - A) = \text{diag}(1/2) * \text{tridiag}([1 \ 0 \ 1]) = \frac{1}{2}\text{tridiag}([1 \ 0 \ 1])$$

We start with the first point of our domain, i.e. the first row of the product  $Bv$ ,

$$(Bv)_1 = \frac{1}{2}v_2 = \frac{1}{2}\sin(2\pi h) = \cos(\pi h)\sin(\pi h).$$

We continue with the interior points, where  $j \in [2, n-1]$ ,

$$\begin{aligned} (Bv)_j &= \frac{1}{2}(v_{j-1} + v_{j+1}) = \frac{1}{2}[\sin(j\pi h)\cos(\pi h) - \sin(\pi h)\cos(j\pi h) + \sin(j\pi h)\cos(\pi h) + \sin(\pi h)\cos(j\pi h)] \\ &= \cos(\pi h)\sin(j\pi h). \end{aligned}$$

Finally, for the last point  $j = n$ ,

$$\begin{aligned} (Bv)_n &= \frac{1}{2}v_{n-1} = \frac{1}{2}\sin((n-1)\pi h) = \frac{1}{2}\sin(n\pi h - \pi h) \\ &= \frac{1}{2}[\sin(n\pi h)\cos(\pi h) - \sin(\pi h)\cos(n\pi h)] \\ &= \frac{1}{2}[\sin(n\pi h)\cos(\pi h) + \sin(n\pi h)\cos(\pi h)] \\ &= \cos(\pi h)\sin(n\pi h), \end{aligned}$$

where we have used the following relations:

$$\begin{aligned} \sin(n\pi h) &= \sin((n+1)\pi h - \pi h) = \sin(\pi - \pi h) = \sin(\pi h) \\ \cos(n\pi h) &= \cos((n+1)\pi h - \pi h) = \cos(\pi - \pi h) = -\cos(\pi h). \end{aligned}$$

Thus, we have obtained that

$$(Bv)_j = \cos(\pi h)\sin(j\pi h),$$

which means that the eigenvalue is  $\lambda = \cos(\pi h)$  and the eigenvector  $v$  has components  $v_j = \sin(j\pi h)$ .

## Problem 2

- a. For SOR/SUR, show that  $\det\{B\} = (1 - \omega)^n$ ,  $0 < \omega < 2$ . *Hint:*  $B$  is the product of triangular matrices ( $A = L + D + U$ ):

$$B = (D + \omega L)^{-1} ((1 - \omega)D - \omega U).$$

**Solution:** It is simply computed that

$$\det\{(D + \omega L)^{-1}\} = \prod_{j=1}^n \frac{1}{d_j} = \frac{1}{\prod_{j=1}^n d_j},$$

since the elements of  $L$  are always multiplied by 0. For the same reason,

$$\det\{(1 - \omega)D - \omega U\} = \prod_{j=1}^n (1 - \omega)d_j = (1 - \omega)^n \prod_{j=1}^n d_j.$$

Hence,

$$\det B = \frac{(1 - \omega)^n \prod_{j=1}^n d_j}{\prod_{j=1}^n d_j} = (1 - \omega)^n.$$

- b. Derive the equation for the SOR  $\omega_{opt}$ , assuming Young's formula applied to the spectral radii:

$$(\lambda + \omega - 1)^2 = \lambda \omega^2 \mu^2.$$

Here  $\mu$  is the spectral radius for the Jacobi iteration method and  $\lambda$  is the spectral radius for the SOR iteration method. *Hint:* Set  $\lambda = \omega - 1$  and minimize  $\lambda$  (using the quadratic formula).

**Solution:** We start from Young's formula and we make  $\lambda = \omega - 1$  and  $\mu = \rho_J$ ,

$$(\lambda + \omega - 1)^2 = \lambda \omega^2 \mu^2$$

$$4\lambda^2 = \lambda(\lambda + 1)^2 \rho_J^2 \Rightarrow$$

$$\Rightarrow -\lambda(\rho_J^2 \lambda^2 + (2\rho_J^2 - 4)\lambda + \rho_J^2) = 0.$$

The solutions to this equation are  $\lambda = 0$  or

$$\lambda = \frac{4 - 2\rho_J^2 \pm \sqrt{(4 - 2\rho_J^2)^2 - 4\rho_J^4}}{2\rho_J^2} = \frac{4 - 2\rho_J^2 \pm 4\sqrt{1 - \rho_J^2}}{2\rho_J^2} = \frac{2 - \rho_J^2 \pm 2\sqrt{1 - \rho_J^2}}{\rho_J^2}.$$

Minimizing  $\rho_{SOR} = \max\{\lambda\} = \omega_{opt} - 1$ :

$$\rho_{SOR} = \frac{2 - \rho_J^2 - 2\sqrt{1 - \rho_J^2}}{\rho_J^2},$$

and,

$$\begin{aligned} \omega_{opt} &= \frac{2 - \rho_J^2 - 2\sqrt{1 - \rho_J^2}}{\rho_J^2} + 1 \\ &= 2 \frac{1 - \sqrt{1 - \rho_J^2}}{\rho_J^2} \end{aligned}$$

- c. Show that for Laplace's equation on the unit square, the SOR  $\lambda = (1 - \sin \pi h)/(1 + \sin \pi h)$ .

**Solution:** Using that  $\rho_J = \cos(\pi h)$ , and the result just proved,

$$\begin{aligned}\rho_{SOR} &= \frac{2 - \rho_J^2 - 2\sqrt{1 - \rho_J^2}}{\rho_J^2} \\ &= \frac{2 - \cos^2(\pi h) - 2\sqrt{1 - \cos^2(\pi h)}}{\cos^2(\pi h)} \\ &= \frac{2 - \cos^2(\pi h) - 2\sin(\pi h)}{\cos^2(\pi h)} \\ &= \frac{1 + \sin^2(\pi h) - 2\sin(\pi h)}{1 - \sin^2(\pi h)} \\ &= \frac{1 - \sin(\pi h)}{1 + \sin(\pi h)}\end{aligned}$$

### Problem 3

- a.  $2 \times 2$  SOR example. Calculate the first two iterates  $x_1$  and  $x_2$  for Jacobi, Gauss-Seidel, and SOR with  $x_0 = (0, 0)$  for  $Ax = b$  with

$$A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ -3 \end{bmatrix}$$

$$M_{SOR} = \begin{bmatrix} \frac{2}{\omega} & 0 \\ -1 & \frac{2}{\omega} \end{bmatrix}, \quad B_{SOR} = \begin{bmatrix} 1 - \omega & \frac{\omega}{2} \\ \frac{\omega}{2}(1 - \omega) & (1 - \frac{\omega}{2})^2 \end{bmatrix}$$

$$\omega_{opt} = 4(2 - \sqrt{3}) \approx 1.0718, \quad \lambda_1 = \lambda_2 = \omega_{opt} - 1 = \rho_{SOR} \approx 0.0718.$$

The exact solution is  $x = (1, -1)$ ,  $x_2^J = (3/4, -3/4)$ , and  $x_2^{GS} = (9/8, -15/16)$ . Calculate  $\|e_2^J\|_1$ ,  $\|e_2^{GS}\|_1$ , and  $\|e_2^{SOR}\|_1$ . Note that the SOR  $x_2$  is much closer to the exact solution.

**Solution:**

- **Jacobi:** We start with the Jacobi iteration,  $x^{(k+1)} = x^{(k)} - D^{-1}r^{(k)}$ , where

$$D = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \Rightarrow D^{-1} = \begin{bmatrix} 1/2 & 0 \\ 0 & 1/2 \end{bmatrix}.$$

Since  $x_0 = 0$ ,  $r_0 = b$ . Then,

$$x^{(1)} = D^{-1}b = \begin{bmatrix} 3/2 \\ -3/2 \end{bmatrix}, \quad r(1) = Ax(1) - b = \begin{bmatrix} 3/2 \\ -3/2 \end{bmatrix}.$$

The second guess is

$$x^{(2)} = x^{(1)} - D^{-1}r^{(1)} = \begin{bmatrix} 3/2 \\ -3/2 \end{bmatrix} - \begin{bmatrix} 1/2 & 0 \\ 0 & 1/2 \end{bmatrix} \begin{bmatrix} 3/2 \\ -3/2 \end{bmatrix} = \begin{bmatrix} 3/4 \\ -3/4 \end{bmatrix}.$$

Lastly, we compute the 1 error norm,  $\|e_J^{(2)}\|_1 = 1/2$ .

- **Gauss Seidel:** Now we proceed with the Gauss-Seidel iteration,  $x^{(k+1)} = M^{-1} (M - A) x^{(k)} - M^{-1}b$ , where

$$M = D + L = \begin{bmatrix} 2 & 0 \\ -1 & 2 \end{bmatrix} \Rightarrow M^{-1} = \frac{1}{4} \begin{bmatrix} 2 & 0 \\ 1 & 2 \end{bmatrix},$$

and

$$M - A = -U = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \Rightarrow M^{-1} (M - A) = \frac{1}{4} \begin{bmatrix} 0 & 2 \\ 0 & 1 \end{bmatrix}$$

Since  $x_0 = 0$ ,

$$x^{(1)} = M^{-1}b = \begin{bmatrix} -3/4 \\ -3/4 \end{bmatrix}.$$

The second guess is

$$x^{(2)} = M^{-1} (M - A) x^{(1)} - M^{-1}b = \frac{1}{4} \begin{bmatrix} 0 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -3/4 \\ -3/4 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 2 & 0 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \end{bmatrix} = \begin{bmatrix} 9/8 \\ -15/16 \end{bmatrix}.$$

Finally, we compute the 1 error norm,  $\|e_{GS}^{(2)}\|_1 = 3/16$ .

- **SOR:** Next, the SOR iteration,  $Mx^{(k+1)} = (M - A) x^{(k)} + b$ , with  $M = \frac{D}{\omega} + L$  and  $M - A = (\frac{1}{\omega} - 1)D - U$ . Therefore,

$$\begin{aligned} \left[ \frac{D}{\omega} + L \right] x^{(k+1)} &= \left[ \left( \frac{1}{\omega} - 1 \right) D - U \right] x^{(k)} + b, \\ [D + \omega L] x^{(k+1)} &= [(1 - \omega) D - U] x^{(k)} + \omega b, \\ x^{(k+1)} &= [D + \omega L]^{-1} [(1 - \omega) D - U] x^{(k)} + \omega [D + \omega L]^{-1} b. \end{aligned}$$

Note that

$$[D + \omega L] = \begin{bmatrix} 2 & 0 \\ -\omega & 2 \end{bmatrix} \Rightarrow [D + \omega L]^{-1} = \frac{1}{4} \begin{bmatrix} 2 & 0 \\ \omega & 2 \end{bmatrix},$$

and

$$[(1 - \omega)D - U] = \begin{bmatrix} 2(1 - \omega) & 1 \\ 0 & 2(1 - \omega) \end{bmatrix} \Rightarrow [D + \omega L]^{-1} [(1 - \omega)D - U] = \frac{1}{4} \begin{bmatrix} 4(1 - \omega) & 2 \\ 2\omega(1 - \omega) & 4(1 - \omega) + \omega \end{bmatrix}.$$

Since  $x_0 = 0$ ,

$$x^{(1)} = \omega [D + \omega L]^{-1} b = \begin{bmatrix} \frac{3\omega}{4} \\ \frac{3\omega^2 - 6\omega}{4} \end{bmatrix}.$$

The second guess is

$$\begin{aligned} x^{(2)} &= [D + \omega L]^{-1} [(1 - \omega) D - U] x^{(1)} + \omega [D + \omega L]^{-1} b \\ &= \frac{1}{4} \begin{bmatrix} 4(1 - \omega) & 2 \\ 2\omega(1 - \omega) & 4(1 - \omega) + \omega \end{bmatrix} \begin{bmatrix} \frac{3\omega}{4} \\ \frac{3\omega^2 - 6\omega}{4} \end{bmatrix} + \frac{\omega}{4} \begin{bmatrix} 2 & 0 \\ \omega & 2 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \end{bmatrix} \\ &= \begin{bmatrix} \frac{-9\omega(\omega-2)}{8} \\ \frac{-3\omega(7\omega^2-18\omega+16)}{16} \end{bmatrix} = \begin{bmatrix} 1.1192 \\ -0.9543 \end{bmatrix}. \end{aligned}$$

Finally, we compute the 1 error norm,  $\|e_{SOR}^{(2)}\|_1 = 0.1649$ . As expected,

$$e_{SOR}^{(2)}\|_1 < \|e_{GS}^{(2)}\|_1 < \|e_J^{(2)}\|_1.$$

## Problem 4

a. Use conjugate gradient on the steepest descent problem we did in class:

$$A = \begin{bmatrix} 4 & -2 \\ -2 & 2 \end{bmatrix}, \quad b = \begin{bmatrix} -2 \\ 2 \end{bmatrix}, \quad x_0 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad x = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

**Solution:** Since  $x_0 = 0$ ,  $r_0 = b$  and  $\beta_1 = 0$ . Then,

$$d_1 = r_0 + \beta_1 d_0 = r_0 = b,$$

and,

$$\alpha_1 = \frac{r_0^T r_0}{d_1^T A d_1} = 1/5.$$

Then new guess is

$$x_1 = x_0 + \alpha_1 d_1 = \begin{bmatrix} -2/5 \\ 2/5 \end{bmatrix},$$

and the new residual is

$$r_1 = r_0 + \alpha_1 A d_1 = \begin{bmatrix} 2/5 \\ 2/5 \end{bmatrix},$$

We repeat the process again,

$$\beta_2 = \frac{r_1^T r_1}{r_0^T r_0} = 1/25,$$

$$d_2 = r_1 + \beta_2 d_1 = \begin{bmatrix} 8/25 \\ 12/25 \end{bmatrix},$$

$$\alpha_2 = \frac{r_1^T r_1}{d_2^T A d_2} = 5/4,$$

$$x_2 = x_1 + \alpha_2 d_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

## Problem 5

- a. Compute the first two conjugate gradient iterates  $x_1$  and  $x_2$  with  $x_0 = (0, 0)$  with and without preconditioning to the solution  $x = (0, 1)$  of  $Ax = b$ :

$$A = \begin{bmatrix} 9 & 1 \\ 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad M = \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix}.$$

Calculate  $\|e_1^{CG}\|_1$  and  $\|e_1^{PCG}\|_1$ . Note that while both CG and PCG give the exact  $x$  in two steps  $x_2 = (0, 1)$ , PCG gives a much better  $x_1$ .

### Solution:

- **CG:** We start without using preconditioning. Since  $x_0 = 0$ ,  $r_0 = b$  and  $\beta_1 = 0$ . Then,

$$d_1 = r_0 + \beta_1 d_0 = r_0 = b,$$

and,

$$\alpha_1 = \frac{r_0^T r_0}{d_1^T A d_1} = 1/6.$$

Then new guess is

$$x_1 = x_0 + \alpha_1 d_1 = \begin{bmatrix} 1/6 \\ 1/6 \end{bmatrix},$$

and the new residual is

$$r_1 = r_0 + \alpha_1 A d_1 = \begin{bmatrix} -2/3 \\ 2/3 \end{bmatrix},$$

We repeat the process again,

$$\beta_2 = \frac{r_1^T r_1}{r_0^T r_0} = 4/9,$$

$$d_2 = r_1 + \beta_2 d_1 = \begin{bmatrix} -2/9 \\ 10/9 \end{bmatrix},$$

$$\alpha_2 = \frac{r_1^T r_1}{d_2^T A d_2} = 3/4,$$

$$x_2 = x_1 + \alpha_2 d_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

- **PCG:** Now we use preconditioning. Since  $x_0 = 0$ ,  $r_0 = b$  and  $\beta_1 = 0$ . We solve

$$M z_0 = r_0 \Rightarrow z_0 = \begin{bmatrix} 1/9 \\ 1 \end{bmatrix}.$$

Then,

$$d_1 = z_0 + \beta_1 d_0 = r_0 = z_0,$$

and,

$$\alpha_1 = \frac{z_0^T r_0}{d_1^T A d_1} = 5/6.$$

Then new guess is

$$x_1 = x_0 + \alpha_1 d_1 = \begin{bmatrix} 5/54 \\ 5/6 \end{bmatrix},$$

and the new residual is

$$r_1 = r_0 + \alpha_1 A d_1 = \begin{bmatrix} -2/3 \\ 2/27 \end{bmatrix},$$

We repeat the process again,

$$M z_1 = r_1 \Rightarrow z_1 = \begin{bmatrix} -2/27 \\ 2/27 \end{bmatrix},$$

$$\beta_2 = \frac{z_1^T r_1}{z_0^T r_0} = 4/81,$$

$$d_2 = z_1 + \beta_2 d_1 = \begin{bmatrix} -50/729 \\ 10/81 \end{bmatrix},$$

$$\alpha_2 = \frac{z_1^T r_1}{d_2^T A d_2} = 27/20,$$

$$x_2 = x_1 + \alpha_2 d_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

To finish we compute

$$\begin{aligned} \|e_{CG}^{(1)}\|_1 &= 1 \\ \|e_{PCG}^{(1)}\|_1 &= 0.2593. \end{aligned}$$

The *PCG* method gives indeed a much better  $x_1$ .