

Another Cone for Integration

Fred J. Hickernell

Room E1-208, Department of Applied Mathematics, Illinois Institute of Technology,
10 W. 32nd St., Chicago, IL 60616

Abstract

Keywords: adaptive, automatic, cones, function recovery, guarantee, integration, quadrature

2010 MSC: 65D05, 65D30, 65G20

1. Introduction

In [1] we considered the problem of integration and the cone of integrands

$$\mathcal{C}_\tau := \{f \in \mathcal{V}^1 : \text{Var}(f') \leq \tau \|f' - f(1) + f(0)\|_1\}, \quad (1)$$

where the total variation and the \mathcal{L}_p norms are defined as

$$\begin{aligned} \text{Var}(f) &:= \sup_{\substack{n \in \mathbb{N} \\ 0=x_0 < x_1 < \dots < x_n=1}} \sum_{i=1}^n |f(x_i) - f(x_{i-1})|, \\ \|f\|_p &:= \begin{cases} \left[\int_0^1 |f(x)|^p dx \right]^{1/p}, & 1 \leq p < \infty, \\ \sup_{0 \leq x \leq 1} |f(x)|, & p = \infty, \end{cases} \\ \mathcal{V}^k &:= \mathcal{V}^k[0, 1] = \{f \in C[0, 1] : \text{Var}(f^{(k)}) < \infty\}. \end{aligned}$$

We derived an algorithm [1, Algorithm 4] that was guaranteed for integrands in \mathcal{C}_τ . In this note we consider another algorithm and other cones.

First we recall some notation and results from [1]. For all $n \in \mathcal{I} := \{2, 3, \dots\}$ we have the linear spline:

$$x_{i,n} := x_i := \frac{i-1}{n-1}, \quad i = 1, \dots, n, \quad (2a)$$

$$\begin{aligned} A_n(f)(x) &:= (n-1) [f(x_i)(x_{i+1} - x) + f(x_{i+1})(x - x_i)] \\ &\quad \text{for } x_i \leq x \leq x_{i+1}. \end{aligned} \quad (2b)$$

Email address: hickernell@iit.edu (Fred J. Hickernell)

The cost of each function value is one and so the cost of A_n is n . The dependence of the nodes, x_i on n is often suppressed for simplicity. Integrating the linear spline gives us the trapezoidal rule based on $n - 1$ trapezoids:

$$T_n(f) := \int_0^1 A_n(f) dx = \frac{1}{2n-2} [f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)].$$

The error of the trapezoidal rule has the following upper bound:

$$\left| \int_0^1 f(x) dx - T_n(f) \right| \leq \frac{\text{Var}(f' - A_n(f'))}{8(n-1)^2} \leq \frac{\text{Var}(f')}{8(n-1)^2}. \quad (3)$$

The variation of the first derivative of f is bounded below by the variation of the first derivative of the linear spline of f :

$$\begin{aligned} \text{Var}(f') &\geq F_n(f) := \text{Var}(A_n(f')) \\ &= \begin{cases} 0, & n = 2, \\ (n-1) \sum_{i=1}^{n-2} |f(x_i) - 2f(x_{i+1}) + f(x_{i+2})|, & n \geq 3. \end{cases} \end{aligned} \quad (4)$$

2. New Cone, New Algorithm

The new cone considered here is defined as

$$\widehat{\mathcal{C}}_{\hat{\tau}} := \{f \in \mathcal{V}^1 : \text{Var}(f' - A_n(f')) \leq \hat{\tau}(n)F_n(f) \ \forall n \in \widehat{\mathcal{I}}\}, \quad (5)$$

Here $\hat{\tau} : \widehat{\mathcal{I}} \rightarrow [0, \infty)$ is some specified function that defines the cone, and $\widehat{\mathcal{I}} = \{N_{\min}, N_{\min} + 1, \dots\}$, where N_{\min} is some integer no smaller than 2.

Algorithm 1 (New Cone Adaptive Univariate Integration). Let the sequence of algorithms $\{T_n\}_{n \in \mathcal{I}}$ $\{F_n\}_{n \in \mathcal{I}}$, and $\widehat{\mathcal{C}}_{\hat{\tau}}$ be as described above. Set $i = 1$, and let $n_1 = N_{\min}$. For any error tolerance ε and input function f , do the following:

Step 1. Bound $\text{Var}(f')$ and check for convergence. Compute $F_{n_i}(f)$ in (4). Check whether n_i is large enough to satisfy the error tolerance, i.e.

$$\hat{\tau}(n_i)F_{n_i}(f) \leq 8(n_i - 1)^2 \varepsilon.$$

If this is true, then return $T_{n_i}(f)$ and terminate the algorithm.

Step 2. Increase the number of trapezoids. If the above condition is false, choose $n_{i+1} = 2n_i$, increment i , and go to Step 1.

3. The New Cone's Relationship to Other Cones

The cone defined in (5) makes Algorithm 1 work. In this section we show that it contains and is contained in other cones that might be more intuitive. One family of cones of interest is defined by replacing $F_n(f)$ by $\text{Var}(f')$ in (5):

$$\tilde{\mathcal{C}}_{\tilde{\tau}} := \{f \in \mathcal{V}^1 : \text{Var}(f' - A_n(f')) \leq \tilde{\tau}(n) \text{Var}(f'), n \in \mathcal{I}\}, \quad (6)$$

where $\tilde{\tau} : \mathcal{I} \rightarrow [0, 2]$ is non-increasing. Another family of cones is related to (1) and is defined as

$$\mathcal{C}_{\overline{\tau}} := \{f \in \mathcal{V}^1 : \text{Var}(f' - A_n(f')) \leq \overline{\tau}(n) \|f' - A_n(f')\|_1, n \in \mathcal{I}\}, \quad (7)$$

where $\overline{\tau} : \mathcal{I} \rightarrow [0, \infty]$. Under this definition $\mathcal{C}_{\overline{\tau}}$ corresponds to defining $\overline{\tau}(2) = \tau$, $\overline{\tau}(n) = \infty$ for $n > 2$.

To facilitate the comparison of $\hat{\mathcal{C}}_{\hat{\tau}}$, $\tilde{\mathcal{C}}_{\tilde{\tau}}$, and $\mathcal{C}_{\overline{\tau}}$ we note several inequalities. For all $f \in \mathcal{V}^1$,

$$\text{Var}(f') \leq \text{Var}(f' - A_n(f')) + \text{Var}(A_n(f')) = \text{Var}(f' - A_n(f')) + F_n(f), \quad (8)$$

$$\text{Var}(f' - A_n(f')) \leq \text{Var}(f') + \text{Var}(A_n(f')) = \text{Var}(f') + F_n(f). \quad (9)$$

prove the following lemma. From (4) and (9) it follows that

$$\text{Var}(f' - A_n(f')) \leq 2 \text{Var}(f') \quad \forall f \in \mathcal{V}^1, \quad (10)$$

which is why $\tilde{\tau}(n) \leq 2$ for all n . Moreover, if $\tilde{\tau}(n) = 2$ for all $n \in \mathcal{I}$, then $\tilde{\mathcal{C}}_{\tilde{\tau}} = \mathcal{V}^1$.

Theorem 1. *Given the function $\hat{\tau} : \hat{\mathcal{I}} \rightarrow [0, \infty)$, suppose that $\tilde{\tau}_j : \mathcal{I} \rightarrow [0, 2]$, $j = 1, 2$ satisfy the inequality*

$$\tilde{\tau}_1(n) \leq \frac{\hat{\tau}(n)}{1 + \hat{\tau}(n)} \leq \min(2, \hat{\tau}(n)) \leq \tilde{\tau}_2(n) \quad \forall n \in \hat{\mathcal{I}}. \quad (11)$$

It follows that $\tilde{\mathcal{C}}_{\tilde{\tau}_1} \leq \hat{\mathcal{C}}_{\hat{\tau}} \subseteq \tilde{\mathcal{C}}_{\tilde{\tau}_2}$.

Proof. First suppose that $f \in \tilde{\mathcal{C}}_{\tilde{\tau}_1}$ where $\tilde{\tau}_1$ satisfies inequality (11). It then follows that

$$\begin{aligned} \text{Var}(f' - A_n(f')) &= (1 + \hat{\tau}(n)) \text{Var}(f' - A_n(f')) - \hat{\tau}(n) \text{Var}(f' - A_n(f')) \\ &\leq [1 + \hat{\tau}(n)] \tilde{\tau}_1(n) \text{Var}(f') - \hat{\tau}(n) \text{Var}(f' - A_n(f')) \quad \text{by (6)} \\ &\leq \hat{\tau}(n) [\text{Var}(f') - \text{Var}(f' - A_n(f'))] \quad \text{by (11)} \\ &\leq \hat{\tau}(n) F_n(f) \quad \text{by (8)}. \end{aligned}$$

Thus, $\tilde{\mathcal{C}}_{\tilde{\tau}_1} \leq \hat{\mathcal{C}}_{\hat{\tau}}$. Now suppose that $f \in \hat{\mathcal{C}}_{\hat{\tau}}$. It follows by (4) and (10) that $f \in \tilde{\mathcal{C}}_{\tilde{\tau}_2}$. \square

To prove the relationship between the cones defined in (5) and (7) the following bound is needed.

Lemma 1. For all $n \in \mathcal{I}$ and all $f \in \mathcal{V}^1$ it follows that

$$F_n(f) \leq 2(n-1) \|f' - A_n(f)'\|_1. \quad (12)$$

Proof. For all $f \in \mathcal{V}^1$ we use the triangle inequality:

$$\begin{aligned} F_n(f) &= (n-1) \sum_{i=1}^{n-2} |f(x_i) - 2f(x_{i+1}) + f(x_{i+2})| \\ &\leq (n-1) \sum_{i=1}^{n-2} \left| f(x_i) - f(x_{i+1}) + \frac{f(1) - f(0)}{n-1} \right| \\ &\quad + (n-1) \sum_{i=1}^{n-2} \left| -f(x_{i+1}) + f(x_{i+2}) - \frac{f(1) - f(0)}{n-1} \right| \\ &\leq 2(n-1) \sum_{i=1}^{n-1} \left| f(x_{i+1}) - f(x_i) - \frac{f(1) - f(0)}{n-1} \right| \\ &= 2(n-1) \|A_n(f)' - f(1) + f(0)\|_1 \\ &\leq 2(n-1) \|f' - f(1) + f(0)\|_1. \end{aligned}$$

□

Theorem 2. Given the function $\hat{\tau} : \hat{\mathcal{I}} \rightarrow [0, \infty)$, suppose that $\bar{\tau}_j : \mathcal{I} \rightarrow [0, \infty)$, $j = 1, 2$ satisfy the inequality

$$\min\{2(n-1)\hat{\tau}(n) : n \in \hat{\mathcal{I}}\} \leq \bar{\tau}_j(n) \quad (13)$$

It follows that $\mathcal{C}_{\bar{\tau}_1} \leq \hat{\mathcal{C}}_{\hat{\tau}} \subseteq \mathcal{C}_{\bar{\tau}_2}$.

Proof. For all $f \in \hat{\mathcal{C}}_{\hat{\tau}}$ it follows from (12) that

$$\text{Var}(f') \leq \hat{\tau}(n) F_n(f) = 2(n-1)\hat{\tau}(n) \|f' - f(1) + f(0)\|_1 \quad \forall n \geq N_{\min}.$$

Applying the definition of τ completes the proof. □

Now we define a cone that is contained in $\hat{\mathcal{C}}_{\hat{\tau}}$. Let

$$\tilde{\mathcal{C}}_{\hat{\tau}} := \{f \in \mathcal{V}^1 : \text{Var}(f') \leq \tilde{\tau}(n) \|f' - f(1) + f(0)\|_1 \quad \forall n \geq 3\}, \quad (14)$$

Theorem 3. For any non-increasing $\hat{\tau} : \mathcal{I} \rightarrow (1, \infty)$, let

$$\tau = \min\{2(n-1)\hat{\tau}(n) : n \geq N_{\min}\}.$$

It follows that $\hat{\mathcal{C}}_{\hat{\tau}} \subseteq \mathcal{C}_{\tau}$.

Proof. For all $f \in \hat{\mathcal{C}}_{\hat{\tau}}$ it follows from (12) that

$$\text{Var}(f') \leq \hat{\tau}(n) F_n(f) = 2(n-1)\hat{\tau}(n) \|f' - f(1) + f(0)\|_1 \quad \forall n \geq N_{\min}.$$

Applying the definition of τ completes the proof. □

References

- [1] N. Clancy, Y. Ding, C. Hamilton, F.J. Hickernell, Y. Zhang, The cost of deterministic, adaptive, automatic algorithms: Cones, not balls, 2013. Submitted for publication, [arXiv.org:1303.2412](https://arxiv.org/abs/1303.2412) [math.NA].