

Construction of many irreducible Sobol' (0,2)-sequences in base $b > 2$

Victor Ostromoukhov

Université Claude Bernard Lyon 1/CNRS

victor.ostromoukhov@liris.cnrs.fr

Coauthor(s): Nicolas Bonneel, David Coeurjolly, Jean-Claude Iehl

When constructing Sobol' (t, s) -sequences from primitive polynomials [10,2,7] or irreducible polynomials [4,5], the t -value is related to the degrees of polynomials ($t = \sum_{i=0}^{s-1} (e_i - 1)$) [9,3,4]. This limits the number of pairs of Sobol' polynomials (and their direction vectors), which produce $(0, 2)$ -sequences in base b . In practice, one can construct (t^*, s) -sequences with $t^* < t$ [3,4].

In this contribution, we explore the construction of irreducible Sobol' $(0,2)$ -sequences in base $b > 2$. It is quite natural to have $(0,2)$ -sequences using irreducible polynomials of degree 1. But, under certain conditions, $(0,2)$ -sequences exist with polynomials of higher degree in base $b > 2$. First, we will demonstrate the evidence of existence of irreducible Sobol' $(0,m,2)$ -nets in bases 3, 5 and 7, up to very large m (we have explored up to $m = 1000$). This allows us to conjecture that such particular polynomials, together with specific direction vectors, form *exceptional* $(0,2)$ -sequences.

Our exploration of $(0,m,2)$ -nets is based on the study of characteristic matrices used in several recent papers [8,6,1]. For a pair of primitive or irreducible polynomials P_i and P_j , which produce generating NUT matrices M_i and M_j , the characteristic matrix C_{ij} can be expressed as $C_{ij} = M_i M_j^{-1}$. t -values of $(t,m,2)$ -nets can be found by studying the characteristic matrix C_{ij} . On the other hand, we have observed self-similar nature of the characteristic matrices produced with exceptional $(0,m,2)$ -nets that we studied. Based on this observation, we conjecture that such $(0,m,2)$ -nets are $(0,2)$ -sequences.

We try to generalise our result and establish a set of rules that would allow to find other irreducible Sobol' $(0,2)$ -sequences of higher degrees, together with associated direction vectors, in different bases.

REFERENCES

- [1] Abdalla GM Ahmed, Mikhail Skopenkov, Markus Hadwiger, and Peter Wonka. 2023. Analysis and synthesis of digital dyadic sequences. *ACM Transactions on Graphics (TOG)* 42, 6 (2023), 1–17.
- [2] Paul Bratley and Bennett L. Fox. 1988. Algorithm 659: Implementing Sobol's quasirandom sequence generator. *ACM Trans. Math. Softw.* 14, 1 (1988), 88–100.
- [3] Josef Dick and Harald Niederreiter. 2008. On the exact t -value of Niederreiter and Sobol' sequences. *Journal of Complexity* 24, 5 (2008), 572–581.
- [4] Henri Faure and Christiane Lemieux. 2016. Irreducible Sobol' sequences in prime power bases. *Acta Arithmetica* 173, 1 (2016), 59–80.
- [5] Henri Faure and Christiane Lemieux. 2019. Implementation of irreducible Sobol' sequences in prime power bases. *Mathematics and Computers in Simulation* 161 (2019), 13–22.
- [6] Roswitha Hofer and Kosuke Suzuki. 2019. A Complete Classification of Digital $(0,$

3)-Nets and Digital $(0, 2)$ -Sequences in Base 2. *Uniform distribution theory* 14, 1 (2019), 43–52.

[7] Stephen Joe and Frances Y Kuo. 2008. Constructing Sobol sequences with better two-dimensional projections. *SIAM Journal on Scientific Computing* 30, 5 (2008), 2635–2654.

[8] Hiroki Kajiura, Makoto Matsumoto, and Kosuke Suzuki. 2018. Characterization of matrices B such that (I, B, B^2) generates a digital net with t -value zero. *Finite Fields and Their Applications* 52 (2018), 289–300.

[9] Harald Niederreiter. 1992. *Random number generation and quasi-Monte Carlo methods*. SIAM.

[10] Il'ya Meerovich Sobol'. 1967. On the distribution of points in a cube and the approximate evaluation of integrals. *Zhurnal Vychislitel'noi Matematiki i Matematicheskoi Fiziki* 7, 4 (1967), 784–802.