

A


kappa
differ
ascites
hepato
creatinine
spiders
Peak FPR
100%
75%
50%
25%
0%
0.0
0.5
1.0
log10(pbaseline) − log10(α)


B


differ
ascites
hepato
spiders
kappa
creatinine
pgr
trig
er
FPR with beta varying, SE fixed at baseline
FPR with SE varying, beta fixed at baseline
0%
25%
50%
75%
100%


lung
pbc
colon
rotterdam
flchain
n
200
1000
6000


Observed peak FPR
10%
50%
90%

C



Peak FPR (%)
Peak width (decades of epsilon)
flchain
pbc
colon
rotterdam
lung
60
40
20
0
0.75
0.5
0.25
200
1000
6000
n (log scale)

D



FPR of the single coefficient
100%
75%
50%
25%
0%
0.1
1
10
100
1k
ε


colon differ (confounding route)
flchain kappa (SE route)

E


Spearman ρ = 0.93
kappa
differ
ascites
hepato
creatinine
spiders
Peak FPR
100%
75%
50%
25%
0%
0
2
4
6
|zeff|

F


differ
kappa
ascites
hepato
creatinine
spiders
Peak FPR
Spearman ρ = −0.52
100%
75%
50%
25%
0%
0.3
0.6
0.9
deff