

Equations and derivations for **ElemCo.jl**

Thomas Schraivogel and Daniel Kats

February 22, 2024

Contents

1	Introduction	2
1.1	General	2
1.2	Notation	2
2	CCSD and DCSD amplitude and Λ equations	3
2.1	Closed-shell CCSD/DCSD Lagrangian	3
2.2	Closed-shell CCSD/DSCD Lagrangian multipliers equations	4
2.3	Perturbative triples for closed-shell CCSD	6
2.4	Open-shell CCSD/DSCD Lagrangian	7
2.4.1	Spin-restricted open-shell CCSD/DSCD	8
2.5	Open-shell CCSD/DSCD Lagrangian multipliers equations	9
2.6	Perturbative triples for unrestricted CCSD	12
3	Two determinant coupled cluster	14
4	Automatically generated UCCSDT and UDC-CCSDT	18
	Bibliography	18

Chapter 1

Introduction

1.1 General

In this document we collect the equations and derivations for methods implemented in the ElemCo.jl package. The final goal is to have a document which can be used as a reference for the equations and derivations. The final equations should also be contained in the code as docstrings or copied to the corresponding Markdown files.

1.2 Notation

We use the following notation throughout the document.

The virtual orbitals are denoted by a, b, c, \dots , the occupied orbitals by i, j, k, \dots , the active (open-shell) orbitals by t, u, v, \dots , and the general orbital indices are denoted by p, q, r, s . The Einstein summation convention is used for repeated indices (repeated lower and upper indices are summed over). The α and β spin orbitals are denoted by p and \bar{p} .

The integrals are **not antisymmetrized** and denoted by v_{pq}^{rs} , where p, q, r, s are indices of orbitals, and the lower indices correspond to the creation and the upper indices to the annihilation operators in the Hamiltonian,

$$\hat{H}_N = f_p^q \{ \hat{a}_p^\dagger \hat{a}_q \}_N + \frac{1}{2} v_{pq}^{rs} \{ \hat{a}_p^\dagger \hat{a}_q^\dagger \hat{a}_s \hat{a}_r \}_N, \quad (1.1)$$

i.e., $f_p^q = \langle p | \hat{f} | q \rangle$ and $v_{pq}^{rs} = \langle pq | rs \rangle$.

Permutation operators:

$$\begin{aligned} \mathcal{P}(ab) X_{ab}^{ij} &= X_{ba}^{ij} \\ \mathcal{P}(ab \leftrightarrow ba) X_{ab}^{ij} &= X_{ba}^{ij} \end{aligned} \quad (1.2)$$

Symmetrization operators:

$$\begin{aligned} \mathcal{S}(ab) X_{ab}^{ij} &= X_{ab}^{ij} + X_{ba}^{ij} \\ \mathcal{S}(ab, ij) X_{ab}^{ij} &= X_{ab}^{ij} + X_{ba}^{ji} \end{aligned} \quad (1.3)$$

Antisymmetrization operators:

$$\begin{aligned} \mathcal{A}(ab) X_{ab}^{ij} &= X_{ab}^{ij} - X_{ba}^{ij} \\ \mathcal{A}(ab, ij) X_{ab}^{ij} &= X_{ab}^{ij} - X_{ab}^{ji} - X_{ba}^{ij} + X_{ba}^{ji} \end{aligned} \quad (1.4)$$

Chapter 2

CCSD and DCSD amplitude and Λ equations

2.1 Closed-shell CCSD/DCSD Lagrangian

The singles-dressed factorization of the closed-shell CCSD and DCSD amplitude equations roughly follows the factorization from Ref. [1]. The closed-shell CCSD and DCSD Lagrangian is given by

$$\begin{aligned}
 \mathcal{L} = & v_{kl}^{cd} \tilde{T}_{cd}^{kl} + \left(\hat{f}_k^c + f_k^c \right) T_c^k + \Lambda_{ij}^{ab} \hat{v}_{ab}^{ij} + \Lambda_{ij}^{ab} \left(\hat{v}_{kl}^{ij} + v_{kl}^{cd} T_{cd}^{ij} \right) T_{ab}^{kl} + \Lambda_{ij}^{ab} \hat{v}_{ab}^{cd} T_{cd}^{ij} \\
 & + \Lambda_{ij}^{ab} v_{kl}^{cd} T_{ad}^{kj} T_{cb}^{il} \\
 & + \Lambda_{ij}^{ab} \mathcal{S}(ab, ij) \left\{ \left(\hat{f}_a^c - 2 \times \frac{1}{2} v_{kl}^{cd} \tilde{T}_{ad}^{kl} \right) T_{cb}^{ij} - \left(\hat{f}_k^i + 2 \times \frac{1}{2} v_{kl}^{cd} \tilde{T}_{cd}^{il} \right) T_{ab}^{kj} \right. \\
 & + \left(\hat{v}_{al}^{id} + \frac{1}{2} v_{kl}^{cd} \tilde{T}_{ac}^{ik} \right) \tilde{T}_{db}^{lj} - \hat{v}_{ka}^{ic} T_{cb}^{kj} - \hat{v}_{kb}^{ic} T_{ac}^{kj} - v_{kl}^{cd} T_{da}^{ki} \left(T_{cb}^{lj} - T_{bc}^{lj} \right) \Big\} \\
 & + \Lambda_i^a \hat{f}_a^i + \Lambda_i^a \hat{f}_j^b \tilde{T}_{ab}^{ij} + \Lambda_i^a \hat{v}_{ak}^{bc} \tilde{T}_{cb}^{ki} - \Lambda_i^a \hat{v}_{jk}^{ic} \tilde{T}_{ca}^{kj}.
 \end{aligned} \tag{2.1}$$

The DCSD Lagrangian is obtained by removing terms in red.

Integrals with hats are dressed integrals, i.e. they are obtained by dressing the integrals with the singles amplitudes, and the Fock matrix is internally dressed, too, e.g.,

$$\begin{aligned}
 \hat{v}_{kl}^{id} &= v_{kl}^{id} + v_{kl}^{cd} T_c^i \\
 \hat{v}_{al}^{ij} &= v_{al}^{ij} - v_{kl}^{ij} T_a^k \\
 \hat{f}_k^c &= h_k^c + 2 \hat{v}_{kl}^{cl} - \hat{v}_{lk}^{cl} = f_k^c + (2 v_{kl}^{cd} - v_{lk}^{cd}) T_d^l.
 \end{aligned} \tag{2.2}$$

Note that only the **lower virtual** and **upper occupied** indices are dressed.

The amplitude equations can be obtained by taking the derivative of the Lagrangian with respect to the Lagrange multipliers Λ and setting the result to zero.

The most efficient version of CCSD/DCSD in **ElemCo.jl** combines the dressed factor-

ization from above with the cckext type of factorization from Ref. [2] and is given by

$$\begin{aligned}
\mathcal{L} = & v_{kl}^{cd} \tilde{T}_{cd}^{kl} + \left(\hat{f}_k^c + f_k^c \right) T_c^k + \Lambda_{ij}^{ab} \left(\hat{v}_{kl}^{ij} + v_{kl}^{cd} T_{cd}^{ij} \right) T_{ab}^{kl} + \Lambda_{ij}^{ab} K_{pq}^{ij} \delta_a^p \delta_b^q \\
& + \Lambda_{ij}^{ab} v_{kl}^{cd} T_{ad}^{kj} T_{cb}^{il} \\
& + \Lambda_{ij}^{ab} \mathcal{S}(ab, ij) \left\{ \left(\hat{f}_a^c - 2 \times \frac{1}{2} v_{kl}^{cd} \tilde{T}_{ad}^{kl} \right) T_{cb}^{ij} - \left(\hat{f}_k^i + 2 \times \frac{1}{2} v_{kl}^{cd} \tilde{T}_{cd}^{il} \right) T_{ab}^{kj} \right. \\
& + \left(\hat{v}_{al}^{id} + \frac{1}{2} v_{kl}^{cd} \tilde{T}_{ac}^{ik} \right) \tilde{T}_{db}^{lj} - \hat{v}_{ka}^{ic} T_{cb}^{kj} - \hat{v}_{kb}^{ic} T_{ac}^{kj} - v_{kl}^{cd} T_{da}^{ki} \left(T_{cb}^{lj} - T_{bc}^{lj} \right) \\
& \left. - K_{pq}^{ij} \left(\delta_k^p \delta_b^q - \frac{1}{2} \delta_k^p \delta_l^q T_b^l \right) T_a^k \right\} + \Lambda_i^a K_{pq}^{ij} (2\delta_a^p \delta_j^q - \delta_j^p \delta_a^q) \\
& - \Lambda_i^a T_a^k K_{pq}^{ij} (2\delta_k^p \delta_j^q - \delta_j^p \delta_k^q) + \Lambda_i^a \hat{h}_a^i + \Lambda_i^a \hat{f}_j^b \tilde{T}_{ab}^{ij} - \Lambda_i^a \hat{v}_{jk}^{ic} \tilde{T}_{ca}^{kj},
\end{aligned} \tag{2.3}$$

where

$$K_{pq}^{ij} = v_{pq}^{rs} \left((T_{ab}^{ij} + T_a^i T_b^j) \delta_r^a \delta_s^b + \delta_r^i T_b^j \delta_s^b + T_a^i \delta_r^a \delta_s^j + \delta_r^i \delta_s^j \right) \tag{2.4}$$

and h is the one-particle part of the Hamiltonian.

2.2 Closed-shell CCSD/DSCD Lagrangian multipliers equations

The Λ equations are obtained by taking the derivative of the Lagrangian Eq. (2.1) with respect to the amplitudes and setting the result to zero, i.e.,

$$\begin{aligned}
\frac{\partial \mathcal{L}}{\partial T_e^m} = & 2(2v_{jm}^{be} - v_{jm}^{eb}) T_b^j + 2f_m^e - \Lambda_{ij}^{eb} \hat{v}_{mb}^{ij} - \Lambda_{ij}^{ae} \hat{v}_{am}^{ij} + \Lambda_{mj}^{ab} \hat{v}_{ab}^{ej} + \Lambda_{im}^{ab} \hat{v}_{ab}^{ie} \\
& + \Lambda_{mj}^{ab} \hat{v}_{kl}^{ej} T_{ab}^{kl} + \Lambda_{im}^{ab} \hat{v}_{kl}^{ie} T_{ab}^{kl} - \Lambda_{ij}^{eb} \hat{v}_{mb}^{cd} T_{cd}^{ij} - \Lambda_{ij}^{ae} \hat{v}_{am}^{cd} T_{cd}^{ij} \\
& - \Lambda_{ij}^{eb} \hat{f}_m^d T_{db}^{ij} - \Lambda_{ij}^{ae} \hat{f}_m^d T_{ad}^{ij} - \Lambda_{mj}^{ab} \hat{f}_k^e T_{ab}^{kj} - \Lambda_{im}^{ab} \hat{f}_k^e T_{ab}^{ik} \\
& + \Lambda_{ij}^{ab} \mathcal{S}(ab, ij) \left\{ (2\hat{v}_{am}^{de} - \hat{v}_{am}^{ed}) T_{db}^{ij} - (2\hat{v}_{km}^{ie} - \hat{v}_{km}^{ei}) T_{ab}^{kj} \right\} \\
& - \Lambda_{ij}^{eb} \hat{v}_{ml}^{id} \tilde{T}_{db}^{lj} - \Lambda_{ij}^{ae} \hat{v}_{ml}^{jd} \tilde{T}_{ad}^{il} + \Lambda_{mj}^{ab} \hat{v}_{al}^{ed} \tilde{T}_{db}^{lj} + \Lambda_{im}^{ab} \hat{v}_{bl}^{ed} \tilde{T}_{ad}^{il} \\
& + \Lambda_{ij}^{eb} \hat{v}_{km}^{ic} T_{cb}^{kj} - \Lambda_{mj}^{ab} \hat{v}_{ka}^{ec} T_{cb}^{kj} + \Lambda_{ij}^{ae} \hat{v}_{km}^{jc} T_{ac}^{ik} - \Lambda_{im}^{ab} \hat{v}_{kb}^{ec} T_{ac}^{ik} \\
& + \Lambda_{ij}^{ae} \hat{v}_{km}^{ic} T_{ac}^{kj} - \Lambda_{mj}^{ab} \hat{v}_{kb}^{ec} T_{ac}^{kj} + \Lambda_{ij}^{eb} \hat{v}_{km}^{jc} T_{cb}^{ik} - \Lambda_{im}^{ab} \hat{v}_{ka}^{ec} T_{cb}^{ik} \\
& - \Lambda_i^e \hat{f}_m^i + \Lambda_m^a \hat{f}_a^e + \Lambda_i^a (2\hat{v}_{am}^{ie} - \hat{v}_{am}^{ei}) \\
& + \Lambda_i^a (2v_{jm}^{be} - v_{jm}^{eb}) \tilde{T}_{ab}^{ij} - \Lambda_i^e v_{mk}^{bc} \tilde{T}_{cb}^{ki} - \Lambda_m^a v_{jk}^{ec} \tilde{T}_{ca}^{kj}.
\end{aligned} \tag{2.5}$$

$$\begin{aligned}
\frac{\partial \mathcal{L}}{\partial T_{ef}^{mn}} = & \frac{1}{2} \mathcal{S}(ef, mn) \left[\Lambda_m^e \hat{f}_n^f \tilde{v}_{mn}^{ef} + \Lambda_{ij}^{ef} (\hat{v}_{mn}^{ij} + v_{mn}^{cd} T_{cd}^{ij}) + \Lambda_{mn}^{ab} v_{kl}^{ef} T_{ab}^{kl} + \Lambda_{mn}^{ab} \hat{v}_{ab}^{ef} \right. \\
& + \Lambda_{in}^{eb} v_{ml}^{cf} T_{cb}^{il} + \Lambda_{mj}^{af} v_{kn}^{ed} T_{ad}^{kj} \\
& + \Lambda_{mn}^{af} \mathcal{S}(af, mn) \left(\hat{f}_a^e - 2 \times \frac{1}{2} v_{kl}^{ed} \tilde{T}_{ad}^{kl} \right) - \Lambda_{ij}^{eb} \mathcal{S}(eb, ij) 2 \times \frac{1}{2} \tilde{v}_{mn}^{cf} T_{cb}^{ij} \\
& - \Lambda_{in}^{ef} \mathcal{S}(ef, in) \left(\hat{f}_m^i + 2 \times \frac{1}{2} v_{ml}^{cd} \tilde{T}_{cd}^{il} \right) - \Lambda_{mj}^{ab} \mathcal{S}(ab, mj) 2 \times \frac{1}{2} \tilde{v}_{kn}^{ef} T_{ab}^{kj} \\
& + 2 \Lambda_{in}^{af} \mathcal{S}(af, in) \left(\hat{v}_{am}^{ie} + \frac{1}{2} v_{km}^{ce} \tilde{T}_{ac}^{ik} \right) - \Lambda_{im}^{af} \mathcal{S}(af, im) \left(\hat{v}_{an}^{ie} + \frac{1}{2} v_{kn}^{ce} \tilde{T}_{ac}^{ik} \right) \\
& + 2 \Lambda_{mj}^{eb} \mathcal{S}(eb, mj) \frac{1}{2} v_{nl}^{fd} \tilde{T}_{db}^{lj} - \Lambda_{nj}^{eb} \mathcal{S}(eb, nj) \frac{1}{2} v_{ml}^{fd} \tilde{T}_{db}^{lj} \\
& - \Lambda_{in}^{af} \mathcal{S}(af, in) \hat{v}_{ma}^{ie} - \Lambda_{in}^{eb} \mathcal{S}(eb, in) \hat{v}_{mb}^{if} \\
& - \Lambda_{nj}^{fb} \mathcal{S}(fb, nj) v_{ml}^{ce} (T_{cb}^{lj} - T_{bc}^{lj}) - \Lambda_{in}^{af} \mathcal{S}(af, in) v_{km}^{ed} T_{da}^{ki} \\
& + \Lambda_{in}^{ae} \mathcal{S}(ab, ij) v_{km}^{fd} T_{da}^{ki} \\
& \left. + \mathcal{T}(mn) \left\{ \Lambda_m^e \hat{f}_n^f + \Lambda_n^a \hat{v}_{am}^{fe} - \Lambda_i^f \hat{v}_{nm}^{ie} \right\} \right], \quad (2.6)
\end{aligned}$$

with a “contravariation” operator,

$$\mathcal{T}(mn) X_{mn}^{ef} = 2X_{mn}^{ef} - X_{nm}^{ef}. \quad (2.7)$$

Now we can introduce useful intermediate quantities, related to the density matrices. The one-body reduced density matrices can be written as

$$\begin{aligned}
D_i^j &= -2\Lambda_{ik}^{cd} T_{cd}^{jk}, \\
D_a^b &= 2\Lambda_{kl}^{bc} T_{ac}^{kl}, \\
D_i^a &= \Lambda_i^a, \\
D_a^i &= \Lambda_k^c \tilde{T}_{ac}^{ik}. \quad (2.8)
\end{aligned}$$

Note that we have excluded here terms coming from the singles amplitudes. Thus, if this density matrix is used to calculate properties, the corresponding integrals should be dressed. Alternatively, one can define “dressed” density matrices which include the singles contributions,

$$\begin{aligned}
\hat{D}_i^j &= D_i^j - D_i^c T_c^j, \\
\hat{D}_a^b &= D_a^b + D_k^b T_a^k, \\
\hat{D}_i^a &= D_i^a, \\
\hat{D}_a^i &= D_a^i + 2T_a^i - D_a^c T_c^i + \hat{D}_k^i T_a^k. \quad (2.9)
\end{aligned}$$

Some parts of the two-body reduced density matrices can be written as

$$\begin{aligned}
D_{ij}^{kl} &= \Lambda_{ij}^{cd} T_{cd}^{kl} \\
D_{ib}^{aj} &= \Lambda_{ik}^{ac} \tilde{T}_{cb}^{kj} \\
\bar{D}_{ib}^{aj} &= \Lambda_{ik}^{ac} T_{cb}^{kj} + \Lambda_{ik}^{ca} T_{bc}^{kj} \quad (2.10)
\end{aligned}$$

Finally, we define the following quantities which correspond to the `cckext` factorization and a doubles-dressing of the Fock matrix,

$$\begin{aligned} K_{mn}^{rs} &= \hat{\Lambda}_{mn}^{pq} v_{pq}^{rs} \\ \hat{\Lambda}_{mn}^{pq} &= \Lambda_{mn}^{ab} \delta_a^p \delta_b^q - \Lambda_{mn}^{ab} T_a^i \delta_i^p \delta_b^q - \Lambda_{mn}^{ab} \delta_a^p T_b^j \delta_j^q + \Lambda_{mn}^{ab} T_a^i T_b^j \delta_i^p \delta_j^q \\ x_m^i &= \tilde{T}_{cd}^{il} v_{ml}^{cd} & x_a^e &= \tilde{T}_{ac}^{kl} v_{kl}^{ec} \end{aligned} \quad (2.11)$$

With these definitions, the Λ equations can be written as

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial T_e^m} &= (2v_{qm}^{pe} - v_{qm}^{ep}) \hat{D}_p^q + 2f_m^e - 2\Lambda_{ij}^{eb} \hat{v}_{mb}^{ij} + 2K_{mj}^{rs} \delta_r^e (\delta_s^j + \delta_s^b T_b^j) \\ &\quad + 2D_{mj}^{kl} \hat{v}_{kl}^{ej} - 2\Lambda_{ij}^{eb} (\hat{v}_{mb}^{cd} T_{cd}^{ij}) - D_d^e \hat{f}_m^d + D_m^k \hat{f}_k^e - 2D_{id}^{el} \hat{v}_{ml}^{id} + 2D_{md}^{al} \hat{v}_{al}^{ed} \\ &\quad + 2\bar{D}_{ic}^{ek} \hat{v}_{km}^{ic} - 2\bar{D}_{mc}^{ak} \hat{v}_{ka}^{ec} - \Lambda_i^e \hat{f}_m^i + \Lambda_m^a \hat{f}_a^e - \Lambda_i^e x_m^i - \Lambda_m^a x_a^e. \end{aligned} \quad (2.12)$$

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial T_{ef}^{mn}} &= \tilde{v}_{mn}^{ef} + \Lambda_{ij}^{ef} (\hat{v}_{mn}^{ij} + \textcolor{red}{v}_{mn}^{cd} T_{cd}^{ij}) + \textcolor{red}{D}_{mn}^{kl} v_{kl}^{ef} + K_{mn}^{rs} \delta_r^e \delta_s^f \\ &\quad + \mathcal{S}(ef, mn) \left\{ \Lambda_{mn}^{af} \left(\hat{f}_a^e - \textcolor{red}{2} \times \frac{1}{2} x_a^e \right) - \Lambda_{in}^{ef} \left(\hat{f}_m^i + \textcolor{red}{2} \times \frac{1}{2} x_m^i \right) \right. \\ &\quad + \mathcal{T}(mn) \left[\textcolor{red}{2} \times \frac{1}{4} v_{kn}^{ef} D_m^k - \textcolor{red}{2} \times \frac{1}{4} v_{mn}^{cf} D_c^e + \Lambda_{in}^{af} (\hat{v}_{am}^{ie} + v_{km}^{ce} \tilde{T}_{ac}^{ik}) \right. \\ &\quad \left. \left. + \frac{1}{2} (\Lambda_m^e \hat{f}_n^f + \Lambda_n^a \hat{v}_{am}^{fe} - \Lambda_i^f \hat{v}_{nm}^{ie}) \right] \right. \\ &\quad \left. - \Lambda_{in}^{af} \hat{v}_{ma}^{ie} - \Lambda_{in}^{eb} \hat{v}_{mb}^{if} - \textcolor{red}{D}_{nc}^{fl} v_{ml}^{ce} + \textcolor{red}{D}_{nd}^{ek} v_{km}^{fd} \right\}. \end{aligned} \quad (2.13)$$

2.3 Perturbative triples for closed-shell CCSD

The perturbative triples equations for CCSD are given by

$$\begin{aligned} E_{[T]} &= \sum_{i \leq j \leq k} p(i, j, k) K_{ijk}^{abc} X_{abc}^{ijk} \\ p(i, j, k) &= \begin{cases} 2 & i \neq j \neq k \\ 1 & i = j \oplus j = k \\ 0 & i = j = k \end{cases} \end{aligned} \quad (2.14)$$

X_{abc}^{ijk} and K_{ijk}^{abc} are calculated for the triangular set of indices $i \leq j \leq k$ (with $k = 1 : n_{occ}$),

$$\begin{aligned} K_{ijk}^{abc} &= K_{abc}^{ijk} = v_{bc}^{dk} T_{ad}^{ij} + v_{ac}^{dk} T_{db}^{ij} + v_{cb}^{dj} T_{ad}^{ik} + v_{ab}^{dj} T_{dc}^{ik} + v_{ca}^{di} T_{bd}^{jk} + v_{ba}^{di} T_{dc}^{jk} \\ &\quad - v_{lc}^{jk} T_{ba}^{li} - v_{lc}^{ik} T_{ab}^{lj} - v_{lb}^{kj} T_{ca}^{li} - v_{lb}^{ij} T_{ac}^{lk} - v_{la}^{ki} T_{cb}^{lj} - v_{la}^{ji} T_{bc}^{lk} \\ X_{abc}^{ijk} &= \frac{4K_{abc}^{ijk} - 2K_{acb}^{ijk} - 2K_{cba}^{ijk} - 2K_{bac}^{ijk} + K_{cab}^{ijk} + K_{bca}^{ijk}}{\epsilon_i + \epsilon_j + \epsilon_k - \epsilon_a - \epsilon_b - \epsilon_c} \end{aligned} \quad (2.15)$$

The (T) correction contains additionally the following terms,

$$\begin{aligned} E_{(T)} &= E_{[T]} + \sum_{i \leq j \leq k} p(i, j, k) \left[v_{jk}^{bc} X_{abc}^{ijk} T_i^{\dagger a} + v_{ik}^{ac} X_{abc}^{ijk} T_j^{\dagger b} + v_{ij}^{ab} X_{abc}^{ijk} T_k^{\dagger c} \right. \\ &\quad \left. + T_{jk}^{\dagger bc} X_{abc}^{ijk} f_i^a + T_{ik}^{\dagger ac} X_{abc}^{ijk} f_j^b + T_{ij}^{\dagger ab} X_{abc}^{ijk} f_k^c \right]. \end{aligned} \quad (2.16)$$

In case of $\Lambda\text{CCSD}(\mathbf{T})$ K_{ijk}^{abc} is different from K_{abc}^{ijk} and is calculated using the Lagrange multipliers,

$$K_{ijk}^{abc} = v_{dk}^{bc} \bar{\Lambda}_{ij}^{ad} + v_{dk}^{ac} \bar{\Lambda}_{ij}^{db} + v_{dj}^{cb} \bar{\Lambda}_{ik}^{ad} + v_{dj}^{ab} \bar{\Lambda}_{ik}^{dc} + v_{di}^{ca} \bar{\Lambda}_{jk}^{bd} + v_{di}^{ba} \bar{\Lambda}_{jk}^{dc} \\ - v_{jk}^{lc} \bar{\Lambda}_{li}^{ba} - v_{ik}^{lc} \bar{\Lambda}_{lj}^{ab} - v_{kj}^{lb} \bar{\Lambda}_{li}^{ca} - v_{ij}^{lb} \bar{\Lambda}_{lk}^{ac} - v_{ki}^{la} \bar{\Lambda}_{lj}^{cb} - v_{ji}^{la} \bar{\Lambda}_{lk}^{bc}, \quad (2.17)$$

where $\bar{\Lambda}_{ij}^{ab}$ are the covariant Lagrange multipliers,

$$\bar{\Lambda}_{ij}^{ab} = \frac{2}{3} \Lambda_{ij}^{ab} + \frac{1}{3} \Lambda_{ij}^{ba}. \quad (2.18)$$

Additionally, the conjugate-transposed amplitudes in Eq. (2.16) are replaced by the covariant Lagrange multipliers $\bar{\Lambda}_{ij}^{ab}$ and $\bar{\Lambda}_i^a = \frac{1}{2} \Lambda_i^a$.

2.4 Open-shell CCSD/DCSD Lagrangian

The factorization of the open-shell CCSD/DCSD amplitude equations roughly follows the factorization of the closed-shell equations, Sec. 2.1. The open-shell CCSD and DCSD Lagrangian – i.e., spin dependent – is given by

$$\mathcal{L} = \mathcal{L}_\alpha + \mathcal{L}_\beta + \mathcal{L}_{\alpha\beta}, \quad (2.19)$$

$$\mathcal{L}_\alpha = \frac{1}{2} \left[v_{kl}^{cd} T_{cd}^{kl} + \left(\hat{f}_k^c + f_k^c \right) T_c^k \right] + \frac{1}{4} \Lambda_{ij}^{ab} \left(\hat{v}_{ab}^{ij} - \hat{v}_{ab}^{ji} \right) + \frac{1}{4} \Lambda_{ij}^{ab} \left(\hat{v}_{kl}^{ij} + \frac{1}{2} v_{kl}^{cd} T_{cd}^{ij} \right) T_{ab}^{kl} \\ + \frac{1}{4} \Lambda_{ij}^{ab} \hat{v}_{ab}^{cd} T_{cd}^{ij} + \frac{1}{4} \Lambda_{ij}^{ab} \mathcal{S}(ab, ij) \left\{ \hat{x}_a^c T_{cb}^{ij} - \hat{x}_k^i T_{ab}^{kj} \right\} \\ + \frac{1}{4} \Lambda_{ij}^{ab} \mathcal{A}(ab; ij) \left\{ \left(\hat{v}_{al}^{id} - \hat{v}_{al}^{di} + \bar{x}_{al}^{id} \right) T_{db}^{lj} + \left(\hat{v}_{al}^{i\bar{d}} + x_{al}^{i\bar{d}} \right) T_{bd}^{j\bar{l}} \right\} \\ + \Lambda_i^a \hat{v}_{al}^{cd} T_{cd}^{il} + \Lambda_i^a \hat{v}_{al}^{c\bar{d}} T_{cd}^{i\bar{l}} - \Lambda_i^a \hat{v}_{jk}^{ic} T_{ac}^{jk} - \Lambda_i^a \hat{v}_{jk}^{i\bar{c}} T_{ac}^{j\bar{k}} \\ + \Lambda_i^a \hat{f}_a^i + \Lambda_i^a \hat{f}_j^b T_{ab}^{ij} + \Lambda_i^a \hat{f}_{\bar{j}}^{\bar{b}} T_{a\bar{b}}^{i\bar{j}}, \quad (2.20)$$

or using the cckext factorization,

$$\mathcal{L}_\alpha = \frac{1}{2} \left[v_{kl}^{cd} T_{cd}^{kl} + \left(\hat{f}_k^c + f_k^c \right) T_c^k \right] + \frac{1}{4} \Lambda_{ij}^{ab} \left(\hat{v}_{kl}^{ij} + \frac{1}{2} v_{kl}^{cd} T_{cd}^{ij} \right) T_{ab}^{kl} \\ + \frac{1}{4} \Lambda_{ij}^{ab} K_{pq}^{ij} \left(\delta_a^p - \delta_k^p T_a^k \right) \left(\delta_b^q - \delta_l^q T_b^l \right) + \frac{1}{4} \Lambda_{ij}^{ab} \mathcal{S}(ab, ij) \left\{ \hat{x}_a^c T_{cb}^{ij} - \hat{x}_k^i T_{ab}^{kj} \right\} \\ + \frac{1}{4} \Lambda_{ij}^{ab} \mathcal{A}(ab; ij) \left\{ \left(\hat{v}_{al}^{id} - \hat{v}_{al}^{di} + \bar{x}_{al}^{id} \right) T_{db}^{lj} + \left(\hat{v}_{al}^{i\bar{d}} + x_{al}^{i\bar{d}} \right) T_{bd}^{j\bar{l}} \right\} \\ + \Lambda_i^a \left(K_{pq}^{ij} \delta_j^q + K_{p\bar{q}}^{i\bar{j}} \delta_{\bar{j}}^{\bar{q}} \right) \left(\delta_a^p - \delta_k^p T_a^k \right) - \Lambda_i^a \hat{v}_{jk}^{ic} T_{ac}^{jk} - \Lambda_i^a \hat{v}_{jk}^{i\bar{c}} T_{ac}^{j\bar{k}} \\ + \Lambda_i^a \hat{h}_a^i + \Lambda_i^a \hat{f}_j^b T_{ab}^{ij} + \Lambda_i^a \hat{f}_{\bar{j}}^{\bar{b}} T_{a\bar{b}}^{i\bar{j}}, \quad (2.21)$$

\mathcal{L}_β is obtained from \mathcal{L}_α by flipping the spins;

$$\mathcal{L}_{\alpha\beta} = v_{kl}^{c\bar{d}} T_{cd}^{k\bar{l}} + \Lambda_{ij}^{a\bar{b}} \hat{v}_{ab}^{i\bar{j}} + \Lambda_{ij}^{a\bar{b}} \left(\hat{v}_{kl}^{i\bar{j}} + v_{kl}^{c\bar{d}} T_{cd}^{i\bar{j}} \right) T_{ab}^{k\bar{l}} + \Lambda_{ij}^{a\bar{b}} \hat{v}_{ab}^{c\bar{d}} T_{cd}^{i\bar{j}} \\ + \Lambda_{ij}^{a\bar{b}} \left\{ \hat{x}_a^c T_{cb}^{i\bar{j}} + \hat{x}_b^{\bar{d}} T_{ad}^{i\bar{j}} - \hat{x}_k^i T_{ab}^{k\bar{j}} - \hat{x}_{\bar{l}}^{\bar{j}} T_{a\bar{b}}^{i\bar{l}} \right\} \\ + \Lambda_{ij}^{a\bar{b}} \left\{ \left(\hat{v}_{al}^{id} - \hat{v}_{al}^{di} + x_{al}^{id} + \bar{x}_{al}^{id} \right) T_{db}^{lj} + \left(\hat{v}_{bl}^{j\bar{d}} - \hat{v}_{bl}^{\bar{d}j} + 2x_{bl}^{j\bar{d}} \right) T_{ad}^{i\bar{l}} \right. \\ \left. + \left(\hat{v}_{al}^{i\bar{d}} + v_{kl}^{c\bar{d}} T_{ac}^{ik} \right) T_{db}^{l\bar{j}} + \hat{v}_{lb}^{d\bar{j}} T_{ad}^{i\bar{l}} - \hat{v}_{ak}^{c\bar{j}} T_{cb}^{i\bar{k}} - \left(\hat{v}_{kb}^{i\bar{d}} - v_{kl}^{c\bar{d}} T_{cb}^{i\bar{l}} \right) T_{ad}^{k\bar{j}} \right\}, \quad (2.22)$$

or using the `ckext` factorization,

$$\begin{aligned}
\mathcal{L}_{\alpha\beta} = & v_{kl}^{cd} T_{cd}^{kl} + \Lambda_{ij}^{ab} \left(\hat{v}_{kl}^{ij} + \textcolor{red}{v}_{kl}^{cd} \textcolor{red}{T}_{cd}^{ij} \right) T_{ab}^{kl} + \Lambda_{ij}^{ab} K_{pq}^{ij} \left(\delta_a^p - \delta_k^p T_a^k \right) \left(\delta_b^q - \delta_l^q T_b^l \right) \\
& + \Lambda_{ij}^{ab} \left\{ \hat{x}_a^c T_{cb}^{ij} + \hat{x}_b^d T_{ad}^{ij} - \hat{x}_k^i T_{ab}^{kj} - \hat{x}_l^j T_{ab}^{il} \right\} \\
& + \Lambda_{ij}^{ab} \left\{ \left(\hat{v}_{al}^{id} - \hat{v}_{al}^{di} + x_{al}^{id} + \bar{x}_{al}^{id} \right) T_{db}^{ij} + \left(\hat{v}_{bl}^{jd} - \hat{v}_{bl}^{dj} + 2x_{bl}^{jd} \right) T_{ad}^{ij} \right. \\
& \left. + \left(\hat{v}_{al}^{id} + v_{kl}^{cd} T_{ac}^{ik} \right) T_{db}^{ij} + \hat{v}_{lb}^{dj} T_{ad}^{il} - \hat{v}_{ak}^{cj} T_{cb}^{ik} - \left(\hat{v}_{kb}^{id} - \textcolor{red}{v}_{kl}^{cd} \textcolor{red}{T}_{cb}^{il} \right) T_{ad}^{kj} \right\}.
\end{aligned} \tag{2.23}$$

The intermediate quantities are defined as follows,

$$\begin{aligned}
K_{pq}^{ij} &= v_{pq}^{rs} D_{rs}^{ij} & K_{p\bar{q}}^{i\bar{j}} &= v_{p\bar{q}}^{r\bar{s}} D_{r\bar{s}}^{i\bar{j}} \\
D_{rs}^{ij} &= \left(T_{ab}^{ij} + T_a^i T_b^j - T_b^i T_a^j \right) \delta_r^a \delta_s^b + \mathcal{A}(ij; rs) \delta_r^i T_b^j \delta_s^b + \delta_r^i \delta_s^j - \delta_s^i \delta_r^j \\
D_{r\bar{s}}^{i\bar{j}} &= \left(T_{ab}^{i\bar{j}} + T_a^i T_b^{\bar{j}} \right) \delta_r^a \delta_{\bar{s}}^b + \delta_r^i T_b^{\bar{j}} \delta_{\bar{s}}^b + T_a^i \delta_s^a \delta_{\bar{s}}^{\bar{j}} + \delta_r^i \delta_{\bar{s}}^{\bar{j}} \\
x_{al}^{id} &= \frac{1}{2} T_{ac}^{ik} \left(v_{kl}^{cd} - \textcolor{red}{v}_{kl}^{dc} \right) \\
\bar{x}_{al}^{id} &= x_{al}^{id} + T_{a\bar{c}}^{i\bar{k}} v_{l\bar{k}}^{d\bar{c}} \\
x_{a\bar{l}}^{i\bar{d}} &= \frac{1}{2} T_{a\bar{c}}^{i\bar{k}} \left(v_{kl}^{\bar{c}\bar{d}} - \textcolor{red}{v}_{kl}^{\bar{d}\bar{c}} \right) \\
\hat{x}_k^i &= \hat{f}_k^i + \textcolor{red}{2} \times \frac{1}{2} \left(v_{kl}^{cd} T_{cd}^{il} + v_{kl}^{cd} T_{cd}^{il} \right) \\
\hat{x}_a^c &= \hat{f}_a^c - \textcolor{red}{2} \times \frac{1}{2} \left(v_{kl}^{cd} T_{ad}^{kl} + v_{kl}^{cd} T_{ad}^{kl} \right)
\end{aligned} \tag{2.24}$$

2.4.1 Spin-restricted open-shell CCSD/DSCD

The spin-restricted versions `rccsd` and `rdcsd` are obtained through spin-projection of the residuals and amplitudes from the spin-dependent equations in each iteration. [3, 4]

In this section we use the following notation:

$$\begin{aligned}
{}^{\alpha\alpha} T_{ab}^{ij} &= T_{ab}^{ij}, \\
{}^{\beta\beta} T_{ab}^{ij} &= \bar{T}_{ab}^{ij}, \\
{}^{\alpha\beta} T_{ab}^{ij} &= \bar{T}_{ab}^{ij},
\end{aligned} \tag{2.25}$$

and the spin-projected amplitudes are denoted by a bar, e.g., ${}^{\alpha\beta} \bar{T}_{ab}^{ij}$. Moreover, the indices i, j, \dots run in the following part of the section over the closed-shell part of occupied orbitals, a, b, \dots over the (doubly) virtual orbitals, and t, u, \dots over the singly occupied (or singly-virtual) orbitals.

The “closed-shell” part of spin-projected $\alpha\beta$ amplitudes is given by

$${}^{\alpha\beta} \bar{T}_{ab}^{ij} = \frac{1}{6} \left({}^{\alpha\alpha} T_{ab}^{ij} + {}^{\beta\beta} T_{ab}^{ij} + 2 {}^{\alpha\beta} T_{ab}^{ij} + {}^{\alpha\beta} T_{ba}^{ij} + 2 {}^{\alpha\beta} T_{ba}^{ji} + {}^{\alpha\beta} T_{ab}^{ji} \right) \tag{2.26}$$

The “open-shell” part of spin-projected $\alpha\beta$ amplitudes is given by

$$\begin{aligned}
{}^{\alpha\beta} \bar{T}_{at}^{ij} &= \frac{1}{3} \left({}^{\beta\beta} T_{at}^{ij} + 2 {}^{\alpha\beta} T_{at}^{ij} + {}^{\alpha\beta} T_{at}^{ji} \right) \\
{}^{\alpha\beta} \bar{T}_{ab}^{tj} &= \frac{1}{3} \left({}^{\alpha\alpha} T_{ab}^{tj} + 2 {}^{\alpha\beta} T_{ab}^{tj} + {}^{\alpha\beta} T_{ba}^{tj} \right) \\
{}^{\alpha\beta} \bar{T}_{au}^{tj} &= {}^{\alpha\beta} T_{au}^{tj} + \frac{\delta_{tu}}{2m_s + 2} \left({}^{\beta\beta} T_a^j - {}^{\alpha\alpha} T_a^j - {}^{\alpha\beta} T_{av}^{uj} \right)
\end{aligned} \tag{2.27}$$

The projection corrections for the remaining amplitudes are defined in terms of the new spin-projected $\alpha\beta$ amplitudes as follows. For singles amplitudes,

$$\begin{aligned}\alpha\bar{T}_a^i &= \frac{1}{2} \left(\alpha T_a^i + \beta T_a^i - \alpha\beta\bar{T}_{av}^i \right), \\ \beta\bar{T}_a^i &= \frac{1}{2} \left(\alpha T_a^i + \beta T_a^i + \alpha\beta\bar{T}_{av}^i \right), \\ \alpha\bar{T}_a^t &= \alpha T_a^t \quad \text{and} \quad \beta\bar{T}_t^i = \beta T_t^i.\end{aligned}\tag{2.28}$$

For the $\alpha\alpha$ and $\beta\beta$ amplitudes,

$$\begin{aligned}\sigma\sigma\bar{T}_{ab}^{ij} &= \alpha\beta\bar{T}_{ab}^{ij} - \alpha\beta\bar{T}_{ba}^{ij}, \\ \alpha\alpha\bar{T}_{ab}^{tj} &= \alpha\alpha\bar{T}_{ba}^{tj} = \alpha\beta\bar{T}_{ab}^{tj} - \alpha\beta\bar{T}_{ba}^{tj}, \\ \beta\beta\bar{T}_{at}^{ij} &= \beta\beta\bar{T}_{ta}^{ji} = \alpha\beta\bar{T}_{at}^{ij} - \alpha\beta\bar{T}_{at}^{ji}, \\ \alpha\alpha\bar{T}_{ab}^{tu} &= \alpha\alpha T_{ab}^{tu} \quad \text{and} \quad \beta\beta\bar{T}_{tu}^{ij} = \beta\beta T_{tu}^{ij}.\end{aligned}\tag{2.29}$$

2.5 Open-shell CCSD/DSCD Lagrangian multipliers equations

The Lagrange multipliers equations for the open-shell CCSD/DSCD Lagrangian can be obtained by taking the derivatives with respect to the amplitudes and setting them to zero.

$$\begin{aligned}\frac{\partial\mathcal{L}_\alpha}{\partial T_e^m} &= (v_{km}^{ce} - v_{km}^{ec}) T_c^k + \frac{1}{2} v_{m\bar{k}}^{e\bar{c}} T_{\bar{c}}^{\bar{k}} + f_m^e - \Lambda_{ij}^{eb} \hat{v}_{mb}^{ij} + \Lambda_{mj}^{ab} \hat{v}_{ab}^{ej} + \frac{1}{4} \Lambda_{mj}^{ab} \hat{v}_{kl}^{ej} T_{ab}^{kl} \\ &+ \frac{1}{4} \Lambda_{im}^{ab} \hat{v}_{kl}^{ie} T_{ab}^{kl} - \frac{1}{4} \Lambda_{ij}^{eb} \hat{v}_{mb}^{cd} T_{cd}^{ij} - \frac{1}{4} \Lambda_{ij}^{ae} \hat{v}_{am}^{cd} T_{cd}^{ij} - \frac{1}{2} \Lambda_{ij}^{eb} \hat{f}_m^c T_{cb}^{ij} - \frac{1}{2} \Lambda_{mj}^{ab} \hat{f}_k^e T_{ab}^{kj} \\ &+ \frac{1}{2} \Lambda_{ij}^{ab} \left\{ (\hat{v}_{am}^{ce} - \hat{v}_{am}^{ec}) T_{cb}^{ij} - (\hat{v}_{km}^{ie} - \hat{v}_{mk}^{ie}) T_{ab}^{kj} \right\} \\ &- \Lambda_{ij}^{eb} \hat{v}_{ml}^{id} T_{db}^{lj} + \Lambda_{mj}^{ab} \hat{v}_{al}^{ed} T_{db}^{lj} + \Lambda_{ij}^{eb} \hat{v}_{ml}^{di} T_{db}^{lj} \\ &- \Lambda_{mj}^{ab} \hat{v}_{al}^{de} T_{db}^{lj} - \Lambda_{ij}^{eb} \hat{v}_{ml}^{i\bar{d}} T_{b\bar{d}}^{j\bar{l}} + \Lambda_{mj}^{ab} \hat{v}_{al}^{e\bar{d}} T_{b\bar{d}}^{j\bar{l}} \\ &- \Lambda_i^e \hat{v}_{ml}^{cd} T_{cd}^{i\bar{l}} - \Lambda_i^e \hat{v}_{ml}^{c\bar{d}} T_{cd}^{i\bar{l}} - \Lambda_m^a \hat{v}_{jk}^{ec} T_{ac}^{jk} - \Lambda_m^a \hat{v}_{jk}^{e\bar{c}} T_{a\bar{c}}^{j\bar{k}} \\ &- \Lambda_i^e \hat{f}_m^i + \Lambda_m^a \hat{f}_a^e + \Lambda_i^a (\hat{v}_{am}^{ie} - \hat{v}_{am}^{ei}) + \Lambda_i^a (v_{jm}^{be} - v_{jm}^{eb}) T_{ab}^{ij} + \Lambda_i^a v_{mj}^{e\bar{b}} T_{ab}^{i\bar{j}}\end{aligned}\tag{2.30}$$

$$\begin{aligned}\frac{\partial\mathcal{L}_\beta}{\partial T_e^m} &= \frac{1}{2} v_{m\bar{k}}^{e\bar{c}} T_{\bar{c}}^{\bar{k}} + \frac{1}{2} \Lambda_{ij}^{\bar{a}\bar{b}} \left\{ \hat{v}_{m\bar{a}}^{e\bar{c}} T_{\bar{c}\bar{b}}^{i\bar{j}} - \hat{v}_{m\bar{k}}^{e\bar{i}} T_{\bar{a}\bar{b}}^{k\bar{j}} \right\} \\ &+ \Lambda_{\bar{i}}^{\bar{a}} \hat{v}_{m\bar{a}}^{e\bar{i}} + \Lambda_{\bar{i}}^{\bar{a}} v_{m\bar{j}}^{e\bar{b}} T_{\bar{a}\bar{b}}^{i\bar{j}} + \Lambda_{\bar{i}}^{\bar{a}} (v_{mj}^{eb} - v_{mj}^{be}) T_{\bar{b}\bar{a}}^{j\bar{i}},\end{aligned}\tag{2.31}$$

$$\begin{aligned}\frac{\partial\mathcal{L}_{\alpha\beta}}{\partial T_e^m} &= -\Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{m\bar{b}}^{i\bar{j}} + \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{ab}^{e\bar{j}} + \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{kl}^{e\bar{j}} T_{ab}^{kl} - \Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{mb}^{cd} T_{cd}^{i\bar{j}} \\ &- \Lambda_{ij}^{\bar{e}\bar{b}} \hat{f}_m^c T_{cb}^{i\bar{j}} - \Lambda_{mj}^{\bar{a}\bar{b}} \hat{f}_k^e T_{ab}^{k\bar{j}} \\ &+ \Lambda_{ij}^{\bar{a}\bar{b}} \left\{ (\hat{v}_{am}^{ce} - \hat{v}_{am}^{ec}) T_{cb}^{i\bar{j}} + \hat{v}_{mb}^{e\bar{d}} T_{ad}^{i\bar{j}} - (\hat{v}_{km}^{ie} - \hat{v}_{mk}^{ie}) T_{ab}^{k\bar{j}} - \hat{v}_{ml}^{e\bar{j}} T_{ab}^{l\bar{i}} \right\} \\ &- \Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{ml}^{id} T_{db}^{l\bar{j}} + \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{al}^{ed} T_{db}^{l\bar{j}} + \Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{ml}^{di} T_{db}^{l\bar{j}} - \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{al}^{de} T_{db}^{l\bar{j}} \\ &- \Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{ml}^{i\bar{d}} T_{b\bar{d}}^{j\bar{l}} + \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{al}^{e\bar{d}} T_{b\bar{d}}^{j\bar{l}} + \Lambda_{ij}^{\bar{e}\bar{b}} \hat{v}_{mk}^{e\bar{j}} T_{cb}^{i\bar{k}} - \Lambda_{mj}^{\bar{a}\bar{b}} \hat{v}_{kb}^{e\bar{d}} T_{ad}^{k\bar{j}},\end{aligned}\tag{2.32}$$

The corresponding equations for the derivatives with respect to the β amplitudes are obtained by flipping the spins.

Derivatives with respect to doubles amplitudes are given by

$$4 \frac{\partial \mathcal{L}_\alpha}{\partial T_{ef}^{mn}} = \mathcal{A}(ef; mn) \left[\frac{1}{2} v_{mn}^{ef} + \frac{1}{4} \Lambda_{ij}^{ef} \left(\hat{v}_{mn}^{ij} + \frac{1}{2} v_{mn}^{cd} T_{cd}^{ij} \right) + \frac{1}{8} \Lambda_{mn}^{ab} v_{kl}^{ef} T_{ab}^{kl} \right. \\ + \frac{1}{4} \Lambda_{mn}^{ab} \hat{v}_{ab}^{ef} + \frac{1}{2} \Lambda_{mn}^{af} \hat{x}_a^e - \frac{1}{2} \Lambda_{in}^{ef} \hat{x}_m^i - 2 \times \frac{1}{4} \Lambda_{ij}^{eb} v_{mn}^{cf} T_{cb}^{ij} - 2 \times \frac{1}{4} \Lambda_{mj}^{ab} v_{kn}^{ef} T_{ab}^{kj} \\ + \Lambda_{in}^{af} (\hat{v}_{am}^{ie} - \hat{v}_{am}^{ei} + \bar{x}_{am}^{ie}) + \frac{1}{2} \Lambda_{mj}^{eb} (v_{nl}^{fd} - v_{nl}^{df}) T_{db}^{lj} \\ \left. + \Lambda_m^a \hat{v}_{an}^{ef} - \Lambda_i^e \hat{v}_{mn}^{if} + \Lambda_m^e \hat{f}_n^f \right], \quad (2.33)$$

$$\frac{\partial \mathcal{L}_\beta}{\partial T_{ef}^{mn}} = 0, \quad (2.34)$$

$$4 \frac{\partial \mathcal{L}_{\alpha\beta}}{\partial T_{ef}^{mn}} = \mathcal{A}(ef; mn) \left[-2 \times \frac{1}{2} \Lambda_{ij}^{e\bar{b}} v_{mn}^{cf} T_{cb}^{i\bar{j}} - 2 \times \frac{1}{2} \Lambda_{mj}^{a\bar{b}} v_{kn}^{ef} T_{ab}^{k\bar{j}} \right. \\ \left. + \Lambda_{m\bar{j}}^{e\bar{b}} (v_{nl}^{fd} - v_{nl}^{df}) T_{db}^{l\bar{j}} + \Lambda_{m\bar{j}}^{e\bar{b}} v_{nl}^{f\bar{d}} T_{db}^{l\bar{j}} + \Lambda_{m\bar{j}}^{e\bar{b}} \hat{v}_{n\bar{b}}^{f\bar{j}} \right], \quad (2.35)$$

$$\frac{\partial \mathcal{L}_\alpha}{\partial T_{ef}^{m\bar{n}}} = -2 \times \frac{1}{4} \Lambda_{ij}^{eb} v_{m\bar{n}}^{c\bar{f}} T_{cb}^{ij} - 2 \times \frac{1}{4} \Lambda_{mj}^{ab} v_{k\bar{n}}^{e\bar{f}} T_{ab}^{kj} \\ + \Lambda_{mj}^{eb} v_{l\bar{n}}^{d\bar{f}} T_{db}^{lj} + \Lambda_{im}^{ae} (\hat{v}_{a\bar{n}}^{i\bar{f}} + x_{a\bar{n}}^{i\bar{f}}) + \frac{1}{2} \Lambda_{mj}^{eb} (v_{n\bar{l}}^{f\bar{d}} - v_{n\bar{l}}^{d\bar{f}}) T_{bd}^{j\bar{l}} \\ + \Lambda_m^a \hat{v}_{a\bar{n}}^{e\bar{f}} - \Lambda_i^e \hat{v}_{m\bar{n}}^{i\bar{f}} + \Lambda_m^e \hat{f}_{\bar{n}}^{\bar{f}}, \quad (2.36)$$

$$\frac{\partial \mathcal{L}_\beta}{\partial T_{ef}^{m\bar{n}}} = -2 \times \frac{1}{4} \Lambda_{ij}^{f\bar{b}} v_{m\bar{n}}^{e\bar{c}} T_{cb}^{i\bar{j}} - 2 \times \frac{1}{4} \Lambda_{n\bar{j}}^{a\bar{b}} v_{m\bar{k}}^{e\bar{f}} T_{ab}^{k\bar{j}} \\ + \Lambda_{n\bar{j}}^{f\bar{b}} v_{m\bar{l}}^{e\bar{d}} T_{db}^{l\bar{j}} + \Lambda_{i\bar{n}}^{a\bar{f}} (\hat{v}_{m\bar{a}}^{e\bar{i}} + x_{a\bar{m}}^{i\bar{e}}) + \frac{1}{2} \Lambda_{n\bar{j}}^{f\bar{b}} (v_{m\bar{l}}^{ed} - v_{m\bar{l}}^{de}) T_{db}^{l\bar{j}} \\ + \Lambda_{n\bar{j}}^{a\bar{f}} \hat{v}_{a\bar{m}}^{e\bar{f}} - \Lambda_{i\bar{n}}^{f\bar{b}} \hat{v}_{n\bar{m}}^{i\bar{e}} + \Lambda_{n\bar{j}}^{f\bar{b}} \hat{f}_m^e, \quad (2.37)$$

$$\frac{\partial \mathcal{L}_{\alpha\beta}}{\partial T_{ef}^{m\bar{n}}} = v_{m\bar{n}}^{e\bar{f}} + \Lambda_{ij}^{e\bar{f}} \left(\hat{v}_{m\bar{n}}^{ij} + v_{m\bar{n}}^{cd} T_{cd}^{ij} \right) + \Lambda_{m\bar{n}}^{ab} v_{kl}^{e\bar{f}} T_{ab}^{kl} + \Lambda_{m\bar{n}}^{ab} \hat{v}_{ab}^{e\bar{f}} \\ + \Lambda_{m\bar{n}}^{a\bar{f}} \hat{x}_a^e - 2 \times \frac{1}{2} \Lambda_{ij}^{e\bar{b}} v_{m\bar{n}}^{c\bar{f}} T_{cb}^{i\bar{j}} + \Lambda_{m\bar{n}}^{e\bar{b}} \hat{x}_b^{\bar{f}} - 2 \times \frac{1}{2} \Lambda_{ij}^{a\bar{f}} v_{m\bar{n}}^{e\bar{d}} T_{ad}^{i\bar{j}} \\ - \Lambda_{i\bar{n}}^{e\bar{f}} \hat{x}_m^i - 2 \times \frac{1}{2} \Lambda_{mj}^{a\bar{b}} v_{k\bar{n}}^{e\bar{f}} T_{ab}^{k\bar{j}} - \Lambda_{mj}^{e\bar{f}} \hat{x}_{\bar{n}}^{\bar{j}} - 2 \times \frac{1}{2} \Lambda_{i\bar{n}}^{a\bar{b}} v_{m\bar{l}}^{e\bar{f}} T_{ab}^{l\bar{i}} \\ + \Lambda_{i\bar{n}}^{a\bar{f}} (\hat{v}_{am}^{ie} - \hat{v}_{am}^{ei} + x_{am}^{ie} + \bar{x}_{am}^{ie}) + \Lambda_{m\bar{j}}^{e\bar{b}} v_{l\bar{n}}^{d\bar{f}} T_{db}^{l\bar{j}} + \Lambda_{m\bar{j}}^{e\bar{b}} (\hat{v}_{b\bar{n}}^{j\bar{f}} - \hat{v}_{b\bar{n}}^{f\bar{j}} + 2x_{b\bar{n}}^{j\bar{f}}) \\ - \Lambda_{m\bar{j}}^{a\bar{f}} \hat{v}_{a\bar{n}}^{e\bar{j}} - \Lambda_{i\bar{n}}^{e\bar{b}} (\hat{v}_{m\bar{b}}^{i\bar{f}} - v_{m\bar{l}}^{c\bar{f}} T_{cb}^{i\bar{l}}) + \Lambda_{m\bar{j}}^{a\bar{f}} v_{k\bar{n}}^{e\bar{d}} T_{ad}^{k\bar{j}}, \quad (2.38)$$

and the derivatives with respect to the $\beta\beta$ amplitudes are obtained by flipping the spins.

The one-body reduced density matrix (without singles contributions) is given by

$$D_i^j = -\frac{1}{2} \Lambda_{ik}^{cd} T_{cd}^{jk} - \Lambda_{ik}^{c\bar{d}} T_{c\bar{d}}^{j\bar{k}} \\ D_a^b = \frac{1}{2} \Lambda_{kl}^{bc} T_{ac}^{kl} + \Lambda_{kl}^{b\bar{c}} T_{a\bar{c}}^{k\bar{l}} \\ D_i^a = \Lambda_i^a \\ D_a^i = \Lambda_k^c T_{ac}^{ik} + \Lambda_k^{\bar{c}} T_{a\bar{c}}^{i\bar{k}} \quad (2.39)$$

and the β 1RDM is obtained by flipping the spins.

The full (dressed) one-body reduced density matrix is given by

$$\begin{aligned}\hat{D}_i^j &= D_i^j - D_i^c T_c^j, \\ \hat{D}_a^b &= D_a^b + D_k^b T_a^k, \\ \hat{D}_i^a &= D_i^a, \\ \hat{D}_a^i &= D_a^i + T_a^i - D_a^c T_c^i + \hat{D}_k^i T_a^k.\end{aligned}\tag{2.40}$$

Additionally, we define intermediates related to the two-body reduced density matrix,

$$\begin{aligned}D_{ij}^{kl} &= \frac{1}{2} \Lambda_{ij}^{cd} T_{cd}^{kl} \\ D_{i\bar{j}}^{k\bar{l}} &= \Lambda_{i\bar{j}}^{cd} T_{cd}^{k\bar{l}} \\ D_{ib}^{aj} &= \Lambda_{ik}^{ac} T_{bc}^{jk} + \Lambda_{ik}^{a\bar{c}} T_{b\bar{c}}^{j\bar{k}} \\ \bar{D}_{i\bar{b}}^{a\bar{j}} &= \Lambda_{ik}^{ac} T_{c\bar{b}}^{k\bar{j}} + \Lambda_{ik}^{a\bar{c}} T_{b\bar{c}}^{j\bar{k}}\end{aligned}\tag{2.41}$$

and doubles-dressed Fock matrix,

$$\begin{aligned}x_k^i &= v_{kl}^{cd} T_{cd}^{il} + v_{k\bar{l}}^{cd} T_{cd}^{i\bar{l}} \\ x_a^c &= v_{kl}^{cd} T_{ad}^{kl} + v_{k\bar{l}}^{cd} T_{a\bar{d}}^{kl}.\end{aligned}\tag{2.42}$$

The intermediates for the `cckext` factorization are given by

$$\begin{aligned}K_{mn}^{rs} &= \hat{\Lambda}_{mn}^{pq} v_{pq}^{rs} \\ \hat{\Lambda}_{mn}^{pq} &= \Lambda_{mn}^{ab} \delta_a^p \delta_b^q - \Lambda_{mn}^{ab} T_a^i \delta_i^p \delta_b^q - \Lambda_{mn}^{ab} \delta_a^p T_b^j \delta_j^q + \Lambda_{mn}^{ab} T_a^i T_b^j \delta_i^p \delta_j^q.\end{aligned}\tag{2.43}$$

$K_{m\bar{n}}^{r\bar{s}}$ and $K_{\bar{m}\bar{n}}^{r\bar{s}}$ are obtained by flipping the spins.

Finally, we define useful intermediates which can be precalculated and reused in the equations,

$$\begin{aligned}\hat{y}_{am}^{ie} &= \hat{v}_{am}^{ie} - \hat{v}_{am}^{ei} + \bar{x}_{am}^{ie} + x_{am}^{ie} \\ \hat{y}_{\bar{b}n}^{\bar{j}f} &= v_{n\bar{l}}^{f\bar{d}} T_{\bar{d}\bar{b}}^{\bar{l}j} + \hat{v}_{n\bar{b}}^{f\bar{j}} + 2x_{\bar{b}n}^{\bar{j}f}\end{aligned}\tag{2.44}$$

With these intermediates the equations for the α Lagrange multipliers are given by

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial T_e^m} &= (v_{pm}^{qe} - v_{pm}^{eq}) \hat{D}_q^p + v_{m\bar{p}}^{e\bar{q}} \hat{D}_{\bar{q}}^{\bar{p}} + f_m^e - \Lambda_{ij}^{eb} \hat{v}_{mb}^{ij} - \Lambda_{ij}^{e\bar{b}} \hat{v}_{m\bar{b}}^{i\bar{j}} \\ &+ K_{mj}^{rs} \delta_r^e (\delta_s^j + \delta_s^b T_b^j) + K_{m\bar{j}}^{r\bar{s}} \delta_r^e (\delta_{\bar{s}}^{\bar{j}} + \delta_{\bar{s}}^{\bar{b}} T_{\bar{b}}^{\bar{j}}) + D_{mj}^{kl} \hat{v}_{kl}^{ej} + D_{m\bar{j}}^{k\bar{l}} \hat{v}_{k\bar{l}}^{e\bar{j}} \\ &- \frac{1}{2} \Lambda_{ij}^{eb} (\hat{v}_{mb}^{cd} T_{cd}^{ij}) - \Lambda_{i\bar{j}}^{e\bar{b}} (\hat{v}_{m\bar{b}}^{cd} T_{cd}^{i\bar{j}}) - D_c^e \hat{f}_m^c + D_m^k \hat{f}_k^e \\ &+ D_{id}^{el} (\hat{v}_{ml}^{di} - \hat{v}_{lm}^{di}) + D_{md}^{al} (\hat{v}_{al}^{ed} - \hat{v}_{al}^{de}) - \bar{D}_{i\bar{d}}^{e\bar{l}} \hat{v}_{m\bar{l}}^{i\bar{d}} + \bar{D}_{m\bar{d}}^{a\bar{l}} \hat{v}_{a\bar{l}}^{e\bar{d}} \\ &+ \Lambda_{i\bar{j}}^{e\bar{b}} \hat{v}_{m\bar{k}}^{c\bar{j}} T_{c\bar{b}}^{i\bar{k}} - \Lambda_{m\bar{j}}^{a\bar{b}} \hat{v}_{k\bar{b}}^{ed} T_{a\bar{d}}^{k\bar{j}} \\ &- \Lambda_i^e \hat{f}_m^i + \Lambda_m^a \hat{f}_a^e - \Lambda_i^e x_m^i - \Lambda_m^a x_a^e\end{aligned}\tag{2.45}$$

$$\begin{aligned}
4 \frac{\partial \mathcal{L}}{\partial T_{ef}^{mn}} = & v_{mn}^{ef} - v_{nm}^{ef} + \Lambda_{ij}^{ef} \left(\hat{v}_{mn}^{ij} + \frac{1}{2} v_{mn}^{cd} T_{cd}^{ij} \right) + D_{mn}^{kl} v_{kl}^{ef} + K_{mn}^{rs} \delta_r^e \delta_s^f \\
& + \mathcal{S}(ef, mn) \left\{ \Lambda_{mn}^{af} \hat{x}_a^e - \Lambda_{in}^{ef} \hat{x}_m^i \right\} \\
& + \mathcal{A}(ef; mn) \left\{ 2 \times \frac{1}{2} D_m^k v_{kn}^{ef} - 2 \times \frac{1}{2} D_c^e v_{mn}^{cf} + \Lambda_{in}^{af} \hat{y}_{am}^{ie} + \Lambda_{mj}^{e\bar{b}} \hat{y}_{bn}^{\bar{j}f} \right. \\
& \left. + \Lambda_m^a \hat{v}_{an}^{ef} - \Lambda_i^e \hat{v}_{mn}^{if} + \Lambda_m^e \hat{f}_n^f \right\}
\end{aligned} \tag{2.46}$$

The equations for the β Lagrange multipliers are obtained by flipping the spins. The equations for the $\alpha\beta$ Lagrange multipliers are given by

$$\begin{aligned}
\frac{\partial \mathcal{L}}{\partial T_{ef}^{m\bar{n}}} = & v_{m\bar{n}}^{e\bar{f}} + \Lambda_{ij}^{e\bar{f}} \left(\hat{v}_{m\bar{n}}^{i\bar{j}} + v_{m\bar{n}}^{c\bar{d}} T_{c\bar{d}}^{i\bar{j}} \right) + D_{m\bar{n}}^{k\bar{l}} v_{k\bar{l}}^{e\bar{f}} + K_{m\bar{n}}^{r\bar{s}} \delta_r^e \delta_{\bar{s}}^{\bar{f}} \\
& + \Lambda_{m\bar{n}}^{a\bar{f}} \hat{x}_a^e + \Lambda_{m\bar{n}}^{e\bar{b}} \hat{x}_{\bar{b}}^{\bar{f}} - \Lambda_{i\bar{n}}^{e\bar{f}} \hat{x}_m^i - \Lambda_{m\bar{j}}^{e\bar{f}} \hat{x}_{\bar{n}}^{\bar{j}} \\
& + 2 \times \frac{1}{2} \left(D_m^k v_{k\bar{n}}^{e\bar{f}} + D_{\bar{n}}^{\bar{k}} v_{m\bar{k}}^{e\bar{f}} - D_c^e v_{m\bar{n}}^{c\bar{f}} - D_{\bar{c}}^{\bar{e}} v_{m\bar{n}}^{e\bar{c}} \right) \\
& + \Lambda_{i\bar{n}}^{a\bar{f}} \hat{y}_{am}^{ie} + \Lambda_{m\bar{j}}^{e\bar{b}} \hat{y}_{b\bar{n}}^{\bar{j}f} + \Lambda_{im}^{ae} \hat{y}_{a\bar{n}}^{i\bar{f}} + \Lambda_{i\bar{n}}^{a\bar{f}} \hat{y}_{am}^{ie} \\
& - \Lambda_{m\bar{j}}^{a\bar{f}} \left(\hat{v}_{a\bar{n}}^{e\bar{j}} - v_{k\bar{n}}^{e\bar{d}} T_{a\bar{d}}^{k\bar{j}} \right) - \Lambda_{i\bar{n}}^{e\bar{b}} \left(\hat{v}_{m\bar{b}}^{i\bar{f}} - v_{m\bar{l}}^{c\bar{f}} T_{c\bar{b}}^{i\bar{l}} \right) \\
& + \Lambda_m^a \hat{v}_{a\bar{n}}^{e\bar{f}} - \Lambda_i^e \hat{v}_{m\bar{n}}^{i\bar{f}} + \Lambda_m^e \hat{f}_{\bar{n}}^{\bar{f}} + \Lambda_{\bar{n}}^{\bar{a}} \hat{v}_{am}^{\bar{f}e} - \Lambda_i^{\bar{f}} \hat{v}_{\bar{n}m}^{\bar{i}e} + \Lambda_{\bar{n}}^{\bar{f}} \hat{f}_m^e.
\end{aligned} \tag{2.47}$$

2.6 Perturbative triples for unrestricted CCSD

The perturbative triples equations for unrestricted CCSD are given by

$$E_{[T]} = \frac{1}{6} \sum_{i < j < k} K_{ijk}^{abc} T_{abc}^{ijk} + \frac{1}{6} \sum_{\bar{i} < \bar{j} < \bar{k}} K_{i\bar{j}\bar{k}}^{\bar{a}\bar{b}\bar{c}} T_{\bar{a}\bar{b}\bar{c}}^{\bar{i}\bar{j}\bar{k}} + \frac{1}{2} \sum_{i < j; \bar{k}} K_{ijk}^{ab\bar{c}} X_{ab\bar{c}}^{ijk} + \frac{1}{2} \sum_{\bar{i} < \bar{j}; k} K_{i\bar{j}k}^{\bar{a}\bar{b}c} X_{\bar{a}\bar{b}c}^{\bar{i}\bar{j}k}. \tag{2.48}$$

K_{ijk}^{abc} and T_{abc}^{ijk} (and the all- β -counterparts) are calculated for a triangular set of indices $i < j < k$ (with $k = 3 : n_\alpha$),

$$\begin{aligned}
K_{ijk}^{abc} = K_{abc}^{ijk} = & \mathcal{A}(abc) \left\{ v_{bc}^{dk} T_{ad}^{ij} + v_{cb}^{dj} T_{ad}^{ik} + v_{ba}^{di} T_{dc}^{jk} - \frac{1}{2} \left(\bar{v}_{cl}^{kj} T_{ab}^{il} + \bar{v}_{cl}^{ki} T_{ab}^{lj} + \bar{v}_{al}^{ij} T_{cb}^{kl} \right) \right\}, \\
\bar{v}_{al}^{ij} = & v_{al}^{ij} - v_{al}^j, \\
T_{abc}^{ijk} = & \frac{K_{abc}^{ijk}}{\epsilon_i + \epsilon_j + \epsilon_k - \epsilon_a - \epsilon_b - \epsilon_c}
\end{aligned} \tag{2.49}$$

$K_{i\bar{j}\bar{k}}^{\bar{a}\bar{b}\bar{c}}$ and $T_{\bar{a}\bar{b}\bar{c}}^{\bar{i}\bar{j}\bar{k}}$ (and the spin-flipped counterparts) are calculated for a triangular set of first two indices $i < j$ (with $j = 2 : n_\alpha$),

$$\begin{aligned}
K_{i\bar{j}\bar{k}}^{\bar{a}\bar{b}\bar{c}} = K_{\bar{a}\bar{b}\bar{c}}^{i\bar{j}\bar{k}} = & \mathcal{A}(ab) \left\{ v_{\bar{b}\bar{c}}^{d\bar{k}} T_{ad}^{i\bar{j}} + v_{ba}^{di} T_{d\bar{c}}^{j\bar{k}} + v_{ab}^{dj} T_{d\bar{c}}^{i\bar{k}} + v_{a\bar{c}}^{i\bar{d}} T_{b\bar{d}}^{j\bar{k}} + v_{b\bar{c}}^{j\bar{d}} T_{ad}^{i\bar{k}} - \bar{v}_{al}^{ij} T_{b\bar{c}}^{l\bar{k}} \right. \\
& \left. - v_{b\bar{l}}^{j\bar{k}} T_{a\bar{c}}^{i\bar{l}} - v_{a\bar{l}}^{i\bar{k}} T_{b\bar{c}}^{j\bar{l}} \right\} - v_{l\bar{c}}^{j\bar{k}} T_{ab}^{il} - v_{l\bar{c}}^{i\bar{k}} T_{ab}^{lj}, \\
T_{\bar{a}\bar{b}\bar{c}}^{i\bar{j}\bar{k}} = & \frac{K_{\bar{a}\bar{b}\bar{c}}^{i\bar{j}\bar{k}}}{\epsilon_i + \epsilon_j + \epsilon_{\bar{k}} - \epsilon_a - \epsilon_b - \epsilon_{\bar{c}}}
\end{aligned} \tag{2.50}$$

The (T) correction contains additionally the following terms,

$$\begin{aligned}
E_{(T)} = E_{[T]} &+ \sum_{i < j < k} \left[v_{jk}^{bc} T_{abc}^{ijk} T_i^{\dagger a} + v_{ik}^{ac} T_{abc}^{ijk} T_j^{\dagger b} + v_{ij}^{ab} T_{abc}^{ijk} T_k^{\dagger c} \right. \\
&+ \frac{1}{2} \left(T_{jk}^{\dagger bc} T_{abc}^{ijk} f_i^a + T_{ik}^{\dagger ac} T_{abc}^{ijk} f_j^b + T_{ij}^{\dagger ab} T_{abc}^{ijk} f_k^c \right) \Big] \\
&+ \sum_{i < j; \bar{k}} \left[v_{j\bar{k}}^{b\bar{c}} T_{ab\bar{c}}^{ij\bar{k}} T_i^{\dagger a} + v_{i\bar{k}}^{a\bar{c}} T_{ab\bar{c}}^{ij\bar{k}} T_j^{\dagger b} + v_{ij}^{ab} T_{ab\bar{c}}^{ij\bar{k}} T_{\bar{k}}^{\dagger \bar{c}} \right. \\
&+ T_{j\bar{k}}^{\dagger b\bar{c}} T_{ab\bar{c}}^{ij\bar{k}} f_i^a + T_{i\bar{k}}^{\dagger a\bar{c}} T_{ab\bar{c}}^{ij\bar{k}} f_j^b + \frac{1}{2} T_{ij}^{\dagger ab} T_{ab\bar{c}}^{ij\bar{k}} f_{\bar{k}}^{\bar{c}} \Big] \\
&+ \text{spin-flipped terms.}
\end{aligned} \tag{2.51}$$

In case of $\Lambda\text{UCCSD}(\mathbf{T})$, K_{ijk}^{abc} and T_{abc}^{ijk} etc are different from K_{ijk}^{abc} and T_{abc}^{ijk} and can be calculated by replacing amplitudes with Lagrange multipliers (and integrals with transpose integrals) in the above equations,

$$\begin{aligned}
K_{ijk}^{abc} &= \mathcal{A}(abc) \left\{ v_{dk}^{bc} \Lambda_{ij}^{ad} + v_{dj}^{cb} \Lambda_{ik}^{ad} + v_{di}^{ba} \Lambda_{jk}^{dc} - \frac{1}{2} (\bar{v}_{kj}^{cl} \Lambda_{il}^{ab} + \bar{v}_{ki}^{cl} \Lambda_{lj}^{ab} + \bar{v}_{ij}^{al} \Lambda_{kl}^{cb}) \right\}, \\
K_{ij\bar{k}}^{ab\bar{c}} &= \mathcal{A}(ab) \left\{ v_{d\bar{k}}^{b\bar{c}} \Lambda_{ij}^{ad} + v_{di}^{ba} \Lambda_{j\bar{k}}^{d\bar{c}} + v_{dj}^{ab} \Lambda_{i\bar{k}}^{d\bar{c}} + v_{id}^{a\bar{c}} \Lambda_{j\bar{k}}^{b\bar{d}} + v_{jd}^{b\bar{c}} \Lambda_{i\bar{k}}^{a\bar{d}} - \bar{v}_{ij}^{al} \Lambda_{l\bar{k}}^{b\bar{c}} \right. \\
&\quad \left. - v_{j\bar{k}}^{b\bar{l}} \Lambda_{i\bar{l}}^{a\bar{c}} - v_{i\bar{k}}^{a\bar{l}} \Lambda_{j\bar{l}}^{b\bar{c}} \right\} - v_{j\bar{k}}^{l\bar{c}} \Lambda_{il}^{ab} - v_{i\bar{k}}^{l\bar{c}} \Lambda_{lj}^{ab},
\end{aligned} \tag{2.52}$$

and the conjugate-transpose of the amplitudes in Eq. (2.51) are replaced with the Lagrange multipliers.

Chapter 3

Two determinant coupled cluster

Amplitudes are normal ordered with respect to the formal reference with two active orbitals t and \bar{u} . The occupied (i, j, \dots) and virtual (a, b, \dots) spaces do not contain the active orbitals. The equations follow the equations presented in Ref.[5]. Differences because of fixed typos or other reasons are coloured **blue**. Terms we have added to ensure energy invariance with respect to the reference choice and which are not explicitly listed in Ref.[5] are coloured **magenta**. Terms we have added to ensure proper antisymmetry and which are not explicitly listed in Ref.[5] are coloured **green**. IAS terms, which are terms including the all internal singles, are coloured **brown**.

$$\begin{aligned} R_a^i &= \langle {}^A\Phi_a^i | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle_C - \left(\langle {}^A\Phi_a^i | e^{\hat{T}_B} | {}^B\Phi \rangle \langle {}^B\Phi | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle \right)_C \\ &\equiv \langle {}^A\Phi_a^i | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle_C + M_a^i W = 0, \end{aligned} \quad (3.1)$$

$$\begin{aligned} R_{ab}^{ij} &= \langle {}^A\Phi_{ab}^{ij} | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle_C - \left(\langle {}^A\Phi_{ab}^{ij} | e^{\hat{T}_B} | {}^B\Phi \rangle \langle {}^B\Phi | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle \right)_C \\ &\quad - \mathcal{A}(ij; ab) \left[\langle {}^A\Phi_a^i | e^{\hat{T}_A} | {}^A\Phi \rangle \left(\langle {}^A\Phi_b^j | e^{\hat{T}_B} | {}^B\Phi \rangle \langle {}^B\Phi | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle \right)_C \right. \\ &\quad \left. - \hat{R}(ia) \langle {}^B\Phi_a^i | e^{\hat{T}_B} | {}^B\Phi \rangle \left(\langle {}^A\Phi_b^j | e^{\hat{T}_B} | {}^B\Phi \rangle \langle {}^B\Phi | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle \right)_C \right] \\ &\equiv \langle {}^A\Phi_{ab}^{ij} | \hat{H}_N e^{\hat{T}_A} | {}^A\Phi \rangle_C + M_{ab}^{ij} W = 0, \end{aligned} \quad (3.2)$$

The operator $\hat{R}(ia)$ excludes the active orbitals from the corresponding orbital spaces. The following intermediates are used:

$$\tau_a^i = T_a^i - T_{\bar{a}}^{\bar{i}}, \quad (3.3)$$

$$\tau_{\bar{a}}^{\bar{i}} = T_{\bar{a}}^{\bar{i}} - T_a^i, \quad (3.4)$$

$$\tau_{ab}^{ij} = T_{ab}^{ij} + T_a^i T_b^j. \quad (3.5)$$

The singles M tensor is built as follows,

$$M_u^i = T_{ut}^{t\bar{i}}, \quad (3.6)$$

$$M_u^i = T_u^t T_{\bar{t}}^{\bar{i}}, \quad (3.7)$$

$$M_a^t = -T_{u\bar{a}}^{t\bar{u}}, \quad (3.8)$$

$$M_a^t = -T_u^t T_{\bar{a}}^{\bar{u}}, \quad (3.9)$$

$$M_a^i = T_{\bar{a}}^{\bar{u}} T_{ut}^{t\bar{i}} + T_{\bar{t}}^{\bar{i}} T_{u\bar{a}}^{t\bar{u}}, \quad (3.10)$$

$$M_a^i = T_u^t T_{\bar{a}\bar{t}}^{\bar{u}\bar{i}}, \quad (3.11)$$

$$M_a^i = T_u^t T_{\bar{a}}^{\bar{u}} T_{\bar{t}}^{\bar{i}}, \quad (3.12)$$

$$M_u^t = T_u^t. \quad (3.13)$$

$$M_{\bar{t}}^{\bar{i}} = T_{ut}^{i\bar{u}}, \quad (3.14)$$

$$M_{\bar{t}}^{\bar{i}} = T_{\bar{t}}^{\bar{u}} T_u^i, \quad (3.15)$$

$$M_{\bar{a}}^{\bar{u}} = -T_{a\bar{t}}^{t\bar{u}}, \quad (3.16)$$

$$M_{\bar{a}}^{\bar{u}} = -T_{\bar{t}}^{\bar{u}} T_a^t, \quad (3.17)$$

$$M_{\bar{a}}^{\bar{i}} = T_a^t T_{ut}^{i\bar{u}} + T_u^i T_{a\bar{t}}^{t\bar{u}}, \quad (3.18)$$

$$M_{\bar{a}}^{\bar{i}} = T_{\bar{t}}^{\bar{u}} T_{au}^{ti}, \quad (3.19)$$

$$M_{\bar{a}}^{\bar{i}} = T_{\bar{t}}^{\bar{u}} T_a^t T_u^i, \quad (3.20)$$

$$M_{\bar{t}}^{\bar{u}} = T_{\bar{t}}^{\bar{u}}. \quad (3.21)$$

The all alpha part of the doubles part is built as follows,

$$M_{ua}^{ij} = \mathcal{A}(ij) \tau_a^i T_{ut}^{t\bar{j}} + \mathcal{A}(ij) T_t^{\bar{i}} T_{u\bar{a}}^{t\bar{j}}, \quad (3.22)$$

$$M_{ua}^{ij} = T_u^t T_{t\bar{a}}^{\bar{i}\bar{j}}, \quad (3.23)$$

$$M_{au}^{ij} = -\mathcal{A}(ij) \tau_a^i T_{ut}^{t\bar{j}} - \mathcal{A}(ij) T_t^{\bar{i}} T_{u\bar{a}}^{t\bar{j}}, \quad (3.24)$$

$$M_{au}^{ij} = -T_u^t T_{t\bar{a}}^{\bar{i}\bar{j}}, \quad (3.25)$$

$$M_{ab}^{ti} = \mathcal{A}(ab) \tau_b^i T_{u\bar{a}}^{t\bar{u}} + \mathcal{A}(ab) T_b^{\bar{u}} T_{u\bar{a}}^{t\bar{i}}, \quad (3.26)$$

$$M_{ab}^{ti} = T_u^t T_{b\bar{a}}^{\bar{u}\bar{i}}, \quad (3.27)$$

$$M_{ab}^{it} = -\mathcal{A}(ab) \tau_b^i T_{u\bar{a}}^{t\bar{u}} - \mathcal{A}(ab) T_b^{\bar{u}} T_{u\bar{a}}^{t\bar{i}}, \quad (3.28)$$

$$M_{ab}^{it} = -T_u^t T_{b\bar{a}}^{\bar{u}\bar{i}}, \quad (3.29)$$

$$\begin{aligned} M_{ab}^{ij} = & -\mathcal{A}(ij; ab) \tau_b^j \left(T_{\bar{a}}^{\bar{u}} T_{ut}^{t\bar{i}} + T_t^{\bar{i}} T_{u\bar{a}}^{t\bar{u}} \right) - \mathcal{A}(\bar{i}\bar{j}; \bar{a}\bar{b}) \left(T_{t\bar{a}}^{\bar{u}\bar{i}} T_{ub}^{t\bar{j}} \right) \\ & - \mathcal{A}(\bar{i}\bar{j}) T_{\bar{a}\bar{b}}^{\bar{u}\bar{i}} T_{ut}^{t\bar{j}} - \mathcal{A}(ab) T_{t\bar{a}}^{\bar{i}\bar{j}} T_{ub}^{t\bar{u}} - \mathcal{A}(\bar{i}\bar{j}; \bar{a}\bar{b}) T_{\bar{a}}^{\bar{u}} T_t^{\bar{j}} T_{ub}^{t\bar{i}}, \end{aligned} \quad (3.30)$$

$$\begin{aligned} M_{ab}^{ij} = & -\mathcal{A}(\bar{i}\bar{j}; \bar{a}\bar{b}) \tau_{\bar{a}}^{\bar{i}} \left(T_{\bar{t}}^{\bar{u}} T_{ub}^{t\bar{j}} + T_u^t T_{t\bar{b}}^{\bar{u}\bar{j}} \right) \\ & - \mathcal{A}(\bar{i}\bar{j}) T_u^t T_{\bar{t}}^{\bar{j}} T_{\bar{a}\bar{b}}^{\bar{u}\bar{i}} - \mathcal{A}(\bar{a}\bar{b}) T_u^t T_{\bar{b}}^{\bar{u}} T_{t\bar{a}}^{\bar{i}\bar{j}}, \end{aligned} \quad (3.31)$$

$$M_{au}^{it} = -T_{u\bar{a}}^{t\bar{i}}, \quad (3.32)$$

$$M_{au}^{ti} = T_{u\bar{a}}^{t\bar{i}}, \quad (3.33)$$

$$M_{ua}^{it} = T_{u\bar{a}}^{t\bar{i}}, \quad (3.34)$$

$$M_{ua}^{ti} = -T_{u\bar{a}}^{t\bar{i}}, \quad (3.35)$$

$$M_{au}^{it} = -\tau_{\bar{a}}^{\bar{i}} T_u^t, \quad (3.36)$$

$$M_{ua}^{it} = +\tau_{\bar{a}}^{\bar{i}} T_u^t, \quad (3.37)$$

$$M_{au}^{ti} = +\tau_{\bar{a}}^{\bar{i}} T_u^t, \quad (3.38)$$

$$M_{ua}^{ti} = -\tau_{\bar{a}}^{\bar{i}} T_u^t. \quad (3.39)$$

The all beta part of the doubles M tensor is obtained from the all alpha part analogously to the presented singles M tensor.

The alpha beta part is calculated as follows,

$$M_{u\bar{a}}^{j\bar{j}} = -\tau_{\bar{a}}^{\bar{j}} T_{u\bar{t}}^{t\bar{i}} - T_u^j T_{a\bar{t}}^{t\bar{i}} - T_a^t \tau_{u\bar{t}}^{j\bar{i}} - T_{\bar{t}}^{\bar{i}} T_{a\bar{u}}^{tj}, \quad (3.40)$$

$$M_{u\bar{a}}^{i\bar{j}} = T_u^t T_{a\bar{t}}^{j\bar{i}}, \quad (3.41)$$

$$M_{a\bar{t}}^{j\bar{i}} = -\tau_a^j T_{u\bar{t}}^{i\bar{u}} - T_{\bar{t}}^{\bar{j}} T_{u\bar{a}}^{i\bar{u}} - T_{\bar{a}}^{\bar{u}} \tau_{u\bar{t}}^{j\bar{i}} - T_u^i T_{\bar{t}\bar{a}}^{j\bar{u}}, \quad (3.42)$$

$$M_{a\bar{t}}^{j\bar{i}} = T_{\bar{t}}^{\bar{u}} T_{u\bar{a}}^{i\bar{j}}, \quad (3.43)$$

$$M_{a\bar{b}}^{t\bar{i}} = \tau_{\bar{b}}^{\bar{i}} T_{u\bar{a}}^{t\bar{u}} + T_b^t T_{u\bar{a}}^{i\bar{u}} + T_u^i \tau_{b\bar{a}}^{t\bar{u}} + T_{\bar{a}}^{\bar{u}} T_{ub}^{it}, \quad (3.44)$$

$$M_{a\bar{b}}^{t\bar{i}} = -T_u^t T_{b\bar{a}}^{i\bar{u}}, \quad (3.45)$$

$$M_{b\bar{a}}^{i\bar{u}} = \tau_b^i T_{a\bar{t}}^{t\bar{u}} + T_b^{\bar{u}} T_{a\bar{t}}^{t\bar{i}} + T_{\bar{t}}^{\bar{i}} \tau_{ab}^{t\bar{u}} + T_a^t T_{\bar{t}\bar{b}}^{i\bar{u}}, \quad (3.46)$$

$$M_{b\bar{a}}^{i\bar{u}} = -T_{\bar{t}}^{\bar{u}} T_{ab}^{t\bar{i}}, \quad (3.47)$$

$$\begin{aligned} M_{ab}^{i\bar{j}} = & -\tau_{\bar{b}}^{\bar{j}} (T_{\bar{a}}^{\bar{u}} T_{u\bar{t}}^{t\bar{i}} + T_{\bar{t}}^{\bar{i}} T_{u\bar{a}}^{t\bar{u}}) - \tau_a^i (T_{u\bar{t}}^{j\bar{u}} T_b^t + T_{b\bar{t}}^{t\bar{u}} T_u^j) \\ & + T_{\bar{t}}^{\bar{i}} T_{\bar{a}}^{\bar{u}} T_{ub}^{tj} + T_b^t T_u^j T_{\bar{t}\bar{a}}^{i\bar{u}} - T_u^j T_{\bar{a}}^{\bar{u}} T_{b\bar{t}}^{t\bar{i}} - T_b^t T_{\bar{t}}^{\bar{i}} T_{u\bar{a}}^{j\bar{u}} \\ & - \tau_{u\bar{t}}^{j\bar{i}} T_{b\bar{a}}^{t\bar{u}} + T_{u\bar{t}}^{j\bar{u}} T_{b\bar{a}}^{t\bar{i}} + T_{u\bar{t}}^{t\bar{i}} T_{b\bar{a}}^{j\bar{u}} + T_{b\bar{t}}^{t\bar{u}} T_{u\bar{a}}^{j\bar{i}} \\ & + T_{u\bar{a}}^{t\bar{u}} T_{b\bar{t}}^{j\bar{i}} - T_{ub}^{tj} T_{\bar{t}\bar{a}}^{i\bar{u}} - T_{u\bar{a}}^{t\bar{i}} T_{b\bar{t}}^{j\bar{u}} - T_{u\bar{a}}^{j\bar{u}} T_{b\bar{t}}^{t\bar{i}}, \end{aligned} \quad (3.48)$$

$$\begin{aligned} M_{ab}^{i\bar{j}} = & -\tau_{\bar{b}}^{\bar{j}} T_u^t T_{a\bar{t}}^{i\bar{u}} - \tau_a^i T_{\bar{t}}^{\bar{u}} T_{ub}^{jt} + T_u^t T_{\bar{t}}^{\bar{i}} T_{b\bar{a}}^{j\bar{u}} + T_u^t T_{\bar{a}}^{\bar{u}} T_{b\bar{t}}^{j\bar{i}} \\ & + T_{\bar{t}}^{\bar{u}} T_b^t T_{u\bar{a}}^{j\bar{i}} + T_{\bar{t}}^{\bar{i}} T_u^t T_{b\bar{a}}^{t\bar{i}}, \end{aligned} \quad (3.49)$$

$$M_{u\bar{t}}^{t\bar{i}} = T_u^i, \quad (3.50)$$

$$M_{u\bar{t}}^{i\bar{u}} = T_{\bar{t}}^{\bar{i}}, \quad (3.51)$$

$$M_{u\bar{a}}^{t\bar{i}} = \tau_{ua}^{it}, \quad (3.52)$$

$$M_{a\bar{t}}^{i\bar{u}} = \tau_{\bar{t}\bar{a}}^{i\bar{u}}, \quad (3.53)$$

$$M_{a\bar{t}}^{t\bar{i}} = \tau_{u\bar{a}}^{i\bar{u}}, \quad (3.54)$$

$$M_{u\bar{a}}^{i\bar{u}} = \tau_{a\bar{t}}^{t\bar{i}}, \quad (3.55)$$

$$M_{u\bar{a}}^{t\bar{u}} = -T_a^t, \quad (3.56)$$

$$M_{a\bar{t}}^{t\bar{u}} = -T_{\bar{a}}^{\bar{u}}, \quad (3.57)$$

$$M_{u\bar{t}}^{i\bar{j}} = -\tau_{u\bar{t}}^{j\bar{i}}, \quad (3.58)$$

$$M_{ab}^{t\bar{u}} = -\tau_{b\bar{a}}^{t\bar{u}}. \quad (3.59)$$

The effective Hamiltonian W is just the all active part of the residuum,

$$W = R_{u\bar{t}}^{t\bar{u}}. \quad (3.60)$$

The all internal doubles $T_{u\bar{t}}^{t\bar{u}}$ coupled cluster amplitude is set to zero at the beginning of every iteration. At the end of every iteration the all internal doubles residuum $R_{u\bar{t}}^{t\bar{u}}$ is set to zero.

IAS contribution to the energy,

$$\Delta E_{\text{IAS}} = -W_{u\bar{t}}^{t\bar{u}} T_u^t T_{\bar{t}}^{\bar{u}}. \quad (3.61)$$

Chapter 4

Automatically generated UCCSDT and UDC-CCSDT

Unrestricted implementations of CCSDT and DC-CCSDT[6, 7, 8] were generated with version 1.0.0 of the Quantwo program[9]. The Quantwo inputs are listed below.

- UCCSDT Quantwo input file:

```
prog,spinintegr=0,nobrafac=1,explspin=1,algo=2
output,level=1,maxlenline=70

\beq
<\Phi^{a}_{i}| \op H (1 + \op T_2 + \op T_3) |0>_C
\eeq
\beq
<\Phi^{ab}_{ij}| \op H (1 + \op T_2 + \half \op T_2 \op T_2 + \op T_3) |0>_C
\eeq
\beq
<\Phi^{abc}_{ijk}| \op H (\op T_2 + \op T_3 + \half \op T_2 \op T_2 + \op T_2 \op T_3) |0>_C
```

- UDC-CCSDT Quantwo input file:

```
%singles and doubles amplitude equations from UCCSDT
%we only modify the triples amplitude equation

prog,spinintegr=0,nobrafac=1,explspin=1,algo=2
output,level=1,maxlenline=70

\beq
<\Phi^{abc}_{ijk}| \op H (\op T_2 + \op T_3 + \frac{1}{2} \op T_2 \op T_2
+ \op T_2 \op T_3) |0>_C
+ (1 - \Perm{IJ}{JI} - \Perm{IK}{KI})(1 - \Perm{AB}{BA} - \Perm{AC}{CA})
(\sum_{LMDE} \tnsr \intg{LE}{MD} \tnsr T^{IL}_{AD} \tnsr T^{MJK}_{EBC})
- \frac{1}{2}(1 - \Perm{KI}{IK} - \Perm{KJ}{JK})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{IJ}_{DE} \tnsr T^{LMK}_{ABC}
- \frac{1}{2}(1 - \Perm{CA}{AC} - \Perm{CB}{BC})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{LM}_{AB} \tnsr T^{IJK}_{DEC}
+ \frac{1}{2}(1 - \Perm{IJ}{JI} - \Perm{IK}{KI})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{LI}_{DE} \tnsr T^{MJK}_{ABC}
+ \frac{1}{2}(1 - \Perm{AB}{BA} - \Perm{AC}{CA})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{LM}_{DA} \tnsr T^{IJK}_{EBC}
+ \frac{1}{2}(1 - \Perm{KI}{IK} - \Perm{KJ}{JK})(1 - \Perm{AB}{BA} - \Perm{AC}{CA})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{IJ}_{AD} \tnsr T^{LMK}_{BEC}
+ \frac{1}{2}(1 - \Perm{IJ}{JI} - \Perm{IK}{KI})(1 - \Perm{CA}{AC} - \Perm{CB}{BC})
\sum_{LMDE} \tnsr \intg{LD}{ME} \tnsr T^{IL}_{AB} \tnsr T^{JMK}_{DEC}
\eeq
```

The program generates TensorOperations code. The generated code used by ElemCo.jl is located in the src/algo directory.

Bibliography

- [1] Kats, D. & Manby, F. R. Sparse tensor framework for implementation of general local correlation methods. *J. Chem. Phys.* **138**, 144101 (2013).
- [2] Hampel, C., Peterson, K. A. & Werner, H.-J. A comparison of the efficiency and accuracy of the quadratic configuration interaction (QCISD), coupled cluster (CCSD), and Brueckner coupled cluster (BCCD) methods. *Chem. Phys. Lett.* **190**, 1 (1992).
- [3] Knowles, P. J., Hampel, C. & Werner, H.-J. Coupled cluster theory for high spin, open shell reference wave functions. *J. Chem. Phys.* **99**, 5219 (1993).
- [4] Knowles, P. J., Hampel, C. & Werner, H.-J. Erratum: “Coupled cluster theory for high spin, open shell reference wave functions” [*J. Chem. Phys.* 99, 5219 (1993)]. *J. Chem. Phys.* **112**, 3106 (2000).
- [5] Szalay, P. G. & Bartlett, R. J. Analytic energy gradients for the two-determinant coupled cluster method with application to singlet excited states of butadiene and ozone. *J. Chem. Phys.* **101**, 4936 (1994).
- [6] Kats, D. & Köhn, A. On the distinguishable cluster approximation for triple excitations. *J. Chem. Phys.* **150**, 151101 (2019).
- [7] Rishi, V. & Valeev, E. F. Can the distinguishable cluster approximation be improved systematically by including connected triples? *J. Chem. Phys.* **151**, 064102 (2019).
- [8] Schraivogel, T. & Kats, D. Accuracy of the distinguishable cluster approximation for triple excitations for open-shell molecules and excited states. *J. Chem. Phys.* **155**, 064101 (2021). URL <https://aip.scitation.org/doi/10.1063/5.0059181>.
- [9] Kats, D. & Schraivogel, T. Quanttwo: second-quantization program (2024). github.com/fkfest/quanttwo [Accessed: 2024-02-22].