



Aalto University

Linear time-invariant systems: Dynamics

Process Automation (CHEM-E7140), 2019-2020

Francesco Corona

Chemical and Metallurgical Engineering
School of Chemical Engineering

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Representation and analysis

LTI systems - Dynamics


Representation and analysis

Consider a linear and stationary system of order n , in **state-space** representation

↪ Let p be the number of outputs

↪ Let r be the number of inputs

$$\begin{cases} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{cases}$$



A ($n \times n$), B ($n \times r$), C ($p \times n$) and D ($p \times r$) are (constant) system matrices

↪ $x(t)$ is the **state vector** (n components)

↪ $\dot{x}(t)$ is the derivative of the state vector (n components)

↪ $u(t)$ is the **input vector** (r components)

↪ $y(t)$ is the **output vector** (p components)

Representation and analysis (cont.)

The analysis problem: Determine the behaviour of state $x(t)$ and output $y(t)$ for $t \geq t_0$

- We are given the input function $u(t)$, for $t \geq t_0$
- We are given the initial state $x(t_0)$

The solution for $t \geq t_0$, for an initial state $x(t_0)$ and an input $u(t \geq t_0)$

$$x(t) = e^{A(t-t_0)}x(t_0) + \int_{t_0}^t e^{A(t-\tau)}Bu(\tau)d\tau$$

$$y(t) = Ce^{A(t-t_0)}x(t_0) + \underbrace{C \int_{t_0}^t e^{A(t-\tau)}Bu(\tau)d\tau + Du(t)}_{Cx(t)}$$

The solution is known as the **Lagrange formula**

- Based on the state transition matrix, e^{At}

Force-free and forced evolution

Note that we can write the state solution $x(t)$, for $t \geq t_0$, as the sum of two terms

$$\begin{aligned} x(t) &= \underbrace{e^{A(t-t_0)} x(t_0)}_{x_u(t)} + \underbrace{\int_{t_0}^t e^{A(t-\tau)} B u(\tau) d\tau}_{x_f(t)} \\ &= x_u(t) + x_f(t) \end{aligned}$$

↪ The **force-free evolution** of the state, $x_u(t)$

↪ The **forced evolution** of the state, $x_f(t)$

The **force-free evolution** of the state, from the initial condition $x(t_0)$

↪ $e^{A(t-t_0)}$ indicates the transition from $x(t_0)$ to $x(t)$

↪ In the absence of contribution from the input

The **forced evolution** of the state, from the contribution of input $u(t)$

↪ In the absence of an initial condition $x(t_0)$

Representation
and analysis

State transition
matrix

Some properties
Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix
Transition matrix

The state transition matrix

LTI systems - Dynamics

The state transition matrix

Consider a square ($n \times n$) matrix A , the exponential e^A is also a square ($n \times n$) matrix

$$\rightsquigarrow e^A = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots = \sum_{k=0}^{\infty} \frac{A^k}{k!}$$

The **state transition matrix** is a matrix exponential e^{At}

\rightsquigarrow Its elements are functions of time

The state transition matrix (cont.)

The exponential function

Let z be some scalar, by definition its exponential is a scalar

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \cdots = \sum_{k=0}^{\infty} \frac{z^k}{k!}$$

The series always converges

The matrix exponential

Let A be a $(n \times n)$ matrix, by definition its exponential is a $(n \times n)$ matrix

$$e^A = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \cdots = \sum_{k=0}^{\infty} \frac{A^k}{k!}$$

The series always converges

The state transition matrix (cont.)

The product of several matrices

The product of A and B is only possible when the matrixes are compatible

- Number of columns of A must equal the number of rows of B

The same applies to the product of several matrixes

$$\underbrace{M}_{(m \times n)} = \underbrace{A_1}_{(m \times m_1)} \underbrace{A_2}_{(m_1 \times m_2)} \cdots \underbrace{A_{k-1}}_{(m_{k-2} \times m_{k-1})} \underbrace{A_k}_{(m_{k-1} \times n)}$$

Powers of a matrix

Let A be an order- n square matrix

The k -th power of matrix A is defined as the n -order matrix A^k

$$A^k = \underbrace{AA \cdots A}_{k \text{ times}}$$

Special cases,

$$\rightsquigarrow A^{k=0} = I, A^{k=1} = A$$

The state transition matrix (cont.)

Definition

The state transition matrix

Consider the state-space model with $(n \times n)$ matrix A

$$\begin{array}{c} u(t) \rightarrow \boxed{\begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array}} \rightarrow y(t) \end{array} \quad \begin{cases} \dot{x}(t) &= Ax(t) + Bu(t) \\ y(t) &= Cx(t) + Du(t) \end{cases}$$

System

The **state transition matrix** is the $(n \times n)$ matrix e^{At}

$$e^{At} = \sum_{k=0}^{\infty} \frac{A^k t^k}{k!}$$

The state transition matrix is well defined for any square matrix A

- (The series always converges)

The state transition matrix (cont.)

Not convenient to determine the state transition matrix starting from its definition

↪ One exception is when A is (block-)diagonal

The matrix exponential of block-diagonal matrixes

Consider any block-diagonal matrix A , we have

$$A = \begin{bmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_q \end{bmatrix} \rightsquigarrow e^A = \begin{bmatrix} e^{A_1} & 0 & \cdots & 0 \\ 0 & e^{A_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e^{A_q} \end{bmatrix}$$

The matrix exponential of diagonal matrixes (as special case)

For any diagonal $(n \times n)$ matrix A , we have

$$A = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} \rightsquigarrow e^A = \begin{bmatrix} e^{\lambda_1} & 0 & \cdots & 0 \\ 0 & e^{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e^{\lambda_n} \end{bmatrix}$$

The state transition matrix (cont.)

Example





Consider the state-space model with the (2×2) diagonal matrix A

$$A = \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$$

We are interested in the corresponding state transition matrix

We have,

$$e^{At} = \begin{bmatrix} e^{(-1)t} & 0 \\ 0 & e^{(-2)t} \end{bmatrix}$$



Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Some properties

State transition matrix

Properties

We state without proof some fundamental results about a state transition matrix e^{At}
↪ They are needed to derive Lagrange formula

Proposition

Derivative of the state transition matrix

Consider the state transition matrix e^{At}

We have,

$$\frac{d}{dt}e^{At} = Ae^{At} = e^{At}A$$

By using the derivative property, we have that A commutes with e^{At}
↪ (This result is important)

Properties (cont.)

Proposition

Composition of two state transition matrices

Consider the two state transition matrices e^{At} and $e^{A\tau}$, we have

$$e^{At}e^{A\tau} = e^{A(t+\tau)}$$

Proposition

Inverse of the state transition matrix

Let e^{At} be a state transition matrix, its inverse $(e^{At})^{-1}$ is matrix e^{-At}

$$e^{At}e^{-At} = e^{-At}e^{At} = I$$

A state transition matrix e^{At} is always invertible (non-singular)

- Even if A were singular

Properties (cont.)

Matrix inverse

Consider a square matrix A of order n

We define the **inverse** of A the square matrix of order n , A^{-1}

$$A^{-1}A = AA^{-1} = I$$

The inverse of matrix A exists if and only if A is non-singular

- When the inverse exists, it is also unique

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Sylvester's formula

The state transition matrix

Sylvester's expansion

We determine the analytical expression of the state transition matrix e^{At}

- The procedure is known as **Sylvester expansion**
- (Does not require computing the infinite series)
- There are also other procedures (later)

Proposition

The Sylvester's expansion

Let A be a $(n \times n)$ matrix and let the corresponding state transition matrix be e^{At}

We have,

$$e^{At} = \sum_{i=0}^{n-1} \beta_i(t) A^i = \beta_0(t)I + \beta_1(t)A + \beta_2(t)A^2 + \cdots + \beta_{n-1}(t)A^{n-1}$$

The coefficients β_i of the expansion are appropriate functions of time

- ↪ They can be determined by solving a set of linear equations
- ↪ There is a finite number (n) of them



Sylvester's expansion (cont.)

We show how to determine the coefficients when A has eigenvalues of multiplicity one

We will not consider the other cases, because rather involved and tedious to derive

- ↪ Matrix A has complex eigenvalues (with multiplicity larger one)
- ↪ Matrix A has complex eigenvalues (with multiplicity one)
- ↪ Eigenvalues of A have multiplicity larger than one

Sylvester's expansion (cont.)

Eigenvalues with multiplicity one

Let matrix A have distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$

$$e^{At} = \sum_{i=0}^{n-1} \beta_i(t) A^i = \beta_0(t)I + \beta_1(t)A + \beta_2(t)A^2 + \dots + \beta_{n-1}(t)A^{n-1}$$

The n unknown functions $\beta_i(t)$ are those that solve the system

$$\leadsto \begin{cases} 1\beta_0(t) + \lambda_1\beta_1(t) + \lambda_1^2\beta_2(t) + \dots + \lambda_1^{n-1}\beta_{n-1}(t) = e^{\lambda_1 t} \\ 1\beta_0(t) + \lambda_2\beta_1(t) + \lambda_2^2\beta_2(t) + \dots + \lambda_2^{n-1}\beta_{n-1}(t) = e^{\lambda_2 t} \\ \dots \\ 1\beta_0(t) + \lambda_n\beta_1(t) + \lambda_n^2\beta_2(t) + \dots + \lambda_n^{n-1}\beta_{n-1}(t) = e^{\lambda_n t} \end{cases}$$

Sylvester's expansion (cont.)

Or, equivalently, in matrix form

$$V\beta = \eta$$

- The vector of unknowns

$$\leadsto \beta = [\beta_0(t) \quad \beta_1(t) \quad \cdots \quad \beta_{n-1}(t)]^T$$

- The coefficients matrix¹

$$\leadsto V = \begin{bmatrix} 1 & \lambda_1 & \lambda_1^2 & \cdots & \lambda_1^{n-1} \\ 1 & \lambda_2 & \lambda_2^2 & \cdots & \lambda_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_n & \lambda_n^2 & \cdots & \lambda_n^{n-1} \end{bmatrix}$$

- The known vector

$$\leadsto \eta = [e^{\lambda_1 t} \quad e^{\lambda_2 t} \quad \cdots \quad e^{\lambda_n t}]^T$$

¹A matrix in this form is known as Vandermonde matrix.

Sylvester's expansion (cont.)

$$\eta = [e^{\lambda_1 t} \quad e^{\lambda_2 t} \quad \dots \quad e^{\lambda_n t}]^T$$

The components of vector η are functions of time, $e^{\lambda t}$

↪ Functions $e^{\lambda t}$ are the **modes** of matrix A

↪ Mode $e^{\lambda t}$ associates with eigenvalue λ

Each element of e^{At} is a linear combination of such modes

Sylvester expansion (cont.)

Example

Consider the (2×2) matrix A , we want to determine e^{At}

$$A = \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix}$$

Matrix A is triangular, the eigenvalues correspond to the diagonal elements

Matrix A has 2 distinct eigenvalues

$$\rightsquigarrow \lambda_1 = -1$$

$$\rightsquigarrow \lambda_2 = -2$$


To determine e^{At} , we write the system

$$\begin{cases} 1\beta_0(t) + \lambda_1\beta_1(t) = e^{\lambda_1 t} \\ 1\beta_0(t) + \lambda_2\beta_1(t) = e^{\lambda_2 t} \end{cases} \rightsquigarrow \begin{cases} \beta_0(t) + (-1)\beta_1(t) = e^{(-1)t} \\ \beta_0(t) + (-2)\beta_1(t) = e^{(-2)t} \end{cases}$$

Sylvester's expansion (cont.)


By simple manipulation, we get

$$\rightsquigarrow \begin{cases} \beta_0(t) = 2e^{-t} - e^{-2t} \\ \beta_1(t) = e^{-t} - e^{-2t} \end{cases}$$



Thus,

$$\begin{aligned} e^{At} &= \beta_0(t)I_2 + \beta_1(t)A = (2e^{-t} - e^{-2t}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + (e^{-t} - e^{-2t}) \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \\ &= \begin{bmatrix} e^{-t} & (e^{-t} - e^{-2t}) \\ 0 & e^{-2t} \end{bmatrix} \end{aligned}$$



Each element of e^{At} is a linear combination of the two modes, e^{-t} and e^{-2t}

Sylvester's expansion (cont.)

Eigenvalues and eigenvectors

Let $\lambda \in \mathcal{R}$ be some scalar and let $v \neq 0$ be some $(n \times 1)$ column vector

Consider a square matrix A of order n , suppose that the identity holds

$$Av = \lambda v$$

The scalar λ is called an **eigenvalue** of A

The vector v is called the associated **eigenvector**

Consider a square matrix A of order n whose elements are real numbers

Matrix A has n (not necessarily distinct) eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$

- They can be real numbers or conjugate-complex pairs
- If $\lambda_i \neq \lambda_j$ for $i \neq j$, A has multiplicity one

Sylvester's expansion (cont.)

Eigenvalues of triangular and diagonal matrices

Let matrix $A = \{a_{i,j}\}$ be a triangular or a diagonal matrix

- The eigenvalues of A are the n diagonal elements $\{a_{i,i}\}$

Sylvester's expansion (cont.)

Characteristic polynomial

The **characteristic polynomial** of a square matrix A of order n

- The n -order polynomial in the variable s

$$P(s) = \det(sI - A)$$

Computing eigenvalues and eigenvectors

The eigenvalues of matrix A of order n solve its characteristic polynomial

\rightsquigarrow The roots of the equation $P(s) = \det(sI - A) = 0$

Let λ be an eigenvalue of matrix A

Each eigenvector v associated to it is a non-trivial solution to the system

$$(\lambda I - A)v = 0$$

Sylvester's expansion (cont.)

Systems of linear equations

Consider a system of n linear equations in n unknowns $Ax = b$

- ↪ A is a $(n \times n)$ matrix of **coefficients**
- ↪ b is a $(n \times 1)$ vector of **known terms**
- ↪ x is a $(n \times 1)$ vector of **unknowns**

If A is non-singular, the system admits one and only one solution

If matrix A is singular, let $M = [A|b]$ be a $[n \times (n + 1)]$ matrix

- If $\text{rank}(A) = \text{rank}(M)$, system has infinite solutions
- If $\text{rank}(A) < \text{rank}(M)$, system has no solutions

Sylvester's expansion (cont.)

Matrix rank

The **rank** of a $(m \times n)$ matrix A is equal to the number of columns (or rows) of the matrix that are linearly independent, $\text{rank}(A)$

Matrix kernel or null space

Consider a $(m \times n)$ matrix A , we define its **null space** or **kernel**

$$\ker(A) = \{x \in \mathcal{R}^n | Ax = 0\}$$

It is the set of all vectors $x \in \mathcal{R}^n$ that left-multiplied by A produce the null vector

The set is a vector space, its dimension is called the **nullity** of matrix A , $\text{null}(A)$

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Lagrange formula

LTI systems - Dynamics

Lagrange formula

We can now prove the solution to the analysis problem for MIMO systems

- Lagrange formula

Theorem

Lagrange formula

Consider the state-space representation of a time-invariant linear system of order n

$$\begin{array}{ccc} u(t) & \xrightarrow{\quad} & \boxed{\begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array}} \xrightarrow{\quad} y(t) \end{array} \quad \left\{ \begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array} \right.$$

System

The solution for $t \geq t_0$, for an initial state $x(t_0)$ and an input $u(t \geq t_0)$

$$x(t) = e^{A(t-t_0)} x(t_0) + \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau$$

$$y(t) = Ce^{A(t-t_0)} x(t_0) + C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t)$$

Lagrange formula (cont.)

Proof

By left-multiplying the state equation $\dot{x}(t) = Ax(t) + Bu(t)$ by e^{-At} , we get

$$e^{-At}\dot{x}(t) = e^{-At}Ax(t) + e^{-At}Bu(t)$$

The resulting state equation can be rewritten,

$$e^{-At}\dot{x}(t) - e^{-At}Ax(t) = e^{-At}Bu(t)$$

Then, by using the result on the derivative of the state transition matrix²,

$$\begin{aligned}\frac{d}{dt}\left[e^{-At}x(t)\right] &= e^{-At}\dot{x}(t) - e^{-At}Ax(t) \\ &= e^{-At}Bu(t)\end{aligned}$$

²Derivative of the state transition matrix

$$\frac{d}{dt}\left[e^{-At}x(t)\right] = e^{-At}\left[\frac{d}{dt}x(t)\right] + \left[\frac{d}{dt}e^{-At}\right]x(t) = e^{-At}\dot{x}(t) - e^{-At}Ax(t).$$

Lagrange formula (cont.)

$$\frac{d}{dt} \left[e^{-At} x(t) \right] = e^{-At} B u(t)$$

By integrating between t_0 and t , we obtain

$$\left[e^{-A\tau} x(\tau) \right]_{t_0}^t = \int_{t_0}^t e^{-A\tau} B u(\tau) d\tau$$

That is,

$$e^{At} x(t) - e^{-At_0} x(t_0) = \int_{t_0}^t e^{-A\tau} B u(\tau) d\tau$$

Thus,

$$e^{-At} x(t) = e^{-At_0} x(t_0) + \int_{t_0}^t e^{-A\tau} B u(\tau) d\tau$$

Lagrange formula (cont.)

$$e^{-At}x(t) = e^{-At_0}x(t_0) + \int_{t_0}^t e^{-A\tau}Bu(\tau)d\tau$$

The first Lagrange formula is obtained by multiplying both sides by e^{At}

$$\rightsquigarrow x(t) = e^{A(t-t_0)}x(t_0) + \int_{t_0}^t e^{A(t-\tau)}Bu(\tau)d\tau$$

The second formula is obtained by substituting $x(t)$ in the output equation

$$y(t) = Cx(t) + Du(t)$$

$$\rightsquigarrow C \underbrace{\left[e^{A(t-t_0)}x(t_0) + \int_{t_0}^t e^{A(t-\tau)}Bu(\tau)d\tau \right]}_{x(t)} + Du(t)$$

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Force-free and forced evolution

Lagrange formula

Force-free and forced evolution

$$x(t) = \underbrace{e^{A(t-t_0)} x(t_0)}_{x_u(t)} + \underbrace{\int_{t_0}^t e^{A(t-\tau)} B u(\tau) d\tau}_{x_f(t)}$$

We can write the state solution (for $t \geq t_0$) as the sum of two terms

$$\rightsquigarrow x(t) = x_u(t) + x_f(t)$$

\rightsquigarrow The **force-free evolution** of the state, $x_u(t)$

\rightsquigarrow The **forced evolution** of the state, $x_f(t)$

Force-free and forced evolution (cont.)

$$x(t) = \underbrace{e^{A(t-t_0)} x(t_0)}_{x_u(t)} + \underbrace{\int_{t_0}^t e^{A(t-\tau)} B u(\tau) d\tau}_{x_f(t)}$$

The **force-free evolution** of the state, from the initial condition $x(t_0)$

$$\rightsquigarrow x_l(t) = e^{A(t-t_0)} x(t_0)$$

$\rightsquigarrow e^{A(t-t_0)}$ indicates the transition from $x(t_0)$ to $x(t)$

\rightsquigarrow In the absence of contribution from the input

The **forced evolution** of the state

$$\rightsquigarrow x_f(t) = \int_{t_0}^t e^{A(t-\tau)} B u(\tau) d\tau = \int_0^{t-t_0} e^{A\tau} B u(t-\tau) d\tau$$

\rightsquigarrow The contribution of $u(\tau)$ to state $x(t)$

\rightsquigarrow Through a weighting function, $e^{A(t-\tau)} B$

Force-free and forced evolution (cont.)

$$y(t) = \underbrace{Ce^{A(t-t_0)}x(t_0)}_{y_u(t)} + \underbrace{C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t)}_{y_f(t)}$$

We can write the output solution (for $t \geq t_0$) as the sum of two terms

$$\rightsquigarrow y(t) = y_l(t) + y_f(t)$$

\rightsquigarrow The **force-free evolution** of the output, $y_u(t)$

\rightsquigarrow The **forced evolution** of the output, $y_f(t)$

Free and forced evolution (cont.)

$$y(t) = \underbrace{Ce^{A(t-t_0)}x(t_0)}_{y_u(t)} + \underbrace{C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t)}_{y_f(t)}$$


The **force-free evolution** of the output, from initial condition $y(t_0) = Cx(t_0)$

$$\rightsquigarrow y_u(t) = Ce^{A(t-t_0)}x(t_0) = Cx_u(t)$$

The **forced-evolution** of the output

$$\rightsquigarrow y_f(t) = C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t) = Cx_f(t) + Du(t)$$

Free and forced evolution (cont.)



Note that for $t_0 = 0$, we have

$$x(t) = e^{At}x(0) + \int_0^t e^{A(t-\tau)}Bu(\tau)d\tau$$

$$y(t) = Ce^{At}x(0) + C \int_0^t e^{A(t-\tau)}Bu(\tau)d\tau + Du(t)$$

Free and forced evolution (cont.)

Example

Consider a linear time-invariant system with the state-space representation,

$$\begin{cases} \begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(t) \\ y(t) = \begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} \end{cases}$$

We want to determine the state and the output evolution for $t \geq 0$

- We consider the input signal $u(t) = 2\delta_{-1}(t)$
- We consider the initial state $x(0) = (3, 4)^T$

The state transition matrix for this state-space representation,

$$e^{At} = \begin{bmatrix} e^{-t} & (e^{-t} - e^{-2t}) \\ 0 & e^{-2t} \end{bmatrix}$$


We computed it earlier

Free and forced evolution (cont.)

The force-free evolution of the state, for $t \geq 0$

$$\leadsto x_u(t) = e^{At}x(0) = \begin{bmatrix} e^{-t} & (e^{-t} - e^{-2t}) \\ 0 & e^{-2t} \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} (7e^{-t} - 4e^{-2t}) \\ 4e^{-2t} \end{bmatrix}$$

That is,




Free and forced evolution (cont.)

The force-free evolution of the output, for $t \geq 0$

$$\rightsquigarrow y_u(t) = Cx_u(t) = \begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} (7e^{-t} - 4e^{-2t}) \\ 4e^{-2t} \end{bmatrix} = 14e^{-t} - 4e^{-2t}$$



That is,



Free and forced evolution (cont.)

The forced evolution of the state, for $t \geq 0$

$$\begin{aligned}
 \rightsquigarrow x_f(t) &= \int_0^t e^{At} B u(t - \tau) d\tau = \int_0^t \begin{bmatrix} e^{-\tau} & (e^{-\tau} - e^{-2\tau}) \\ 0 & e^{-2\tau} \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} 2 d\tau \\
 &= 2 \int_0^t \begin{bmatrix} (e^{-\tau} - e^{-2\tau}) \\ e^{-2\tau} \end{bmatrix} d\tau = 2 \begin{bmatrix} \int_0^t (e^{-\tau} - e^{-2\tau}) d\tau \\ \int_0^t e^{-2\tau} d\tau \end{bmatrix} \\
 &= 2 \begin{bmatrix} (1 - e^{-t}) - 1/2(1 - e^{-2t}) \\ 1/2(1 - e^{-2t}) \end{bmatrix} = \begin{bmatrix} (1 - 2e^{-t} + e^{-2t}) \\ (1 - e^{-2t}) \end{bmatrix}
 \end{aligned}$$




Free and forced evolution (cont.)

Since $D = 0$, the forced evolution of the output for $t \geq 0$

$$\rightsquigarrow y_f(t) = Cx_f(t) = \begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} (1 - 2e^{-t} + e^{-2t}) \\ (1 - e^{-2t}) \end{bmatrix} = 3 - 4e^{-t} + e^{-2t}$$

That is,



Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Similarity transformation

LTI systems - Dynamics

Similarity transformation

The form of the state space representation depends on the choice of states

- The choice is not unique

There is an infinite number of different representations of the same system

- They are all related by a **similarity transformation**
- These transformations allow flexibility in the analysis
- We can change to easier system representations

The state matrix can be set to a **canonical form**

↪ **Diagonal form**

↪ **Jordan form**

↪ ...

Similarity transformation (cont.)

Definition

Similarity transformation

Consider the state-space representation of a linear time-invariant system of order n

$$\begin{array}{ccc}
 u(t) & \xrightarrow{\quad} & \boxed{\begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array}} \xrightarrow{\quad} y(t) \\
 & & \text{System}
 \end{array}
 \qquad
 \begin{cases} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{cases}$$

- $x(t)$ and $\dot{x}(t)$, state vector and its derivative (n components)
- $u(t)$, input vector (r components)
- $y(t)$, output vector (p components)

Let vector $z(t)$ be related to $x(t)$ by a linear transformation P , $x(t) = Pz(t)$

P is any $(n \times n)$ non-singular matrix of constants, its inverse always exists

- We have $z(t) = P^{-1}x(t)$

Transformation/matrix P is called **similarity transformation/matrix**

Similarity transformation (cont.)

Proposition

Similar representation

Consider the state-space representation of a linear time-invariant system of order n

$$\begin{array}{ccc} u(t) & \xrightarrow{\quad} & \boxed{\begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array}} \xrightarrow{\quad} y(t) \end{array} \quad \left\{ \begin{array}{l} \dot{x}(t) = Ax(t) + Bu(t) \\ y(t) = Cx(t) + Du(t) \end{array} \right.$$

System

Let P be some transformation matrix such that $x(t) = Pz(t)$

Vector $z(t)$ satisfies the new state-space representation

$$\begin{array}{ccc} u(t) & \xrightarrow{\quad} & \boxed{\begin{array}{l} \dot{x}(t) = A'x(t) + B'u(t) \\ y(t) = C'x(t) + D'u(t) \end{array}} \xrightarrow{\quad} y(t) \end{array} \quad \left\{ \begin{array}{l} \dot{z}(t) = A'z(t) + B'u(t) \\ y(t) = C'z(t) + D'u(t) \end{array} \right.$$

System

$$\begin{aligned} &\rightsquigarrow A' = P^{-1}AP \\ &\rightsquigarrow B' = P^{-1}B \\ &\rightsquigarrow C' = CP \\ &\rightsquigarrow D' = D \end{aligned}$$

Similarity transformation (cont.)

Proof

By taking the time-derivative of $x(t) = Pz(t)$,

$$\rightsquigarrow \dot{x}(t) = P\dot{z}(t)$$


By substituting $x(t)$ and $\dot{x}(t)$ into the state-space representation,

$$\rightsquigarrow \begin{cases} P\dot{z}(t) = APz(t) + Bu(t) \\ y(t) = CPz(t) + Du(t) \end{cases}$$

Pre-multiply the state equation by P^{-1} , to complete the proof



Similarity transformation (cont.)



We obtained a different state-space representation of the same dynamical system

- Input $u(t)$ and output $y(t)$ are left unchanged
- The new state is indicated by $z(t)$

There is an infinite number of non-singular matrixes P that could be used

↪ Thus, there is also an infinite number of equivalent representations

$$\rightsquigarrow A' = P^{-1}AP$$

$$\rightsquigarrow B' = P^{-1}B$$

$$\rightsquigarrow C' = CP$$

$$\rightsquigarrow D' = D$$

Similarity transformation (cont.)

Example

Consider a linear time-invariant system with state-space representation $\{A, B, C, D\}$

$$\begin{cases} \begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = \underbrace{\begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix}}_A \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \underbrace{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}_B u(t) \\ \begin{bmatrix} y_1(t) \\ y_2(t) \end{bmatrix} = \underbrace{\begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}}_C \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \underbrace{\begin{bmatrix} 1.5 \\ 0 \end{bmatrix}}_D u(t) \end{cases}$$

Consider the similarity transformation of the state using the matrix P

$$\begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} = \underbrace{\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}}_P \begin{bmatrix} z_1(t) \\ z_2(t) \end{bmatrix}$$

What is the $\{A', B', C', D'\}$ state-space representation for state $z(t)$

Similarity transformation (cont.)

We have,

$$P = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \rightsquigarrow P^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$$

Since $z(t) = P^{-1}x(t)$, we have

$$\begin{bmatrix} z_1(t) \\ z_2(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} = \begin{bmatrix} x_2(t) \\ x_1(t) - x_2(t) \end{bmatrix}$$

The second component of z is the difference between first and second component of x

\rightsquigarrow The first component of z simply equals the second component of x

Similarity transformation (cont.)

In addition, we can calculate the state-space representation

$$A' = P^{-1}AP = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -2 & 0 \end{bmatrix} = \begin{bmatrix} -2 & 0 \\ 2 & -1 \end{bmatrix}$$

$$B' = P^{-1}B = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$C' = CP = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 2 & 0 \end{bmatrix}$$

$$D' = D = \begin{bmatrix} 1.5 \\ 0 \end{bmatrix}$$



Similarity transformation (cont.)

Proposition

Similarity and state transition matrix

Consider the state matrix $A' = P^{-1}AP$ from some similarity transformation P

The corresponding state transition matrix,

$$e^{A't} = P^{-1}e^{At}P$$

Proof

Note that

$$(A')^k = \underbrace{(P^{-1}AP) \cdot (P^{-1}AP) \cdots (P^{-1}AP)}_{k \text{ times}} = P^{-1} \underbrace{AA \cdots A}_{k \text{ times}} P = P^{-1}A^kP$$

Thus, by definition

$$e^{A't} = \sum_{k=0}^{\infty} \frac{(A')^k t^k}{k!} = \sum_{k=0}^{\infty} \frac{(P^{-1}A^kP)t^k}{k!} = P^{-1} \left(\sum_{k=0}^{\infty} \frac{A^k t^k}{k!} \right) P = P^{-1}e^{At}P$$

Similarity transformation (cont.)

We show how two similar state-space representations describe the same IO relation

Proposition

Invariance of the IO relationship by similarity

Consider two similar state-space representations of a linear time-invariant system

$$\rightsquigarrow \{A, B, C, D\} \text{ and } \{A', B', C', D'\}$$

$$\rightsquigarrow P \text{ is the transformation matrix}$$

Suppose that the system be subjected to some known input $u(t)$

The two representations produce the same forced response

$$\rightsquigarrow y_f(t)$$

Similarity transformation (cont.)

Proof

Consider the Lagrange formula

The forced response of the second representation due to input $u(t)$

$$\begin{aligned} y_f(t) &= C' \int_{t_0}^t e^{A'(t-\tau)} B' u(\tau) d\tau + Du(t) \\ &= CP \int_{t_0}^t \underbrace{P^{-1} e^{A(t-\tau)} P}_{e^{A'(t-\tau)}} \underbrace{P^{-1} B}_{B'} u(\tau) d\tau + Du(t) \\ &= C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t) \end{aligned}$$

This response corresponds to that of the first SS representation

$$y_f(t) = C \int_{t_0}^t e^{A(t-\tau)} Bu(\tau) d\tau + Du(t)$$



Similarity transformation (cont.)

Proposition

Invariance of the eigenvalues under similarity transformations

Matrix A and $P^{-1}AP$ have the same characteristic polynomial

Proof

The characteristic polynomial of matrix A'

$$\begin{aligned}\det(\lambda I - A') &= \det(\lambda I - P^{-1}AP) = \det(\lambda \underbrace{P^{-1}P}_I - P^{-1}AP) \\ &= \det[P^{-1}(\lambda I - A)P] = \det(P^{-1}) \det(\lambda I - A) \det(P) \\ &= \det(\lambda I - A)\end{aligned}$$

The last equality is obtained from $\det(P^{-1})\det(P) = 1$

A and A' share the same characteristic polynomial

~> Thus, also the eigenvalues are the same

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Similarity transformation (cont.)

Two similar representations have the same modes, the modes characterise the dynamics

The modes are therefore independent of the representation

↪ This is important

Similarity transformation (cont.)

Example

Consider two similar state-space representations of a linear time-invariant system

$$A = \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix}$$

$$A' = \begin{bmatrix} -2 & 0 \\ 2 & -1 \end{bmatrix}$$

The similarity transformation matrix

$$P = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

We are interested in the eigenvalues and modes of the system

Matrix A and A' have two eigenvectors

- $\lambda_1 = -1$
- $\lambda_2 = -2$

The system modes are e^{-t} and e^{-2t}

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Diagonalisation

LTI systems - Dynamics

Diagonalisation

We consider a special similarity transformation \mathbf{P} , we seek for a diagonal matrix \mathbf{A}'

↪ A state-space representation with a diagonal state matrix

↪ **Diagonal canonical form**

↪ $\mathbf{\Lambda} = \mathbf{A}' = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$

Consider the linear time-invariant system with a single input (and, say, single output)

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \\ \vdots \\ \dot{x}_n(t) \end{bmatrix} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} u(t)$$

The evolution of the i -th component of the state vector

$$\rightsquigarrow \dot{x}_i(t) = \lambda_i x_i(t) + b_i u(t)$$

State derivatives are not related to other components

Diagonalisation (cont.)

We think of a system with diagonal matrix A as a collection of sub-systems

- ↪ Each sub-system is described by a single state component
- ↪ Each state component evolves independently
- ↪ The representation is **decoupled**
- ↪ n first-order subsystems

The characteristic polynomial of the system for the i -th component

$$\rightsquigarrow P_i(s) = (s - \lambda_i)$$

This subsystem has mode $e^{-\lambda_i t}$

We show how to determine a similarity transformation into a diagonal form

- A somehow special similarity transformation matrix

Representation
and analysis

State transition
matrix

Some properties

Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

Modal matrix

Diagonalisation

Diagonalisation (cont.)

Definition

Modal matrix

Consider a system in state-space representation with $(n \times n)$ matrix A

- Let v_1, v_2, \dots, v_n be a set of all the eigenvectors of matrix A
- Suppose that they correspond to eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$

Suppose that eigenvectors in this set are linearly independent

We define the **modal matrix** of A as the $(n \times n)$ matrix V

$$V = [v_1 | v_2 | \dots | v_n]$$

Diagonalisation (cont.)

If a matrix A has n distinct eigenvalues λ , then the modal matrix A always exists

- As its n eigenvectors \mathbf{v} are linearly independent

Distinct eigenvalues

Let A be a n -order matrix whose n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ are distinct

Then, there is a set of n linearly independent eigenvectors

- Vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ form a basis for \mathcal{R}^n

Diagonalisation (cont.)

Example

Consider a state-space representation of a linear time-invariant system with matrix A

$$A = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$$

We are interested in the modal matrix V of A

The eigenvalues and eigenvectors of A

$$\rightsquigarrow \lambda_1 = 1 \text{ and } v_1 = \begin{bmatrix} 1 & -1 \end{bmatrix}^T$$

$$\rightsquigarrow \lambda_2 = 5 \text{ and } v_2 = \begin{bmatrix} 1 & 3 \end{bmatrix}^T$$

The modal matrix V ,

$$V = [v_1 | v_2] = \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix}$$

Diagonalisation (cont.)

$$V = [v_1|v_2] = \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix}$$

It is important to note that the eigenvectors are determined up to a scaling constant

- (Plus, the ordering of the eigenvalues is arbitrary)
- There can be more than one modal matrix

These two modal matrices of matrix A are equivalent

$$V' = [2v_1|3v_2] = \begin{bmatrix} 2 & 3 \\ -2 & 9 \end{bmatrix}$$

$$V'' = [v_2|v_1] = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$$



Diagonalisation (cont.)

Consider a matrix A with some eigenvalues λ that have multiplicity ν larger than one

- The modal matrix V exists if and only if to each eigenvalue λ_i with multiplicity ν_i is possible to associate ν_i linearly independent eigenvectors $\{v_{i,1}, v_{i,2}, \dots, v_{i,\nu_i}\}$

This is not always possible

But, ...

If a matrix admits a modal matrix, then it can be diagonalised

- (This is what matters to us)

Diagonalisation (cont.)

Example

Consider a state space representation of a linear time-invariant system with matrix A

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

Its only eigenvalue $\lambda = 2$ has multiplicity $\nu = 2$

Its eigenvectors are obtained by solving the system $[\lambda I - A]v = 0$

$$[2I - A]v = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightsquigarrow \begin{cases} 0 = 0 \\ 0 = 0 \end{cases}$$

We can choose any two linearly independent eigenvectors for λ

- As the equation is satisfied for any value of a and b

A modal matrix with the eigenvectors from the canonical basis

$$\rightsquigarrow V = [v_1 | v_2] = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Diagonalisation (cont.)

Example

Consider a state space representation of a linear time-invariant system with matrix A

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$$

Its only eigenvalue $\lambda = 2$ has multiplicity $\nu = 2$

Its eigenvectors are obtained by solving the system $[\lambda I - A]v = 0$

$$[2I - A]v = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightsquigarrow \begin{cases} -b = 0 \\ 0 = 0 \end{cases}$$

As $b = 0$, we can choose only one linearly independent eigenvector for λ

$$v_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Matrix A does not admit a modal matrix

Diagonalisation (cont.)

Proposition

Diagonalisation

Consider a state-space representation of a linear time-invariant system with matrix A

Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be its eigenvalues and $V = [v_1 | v_2 | \dots | v_n]$ one of its modal matrices

Let Λ be the state matrix M transformed according to $\Lambda = V^{-1}AV$

$\leadsto \Lambda$ is diagonal



Diagonalisation (cont.)

Example

Consider a linear time-invariant system with matrixes $\{A, B, C, D\}$

$$\begin{cases} \begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(t) \\ \begin{bmatrix} y_1(t) \\ y_2(t) \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 1.5 \\ 0 \end{bmatrix} u(t) \end{cases}$$

We are interested in a diagonal representation by similarity

We can compute the eigenvalues and eigenvectors of A

- $\lambda_1 = -1$ and $v_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
- $\lambda_2 = -2$ and $v_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$

Diagonalisation (cont.)

Then, we can determine a modal matrix and its inverse

$$V = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$
$$V^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

From the similarity transformation expressions, we get

$$\begin{aligned} A' &= V^{-1}AV = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & -2 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix} = \Lambda \\ B' &= V^{-1}B = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\ C' &= CV = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & -2 \end{bmatrix} \\ D' &= D = \begin{bmatrix} 1.5 \\ 0 \end{bmatrix} \end{aligned}$$

Representation
and analysis

State transition
matrix

Some properties
Sylvester's formula

Lagrange
formula

Force-free and
forced evolution

Similarity
transformation

Diagonalisation

Modal matrix

Transition matrix

State transition matrix by diagonalisation

Diagonalisation

State transition matrix by diagonalisation

We show a procedure alternative to Sylvester's formula for the state transition matrix

- We assume a linear time-invariant state-space system representation
- We assume that the state matrix A can be diagonalised

Transition matrix by diagonalisation (cont.)

Proposition

State transition matrix by diagonalisation

Consider a $(n \times n)$ state matrix A and let $\lambda_1, \lambda_2, \dots, \lambda_n$ be its eigenvalues

Suppose that A admits the modal matrix V

We have for the state transition matrix

$$e^{At} = V e^{\Lambda t} V^{-1} = V \begin{bmatrix} e^{\lambda_1 t} & 0 & \cdots & 0 \\ 0 & e^{\lambda_2 t} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e^{\lambda_n t} \end{bmatrix} V^{-1}$$

Because we have a diagonal state matrix

$$\Lambda = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$

State transition matrix by diagonalisation (cont.)

Proof

We have shown that the identity holds (see similarity and state transition matrices³)

$$e^{At} = V^{-1} e^{A_d t} V$$

To complete the proof, multiply both sides by V on the left and by V^{-1} on the right



³Given $A' = P^{-1}AP$, we have $e^{A't} = P^{-1}e^{At}P$.

State transition matrix by diagonalisation (cont.)

Example

Consider a linear time-invariant system with matrixes $\{A, B, C, D\}$

$$\begin{cases} \begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(t) \\ \begin{bmatrix} y_1(t) \\ y_2(t) \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 1.5 \\ 0 \end{bmatrix} u(t) \end{cases}$$

We are interested in computing the state transition matrix e^{At}

State transition matrix by diagonalisation (cont.)

We have already computed the modal matrix of A and its inverse, V and V^{-1}

$$V = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

$$V^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

Thus, we have

$$\begin{aligned} e^{At} &= V \begin{bmatrix} e^{\lambda_1 t} & 0 \\ 0 & e^{\lambda_2 t} \end{bmatrix} V^{-1} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} e^{-t} & 0 \\ 0 & e^{-2t} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} e^{-t} & e^{-t} \\ 0 & -e^{-2t} \end{bmatrix} = \begin{bmatrix} e^{-t} & (e^{-t} - e^{-2t}) \\ 0 & e^{-2t} \end{bmatrix} \end{aligned}$$

This is the same result we determined by using the Sylvester expansion

