
Calculus 1

March 4, 2025

Contents

Differentiable Functions	5
1 Differentiability	5
2 Using Algebra to Compute Derivatives	8
3 Applications	11
Differentiable Functions, Part 2	13
1 Using Limits to Compute Derivatives	13
Free Fall	16
Purchasing Power	19
1 Purchasing Power	19
2 Weight in Space	23
3 A Rock in Free Fall	24
4 Distance to the Horizon	25
5 Intensity of Sound	26
Practice Quiz 1	27
1 Part 1	27
Introduction to Motion	30
Introduction	31
Small Changes	38
1 Odometer Readings	38
2 A Projectile	41
3 The Falling Ladder, Part 1	42
4 The Falling Ladder, Part 2	43
5 Riding a Ferris Wheel	45
Limits	46
1 Limits and Tangent Lines	46
2 Limits and Gas Mileage	46
3 Limits, Gas Mileage and Speed	47
4 Limits, Speed and Altitude	48
5 Limits and Gas Mileage	49
6 Limits and Purchasing Power	49

The Derivative	50
1 Limits, Speed and Altitude	55
Derivatives of Polynomials	57
1 Differentiating Polynomials	57
2 Stock Price	59
3 Homework Solution	68
Gas Consumption	69
The Quotient and Product Rules	71
1 Relative Rates of Change	71
2 Exercises	72
Test 1	75
Derivatives of Exponential Functions	79
Circular Interpolation	91
Derivatives of Trigonometric Functions	96
1 Exercises	96
2 Visualizing Derivatives	98
3 The Derivative of the Sine Function	99
4 Applications 1	100
5 Transformations of the Sine Function	100
6 Applications 2	101
7 MATH 142	101
The Chain Rule	102
Exponential Functions, Part 2	117
Practice Quiz 2	122
Derivatives of Inverse Functions, Short Version	126
1 The Main Idea	126
2 Derivatives of Inverse Trig Functions	127
3 Exercises	128
Derivatives of Inverse Functions	131
The Quotient Rule	147
Implicit Differentiation	157
The Ladder and the Tree	165

The Ladder and the Tree, Part 2	167
More Related Rates Problems	170
1 Kite Flying	170
2 A Crawling Beetle	171
3 A Mechanical Motion	171
4 Tracking a Helicopter	171
5 Distance to the Horizon	172
6 Jar Lid and Rubber Band	172
7 Disco Dancing	173
The Natural Log Function	174
1 The Derivative of the Natural Log Function	174
2 Relative Rates of Change	176
Derivatives and the Shapes of Curves	178
Optimization	179
Hours of Daylight	181
Motion	184
Review of Differentiation	186
Review Problems	192
Hanging Chains	194
Electron in a Crossed Field	195
1 Newton's Law of Cooling	197
Short Quizzes Math 151	198
Short Quizzes Math 142	200
Sinusoidal Modeling, Math 142	202
Simple Harmonic Motion	204

Differentiable Functions

An introduction to what it means for a function to be differentiable.

1 Differentiability

Zoom in closer and closer near a point $(a, f(a))$ on the graph of a common function f and you'll most likely notice that the graph looks more and more like the graph of a linear function (ie. like a non-vertical line). If so, we say the function $y = f(x)$ is *differentiable* at $x = a$. And the *derivative* of the function, evaluated at $x = a$, written as

$$\left. \frac{dy}{dx} \right|_{x=a},$$

is just the slope of that line or the rate of change of the linear function.

Exploration 1 Use the graph of the function $y = f(x)$ below to answer the following questions.

Desmos link: <https://www.desmos.com/calculator/ojdj4r3r9v>

151: Diff0

- (a) Zoom in toward the point A to determine if the function $y = f(x)$ is differentiable at $x = 1$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=1} = \left. \frac{d}{dx} (f(x)) \right|_{x=1}.$$

The derivative is

$$\left. \frac{dy}{dx} \right|_{x=1} = 3.$$

- (b) Zoom in toward the point B to determine if the function $y = f(x)$ is differentiable at $x = 0$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0} = \left. \frac{d}{dx} (f(x)) \right|_{x=0}.$$

The derivative is

$$\left. \frac{dy}{dx} \right|_{x=0} = -2.$$

Learning outcomes:
Author(s):

Exploration 2 Use the graph of the function

$$y = f(x) = |x|$$

below to evaluate the following derivatives if possible. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/us2fruzbra>

151: Diff1

(a)

$$\left. \frac{d(|x|)}{dx} \right|_{x=2} = 1$$

(b)

$$\left. \frac{d(|x|)}{dx} \right|_{x=0} =$$

(c)

$$\left. \frac{d(|x|)}{dx} \right|_{x=-3} = -1$$

Exploration 3 Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0$. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/ov8qt938ot>

151: Diff2

Exploration 4 (a) Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0.2$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0.2} = \left. \frac{d}{dx} (f(x)) \right|_{x=0.2}.$$

Explain your reasoning.

- (b) Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0} = \left. \frac{d}{dx} (f(x)) \right|_{x=0}.$$

Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/tvwtbx9hco>

151: Not Differentiable

Exploration 5 Use the graph of $y = f(x)$ below to determine whether each of the following derivatives are negative, positive, or zero. Explain your reasoning.

(a)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=0.5}$$

(b)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=1}$$

(c)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=1.5}$$

(d)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=2.29}$$

Desmos link: <https://www.desmos.com/calculator/ks2yui6ofs>

151:Diff 7

Question 6 (a) Summarize your understanding of the main ideas of this section.

(b) What questions do you have about this section?

Free Response:

2 Using Algebra to Compute Derivatives

Exploration 7 The aim of this problem is to use numerical methods to approximate and algebra to evaluate the derivative

$$\left. \frac{d}{dx} (x^3) \right|_{x=1}.$$

- (a) First use the graph of the function $y = f(x) = x^3$ below to approximate or guess the value of the above derivative by zooming in on the appropriate point.

$$\left. \frac{d}{dx} (x^3) \right|_{x=1} = 3.$$

Desmos link: <https://www.desmos.com/calculator/rto22qzlvw>

151: Cubing Function

- (b) The idea to compute the derivative algebraically is this: Fix the point $P(1,1)$ on the graph of $y = f(x) = x^3$. Then choose a variable point Q on the graph, different from P , with coordinates (v, v^3) . When Q is sufficiently close to P , the line PQ approximates the curve $y = x^3$ near $P(1,1)$ and the slope of this line approximates the derivative above.
- (i) Our first step is to find an expression for the slope of line PQ as a function of v . The slope is

$$\begin{aligned} m(v) &= \frac{\Delta y}{\Delta v} \\ &= \frac{f(v) - f(1)}{v - 1} \\ &= \frac{v^3 - 1}{v - 1} \end{aligned}$$

all assuming $v \neq 1$.

- (ii) Now we'll use the slope (or average rate of change) function $m(x)$ to create a table of slopes for the lines PQ . Reveal the contents of the Table folder in Line 1 of the worksheet above by clicking the Right Arrow just to the left of "Table".
- Add a few entries to the table to get better approximations to the above derivative.
 - The slopes $m(v)$ should appear should appear to approach some number as v approaches 1. What is that number?

iii. This suggests that

$$\left. \frac{d}{dx} (x^3) \right|_{x=1} = \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} = 3.$$

(iii) Another approach to approximating the derivative is to graph the slope function

$$m(v) = \frac{v^3 - 1}{v - 1}, v \neq 1.$$

Activate the Average Rate of Change folder on Line 11 by clicking the camera icon to the left of the line to see the graph of this function. Activate also the Table folder in Line 1.

- i. Drag the slider v on Line 4. Describe the relationship between the graph of the slope function and the line PQ .
 - ii. There is a hole in the graph of the function $y = m(x)$. Where is it? Why is it there?
- (iv) To verify the numerical and graphical evidence that

$$\left. \frac{d}{dx} (x^3) \right|_{x=1} = \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} = 3,$$

we'll use algebra to evaluate the above limit.

The idea is to factor the numerator $v^3 - 1$. Since

$$(v^3 - 1) \Big|_{v=1} = 1^3 - 1 = 0,$$

we know that $v - 1$ is a factor of $v^3 - 1$. To simplify the quotient

$$\frac{v^3 - 1}{v - 1}, v \neq 1,$$

we could use long division or factor the difference of two cubes:

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2).$$

Either way, the result is that

$$v^3 - 1 = (v - 1)(v^2 + v + 1).$$

So

$$\frac{v^3 - 1}{v - 1} = v^2 + v + 1, v \neq 1.$$

Putting this all together, we get

$$\begin{aligned}
 \left. \frac{d}{dx} (x^3) \right|_{x=1} &= \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} \\
 &= \lim_{v \rightarrow 1} \frac{(v - 1)(v^2 + v + 1)}{v - 1} \\
 &= \lim_{v \rightarrow 1} (v^2 + v + 1) \\
 &= 1^2 + 1 + 1 \\
 &= 3.
 \end{aligned}$$

Exploration 8 Repeat all parts of Exploration 2 for the following derivatives.

(a)

$$\left. \frac{d}{dx} (x^3) \right|_{x=2} = 12.$$

(b)

$$\left. \frac{d}{dx} (x^3) \right|_{x=a} = 3a^2.$$

Use the demonstration below to check your work by dragging the sliders v (Line 4) and a (Line 16). Then activate the Derivative Function folder on Line 17.

Desmos link: <https://www.desmos.com/calculator/ewy7jqij6s>

151: Cubing Function 2

(c)

$$\left. \frac{d}{dx} \left(\frac{1}{x} \right) \right|_{x=1} = -1.$$

(d)

$$\left. \frac{d}{dx} \left(\frac{1}{x} \right) \right|_{x=a} = -1/a^2.$$

(e)

$$\left. \frac{d}{dx} \left(\frac{1}{x^2} \right) \right|_{x=3} = -2/27.$$

(f)

$$\frac{d}{dx} \left(\frac{1}{x^2} \right) \Big|_{x=a} = -2/a^3.$$

(g)

$$\frac{d}{dx} \left(\frac{1}{1+x^2} \right) \Big|_{x=1} = -\frac{1}{2}.$$

(h)

$$\frac{d}{dx} \left(\frac{1}{1+x^2} \right) \Big|_{x=a} = -\frac{2a}{(1+a^2)^2}.$$

3 Applications

Question 9 The function

$$h = f(t), \quad 0 \leq t \leq 2.2,$$

expresses the height of a balloon (in hundreds of feet) in terms of the number of minutes past noon.

The graph of the function $h = f(t)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/yd4xm6x6ub>

151: Balloon

- Find an expression for the function $r = m(v)$ that gives the balloon's average rate of ascent (measured in hundreds of feet per minute) between time v minutes past noon and 12:02 pm. Include also the function's domain.
- How is the average rate of ascent function in part (a) related to the line PQ in the demonstration above?
- Use the graph of the function $h = f(t)$ above to sketch a rough graph of the function $r = m(v)$.
- Activate the folder (avg. rate of change function) in Line 11 to check your graph of the function $r = m(v)$.
- Open the Table in Line 1 by clicking the right arrow at the left of the line.

- (i) What is the balloon's average rate of ascent between 12:02:00 pm and 12:02:06 pm?
 - (ii) Use the graph of the function $h = f(t)$ above to estimate the balloon's rate of ascent at 12:02pm by zooming in on the appropriate point.
 - (iii) What does the table suggest about the balloon's rate of ascent at 12:02pm? Explain.
- (f) Now suppose
- $$f(t) = -2t^3 + 7t^2 - 8t + 8.$$
- (i) Find a simplified expression for the function $r = m(v)$. Include the appropriate domain.
 - (ii) Use your simplified expression to compute the balloon's rate of ascent at 12:02pm.
-

Differentiable Functions, Part 2

Using limits to compute the derivative of a function at a general input.

1 Using Limits to Compute Derivatives

Exploration 10 In part (b) of Exploration 7 from the previous chapter where we computed

$$\left. \frac{d}{dx} (x^3) \right|_{x=1}.$$

Now we'll use limits to evaluate the derivative

$$\left. \frac{d}{dx} (x^3) \right|_{x=a}.$$

In other words, we'll compute the derivative of the function $f(x) = x^3$ at a general input $x = a$. Think about zooming in on the graph of $y = x^3$ sufficiently close to the point (a, a^3) so that the graph looks like a straight line. We'll compute the slope of that line.

The algebra here is nearly identical to what we did earlier. You should compare the two computations.

Desmos link: <https://www.desmos.com/calculator/8eiffwbgt5>

151: Cubing Function 3

The idea to compute the derivative algebraically is this: Fix the point $P(a, a^3)$ on the graph of $y = f(x) = x^3$. Then choose a variable point Q on the graph, different from P , with coordinates (v, v^3) . When Q is sufficiently close to P , the line PQ approximates the curve $y = x^3$ near $P(a, a)$ and the slope of this line is the derivative.

(a) Our first step is to find an expression for the slope of line PQ as a function

Learning outcomes:
Author(s):

of v . The slope is

$$\begin{aligned} m(v) &= \frac{\Delta y}{\Delta v} \\ &= \frac{f(v) - f(a)}{v - a} \\ &= \frac{v^3 - a^3}{v - a} \end{aligned}$$

all assuming $v \neq a$.

- (b) The next step is to factor the numerator $v^3 - a^3$. Since

$$(v^3 - a^3) \Big|_{v=a} = a^3 - a^3 = 0,$$

we know that $v - a$ is a factor of $v^3 - a^3$. To simplify the quotient

$$\frac{v^3 - a^3}{v - a}, \quad v \neq a,$$

we could use long division or factor the difference of two cubes:

$$A^3 - B^3 = (A - B)(A^2 + AB + B^2).$$

Either way, the result is that

$$v^3 - a^3 = (v - a)(v^2 + va + a^2).$$

So

$$\frac{v^3 - a^3}{v - a} = v^2 + va + a^2, \quad v \neq a.$$

Putting this all together, we get

$$\begin{aligned} \frac{d}{dx} (x^3) \Big|_{x=a} &= \lim_{v \rightarrow a} \frac{v^3 - a^3}{v - a} \\ &= \lim_{v \rightarrow a} \frac{(v - a)(v^2 + va + a^2)}{v - a} \\ &= \lim_{v \rightarrow a} (v^2 + va + a^2) \\ &= a^2 + a^2 + a^2 \\ &= 3a^2. \end{aligned}$$

- (c) More simply put, we just write the derivative

$$\frac{d}{dx} (x^3) = 3x^2$$

as a function of x .

(d) We can check our result using the desmos worksheet above. Do this as follows:

- (i) Input the correct expression for the derivative $d(x^3)/dx$ on Line 17.
- (ii) For a differentiable function $f(x)$ and values of h near zero, we can approximate the derivative $d(f(x))/dx$ as the slope

$$\frac{d}{dx}(f(x)) = \frac{f(x+h) - f(x-h)}{2h}$$

of the line through the points $(x-h, f(x-h))$ and $(x+h, f(x+h))$. Now activate the folder Approximation to the derivative function on Line 19. Then drag the slider h on Line 21 to get a better approximation to the derivative.

Free Response: Describe what happens to the above approximation as $h \rightarrow 0$.

Exercise 11 (a) Follow the method of Exploration 1 for the function $f(x) = x^4$ to compute the derivative

$$\left. \frac{d}{dx}(x^4) \right|_{x=a}.$$

- (b) Modify the desmos worksheet below for the function $f(x) = x^4$.

151: Cubing Function 3

Then to check your work, do part (d) of Exploration 1 for $f(x) = x^4$. Describe also what happens to the approximation in part (d) as $h \rightarrow 0$. Include a screenshot to help with your description.

Exercise 12 Repeat all parts of Exercise for the function $f(x) = 1/x^4$.

Exercise 13 Repeat all parts of Exercise for the function $f(x) = 1 + x^2$.

Exercise 14 Repeat all parts of Exercise for the function $f(x) = 1/(1 + x^2)$.

Free Fall

Rocks in free fall.

Question 15 Part (a) of the following question is an important takeaway from this chapter. It is not necessary to answer this question now.

A rock is dropped near the surface of a planet without an atmosphere.

- (a) Through what fraction of its initial height does the rock fall during the last $1/10$ of the time it takes to hit the surface? Give a quick estimate.
- (b) What is the exact fraction?

Exploration 16 Play the slider u in Line 1 of the desmos worksheet below to watch two motions. One is a rock falling to the surface of Mars when dropped from a height of 190 meters above the surface. The other is a constant-speed motion.

Desmos link: <https://www.desmos.com/calculator/j2ciuemii7>

151: Free Fall 1

- (a) Use the animation to sketch a graph of the function $s = g(t)$ that expresses the distance (in meters) of the rock from its starting point in terms of the number of seconds since the rock was released. Include the appropriate units and variable names on the axes.
- (b) Activate the folder Graph of distance function in Line 17 above to see how you did.
- (c) Our aim now is to compute the (instantaneous) speed of the rock (in m/s) at time $t = u$ seconds. We'll assume that

$$s = g(t) = 1.9t^2, \quad 0 \leq t \leq 10.$$

Do this as follows.

- (i) Find an expression that gives the rock's average speed between times $t = u$ and $t = v$.

Learning outcomes:
Author(s):

- (ii) Evaluate the appropriate limit to find the speed of the rock at time $t = u$ seconds. Click the Hint tab above for help.

Hint: The speed at time $t = u$ is

$$\begin{aligned}\frac{ds}{dt}\bigg|_{t=u} &= \lim_{v \rightarrow u} \frac{1.9v^2 - 1.9u^2}{v - u} \\ &= \lim_{v \rightarrow u} 1.9(v + u) \\ &= 3.8u.\end{aligned}$$

- (iii) Use the result of part (ii) to find the rock's speed as it hits the ground. Assume the rock is dropped from a height of 190 meters.
- (iv) Compare the rock's speed as it hits the ground with its average speed during the entire time of its fall.
- (v) Use the result of part (iv) to approximate the fraction of the total distance of 190 meters the rock falls through during the last second of its fall?
- (vi) Compare the rock's speed at time $t = u$ with its average speed during the first u seconds of its fall.
- (vii) Use point-slope to find an equation of the tangent line to the curve $s = 1.9t^2$ at the point $(u, 1.9u^2)$. Enter this equation in Line 21 of the worksheet above.
The point-slope equation is

$$s = 1.9u^2 + 3.8u(t - u).$$

- (viii) Activate the folder Average speed in Line 23 above. How is the slope of the tangent line related to the slope of the line OP ?
- (ix) Find the coordinates of the point where the tangent line intersects the horizontal axes. Enter these in Line 30 above.
The coordinates are

$$(t, s) = \left(\frac{u}{2}, 0\right).$$

Question 17 Play the slider u in Line 2 of the desmos worksheet below to watch two motions. One shows a rock falling to the surface of a planet when it is dropped from rest near the surface. The other is a constant-speed motion.

Desmos link: <https://www.desmos.com/calculator/dmlrxahkld>

- (a) What does the animation suggest about the speed of the rock as it hits the surface compared with its average speed during the time it takes to hit the ground?
- (b) What does the animation suggest about the fraction of its initial height the rock falls through during the last $1/20$ of the time it takes to hit the surface? Give a quick estimate.
- (c) Now suppose the rock is dropped from a height of H meters and that it takes T seconds to hit the surface. Assume also that the planet has no atmosphere so that the function expressing the rock's distance from its starting point (in meters) in terms of the number of seconds since it was dropped is of the form

$$s = f(t) = at^2$$

for some constant a .

- (i) Express the constant a in terms of H and T .
 - (ii) What are the units of the constant a ?
 - (iii) Explain the meaning of the constant a . Do not use the word acceleration.
 - (iv) Find the domain of the function f .
 - (v) Use calculus to express the speed of the rock (in metes/sec) in terms of H and T . Check that your expression has the correct units.
 - (vi) Compare the speed of the rock as it hits the ground with its average speed during the time it falls.
 - (vii) Through what exact fraction of its initial height the rock fall through during the last $1/20$ of the time it takes to hit the surface? Compare this with your earlier estimate.
-

Purchasing Power

How a small change in price affects the quantity of an item you can buy.

1 Purchasing Power

Question 18 This problem investigates how a small change in the price of apples would affect the number of pounds we can buy with \$10.

- (a) We'll start by finding a function

$$n = f(p), p > 0,$$

that expresses the number of pounds of apples we can buy with \$10 in terms of the price (measured in dollars/pound).

The function is

$$n = f(p) = 10/p, p > 0.$$

- (b) Would you expect the derivative dn/dp to be positive, negative, or zero? Explain.
- (c) Use limits to find an expression for the derivative

$$\left. \frac{dn}{dp} \right|_{p=a}.$$

The derivative is (click the Hint tab above for help)

Hint:

$$\begin{aligned} \left. \frac{dn}{dp} \right|_{p=a} &= \lim_{p \rightarrow a} \frac{1}{p-a} (f(p) - f(a)) \\ &= \lim_{p \rightarrow a} \frac{1}{p-a} \left(\frac{10}{p} - \frac{10}{a} \right) \\ &= \lim_{p \rightarrow a} \frac{10}{p-a} \left(\frac{a-p}{pa} \right) \\ &= \lim_{p \rightarrow a} \frac{-10}{pa} \\ &= \frac{-10}{a^2}. \end{aligned}$$

Learning outcomes:
Author(s):

- (d) Evaluate the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

- (i) What are the units of the above derivative?
 - (ii) What do you get by “simplifying” the units? Explain how simplifying the units of this derivative gives you insight into its meaning.
- (e) Interpret the meaning of the above derivative. Click the arrow to the lower right for help.

For a way to interpret the meaning of the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

we can ask ourselves what happens to the number of pounds of apples we can buy with \$10 if the price changes by a small amount from \$2/lb. For this, let Δp be a small change in price from \$2/lb (measured in dollars/lb) and let Δn be the corresponding change in the number of pounds we can buy with \$10.

For $\Delta p \sim 0$, we have

$$-2.5 = \left. \frac{dn}{dp} \right|_{p=2} \sim \frac{\Delta n}{\Delta p}$$

and (type “Delta” for Δ)

$$\Delta n \sim -2.5 \Delta p.$$

This tells us that if the price increases from \$2/lb to say \$2.10/lb, then

$$\Delta n \sim \left(-2.5 \frac{\text{lb}}{\$/\text{lb}} \right) \left(0.10 \frac{\$}{\text{lb}} \right) = -0.25 \text{ lbs.}$$

So if the price increases from \$2/lb to say \$2.10/lb, we can buy about 0.25 fewer pounds of apples with ten dollars.

- (f) Next we’ll use the worksheet below to visualize the approximate change

$$\begin{aligned} \Delta n &\sim \left(\left. \frac{dn}{dp} \right|_{p=2} \right) \Delta p \\ &= \left(-2.5 \frac{\text{lb}}{\$/\text{lb}} \right) \left(0.10 \frac{\$}{\text{lb}} \right) \\ &= -0.25 \text{ lbs.} \end{aligned}$$

in the number of pounds of apples we can buy with \$10 as the price increases from \$2/lb to \$2.10/lb.

Desmos link: <https://www.desmos.com/calculator/qw7wis1q0c>

151: Apples

- (i) To get started, find an equation (in point-slope form) of the tangent line to the curve $n = f(p)$ at the point P with coordinates $(2, 5)$. The tangent line has a slope equal to the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

So its equation is

$$n = 5 + -2.5(p - 2).$$

- (ii) Enter your equation of the tangent line on Line 17 of the desmos worksheet.
- (iii) Activate the folders tangent line and linear approximation on Lines 18 and 22.
- (iv) Explain why the difference in the n -coordinates of points P and R' (ie. n -coordinate of R' minus n -coordinate of P) is equal to our approximation of Δn above when Q as coordinates $(2.10, f(2.10))$.
- (v) Drag Slider v on Line 2 to make Q approach P . What so you think happens to the ratio of Δn to our approximation of Δn as $v \rightarrow 2$?
- (g) We can get a better understanding of the changes

$$\Delta p = 2.1 - 2 = 0.10 \text{ dollars/pound}$$

and

$$\Delta n = f(2.1) - f(2) \sim 0.25 \text{ pounds}$$

by thinking about relative instead of absolute change.

An increase in price from \$2/lb to \$2.10/lb is a relative change of

$$\frac{\Delta p}{p} = \frac{\$0.1/\text{lb}}{\$2/\text{lb}} = 0.05 = 5\%.$$

This causes a relative change in the number of pounds we can buy of approximately

$$\frac{\Delta n}{n} \sim \frac{-0.25\text{lbs}}{5\text{lbs}} = -0.05 = -5\%.$$

- (h) We can get a the same result relating the two relative changes of the previous question by working in general.

Suppose we increase the price of apples by $Q\%$, where $Q \sim 0$. What can we say about the relative change in the number of pounds of apples we can buy with \$10?

Well, if the change in price Δp is near zero and Δn is the corresponding change in the number of pounds we can buy, then

$$\frac{\Delta n}{\Delta p} \sim \frac{dn}{dp} = -10/p^2.$$

So

$$\Delta n \sim \left(\frac{-10}{p^2} \right) \Delta p.$$

Dividing both sides by $n = 10/p$ tells us that

$$\frac{\Delta n}{n} \sim \left(\frac{-10}{p^2} \right) \left(\frac{\Delta p}{\frac{10}{p}} \right) = -\frac{\Delta p}{p} = -Q\%.$$

So if we increase the price by $Q\% \sim 0$, then the number of apples we can buy decreases by approximately that same $Q\%$.

- (i) We can think about this relationship between the relative changes geometrically. To do this, remove your equation of the tangent line on Line 17 and activate the folder tangent line in the worksheet above. Then drag slider a on Line 4 to move point P . What do you notice about the ratio $PB : PA$ of the distances PA and PB as P moves? More on this later.

Question 19 Due to a printing error, the graph of the function

$$n = f(p), p > 0,$$

expressing the number of pounds of an item we can buy with \$100 in terms of the price (measured in \$/lb) is not shown below. All we can see is a point A on the graph.

Desmos link: <https://www.desmos.com/calculator/y5nqs8fkvj>

151: Printers Error 1

The problem is to draw the tangent line to the graph at A without sketching the graph or doing any kind of computation. Click the arrow below for a hint.

The key idea is to relate the slope of the tangent line at A to the slope of the line OA through the origin and A .

We know that

$$n = f(p) = \frac{100}{p}, p > 0,$$

and

$$\frac{dn}{dp} = -100/p^2.$$

Now write this derivative in terms of both n and p ,

$$\frac{dn}{dp} = -\frac{100}{p^2} = -\frac{n}{p}.$$

Now relate this to the slope of line OA and draw the tangent line at A .

2 Weight in Space

Question 20 The weight of an object is the gravitational force that the earth exerts on the object's mass and varies with the object's height above the surface.

The function

$$W = f(h) = \frac{k}{(h + 4)^2}, \quad h \geq 0,$$

expresses the weight (in pounds) of an object in terms of its height (in thousands of miles) above the surface of the earth. We'll suppose the object weighs 200 pounds on the surface.

Desmos link: <https://www.desmos.com/calculator/zklsucctjp>

151: Weight in Space

- Find the value of the constant k . What are its units?
- Find the average rate of change of W with respect to h over the interval between heights of h and v thousands of miles.
- Use part (b) to find an expression for the derivative dW/dh .
- Use the graph of the function $W = f(h)$ above to approximate the derivative

$$\left. \frac{dW}{dh} \right|_{h=4}.$$

Include units

- Evaluate the derivative

$$\left. \frac{dW}{dh} \right|_{h=4}$$

using your expression for dW/dh and compare this with your estimate.

- What are the units of the derivative above? Interpret its meaning.

- (i) At approximately what altitude does your weight decrease by 4 pounds when your altitude increases by 200 miles? Use your expression for the derivative dW/dh to help.
- (ii) Compute the exact change in your weight over the interval you found in part (i).

3 A Rock in Free Fall

Question 21 A rock is dropped from a height of 100 meters on the planet Krypton.

The function

$$v = f(h), 0 \leq h \leq 100,$$

expresses the speed (in meters/sec) of the rock in terms of its height (in meters) above the surface.

Desmos link: <https://www.desmos.com/calculator/hcqb5nxc18>

151: Free Fall Speed and Height

- (a) Would you expect the derivative

$$\left. \frac{dv}{dh} \right|_{h=36}$$

to be positive or negative? Why?

- (b) Suppose

$$v = f(h) = 4\sqrt{100 - h}, 0 \leq h \leq 100$$

and find an expression for the average rate of change of the rock's speed (measured in meters/sec) with respect to its height (in meters) between heights of v meters and a meters.

- (c) Use your expression from part (b) to find an expression for the derivative

$$\left. dv/dh \right|_{h=a}.$$

- (d) Use the graph above to approximate the derivative

$$\left. dv/dh \right|_{h=64}.$$

Include units.

- (e) Evaluate the derivative

$$dv/dh \Big|_{h=64}$$

and compare the exact value with your estimate.

- (f) What are the units of the derivative above? What is its meaning?
- (g) “Simplify the units of the derivative”. What insight does this give you into its meaning?
- (h) Use part (e) to approximate the rock’s speed at a height of 61 feet. Compare your estimate with the actual speed.
- (i) Use part (c) to approximate the interval over which the rock falls two meters as its speed increases by 0.5m/s.

4 Distance to the Horizon

Question 22 The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 10,000$$

expresses the distance to the horizon (measured in miles) in terms of your altitude (measured in feet).

- (a) Find an expression for the average rate of change in the distance to the horizon with respect to altitude between altitudes h feet and v feet.
- (b) Use your expression from part (a) to find an expression for the derivative ds/dh .
- (c) Evaluate the derivative

$$\frac{ds}{dh} \Big|_{h=25}.$$

- (d) What are the units of the derivative above? Interpret its meaning.
- (e) Use the result of part (c) to approximate the distance to the horizon at an altitude of 24 feet. Then compare this approximation with the actual distance.
- (f) Approximate the relative change in the distance to the horizon in terms of a small relative change in altitude.
- (g) Use your result from part (b) to approximate the altitude at which moving 10 feet higher increase the distance to the horizon by 0.5 miles.

5 Intensity of Sound

Question 23 The intensity of sound is measured in Watts per square meter and is a function of the distance from the source.

Suppose for a jet taking off, this function is

$$I = f(r) = \frac{100}{r^2}, \quad r \geq 10,$$

where r is the distance (in meters) from the jet.

- (a) Find an expression for the average rate of change of the sound intensity (in Watts/m²) with respect to the distance from the source (in meters) between distances r and a meters.
- (b) Use your expression from part (a) to find an expression for the derivative

$$dI/dr \Big|_{r=a}$$

- (c) Evaluate the derivative

$$dI/dr \Big|_{r=20}.$$

- (d) What are the units of the derivative above? What is its meaning?
- (e) Approximate the relative change in the sound intensity in terms of a small relative change in the distance to the source.



Practice Quiz 1

First practice quiz, Weeks 1-2

Directions:

- (a) Show all work.
- (b) Give brief explanations for each problem. Include these explanations in the flow of the solution.
- (c) Show all units in all computations.

1 Part 1

Question 24 Explain what it means intuitively for a function to be differentiable at some input in terms of the graph of that function.

Question 25 The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 10,000,$$

expresses the distance to the horizon (measured in miles) in terms of your altitude (measured in feet).

- (a) Find an expression for the average rate of change in the distance to the horizon with respect to altitude between altitudes b feet and v feet.
- (b) Use your expression from part (a) to find an expression for the derivative

$$\left. \frac{ds}{dh} \right|_{h=b}.$$

- (c) Evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=25}.$$

- (d) What are the units of the derivative above? Interpret its meaning.

Learning outcomes:
Author(s):

- (e) Use the result of part (c) to approximate the distance to the horizon at an altitude of 24 feet. Then compare this approximation with the actual distance.
- (f) Approximate the relative change in the distance to the horizon in terms of a small relative change in altitude.
- (g) Use your result from part (b) to approximate the altitude at which moving 10 feet higher increase the distance to the horizon by 0.5 miles.

Question 26 The intensity of sound is measured in Watts/(square meter) and is a function

$$I = f(r) = \frac{100}{r^2}, \quad r \geq 0.5,$$

of the distance from the source.

Suppose r is measured in meters and that the intensity of the sound emitted by a vacuum cleaner is 10^{-4} watts/ m^2 at a distance of 0.5 meters.

- (a) Compute the value of the constant k . What are its units?
- (b) Find an expression for the average rate of change of the sound intensity (in Watts/ m^2) with respect to the distance from the source (in meters) between distances b and r meters.
- (c) Use your expression from part (b) to find an expression for the derivative

$$\left. \frac{dI}{dr} \right|_{r=b}.$$

- (d) Evaluate the derivative

$$\left. \frac{dI}{dr} \right|_{r=2}.$$

- (e) What are the units of the derivative above? What is its meaning?
- (f) Approximate the relative change in the sound intensity in terms of a small relative change in the distance to the source.

Question 27 Between speeds of 70 miles/hr and 84 miles/hr, the gas mileage of a car (in miles/gal) is a one-to-one function $G = f(v)$ of its speed (in miles/hour). The car gets 10 miles/gal at a speed of 80 miles/hour.

- (a) Which of the following is more likely to be true? Explain your reasoning.

(i) $\left. \frac{dG}{dv} \right|_{v=80} = 0.25$ or

(ii) $\left. \frac{dG}{dv} \right|_{v=80} = -0.25$

(b) What are the units of the correct derivative above? Explain its meaning.

(c) Assuming the correct choice in part (b), evaluate the derivative

$$\left. \frac{dv}{dG} \right|_{G=10}.$$

(d) Simplify the units of the derivative in part (c). What does this tell you about its meaning?

(e) At what rate (in gal/hour) does the car burn gas at a speed of 80 miles/hour?

Question 28 Between speeds of 55 miles/hour and 70 miles/hour the gas mileage of a car is a linear function of its speed. The car gets 40 miles/gal at a speed of 55 miles/hour and 30 miles/gal at a speed of 70 miles/hour.

(a) Find a function

$$r = f(v), \quad 55 \leq v \leq 70,$$

that express the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hour). Note: This function is not linear

(b) Find an expression for the average rate of change of r with respect to v between speeds of b miles/hour and w miles/hour.

(c) Use your expression from part (b) to find an expression for the derivative

$$\left. \frac{dr}{dv} \right|_{v=b}.$$

(d) Evaluate the derivative

$$\left. \frac{dr}{dv} \right|_{v=60}.$$

(e) What are the units of the derivative above? Interpret its meaning.

Introduction to Motion

An introduction to motion

Exploration 29 *Desmos link:*

<https://www.desmos.com/calculator/bk1z9cwbhb>

151: Sound Squaring Function

Learning outcomes:
Author(s):

Introduction

Introduction to Differential Calculus

This course is listed as Calculus I in the catalogue, but it really should be called Differential Calculus. *Differential* has the same root as *difference* and *calculus* the same root as *calculate*. So this class is really about calculating differences, or more simply put it's about subtracting. But it's about subtracting in the context of functions.

Pick a specific input to specific function and you'll likely find that the changes in the function's output are approximately proportional to *small* changes in the input. If so, we say that the function is *differentiable* at that input and we call the proportionality constant the *derivative*.

For example, let's look at the behavior of the function

$$A = f(s) = s^2, s \geq 0,$$

near the input $s = 5$. To emphasize the importance of units, let's define the input s to be the side length of a square measured in feet and the output $f(s)$ to be the area of that square, measured in square feet. The problem before us is to describe a simple relationship between a small change in the side length

$$\Delta s = s - 5$$

of the square and the change

$$\Delta A = f(s) - f(5)$$

in its area.

Question 30 (a) We'll first take a numerical approach and compute some small changes and their ratios. Fill in the missing entries in the table below.

s (ft)	$A = s^2$ (ft ²)	$\Delta s = s - 5$ (ft)	$\Delta A = s^2 - 25$ (ft ²)	$\Delta A / \Delta s$ (ft ² /ft)
4.9	24.01	-0.1	-0.99	9.9
4.99	24.9001	-0.01	-0.0999	9.99
5	25	0	0	—
5.01	25.1001	0.01	0.1001	10.01
5.1	26.01	0.1	1.01	10.1

Learning outcomes:
Author(s):

(b) The data in the table above suggests an approximate proportional relationship between ΔA and Δs . We can guess the constant of proportionality from the fifth column. As $s \rightarrow 5$ (as s approaches 5), it looks like the ratio $\Delta A/\Delta s$ approaches some number, the constant of proportionality.

i) What is that number? 10

ii) What are its units?

Free Response:

So for $\Delta s \sim 0$, we suspect that

$$\Delta A \sim 10\Delta s.$$

(c) The constant of proportionality is called the derivative, in this case of the function $A = s^2$, at the input $s = 5$. We write this as

$$\left. \frac{dA}{ds} \right|_{s=5} = 10.$$

(d) We could have taken an algebraic approach to determine this constant of proportionality instead. The idea is to first simplify the quotient $\Delta A/\Delta s$ as

$$\begin{aligned} \frac{\Delta A}{\Delta s} &= \frac{s^2 - 25}{s - 5} \\ &= \frac{(s + 5)(s - 5)}{s - 5} \\ &= s + 5 \text{ if } s \neq 5. \end{aligned}$$

So, for example, if $s = 4.99$, then

$$\frac{\Delta A}{\Delta s} = 4.99 + 5 = 9.99$$

as shown in the last column of the second row of the above table.

The advantage of this algebraic approach is that we can now compute the proportionality constant as a limit:

$$\begin{aligned} \left. \frac{dA}{ds} \right|_{s=5} &= \lim_{s \rightarrow 5} \frac{\Delta A}{\Delta s} \\ &= \lim_{s \rightarrow 5} (s + 5) \\ &= 5 + 5 \\ &= 10. \end{aligned}$$

(d) We can also use the graph of the function $A = f(s) = s^2$ to interpret the ratios

$$\frac{\Delta A}{\Delta s} = \frac{f(s) - f(5)}{s - 5}$$

geometrically. Move the slider s in the demonstration below and describe

- (i) how the line through the points P and Q is related to the ratio $\Delta A/\Delta s$ show on Line 2,
- (ii) what happens to the line PQ as $s \rightarrow 5$, and
- (iii) what happens to the line PQ when $s = 5$.

Free Response:

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/vz9ud5txva>

Continuing with the above demonstration,

- (i) Open the Code folder in Line 3 and turn off the line PQ in Line 7.
- (ii) Write an equation for the line through the point P with slope equal to the proportionality constant in the line below and on Line 8 in the desmos worksheet:

$$A = L(s) = 25 + 10(s - 5).$$

- (iii) Zoom in close enough to the point P to make the graph of the function $A = f(s)$ look like a line. How do the graph of the function and the graph of the line $A = L(s)$ compare in this close-up view?

Free Response:

(e) **Summary:**

- If we change the side of a square from a length of 5 feet to a length of $s \sim 5$ feet, then the area of the square changes by approximately

$$\Delta A = s^2 - 25 \sim 10\Delta s = 10(s - 5)$$

square feet. The proportionality constant 10 has units $ft^2/ft = ft$.

- Zoom in close enough to the graph of the function $A = f(s) = s^2$ near the point $P(5, 25)$ and the graph looks like a line with slope equal to the proportionality constant.

- We can compute the proportionality constant as the limit

$$\left. \frac{dA}{ds} \right|_{s=5} = \lim_{s \rightarrow 5} \frac{f(s) - f(5)}{s - 5}.$$

- Suppose for example, we wanted to approximate the side length s of a square with area 25.06 ft^2 . Then

$$\Delta A = s^2 - 25 = 25.06 - 25 = 0.06.$$

And since

$$\Delta A \sim 10\Delta s = 10(s - 5),$$

$$0.06 \sim 10\Delta s.$$

So

$$\Delta s \sim 0.006$$

and a square with area 25.06 ft^2 has an approximate side length (measured in feet) of

$$s = 5 + \Delta s \sim 5.006.$$

Question 31 On a clear day with an unobstructed view (like you might have at the beach or in a hot air balloon), the distance to the horizon is limited by the curvature of the earth as illustrated in the demonstration below.

In fact, as long as you are not too high above the surface of the earth, the function

$$s = f(h) = 1.22\sqrt{h}, 0 \leq h \leq 20,000,$$

gives a good approximation to the distance to the horizon (the length of the red arc AT below, measured in miles) in terms of your height above the ground (the distance AP below, measured in feet).

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

Desmos activity available at

[151:Distance to Horizon 1](#)

Our aim is to approximate the change in the distance to the horizon (in miles) in terms of a small change in height (in feet) from a height of 25 feet.

(i) To start, what are the units of the constant 1.22 above? Explain how you know.

Free Response:

(ii) Go through a similar analysis as in parts (a)-(e) of Example 1, to approximate the change $\Delta s = s - f(25)$ in the distance to the horizon in terms of the change $\Delta h = h - 25$ in your height above the ground. Start by completing the column headings (with units) and the missing entries in the table below.

h (ft)	$s = 1.22\sqrt{h}$ (miles)	$\Delta h = h - 25$ (ft)	$\Delta s = f(h) - f(25)$ (miles)	$\Delta s/\Delta h$ (units?)
4.9^2				
4.99^2				
25		0	0	—
5.01^2				
5.1^2				

Question 32 This question is similar to the last, but suppose instead we are looking down on the earth from the space station or a rocket. Then the approximation to the distance to the horizon from the previous problem will not work.

So our first step is to find a function

$$s = f(h), h \geq 0,$$

that expresses the distance to the horizon (still measured in miles) in terms of our height above the earth's surface, now measured in miles instead of feet. We'll suppose the earth to be a perfect sphere of radius 3960 miles. The distance to the horizon is the arclength AT below, measured along the surface of the earth (you can think of this distance as the radius of the spherical disk visible to us). Our height is the distance AP .

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

(a) Find an expression for the above function.

Hint: Use right triangle $\triangle OTP$ to find an expression for the radian measure of angle $\angle POT$. Then use this angle to find an expression for the arclength AT .

Here are more details.

(i) Enter the two side lengths, measured in feet, in right triangle $\triangle OPT$ below.

$$OT = 3960$$

and

$$OP = h + 3960.$$

(ii) Let θ be the radian measure of $\angle TOP$. Write an equation with a trigonometric function of θ that relates the two lengths in part (i). Use the Math Editor tab to enter the trig function and the angle θ .

$$\cos \theta = \frac{3960}{h + 3960}.$$

(iii) Now solve the equation from part (ii) for θ in terms of h . Then use what you know about measuring arclength along a circle to find an expression for the function f . Use the Math Editor tab to help.

$$s = f(h) = 3960 \arccos\left(\frac{3960}{h + 3960}\right).$$

(b) Now suppose we are 165 miles above the surface of the earth and we wish to approximate how a small change in our altitude changes the distance to the horizon.

To do this, fill in the missing entries in the table below.

h (miles)	$s = f(h)$ (miles)	$\Delta h = h - 165$ (miles)	$\Delta s = f(h) - f(165)$ (miles)	$\Delta s / \Delta h$ (units?)
162				
163				
164				
165				
166				
167				
168				

(c) Do the data above suggest that the quotients $\Delta s / \Delta h$ approach some number as h approaches 165? If so, use the data to approximate that number. If not, explain why not.

(d) Make your own table similar to the one above to get a better approximation, correct to the nearest thousandth, to

$$\lim_{h \rightarrow 165} \frac{f(h) - f(165)}{h - 165}.$$

(e) Use your result from part (d), rounded to the nearest thousandth, to approximate Δs in terms of Δh and enter your result below.

$$\Delta s \sim 3.291 \Delta h, \text{ for } \Delta h \sim 0.$$

(f) Explain the meaning of the proportionality constant in parts (d) and (e). Be sure to include units in your explanation.

Question 33 The function

$$h = f(v), 80 \leq v \leq 120,$$

expresses the height of a helicopter (measured in feet) in terms of its speed (measured in ft/sec).

Suppose

$$\left. \frac{dh}{dv} \right|_{v=100} = -20.$$

- (a) What are the units of the above derivative?
 - (b) Explain the meaning of the derivative using the language of small changes.
-

Small Changes

We explore how small changes to the input of a function change the output.

The main idea of differential calculus is to approximate the change in the output of a function in terms of a small change in the input. For some functions, called *differentiable*, the change in the output is approximately proportional to the (small) change in the input. The proportionality factor is called the derivative. In this chapter we explore this idea.

1 Odometer Readings

Example 1. The graph of the function

$$s = f(t), 0 \leq t \leq 2,$$

that expresses the trip odometer reading (measured in miles) on your car in terms of the number of hours past noon during a two-hour trip is shown below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/iw69lr1bc>

Desmos activity available at

151: Odometer

Our goal is to approximate the car's speed at 12:30pm in three ways:

- (1) geometrically, using the above graph as is.
- (2) geometrically, by zooming in on the above graph.
- (3) arithmetically, using the specific expression for the function f .

(a) Start by using the graph above to describe how the speed of the car varies over the two-hour period. Explain your reasoning. Then play the Slider u in Line 2 and use the animation of the motion to check if your description was accurate. Explain.

Learning outcomes:
Author(s):

Small Changes

(b) Set the slider in Line 6 to $n = 20$. Use *only* the graph (with $u = 2$) to create a table with five columns showing the values of t , s (approximate this), $\Delta t = t - 0.5$, $\Delta s = f(t) - f(0.5)$, and $\Delta s/\Delta t$. Include units in the heading of each column. The table should include five rows, with $t = 0.3, 0.5, \dots 0.7$.

t (hours)	$s = f(t)$ (miles)	$\Delta t = t - 0.5$ (hrs)	$\Delta s = f(t) - f(0.5)$ (miles)	$\Delta s/\Delta t$ (miles/hr)
0.3				
0.4				
0.5				
0.6				
0.7				

(c) Explain the meaning of the fifth column in the table of part (b). What do the entries in this column suggest about the speed of the car at 12:30pm?

(d) Now we'll use the fact that

$$s = f(t) = 48t^2 - 16t^3, 0 \leq t \leq 2,$$

to construct another table like the one in part (b). Do this by finding expressions for

$$\Delta s = f(t) - f(0.5)$$

and

$$r = g(t) = \frac{\Delta s}{\Delta t} = \frac{f(t) - f(0.5)}{t - 0.5},$$

both in terms of t (and *not* Δt). Use these functions to fill in the missing entries in the table below.

t (hours)	$s = f(t)$ (miles)	$\Delta t = t - 0.5$ (hrs)	$\Delta s = f(t) - f(0.5)$ (miles)	$\Delta s/\Delta t$ (miles/hr)
0.49				
0.499				
0.4999				
0.5				
0.5001				
0.501				
0.51				

(e) Does your table from part (d) suggest that the ratios $\Delta s/\Delta t$ approach some number as $t \rightarrow 0.5$? If so, what would be your guess for the exact value of this number? What are its units? What is its meaning?

(f) Activate the folder "Graph of average speed" on Line 9.

(i) Use the graph to check some your entries in the fifth column of your table from part (d). Explain.

- (ii) How is the line PQ , through the fixed point $P(0.5, 10)$ and the variable point $Q(t, s)$ on the graph of the function $r = g(t)$ related to the ratio $\Delta s / \Delta t$?
- (iii) Describe what happens to the line PQ as point Q approaches point P .
- (g) For a quicker way to approximate car's speed at 12:30pm, zoom in sufficiently close to point P in the graph above to make the graph of $s = f(t)$ look like a line. Use the coordinates of point P and a second point in the window far away from P to estimate the car's speed at 12:30pm. Explain your method.
- (h) Summarize your conclusions by comparing your three estimates for the car's speed at 12:15pm. Which estimate do you think is most accurate? Least accurate?

Example 2. This is a continuation of the previous example where we'll algebraically compute the exact speed of the car at 12:30pm, using the odometer function

$$s = f(t) = 48t^2 - 16t^3, 0 \leq t \leq 2.$$

The idea is to first find an algebraic expression for the car's average speed between time t and time $t = 0.5$ hours past noon. Then we'll evaluate the limit of this average speed as $t \rightarrow 0.5$ to find the (instantaneous) speed at 12:30pm.

Question 34 First we'll find the average speed between time t and time $t = 0.5$.

(a) Explain in general how to compute a car's average speed over some time interval. What do you need to know? What is the computation? Make up your own specific example.

(b) Now for our particular odometer function above, the average speed $v_{\text{avg}}(t)$, measured in miles/hour, between time t and time $t = 0.5$ is

$$\begin{aligned} v_{\text{avg}}(t) &= \frac{f(t) - f(0.5)}{t - 0.5} \\ &= \frac{48t^2 - 16t^3 - 10}{t - 0.5} \\ &= \frac{(2t - 1)(-8t^2 + 20t + 10)}{t - 0.5} \\ &= -16t^2 + 40t + 20 \text{ if } t \neq 0.5. \end{aligned}$$

The key step in the computation above is in the third line. How did we know $2t - 1$ was a factor of

$$f(t) - f(0.5) = 48t^2 - 16t^3 - 10?$$

The reason is that $t = 0.5$ is a root of the polynomial $f(t) - f(0.5)$ and therefore $t - 0.5$ is a factor. And so

$$2(t - 0.5) = 2t - 1$$

is also a factor. Then we can use long division to find the quotient.

(c) Show the steps in the long division.

(d) The final step in computing the car's speed v (in miles/hour) at 12:30pm is to evaluate the limit of these average speeds as $t \rightarrow 0.5$. We get

$$\begin{aligned} v &= \lim_{t \rightarrow 0.5} (-16t^2 + 40t + 20) \\ &= 36. \end{aligned}$$

(e) Here's another way to simplify the average speed in part (b). Fill in the missing steps.

$$\begin{aligned} v_{\text{avg}}(t) &= \frac{f(t) - f(0.5)}{t - 0.5} \\ &= \frac{(48t^2 - 16t^3) - (48(0.5)^2 - 16(0.5)^3)}{t - 0.5} \\ &= \frac{(48t^2 - 48(0.5)^2) - (16t^3 - 16(0.5)^3)}{t - 0.5} \\ &= \frac{48(t^2 - (0.5)^2) - 16(t^3 - (0.5)^3)}{t - 0.5} \\ &= \frac{48(t - 0.5)(t + 0.5) - 16(t - 0.5)(t^2 + 0.5t + 0.25)}{t - 0.5} \\ &= 48(t + 0.5) - 16(t^2 + 0.5t + 0.25) \text{ if } t \neq 0.5. \end{aligned}$$

(f) Use the above expression for the average speed function to compute the (instantaneous) speed of the car at 12:30pm by evaluating the appropriate limit.

(g) Sketch by hand a graph of the average speed function $y = v_{\text{avg}}(t)$ over the appropriate domain. Be sure also to state this function's domain.

2 A Projectile

Example 3. Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/l4fknr0hpl>

Desmos activity available at

151: Projectile

3 The Falling Ladder, Part 1

Example 4. The top end of a ten-foot ladder leans against a vertical wall and the bottom end rests on the horizontal floor. We analyze how a small change in the distance between the wall and the bottom of the ladder affects the height of the ladder's top above the floor.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/dvyuifyyg4>

Desmos activity available at

151: Ladder 1B

Question 35 (a) The slider s in Line 1 of the demonstration above controls the distance between the wall and the bottom of the ladder, measured in feet. Use the slider s to describe qualitatively how a small change in s changes the height h (also measured in feet) of the ladder's top end above the floor.

(i) Do the small changes have the same or opposite signs?

(ii) At what positions of the ladder does a small change in s result in a comparatively large change in h ?

(b) Now let's focus on the particular position of the ladder where the bottom end is $s = 8$ feet from the wall. For this, turn on the "one position" folder in Line 3.

(i) Drag the slider s close to $s = 8$ and use the coordinates of the endpoints of the ladder to construct a table with five columns showing the values of s , h , $\Delta s = s - 8$, $\Delta h = f(s) - f(8)$, and $\Delta h/\Delta s$. Include units in the heading of each column. The table should include seven rows, with $s = 7.7, 7.8, \dots, 8.2, 8.3$. Here $h = f(s)$ is the function described in part (ii) below.

(ii) Find a function

$$h = f(s), 0 \leq s \leq 10,$$

that expresses the height of the ladder's top end above the ground (in feet) in terms of the distance of its bottom end from the wall (in feet).

(iii) Use your function f to construct another table, like the one in part (i), with $s = 7.9, 7.99, 7.999, 8, 8.0001, 8.01, 8.1$. Do this by finding expressions for

$$\Delta h = f(s) - f(8)$$

and

$$r = g(s) = \frac{\Delta h}{\Delta s} = \frac{f(s) - f(8)}{s - 8},$$

both in terms of s (and not Δs).

(iv) Does your table from part (iii) suggest that the ratios $\Delta h/\Delta s$ approach some number as $s \rightarrow 8$? If so, what would be your guess for the exact value of this limit? What are its units? What is its meaning?

(v) Activate the folder “graph of function” on Line 8. How is the line PQ , through the fixed point $P(8, 6)$ and the variable point $Q(s, h)$ on the graph of the function $h = f(s)$ related to the ratios $\Delta h/\Delta s$?

(vi) Change the bounds for s in Line 1 to run between $s = 7.9$ and $s = 8.1$. Then activate the folder “graph: average rate of change function” on Line 18. Move the slider s and use the graph of the function $r = g(s)$ to check your computations in part (iii).

(vii) Use the result of part (iv) to write an approximation for the change in height

$$\Delta h = h - 8$$

in terms of the change

$$\Delta s = s - 8.$$

(viii) The graph of the function $h = f(s)$ suggests another, geometric way to find the proportionality constant (of part (vii)) that relates Δh to Δs . Explain how.

4 The Falling Ladder, Part 2

Example 5. A tree leans precariously with its trunk making an angle of $\phi = \pi/6$ radians with the ground. One end of a ten-foot ladder leans against the trunk, the other rests on the horizontal ground. We analyze how a small change in the distance between the bottom of the ladder and the base of the trunk changes the distance between the top of the ladder and the base of the trunk.

Access Geogebra interactives through the online version of this text at

Geogebra link: <https://www.geogebra.org/m/qmke5y7x>

We'll let t be the distance between the top of the ladder and the base of the trunk (measured in feet) and s the distance between the bottom of the ladder and the base of the trunk (also measured in feet).

The slider θ in the demonstration above controls the angle that the ladder makes with the ground, but this angle does not come into play in our problem.

(a) Use the slider θ to describe qualitatively how a small change in s (the length of segment GC) changes t (the length of segment GB):

(i) For what positions of the ladder do these small changes have the same signs? Opposite signs?

(ii) For what positions of the ladder does a small change in s result in a comparatively large change in t ?

(b) Now let's focus on the particular position of the ladder when the bottom end C is 16 feet from the trunk's base and the top end D is about 8 feet from the base *as illustrated above*. Our first step is to find a function

$$t = f(s)$$

that expresses t in terms of s for values of s and t near $s = 16$ and $t = 8$ respectively.

To do this, first use the law of cosines to write an equation relating s and t . Then *complete the square* to solve this equation for t in terms of s to find the function $t = f(s)$. Keep in mind that when $s = 16$, we must have $t \sim 8$.

Hint: Here is an outline of the steps to check your work:

(i) Use the law of cosines to relate s and t .

$$s^2 + t^2 - \sqrt{3}st = 100.$$

(ii) Solve the above equation for t in terms of s as follows:

First rewrite the equation with the two terms with t as a factor on the left side and the other two terms on the right.

$$t^2 - \sqrt{3}st = 100 - s^2.$$

Then complete the square by adding the same perfect square to each side.

$$t^2 - \sqrt{3}st + \left(\frac{\sqrt{3}s}{2}\right)^2 = 100 - s^2 + \left(\frac{\sqrt{3}s}{2}\right)^2.$$

Then factor the LHS and simplify the RHS.

$$\left(t - \frac{\sqrt{3}s}{2}\right)^2 = 100 - \frac{s^2}{4}.$$

Next, solve for t in terms of s .

$$t = \frac{\sqrt{3}}{2}s \pm \sqrt{100 - \frac{s^2}{4}}$$

Finally, make the correct choice of \pm to solve for t in terms of s , when s is near 16 and t near 8.

$$t = \frac{\sqrt{3}}{2}s - \sqrt{100 - \frac{s^2}{4}}.$$

(c) Use your function from part (b) to find an expressions for

$$\Delta t = f(s) - f(16)$$

and for the function

$$r = g(s) = \frac{\Delta t}{\Delta s} = \frac{f(s) - f(16)}{s - 16}.$$

Explain what the output of the function g measures. What are its units?

(d) Use the results of part (c) to construct a table with five columns showing the values of s , t , $\Delta s = s - 16$, $\Delta t = f(s) - f(16)$, and $\Delta t/\Delta s$. Include units in the heading of each column. The table should include seven rows, with $s = 15.9, 15.99, 15.999, 16, 16.001, 16.01, 16.1$.

(e) Does your table from part (d) suggest that the ratio $\Delta t/\Delta s$ approaches some number as $s \rightarrow 16$? If so, approximate the value of this number. What are its units?

(f) Check the box “GraphofRelation” in the demonstration above and explain how the line EF is related to part (d).

(g) Use the result of part (e) to write an approximation for the change

$$\Delta t = f(s) - f(16)$$

in terms of the change

$$\Delta s = s - 16$$

for values of s near 16. Use this approximation to estimate the distance between the top of the ladder and the base of the trunk when the bottom of the ladder is 16.4 feet from the trunk’s base. Compare your approximation with the exact distance.

5 Riding a Ferris Wheel

Suppose you ride a ferris wheel

Limits

Limits in context.

1 Limits and Tangent Lines

Example 6. Let

$$g(x) = \frac{x^2 - 9}{3x - 9}.$$

(a) Evaluate each of the following expressions.

(i) $g(7)$

(ii) $\lim_{x \rightarrow 7} g(x)$

(iii) $g(3)$

(iv) $\lim_{x \rightarrow 3} g(x)$

(b) Simplify and then graph the function $g(x)$.

(c) Interpret the expressions in part (a) geometrically by considering the graph of the function $f(x) = x^2/3$ as in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/u0uvuchnrk>

Desmos activity available at 151: Parabola Basic

2 Limits and Gas Mielage

Example 7. The function

$$G = f(s) = \frac{2}{5} + \frac{1}{5000}(40(s+2)^2 - (s+2)^3), \quad 0 \leq s \leq 23,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/xzknfkw3>

Learning outcomes:
Author(s):

Desmos activity available at 151: Gas as a Function of Distance

- (a) Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
- (b) Find your average gas mileage (in miles/gallon) over the interval $s \in [8, 18]$.
- (c) Use the graph to approximate your gas mileage at the moment you are 18 miles from home. Do this by zooming in on the appropriate point.
- (d) Use the algebra of limits to determine your exact gas mileage at the moment you are 18 miles from home.
- (e) Use the result of part (d) to approximate the change in the volume of gas

$$\Delta G = f(s) - f(18)$$

in terms of the change

$$\Delta s = s - 18$$

in your distance from home for values of s near 18 miles. What are the units of the proportionality constant?

- (f) Use the result of part (e) to approximate your distance from home when there are 1.9 gallons of gas in your tank.

3 Limits, Gas Mileage and Speed

Example 8. Suppose that between speeds of 30 miles/hour and 70 miles/hour the gas mileage of a car is a quadratic function of its speed. Suppose also that the car gets a maximum of 42 miles/gal at a speed of 50 miles/hour and that the car gets 38 miles/gallon at a speed of 40 miles/hour.

- (a) Find an expression for the function

$$G = f(v), \quad 30 \leq v \leq 70,$$

that gives the gas mileage (in miles/gal) in terms of the speed (in miles/hour).

- (b) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the car's speed at 60 miles/hour gives an approximately proportional change in its gas mileage.
- (c) Use the results of part (b) to approximate the proportionality constant. What are its units?
- (d) Use the algebra of limits to find the exact value of the proportionality constant.
- (e) Explain the meaning of the proportionality constant.
- (f) Approximate the change

$$\Delta G = g - f(60)$$

in gas mileage in terms of a small change

$$\Delta v = v - 60$$

in the car's speed.

- (g) Use part (f) to approximate the speed at which the car gets 36 miles/gallon.
- (h) Would you expect your approximation in part (g) to be greater or less than the actual speed? Explain your reasoning with a graph.
- (i) Simplify the units of the proportionality constant. What might these units suggest about a way to interpret the constant?
- (j) At what rate (in gal/hr) does the car burn gas at a speed of 60 miles/hour?
- (k) How is the rate in part (j) related to the proportionality constant?

4 Limits, Speed and Altitude

Question 36 A rock dropped from a height of 100 feet falls to the surface of Planet Krypton without air resistance.

(a) By considering only the physical situation and without doing any computations, sketch a graph of the function

$$v = g(h), 0 \leq h \leq 100$$

that expresses the rock's speed (in ft/sec) in terms of its height (in feet).

(b) Use the results from part (a) to choose a reasonable expression for the function g from the list below.

Multiple Choice:

- (a) $g(t) = 100 - 9t^2, 0 \leq t \leq 10/3$
- (b) $g(h) = 100 - 9h^2, 0 \leq h \leq 100$
- (c) $g(h) = 0.005(100 - h)^2, 0 \leq h \leq 100$
- (d) $g(h) = 6\sqrt{100 - h}, 0 \leq h \leq 100$ ✓

(c) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the rock's height from 64 feet gives an approximately proportional change in its speed.

(d) Use the results of part (c) to approximate the proportionality constant. What are its units?

(e) Use the algebra of limits to find the exact value of the proportionality constant.

(f) Explain the meaning of the proportionality constant.

(g) Approximate the change

$$\Delta v = v - g(64)$$

in the rock's speed in terms of a small change

$$\Delta h = h - 64$$

in its height.

(h) Use part (g) to approximate the rock's speed at a height of 63 feet.

(i) Would you expect your approximation in part (h) to be greater or less than the actual speed? Explain your reasoning with a graph.

(j) Simplify the units of the proportionality constant. Does this simplification help to understand or obscure the meaning of the proportionality constant?

5 Limits and Gas Mileage

6 Limits and Purchasing Power

The Derivative

Computing derivatives with limits.

Example 9. Suppose

$$y = f(x) = x^2$$

and let's use limits to evaluate

$$f'(3) = \left. \frac{dy}{dx} \right|_{x=3} = \left. \frac{d(x^2)}{dx} \right|_{x=3}.$$

We have

$$\begin{aligned} f'(3) &= \left. \frac{d(x^2)}{dx} \right|_{x=3} = \lim_{v \rightarrow 3} \frac{f(v) - f(3)}{v - 3} \\ &= \lim_{v \rightarrow 3} \frac{v^2 - 9}{v - 3} \\ &= \lim_{v \rightarrow 3} \frac{(v - 3)(v + 3)}{v - 3} \\ &= \lim_{v \rightarrow 3} (v + 3) \\ &= (3 + 3) \\ &= 6. \end{aligned}$$

Next let's do almost the same thing and compute

$$f'(x) = \frac{dy}{dx}$$

for the function

$$y = f(x) = x^2$$

by replacing 3 in the above computation with x .

Learning outcomes:
Author(s):

We get

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d(x^2)}{dx} = \lim_{v \rightarrow x} \frac{f(v) - f(x)}{v - x} \\
 &= \lim_{v \rightarrow x} \frac{v^2 - x^2}{v - x} \\
 &= \lim_{v \rightarrow x} \frac{(v - x)(v + x)}{v - x} \\
 &= \lim_{v \rightarrow x} (v + x) \\
 &= (x + x) \\
 &= 2x.
 \end{aligned}$$

Just as a check, when $x = 3$,

$$\left. \frac{dy}{dx} \right|_{x=3} = (2x) \Big|_{x=3} = 2(3) = 6.$$

Question 37 Use the result of the previous example to solve each of the following problems. Do not use limits.

- (a) Find an equation of the tangent line to the parabola $y = x^2$ at the point $(-4, 16)$.
- (b) Find an equation of the tangent line to the parabola perpendicular to the tangent line in part (a).
- (c) Find the coordinates of the point where the lines in parts (a) and (b) intersect.
- (d) Let \mathcal{L} be the line through the points of tangency of the lines in parts (a) and (b). Find the coordinates of the point where \mathcal{L} intersects the y -axis.
- (e) Repeat parts (a)-(d) above for the tangent line to the parabola $y = x^2$ at the general point (a, a^2) . What do you notice?

Question 38 (a) Use the method of Example 1 for the function

$$y = g(x) = 1/x^2$$

to compute

$$g'(3) = \left. \frac{dy}{dx} \right|_{x=3} = \left. \frac{d(1/x^2)}{dx} \right|_{x=3}$$

and

$$g'(x) = \frac{dy}{dx}.$$

(b) Use the result of part (a) to find an equation of the tangent line to the curve $y = 1/x^2$ at the point $(3, 1/9)$.

Question 39 (a) Use numerical methods to estimate the slope of the tangent line to the curve

$$y = f(x) = x^3$$

at the point $(2, 8)$. Include enough data to suggest a progression toward a limit.

(b) Use the algebra of limits to find the exact slope of the tangent line in part (a).

(c) Use algebra to find the coordinates of the all pointw where the tangent line in part (a) intersects the curve $y = x^3$.

(d) Suppose you measure the edge length of a cube to be 2cm and then use this measurement to compute the volume of the cube. Use the result of part (b) to approximate your error in computing the volume in terms of your error in measuring the edge length. Assume the latter error is small.

Then compare your exact error in computing the volume with your approximation for some specific edge length near 2cm. You should start this problem defining a function with meaningful variable names (do not use x and y).

Question 40 (a) Find a function

$$s = g(V), V \geq 0$$

that expresses the edge length (measured in cm) of a cube in terms of its volume (measured in cubic centimeters).

(b) Use the algebra of limits to evaluate the derivative

$$g'(V_0) = \left. \frac{ds}{dV} \right|_{V=V_0}.$$

(c) What are the units of the derivative in part (b)? Explain how you know.

(d) Suppose you submerge the cube in water and measure its volume to be 8 cm^3 . You then use this measurement to compute the edge length of the cube. Use the result of part (b) to approximate your error in computing the edge length in terms of your error in measuring the volume. Assume the latter error is small.

Start this problem by defining the errors Δs and ΔV in terms of the actual edge length s of the cube.

(e) Use the result of part (d) to approximate your error in computing the edge length of the cube if the cube's actual volume is 7.7 cm^3 .

(f) Explain how this question is related to Question 4.

Question 41 Suppose that between speeds of 60 miles/hr and 72 miles/hr, the gas mileage of a car is a linear function of its speed. Suppose also that the car gets 36 miles/gallon at a speed of 60 miles/hour and 32 miles/gallon at a speed of 72 miles/hour.

(a) Find a function

$$r = f(v), 60 \leq v \leq 72,$$

that expresses the rate (measured in gal/hr) at which the car burns gas in terms of its speed (measured in miles/hour). Explain your reasoning. This function is not linear.

Hint:

$$r = f(v) = \frac{3v}{168 - v}, 60 \leq v \leq 72.$$

(b) Use numerical methods to estimate the value of the derivative

$$f'(63) = \left. \frac{dr}{dv} \right|_{v=63}.$$

Make a table that shows enough data to suggest a progression toward a limit. Include units in all column headings.

(c) Use the algebra of limits to find an expression for the derivative

$$f'(v) = \frac{dr}{dv}.$$

Then use this expression to find the exact value of the derivative in part (b).

(d) What are the units of the derivative in part (b)? Explain its meaning.

(e) Use the result of part (c) to approximate the change

$$\Delta r = f(v) - f(63)$$

in the rate at which your car burns gas in terms of the change

$$\Delta v = v - 63$$

of the car's speed. Assume $\Delta v \sim 0$.

Question 42 Suppose that between speeds of 30 miles/hour and 70 miles/hour the gas mileage of a car is a quadratic function of its speed. Suppose also that

the car gets a maximum of 42 miles/gal at a speed of 50 miles/hour and 34 miles/gallon at a speed of 30 miles/hour.

(a) Find a function

$$r = h(v), \quad 30 \leq v \leq 70,$$

that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hour).

Hint: (i) At what rate does the car burn gas at a speed of 50 miles/hour? 25/21 gal/mile

(ii) Find a function that expresses the gas mileage G (measured in miles/gallon) in terms of the speed (measured in miles/hr).

$$G = 42 - 0.02(v - 50)^2, \quad 30 \leq v \leq 70.$$

(ii) The rate (in gal/hr) at which the car burns gas as a function of its speed (in miles/hr) is

$$r = h(v) = \frac{v}{42 - 0.02(v - 50)^2}, \quad 30 \leq v \leq 70.$$

(b) Use the algebra of limits to evaluate the derivative

$$h'(40) = \left. \frac{dr}{dv} \right|_{v=40}.$$

(c) What are the units of the above derivative? How do you know?

(d) Express the meaning of the derivative in the context of small changes.

Question 43 At 10:00am on April 18, the wholesale price of Cosmic Crisp apples is \$2.00/lb and is decreasing at the rate of \$0.10/lb/hour.

Use the algebra of limits to determine the rate (in pounds/hour) at which the number of pounds of apples a store can purchase with \$1000 is changing at this time.

Start this question by defining a function that expresses the number of pounds of apples the store can buy with \$1000 in terms of the price (in \$/lb). Choose meaningful variable names (not x and y). Do not assume the price is decreasing at a constant rate.

1 Limits, Speed and Altitude

Question 44 A rock dropped from a height of 100 feet falls to the surface of Planet Krypton without air resistance.

(a) By considering only the physical situation and without doing any computations, sketch a graph of the function

$$v = g(h), 0 \leq h \leq 100$$

that expresses the rock's speed (in ft/sec) in terms of its height (in feet). Explain your reasoning.

(b) Use the results from part (a) to choose a reasonable expression for the function g from the list below.

Multiple Choice:

- (a) $g(t) = 100 - 9t^2, 0 \leq t \leq 10/3$
- (b) $g(h) = 100 - 9h^2, 0 \leq h \leq 100$
- (c) $g(h) = 0.005(100 - h)^2, 0 \leq h \leq 100$
- (d) $g(h) = 6\sqrt{100 - h}, 0 \leq h \leq 100$ ✓

(c) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the rocks height from 64 feet gives an approximately proportional change in its speed. Then approximate the proportionality constant. What are its units?

(d) Use the algebra of limits to find an expression for the derivative

$$g'(h) = \frac{dv}{dh}.$$

Then use this expression to find the exact value of the proportionality constant in part (c).

(e) Explain the meaning of the proportionality constant.

(f) Approximate the change

$$\Delta v = v - g(64)$$

in the rock's speed in terms of a small change

$$\Delta h = h - 64$$

in its height.

- (g) Use part (g) to approximate the rock's speed at a height of 63 feet.
 - (h) Would you expect your approximation in part (h) to be greater or less than the actual speed? Explain your reasoning with a graph.
 - (i) Simplify the units of the proportionality constant. Does this simplification help to understand or obscure the meaning of the proportionality constant?
-

Derivatives of Polynomials

Working with polynomials and their derivatives.

1 Differentiating Polynomials

Question 45 Let $f(x) = x^5$.

- (a) Use the algebra of limits to find an expression for the derivative

$$\frac{d}{dx}(f(x)) = \frac{d}{dx}(x^5).$$

- (b) Use the result of part (a) to find an expression for the derivative

$$\frac{d}{dx}(f^{-1}(x)) = \frac{d}{dx}(x^{1/5}).$$

Use set-builder notation to state the domains of $f^{-1}(x)$ and its derivative.

- (c) Use the result of part (a) and the algebra of limits to find an expression for the derivative

$$\frac{d}{dx}\left(\frac{1}{f(x)}\right) = \frac{d}{dx}(x^{-5}).$$

- (d) Use the results of parts (a)-(c) to find equations of the tangent lines to the three curves $y = f(x)$, $y = f^{-1}(x)$, and $y = 1/f(x)$ at the point $(1, 1)$. Graph the curves and their tangent line in Desmos to check your work.

Question 46 (a) Use the results of Question 1 to make a conjecture about the derivative

$$\frac{d}{dx}(f(x)) = \frac{d}{dx}(x^n)$$

of the function $f(x) = x^n$.

- (b) What do you get for the derivative when $n = 0$? When $n = 1$? Are these results correct? Explain.

Learning outcomes:
Author(s):

Question 47 Due to a printing error, the graph of the function $y = f(x) = 234x^5$ is missing below. All we see is a point A on the graph of the function.

Desmos link: <https://www.desmos.com/calculator/pjyqjtdcxm>

151: Printers Error 2

Without sketching a graph, drag the slider m to draw the tangent line to the curve $y = 234x^5$. Explain your reasoning.

Question 48 (a) Suppose one giraffe is always twice as tall as another. What can you say about their growth rates at any instant?

(b) Suppose one giraffe is always two feet taller than another. What can you say about their growth rates at any instant?

(c) What do parts (a) and (b) suggest about how to compute the derivatives

$$\frac{d}{dx}(f(x) + b),$$

$$\frac{d}{dx}(af(x)),$$

and

$$\frac{d}{dx}(af(x) + b)$$

for constants $a, b \in \mathbb{R}$?

(d) Make up your own scenario that suggests how to compute the derivative

$$\frac{d}{dx}(f(x) + g(x))$$

of the sum of two functions.

Question 49 Find an equation of the tangent line to the curve

$$y = -x^3 + 4x^2 - 3x + 1$$

at the point $(2, 3)$. Graph the curve and its tangent line on Desmos.

2 Stock Price

Question 50 The function

$$P = f(t) = 0.1t^3 - t^2 + t + 20, \quad 0 \leq t \leq 8,$$

expresses the price (in dollars/share) of a stock in terms of the number of hours past 9am. The graph of the function is shown below.

Desmos link: <https://www.desmos.com/calculator/gpqicgxvk3>

151: Stock Price

- (a) Is the price increasing or decreasing at 10am? Use the graph to estimate this rate as follows:
 - (i) First drag the slider u (another name for t) on Line 5 to the appropriate value.
 - (ii) Then drag the slider m (the slope of the orange line) so that the orange line looks parallel to the (red) tangent line to the curve $P = f(t)$.
- (b) Use the function to compute the exact rate at which the price is changing at 10am. Include units.
- (c) At what relative rate is the price changing at 10am? Include units.
- (d) At what rate is the number of shares you can buy with \$1000 changing at 10am? Do not use the quotient rule or the chain rule (we have not learned these yet). Use limits instead, but work in general (ie. not with this specific price function). See the Explanation below for help.
- (e) Use the sliders m and b to control the line $P = b + mt$ and approximate when the price is decreasing at the rate of (\$1.8/share)/hr. Then use calculus and algebra to compute the exact time(s).
- (f) Use calculus and technology to approximate the time(s) when the price is decreasing at the relative rate of 10%/hour.
- (g) Use the graph to approximate the minimum and maximum prices of the stock during the eight-hour period. Then use calculus and algebra to compute these exact prices. Click the arrow below for a hint.

What is the value of the derivative dP/dt when the price is a maximum/minimum?

- (h) Use the slider u (another name for the variable t) to approximate the time(s) when the price is decreasing at the fastest rate. Then use calculus to compute the exact time(s). Click the arrow below for a hint.

It helps to graph the function

$$dP/dt = 0.03t^2 - 2t + 1, 0 \leq t \leq 8,$$

Then think about how you would use calculus to compute the exact time when the price is decreasing at the fastest rate. This is similar to the previous question.

- (i) Use the graph above to approximate the end of the time interval beginning at 9am over which the price decreases at the greatest average rate. Do this by activating the folder secant line on Line 6 (click the circle on the left) and dragging the slider u .
- (j) Use calculus and algebra to find the end of the time interval beginning at 9am over which the price decreases at the greatest average rate. Click the arrow below for a hint.

Think about the function

$$m(t) = \frac{f(t) - f(0)}{t - 0}, 0 < t \leq 8,$$

that expresses the average rate of change of the price (with respect to time) over the time interval between 9am and time t hours past 9am. Then use this function and the ideas of parts (g) and (h).

Explanation. (d) Here's the solution to part (d).

- Start by finding a function

$$S = g(t), 0 \leq t \leq 8,$$

that expresses the number of shares we can buy with \$1,000 in terms of the number of hours past 9am. Click the arrow to the lower right to check.

This function is

$$S = g(t) = \frac{1000}{f(t)}, 0 \leq t \leq 8$$

- Next we'll use limits to find an expression for the derivative

$$\left. \frac{dS}{dt} \right|_{t=a}$$

as follows.

$$\begin{aligned}
 \left. \frac{dS}{dt} \right|_{t=a} &= \lim_{v \rightarrow a} \frac{g(v) - g(a)}{v - a} \\
 &= \lim_{v \rightarrow a} \left(\frac{1000}{v - a} \right) \left(\frac{1}{f(v)} - \frac{1}{f(a)} \right) \\
 &= \lim_{v \rightarrow a} \left(\frac{1000}{v - a} \right) \left(\frac{f(a) - f(v)}{f(v)f(a)} \right) \\
 &= \lim_{v \rightarrow a} \left(\frac{-1000}{f(v)f(a)} \right) \left(\frac{f(v) - f(a)}{v - a} \right) \\
 &= \lim_{v \rightarrow a} \frac{-1000}{f(v)f(a)} \cdot \lim_{v \rightarrow a} \frac{f(v) - f(a)}{v - a}
 \end{aligned}$$

Let's pause here for a moment. The last equality above follows from the fact that both limits in the product exist. The first limit is easy enough to evaluate. The second limit exists because *it is equal to the derivative*

$$\left. \frac{dP}{dt} \right|_{t=a}.$$

That's the key point. Now we know that

$$\begin{aligned}
 \left. \frac{dS}{dt} \right|_{t=a} &= \lim_{v \rightarrow a} \frac{-1000}{f(v)f(a)} \cdot \lim_{v \rightarrow a} \frac{f(v) - f(a)}{v - a} \\
 &= \left(\frac{-1000}{(f(a))^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right)
 \end{aligned}$$

This is enough to compute the rate at which the number of shares we can buy with \$1000 is changing at 10am.

But we can get a better understanding of this by writing the above expression in a more meaningful way. Remember that $f(a) = P$ is the price at time t . So

$$\begin{aligned}
 \left. \frac{dS}{dt} \right|_{t=a} &= - \left(\frac{1000}{(f(a))^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right) \\
 &= - \left(\frac{1000}{P^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right) \\
 &= - \left(\frac{1000}{P} \right) \left(\frac{1}{P} \cdot \left. \frac{dP}{dt} \right|_{t=a} \right) \\
 &= -S \left(\frac{1}{P} \cdot \left. \frac{dP}{dt} \right|_{t=a} \right),
 \end{aligned}$$

where $S = 1000/P$ is the number of shares we can buy with \$1,000 at a price of P dollars/share.

Now remember that the product

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=a}$$

is the *relative* rate of change in the price (with respect to time) at time $t = a$ hours past 9am.

From part (c) we know at 10am this relative rate is equal to

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=1} = -\frac{0.7}{20.1} \text{hr}^{-1} \sim -3.4826\%/\text{hour}$$

So at 10am we can buy

$$S = \frac{\$1000}{\$20.1/\text{share}} \sim 47.7512 \text{ shares}$$

with \$1,000. And the number of shares we can buy with is increasing at the rate of approximately

$$(47.7512 \text{ shares}) (-3.4826\%/\text{hour}) \sim 1.733 \text{ shares}/\text{hour}.$$

at 10am.

Thinking about Parabolas

Question 51 (a) Find an equation of the tangent line to the parabola $y = x^2$ at the point $(-3, 9)$.

(b) Find an equation of the tangent line to the parabola perpendicular to the tangent line in part (a).

(c) Find the coordinates of the point where the lines in parts (a) and (b) intersect.

(d) Let \mathcal{L} be the line through the points of tangency of the lines in parts (a) and (b). Find the coordinates of the point where \mathcal{L} intersects the y -axis.

(e) Repeat parts (a)-(d) above for the tangent line to the parabola $y = x^2$ at the general point (b, b^2) . What do you notice? Enter your work in the Desmos activity below.

Access Desmos interactives through the online version of this text at

Desmos link: <https://www.desmos.com/calculator/qe7mgnu5sv>

Question 52 Let

$$y = f(x) = ax^2$$

where $a \in \mathbb{R}$ is a constant, and the variables x, y are measured in meters.

- (a) What are the units of the constant a ? How do you know?
- (b) Answer part (e) of the previous question for this function. Modify the desmos activity in the previous question to check your work.

Question 53 In the absense of air resistance, a rock released from rest near the surface of the earth falls $s = 16t^2$ feet during the first t seconds of its fall.

Compare the speed of the rock when it hits the ground with the average speed of the rock during the entire time interval of its fall.

Question 54 (a) The demonstration below shows two normal lines to a parabola and their point of intersection P . What do you think happens to P as point B approaches A ? Answer this question without dragging the slider b .

- (b) Now drag the slider b near $a = 2$ and observe what happens to point P . Were you correct?

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/ybaivhc2t1>

Access this activity online at 151: Normals to Parabola

The parabola has equation $y = x^2/4$, point A has coordinates $(2, 1)$, and point B has coordinates $(b, b^2/4)$.

- (c) Find an equation of the normal lines to the parabola at A and B .
- (d) Use algebra to find an equation of the point P where the normal lines intersect.
- (e) The point P approaches some point Q as B appraoches A . Use the algebra of limits to find the coordinates of Q .
- (f) Find an equation of the circle centered at Q through A . Do this by first using vector algebra to find the coordinates of the center of the circle.
- (g) Repeat parts (c)-(g), replacing the point $A(2, 1)$ with the point $A(a, a^2)$.

Applications

Question 55 Play the slider u in the animation below to watch the motion of a balloon as it leaves behind track's at equal time intervals. Ignore the balloon's rightward jogs. Their purpose is to prevent the tracks from overlapping.

Desmos link: <https://www.desmos.com/calculator/h91txxjcmi>

151: Balloon Motion 7

- (a) Use the animation to sketch by hand the graph of the function

$$h = f(t), \quad -1 \leq t \leq 1.8,$$

that expresses the balloon's height (in thousands of feet) in terms of the number of hours past noon.

- (b) Use the animation or your graph from part (a) to sketch by hand the graph of the function

$$r = g(t), \quad -1 \leq t \leq 1.8,$$

that expresses the balloon's rate of ascent (in thousands of feet/hour in terms of the number of hours past noon.

- (c) Use your graph(s) or the animation to approximate when the balloon descends at the fastest rate.

Question 56 The function

$$h = f(t) = 5 - 2t - \frac{t^2}{4} + t^3, \quad -1 \leq t \leq 1.8,$$

expresses the height (in thousands of feet) of a balloon in terms of the number of hours past noon.

- (a) Find the balloon's average rate of ascent between 11:00am and 11:30am.
- (b) Is the balloon rising or falling at 1:00pm? At what rate? Use the graph of the function $h = f(t)$ below to approximate the rate. Then compute the exact rate.
- (c) When is the balloon descending at the rate of 1000 ft/hour? Use the sliders m and b below to approximate the time(s). Then compute the exact time(s).
- (d) Use the graph below to approximate when the balloon is descending at the fastest rate. Approximate this rate from the graph. Then compute the exact time and rate.

- (e) Use the graph below to approximate when the balloon is at its lowest point. Then compute the exact time.
- (f) Use the graph below to approximate when the balloon is at its highest point between 11am and 1:36pm. Then compute the exact time.
- (g) Use algebra to find all half-hour time intervals during which the balloon descends at an average rate of 500 ft/hour.

Desmos link: <https://www.desmos.com/calculator/bvtukd0v1c>

Access this activity online at 151: Height of Balloon

Question 57 The function

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), \quad 3 \leq s \leq 28,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/cphmgnr7m7>

Desmos activity available at 151: Gas as a Function of Distance 2c

- (a) Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
- (b) Zoom in on the graph to approximate your gas mileage at the moment you are 20 miles from home. Show a screenshot to help explain how you got your approximation. Then compute the exact gas mileage.
- (c) Use the sliders m and b_1 in the graph to approximate your distance from home at the moment your car gets 30 miles/gallon. Show a screenshot to help explain how you got your approximation. Then compute the exact distances.
- (d) Use the sliders m and b_1 in the graph to approximate an interval beginning or ending when you are 20 miles from home over which your average gas mileage is equal to you gas mileage at the moment you are 20 miles from home. Show a screenshot to help explain how you got your approximation. Then compute the exact interval.
- (e) Sensors on your car measure both the (instantaneous) gas mileage and the number of gallons of gas in your tank at each instant. A computer then uses these measurements to estimate the number of additional miles you can drive before running out of gas. Use this idea to find a function

$$m = g(s), \quad 3 \leq s \leq 28,$$

that expresses the number of miles you can drive before running out of gas (assuming your gas mileage remains constant for the remainder of your trip) in terms of your distance from home. Explain your reasoning.

Question 58 (a) Make up your own quadratic function

$$v = f(G) = aG^2 + bG + c,$$

with a, b , and c all not equal to zero, that expresses the speed of a car (measured in miles/hour) in terms of its gas mileage (measured in miles/gallon). Be sure to include a domain. Explain why you think your function is reasonable.

(b) Compute the derivative dv/dG and evaluate it at a specific gas mileage. Include units.

(c) Evaluate the derivative dv/dG at a specific gas mileage and its meaning. Include units in your explanation.

(d) Use your function from part (a) to find a function

$$r = h(G)$$

that expresses the rate (in gal/hr) at which the car burns gas in terms of its gas mileage (in miles/gal). Explain your logic thoroughly.

(e) Evaluate $h(G)$ at the same gas mileage, say G_0 , you used in part (c). Compare the units of $h(G_0)$ and the derivative $dv/dG|_{G=G_0}$. Are these two numbers related? What does this tell you about simplifying the units of a derivative?

(f) Use the ideas of this chapter (ie. derivatives of polynomials, and nothing beyond) to find an expression for the derivative dr/dG .

(g) What are the units of the derivative dr/dG ?

(h) Evaluate the derivative dr/dG at a specific gas mileage and explain its meaning. Include units in your explanation.

(i) Make up and answer your own question about the derivative dr/dG at a specific gas mileage.

A Few More Problems

Question 59 The function

$$h = f(t) = kt^3, t \geq 0,$$

expresses the height of a balloon (in thousands of feet) in terms of the number of hours past noon. Here $k > 0$ is a positive constant.

- (a) What are the units of the constant k ? Explain how you know.
- (b) Find the balloon's rate of ascent (measured in thousands of ft/hr) at time $t = 3$ hours past noon. Then find its average rate of ascent (measured in thousands of ft/hr) between noon and 3pm. How are these rates related? Note that both will be expressed in terms of k .
- (c) Find a function
- $$r = g(a)$$
- that expresses the balloon's rate of ascent (measured in thousands of ft/hr) at time $t = u$ hours past noon in terms of its average rate of ascent (measured in thousands of ft/hr) over the time interval $t \in [0, u]$. Assume $u > 0$.
- (d) Interpret your result from part (b) geometrically on the graph of the function f . Follow the directions on Lines 4, 6, and 8 in the demonstration below to help explain your interpretation.
- (e) Which of the following expresses the balloon's rate of ascent at time t hours past noon in terms of t and $h = f(t)$?

Multiple Choice:

- (a) $\frac{h}{t}$
- (b) $\frac{h}{3t}$
- (c) $\frac{3h}{t}$ ✓

Desmos link: <https://www.desmos.com/calculator/0byjcy77yw>

Desmos activity available at 151: Subtangent 1

Question 60 The function

$$v = f(G)$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon) for speeds between 55 miles/hour and 70 miles/hour.

Suppose $f(30) = 60$.

- (a) Which of the following is more likely to be true?

$$\left. \frac{dv}{dG} \right|_{G=30} = 4$$

or

$$\left. \frac{dv}{dG} \right|_{G=30} = -4?$$

Explain your reasoning.

- (b) What are the units of the above derivative? Do not simplify the units and do not write “per” in place of “/”.
- (c) Explain the meaning of the derivative in part (a). It is not enough to say “the rate of change of something with respect to something else.” Remember this class is all about small changes and your explanation should be about an approximate relationship between small changes in this setting.
- (d) Simplify the units of the derivative. What does this suggest about its meaning?
- (e) At what rate does the car burn gas (in gal/hour) at a speed of 60 miles/hour?
- (f) What does this problem suggest about simplifying the units of a derivative?

Question 61 The function

$$v = f(G), 20 \leq G \leq 40,$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon). Use the graph of the function f to find approximate answers to the following questions. Change the position and slope of line AB by dragging either the line or the points A or B . Change the position of the tangent line by dragging the slider G .

- (a) Label the axes with the appropriate variable names and units.
- (b) At what speed does the car burn gas as the fastest rate?

Hint: One of questions (b), (c) is related to the tangent lines to the curve, the other is related to the lines through the origin and the points of the curve.

- (c) At what speed does increasing the speed by 0.1 miles/hour result in the greatest change in the gas mileage? Approximate that change.

Geogebra link: <https://www.geogebra.org/m/vjdf6x6z>

Geogebra activity available at [151: Gas Mileage](#)

3 Homework Solution

Desmos link: <https://www.desmos.com/calculator/jku58rp2ve>

Gas Consumption

Gas consumption and derivatives.

Question 62 The function

$$\begin{aligned} G &= f(v) \\ &= -\frac{v^2}{20} + 5v - 90, \quad 35 \leq v \leq 70, \end{aligned}$$

graphed below expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

Desmos link: <https://www.desmos.com/calculator/fapdhcqml>

151: Burning Gas

- Use the graph to determine the rate (in gal/hr) at which the car burns gas at a speed of 50 miles/hour.
- Drag the slider v in Line 1 to approximate the speeds between 35 miles/hour and 70 miles/hour at which the car burns gas at the maximum and minimum rates (measured in gal/hr). Explain your reasoning.
- Use calculus to determine the exact speeds in part (b). Find a way that avoids using the quotient rule.

Question 63 The function

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), \quad 3 \leq s \leq 28,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/pb8v4t3cxg>

Desmos activity available at 151: Gas as a Function of Distance 33

Learning outcomes:
Author(s):

- (a) Use the graph above to determine if you are driving toward or away from home. Explain your reasoning.
- (b) Sensors on your car measure both the (instantaneous) gas mileage and the number of gallons of gas in your tank at each instant. A computer then uses these measurements to estimate the number of additional miles you can drive before running out of gas.
- (i) Use the graph and the slider s_0 above to approximate the reading for the number of additional miles you can drive when you are 10 miles from home and 20 miles from home. Explain your reasoning.
- (ii) Find a function

$$m = g(s), 3 \leq s \leq 28,$$

that expresses the number of miles you can drive before running out of gas (assuming your gas mileage remains constant for the remainder of your trip) in terms of your distance from home. Explain your reasoning.

- (iii) Find a function

$$r = h(s), 3 \leq s \leq 28,$$

that expresses the relative rate of change in the function f with respect to and in terms of the number of miles from home.

- i. Evaluate $h(10)$ and interpret its meaning. Include units.
- ii. Compare the functions $r = h(s)$ and $m = g(s)$. How are they related? Explain the logic behind this relation.

The Quotient and Product Rules

Quotient and product rules.

Here are examples of how to use the Leibniz notation correctly in computing the derivative of a product and quotient of functions.

Example 10. Find expressions for the derivatives

(a)

$$\frac{d}{dx} ((5x^3 - 2)(4 - 6x))$$

(b)

$$\frac{d}{dt} \left(\frac{5t^3 - 2}{4 - 6t} \right)$$

Explanation. (a)

$$\begin{aligned} \frac{d}{dx} ((5x^3 - 2)(4 - 6x)) &= (4 - 6x) \frac{d}{dx} (5x^3 - 2) + (5x^3 - 2) \frac{d}{dx} (4 - 6x) \\ &= (4 - 6x)15x^2 + (5x^3 - 2)(-6). \end{aligned}$$

(b)

$$\begin{aligned} \frac{d}{dt} \left(\frac{5t^3 - 2}{4 - 6t} \right) &= \frac{1}{(4 - 6t)^2} \left((4 - 6t) \frac{d}{dt} (5t^3 - 2) - (5t^3 - 2) \frac{d}{dt} (4 - 6t) \right) \\ &= \frac{(4 - 6t)15t^2 - (5t^3 - 2)(-6)}{(4 - 6t)^2}. \end{aligned}$$

1 Relative Rates of Change

Suppose $P = f(t)$ and $Q = g(t)$ are differentiable functions of t and $g(t) \neq 0$. Then the quotient and product rules are better written in the forms

$$\frac{d}{dt} \left(\frac{P}{Q} \right) = \frac{P}{Q} \left(\frac{1}{P} \cdot \frac{dP}{dt} - \frac{1}{Q} \cdot \frac{dQ}{dt} \right) \quad (1)$$

and

$$\frac{d}{dt} (PQ) = PQ \left(\frac{1}{P} \cdot \frac{dP}{dt} + \frac{1}{Q} \cdot \frac{dQ}{dt} \right).$$

Learning outcomes:
Author(s):

Question 64 (a) Verify the statements above.

(b) What do they say about the relative rate of change in the quotient of two functions? In their product?

2 Exercises

Exercise 65 Between speeds of 45 miles/hr and 75 miles/hr, the function

$$G = f(v), 45 \leq v \leq 75,$$

expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

Suppose $f(50) = 25$ and

$$\left. \frac{dG}{dv} \right|_{v=50} = 0.8.$$

- What are the units of the derivative above? What is the meaning? Explain in terms of small changes.
- What are the simplified units of the derivative above? What insight do these units give you about the derivative's meaning?
- Approximate the gas mileage at a speed of 48 miles/hour.
- Let

$$r = g(v), 45 \leq v \leq 75,$$

be the function that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hr).

- Use Equation (1) above to evaluate the derivative

$$\left. \frac{dr}{dv} \right|_{v=50}.$$

Include units for all numbers.

- What are the units of the derivative above? What is the meaning? Explain in terms of small changes.
 - What are the simplified units of the derivative above? What insight do these units give you about the derivative's meaning?
 - Approximate the rate (in gal/hr) at which the car burns gas at a speed of 48 miles/hr.
-

Exercise 66 At 9:00am on February 23, 2023, the price of oil was decreasing at the relative rate of 2%/hour. At what relative rate was the number of gallons of oil you could buy with \$100,000 changing at that time? Use calculus to justify your assertion.

Exercise 67 The function

$$P = f(t) = 5 - 3t + t^2, 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) Use the graphs of the function $P = f(t)$ and the function $r = f'(t)/f(t)$ to estimate when the stock price is increasing at the greatest relative rate.
- (b) Use algebra to find the exact time when the stock price is increasing at the greatest relative rate.

Hint: What is the value of the derivative dr/dt at this time? But start by finding an expression for the instantaneous relative rate of change in the stock price.

Desmos link: <https://www.desmos.com/calculator/xuupp3srqv>

Desmos activity available at 151: Stock Price 4

Exercise 68 An h -foot tall giraffe walks directly toward a spotlight on the ground as the light casts its shadow on a vertical wall as illustrated below. The wall is b feet from the light.

Desmos link: <https://www.desmos.com/calculator/2eiyjjpu9n>

Desmos activity available at 151: Spotlight

Suppose at a certain instant the giraffe is w feet from the spotlight and is walking at a speed of v ft/sec.

- (a) Is the length of the giraffe's shadow increasing or decreasing at this instant?
- (b) At what rate?
- (c) Check that your answer in part (b) has the correct units.

- (d) Assume v is constant and describe what happens to the rate in part (b) as the giraffe approaches the spotlight. How is your conclusion revealed in the animation?
-

Exercise 69 (a) On January 1, 2024 the national debt of a country was decreasing at the rate of 3%/yr and the population was increasing at the rate of 2%/yr. Was the per-capita (ie. per person) share of the national debt increasing or decreasing at this time? At what relative rate?

- (b) During the year 2024 the national debt of a country decreased 3% and the population increased 2%. Did the per-capita share of the national debt increase or decrease during the year? By what percent?

- (c) Compare the two questions above and their answers.
-

Test 1

Test 1.

- (a) (6 points) The function

$$W = f(r) = \frac{2000}{r^2}, r \geq 4,$$

expresses the weight of an astronaut (measured in pounds) in terms of her distance from the center of the earth (measured in thousands of miles).

- (i) (2 points) Find an expression for the average rate of change in the astronaut's weight with respect to her distance from the earth's center between distances $r = b$ and $r = c$ thousands of miles from the center. Assume $b, c \geq 4$ and that $b \neq c$.
- (ii) (4 points) Use your *expression from part (a) directly and the definition of the derivative* to find an expression (fully simplified) for the derivative

$$\left. \frac{dW}{dr} \right|_{r=b}.$$

In particular, do *not* use the power rule. Show all work in its mathematically correct form. Write vertically, one equal sign per line. No need to explain algebra.

Explanation. (i) The average rate of change is

$$\begin{aligned} \frac{\Delta W}{\Delta h} &= \frac{f(c) - f(b)}{c - b} \\ &= \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \end{aligned}$$

- (ii) (Some algebra omitted, left for you to fill in).

$$\begin{aligned} \left. \frac{dW}{dr} \right|_{r=b} &= \lim_{c \rightarrow b} \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \\ &= \lim_{c \rightarrow b} \frac{-2000(c + b)}{c^2 b^2} \\ &= -\frac{2000(b + b)}{b^2 b^2} \\ &= -\frac{4000}{b^3}. \end{aligned}$$

Learning outcomes:
Author(s):

- (b) (5 points) At 3pm, you can buy 5 pounds of cod with \$100. And at 3pm the number of pounds of cod you can buy with \$100 is decreasing at the rate of 0.4 lbs/hour.

At what rate (with respect to time) is the price of cod changing at 3pm? Explain your reasoning and be sure to end with a concluding sentence.

Explanation. Let $P = f(t)$ be the function that expresses the price of cod (in dollars/pound) in terms of the number of hours since 3pm.

Let $Q = g(t)$ be the function that expresses the number of pounds of cod you can buy with \$100 in terms of the number of hours since 3pm.

Then

$$P = \frac{100}{Q}$$

and the quotient rule gives

$$\begin{aligned} \frac{dP}{dt} &= \frac{1}{Q^2} \left(Q \frac{d}{dt} (100) - 100 \frac{dQ}{dt} \right) \\ &= -\frac{100}{Q^2} \cdot \frac{dQ}{dt} \end{aligned}$$

Since at 3pm the number of pounds we can buy with \$100 is decreasing at the rate of 0.4 lbs/hour, we know that

$$\left. \frac{dQ}{dt} \right|_{t=0} = -0.4 \text{ lb/hr.}$$

And since we can buy $Q = g(0) = 5$ pounds with \$100 at 3pm,

$$\begin{aligned} \left. \frac{dP}{dt} \right|_{t=0} &= - \left(\frac{100}{Q^2} \cdot \frac{dQ}{dt} \right) \Big|_{t=0} \\ &= \left(-\frac{100 \text{ dollars}}{25 \text{ lb}^2} \right) \left(-0.4 \frac{\text{lb}}{\text{hr}} \right) \\ &= (1.6 \text{ dollars/lb})/\text{hr.} \end{aligned}$$

So at 3pm the price of cod is increasing at the rate of (1.6 dollars/lb)/hr.

- (c) (5 points) Between speeds of 60 miles/hour and 80 miles/hour, the function

$$v = f(G), \quad 8 \leq G \leq 20,$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon).

Suppose $f(12) = 72$.

- (i) (1 point) Which would be more likely, that

$$\left. \frac{dv}{dG} \right|_{G=12} = 4 \quad \text{or that} \quad \left. \frac{dv}{dG} \right|_{G=12} = -4?$$

Explain your reasoning.

- (ii) (1 point) What are the units of the correct derivative above?
 (iii) (2 points) Explain the meaning of the correct derivative above using the language of *small changes*.
 (iv) (1 point) Simplify the units of the correct derivative above. What do these simplified units tell you about the derivative's meaning?

Explanation. (i) At the high speed of 72 miles/hour it is likely that a small increase in the speed will decrease the gas mileage. So we would expect the derivative to be negative, so

$$\left. \frac{dv}{dG} \right|_{G=12} = -4.$$

- (ii) Since v has units miles/hour and G has units miles/gal, the derivative dv/dG has units

$$\frac{\text{miles/hr}}{\text{miles/gal}}.$$

- (iii) We know that

$$\left. \frac{dv}{dG} \right|_{G=12} = -\frac{0.4 \text{ miles/hr}}{0.1 \text{ miles/gal}} \sim \frac{\Delta v}{\Delta G}$$

if $\Delta G \sim 0$. So increasing the speed by 0.4 mile/hour (from 72 miles/hr to 72.4 miles/hr) would decrease the gas mileage by about 0.1 miles/gal (from 12 miles/gal to about 11.9 miles/gal).

- (iv) The units of the simplified derivative are

$$\frac{\text{miles/hr}}{\text{miles/gal}} = \frac{\text{gal}}{\text{hr}}.$$

Although this suggests the derivative is related to the rate at which the car burns gas, this is *not* true. Our interpretation of the derivative had nothing to do with this rate and in fact the car burns gas at the rate of 6 gal/hour at a speed of 72 miles/hour. So the simplified units are misleading and tell us *nothing* about the derivative's meaning.

- (d) (5 points) The function

$$G = f(s) = \frac{3s + 25}{s + 5}, \quad 0 \leq s \leq 10,$$

expresses the number of gallons of gas in a car in terms of the trip odometer reading (measured in miles).

Sensors on the car measure both the number of gallons of gas in the tank and the current (instantaneous) gas mileage (measured in miles/gallon).

A computer then calculates the number of miles you have left to drive assuming the car's gas mileage remains constant for the remainder of your trip (and equal to the current gas mileage). This number is displayed on the dashboard.

Determine the exact trip odometer reading when the dashboard reading indicates that you have 13 miles left to drive. Explain your reasoning and be sure to end with a concluding sentence.

Explanation. I will not solve this problem, but will give you a hint instead by solving a closely related problem.

Suppose, for example, that we wish to find the dashboard reading (for the number of miles left to drive) at the odometer reading $s = 5$ miles.

First, we would find the number of gallons in the tank at this odometer reading. At this time, the tank has

$$f(5) = \frac{3(5) + 25}{5 + 5} = 4$$

gallons of gas.

Now at $s = 5$, the derivative dG/ds is (left for you to verify)

$$\left. \frac{dG}{ds} \right|_{s=5} = -0.10 \frac{\text{gal}}{\text{mile}}.$$

This means that five miles into our trip the car is burning gas at the rate of 0.10 gallons/mile. So if this rate were to remain constant for the remainder of the trip, we would be able to drive an additional

$$\frac{4 \text{ gallons}}{0.10 \frac{\text{gal}}{\text{mile}}} = 40 \text{ miles}$$

before running out of gas.

Derivatives of Exponential Functions

Working with exponential functions and their derivatives.

Relative Changes and Relative Rates of Change

Relative changes and relative errors are often more meaningful than absolute changes and errors. For example, I might measure the distance from Shoreline's Central Market to the Richmond beach library to be 5 km with an error of at most 0.2 km, while NASA might measure the between the earth and the moon on the first day of spring to be 384,400 km with an error of at most 100 km. The relative error in my measurement is at most

$$\frac{0.2 \text{ km}}{5 \text{ km}} = 0.04 = 4\%,$$

while the relative error in NASA's measurement is at most

$$\frac{100 \text{ km}}{384,400 \text{ km}} \sim 0.00026 = 0.026\%.$$

Relatively speaking, NASA's measurement was about 150 times more accurate than mine.

Question 70 At 10:00am the prices of Stock A and Stock B are both increasing at the rate of (\$2/share)/hour. At 10:00am Stock A sells for \$50/share and Stock B for \$10/share. Compare the relative rates at which the share prices are changing at 10:00am.

Question 71 The function

$$P = f(t) = 5 - 3t + t^2, \quad 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) At what relative rate is the price of the stock changing at 10am?
- (b) When is the share price increasing at a rate of 60%/hr?
- (c) During what time interval is the price of the stock increasing?

Learning outcomes:
Author(s):

(d) During what time interval is the relative rate of change in the price of the stock increasing?

Desmos link: <https://www.desmos.com/calculator/hhkveu6lxp>

Desmos activity available at 151: Stock Price

Question 72 The function

$$P = f(t), 1 \leq t \leq 3.6,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am. Use the graph below to approximate the answers to the following questions without putting a scale on the P axis.

- (a) At what relative rate is the price changing at 11am? At 12:30pm?
- (b) When is the stock price increasing at its maximum relative rate? At its minimum relative rate? Approximate these rates.

Desmos link: <https://www.desmos.com/calculator/jyebaj5jif>

Desmos activity available at 151: Stock Price 2

Question 73 The function

$$W = f(t) = 200 + 4t + 2t^2, 0 \leq t \leq 12,$$

expresses the weight (in pounds) of a baby elephant in terms of its age (in months).

- (a) Find the average rate at which the elephant gained weight between ages 4 and 10 months.
- (b) Find the relative average rate at which the elephant gained weight between ages 4 and 10 months.
- (c) Find the relative instantaneous rate at which the elephant is gaining weight at age 4 month.
- (d) Find the relative instantaneous rate at which the elephant is gaining weight at age 10 months.
- (e) Use the graph below to interpret your answers to parts (b)-(d) geometrically.

Desmos link: <https://www.desmos.com/calculator/2xj6xy7ggo>

Desmos activity available at 151: Elephant

Exponential Growth

Question 74 (a) What does it mean for a population to grow exponentially?
 (b) Is it possible for a population to increase by 20% every year and not grow exponentially?

Question 75 Suppose between noon and 10pm a colony of bacteria grows exponentially. The population is 200,000 at noon and 242,200 at 1pm.

- (a) Describe how the population grows. Keeping Question 2(b) in mind, is your description sufficient?
- (b) How might we find a complete description of the exponential growth?
- (c) Determine the relative average growth rate between noon and 12:30pm. Between 1pm and 1:30pm. Over any half-hour time period. Use the slider u in the graph below to interpret these rates geometrically.
- (d) Approximate the instantaneous relative growth rates in the population at noon, at 1pm, and at 2pm. Modify the definition of $v = u + 1/2$ in the demonstration below and interpret these rates geometrically.
- (e) Use limits to write an expression that gives the instantaneous growth rate at time t hours past noon. What can you conclude?
- (f) Try to answer part (b) again.

Desmos link: <https://www.desmos.com/calculator/wvpsotdhby>

Desmos activity available at 151: [Exp Growth 1](#)

Question 76 The function

$$P = f(t) = 3(2)^t, -2 \leq t \leq 4,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

- (a) Describe how the population grows. Is your description sufficient?

The population *doubles* every *hour*.

- (b) Find an expression for the relative growth rate between time $t = u$ hours past noon and time $t = u + h$ hours past noon. Measure the rate relative to the population at time $t = u$. Is this question asking about an average or an instantaneous relative growth rate?

Derivatives of Exponential Functions

The relative growth rate is

$$\begin{aligned}\frac{1}{P} \left(\frac{\Delta P}{\Delta t} \right) &= \frac{1}{f(u)} \left(\frac{f(u+h) - f(u)}{h} \right) \\ &= \frac{1}{3(2)^u} \left(\frac{3(2)^{u+h} - 3(2)^u}{h} \right) \\ &= \frac{1}{3(2)^u} \left(\frac{3(2)^u(2^h - 1)}{h} \right) \\ &= \frac{2^h - 1}{h}\end{aligned}$$

- (c) What are the units of the relative average growth rate in part (b)?
- (d) Input your function from part (b) on Line 5 in the worksheet below.
- (e) What do you notice about the distance between points R and S as you drag the slider u below. How is this distance related to the relative average growth rate in part (b)?
- (f) Use limits to write an expression for the relative growth rate at time $t = u$ hours past noon. Simplify this expression as much as possible. What can you conclude about how the relative instantaneous growth rate varies with time?

The relative growth rate is

$$\lim_{h \rightarrow 0} \frac{2^h - 1}{h}.$$

- (g) Interpret your expression from part (f) as the derivative of a specific function evaluated at a specific input. What does this tell you about the relative growth rate of this particular population?

The relative instantaneous growth rate of the population is equal to the derivative of the function

$$f(x) = 2^x$$

evaluated at $x = 0$.

- (h) Use part (f) to numerically approximate the relative (instantaneous) growth rate of the population. Show a table that suggests a progression toward a limit.
- (i) Use a similar method to approximate the relative instantaneous growth rate of the population

$$P = f(t) = 5(3)^t.$$

Desmos link: <https://www.desmos.com/calculator/omjbec2hpu>

Desmos activity available at 151: Exponential Growth 1

Question 77 Parts (h) and (i) of the previous question suggest that there is a number e between 2 and 3 that makes the relative growth rate of the function

$$P = f(t) = P_0 e^t, -3 \leq t \leq 5$$

equal to 100%/hr, where we assume here that t is measured in hours.

- (a) What is the one-hour growth factor for this population?
- (b) Describe what happens to the population every hour.
- (c) At what relative rate is the population increasing at 1:00pm?
- (d) Suppose at 1:00pm the population is 500,000. Approximate the population at 1:03pm and compare your approximation to the actual population at that time.

Exponential Functions with Bases other than e

Question 78 The function

$$P = g(t) = P_0 e^{t/2}, -6 \leq t \leq 10$$

expresses the population (Colony B) of bacteria in terms of the number of hours past noon.

- (a) Describe a transformation that takes the graph of the population function

$$P = f(t) = P_0 e^t, -3 \leq t \leq 5$$

for Colony A (where t is also the number of hours past noon) to the graph of $P = g(t)$.

- (b) Suppose that the population of Colony A is 400,000 at 4:00pm.
 - (i) When is the population of Colony B equal to 400,000?
 - (ii) What are the growth rates of the two populations when they each have respective populations of 400,000 bacteria?
 - (iii) What are the relative growth rates of the two populations when they each have respective populations of 400,000 bacteria?

Question 79 Here's another way to think about differentiating the function

$$P = g(t) = P_0 e^{t/2}, -6 \leq t \leq 10$$

that expresses the population of a colony of bacteria in terms of the number of hours past noon.

We'll let $u = t/2$ be the number of two-hour periods since noon.

- (a) Express the population in terms of u .
- (b) Use what you know about the exponential function base e to express the growth rate of the population in terms of u .
- (c) Use part (b) to find the growth rate of the population at 6pm. Pay careful attention to units.
- (d) Use the idea of part (c) to express the growth rate

$$\frac{dP}{dt} = g'(t)$$

in terms of t .

- (e) Suppose instead that the population grows exponentially and doubles every hour. Find the relative instantaneous growth rate of the population.

Question 80 Between 11am and 8pm, a population of bacteria grows exponentially. The population is 4 million at noon and 5 million at 1pm.

- (a) What is the one-hour growth factor? 1.25
- (b) Describe how the population grows.

The population increases by 25% every hour.

- (c) Use your description from part (b) to find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon. Do not use e in your function. Define meaningful variables and include a domain.

The function

$$P = f(t) = 4(5/4)^t, -1 \leq t \leq 8$$

expresses the population (in millions of bacteria) in terms of the number of hours past noon.

- (d) Use the fact that $k = e^{\ln k}$ for $k > 0$ to express your function from part (c) using an exponential function with base e .

$$P = f(t) = 4e^{t \ln(1.25)}, -1 \leq t \leq 8$$

(e) Use u -substitution and the chain rule (show all steps) along with the fact that

$$\frac{d}{du}(e^u) = e^u$$

to find the relative instantaneous growth rate of the population. Include units in your conclusion.

The relative instantaneous growth rate of the population is

$$\frac{1}{P} \left(\frac{dP}{dt} \right) = \ln(5/4) \sim 22.314\%/hour.$$

(f) At what rate is the population growing when there are 10 million bacteria? Do this the easy way. Note that the question is not asking for a relative rate.

(g) Find the population when the population is increasing at the rate of 3 million bacteria/hour. Do this the easy way.

Question 81 Between 11am and 8pm, a population of bacteria decreases exponentially. The population is 5 million at noon and 4 million at 4pm.

(a) What is the four-hour growth factor? (This is the number you multiply the current population by to get the population four hours later). 0.8

(b) What is the one-hour growth factor? $(4/5)^{1/4}$

(c) Describe how the population decreases.

The population decreases by 20% every 4 hours.

(d) Use your description from part (c) to find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon. Do not use e in your function. Start by define meaningful variables. Include a domain with your function.

The function

$$P = f(t) = 5(4/5)^{t/4}, -1 \leq t \leq 8$$

expresses the population (in millions of bacteria) in terms of the number of hours past noon.

(e) Express your function from part (c) using an exponential function with base e . Then use u -substitution and the chain rule to find the relative instantaneous growth rate of the population. Show all steps in using the chain rule.

The population function is

$$P = f(t) = 5e^{0.25 \ln(0.8)t}, -1 \leq t \leq 8.$$

The relative instantaneous growth rate is

$$\frac{1}{P} \left(\frac{dP}{dt} \right) = 0.25 \ln(0.8) \sim -5.58\%/hour.$$

(f) At what rate is the population decreasing when there are 2 million bacteria? Do this the easy way.

(g) Find the population when the population is decreasing at the rate of 300,000 bacteria/hour. Do this the easy way.

(h) Use the result of part (f) to approximate the population 4 minutes after there are 2 million bacteria.

Relative Rates Again

Question 82 The function

$$P = f(t), 0 \leq t \leq 2,$$

expresses the balance (in dollars) in an account in terms of the number of years since the start of 2022. Suppose

$$\left. \frac{1}{P} \frac{dP}{dt} \right|_{P=5000} = 0.08.$$

- (a) What are the units of the above derivative? How do you know?
- (b) Interpret the meaning of the above derivative.
- (c) Approximate the balance in the account four days after the account has \$5,000. Explain your reasoning.

Question 83 The function

$$q = f(p) = 0.5(p - 18)^2, 6 \leq p \leq 15,$$

expresses the average number of burgers/day sold at Five Guys of Edmonds in terms of the price (in \$/burger).

- (a) At what relative rate does the quantity sold (q) change with respect to the price (p) at a price of \$10/burger?
- (b) What are the units of the above relative rate of change?
- (c) Explain the meaning of the relative rate of change in part (a).

(d) Use the graph of the function $q = f(p)$ and the slider u in the desmos activity below to interpret the relative rate of change in part (a) geometrically. Explain your reasoning.

(e) Use the result of part (a) to approximate the relative change in the average number of burgers sold per day if the Five Guys increases the price from \$10/burger to \$10.25/burger. Explain your reasoning.

(f) Use the result of part (a) to approximate the relative change $\Delta a\%$ in the average number of burgers sold per day in terms of a small relative change $\Delta r\%$ in the price from \$10/burger. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/ylgk03oaza>

Desmos activity available at 151: Burgers 1

Question 84 The function

$$P = f(t), -2 \leq t \leq 5,$$

expresses the population of a colony (call it Colony A) of bacteria in terms of the number of hours past noon.

The function

$$P = g(t) = f(t/2), -4 \leq t \leq 10,$$

expresses the population of Colony B in terms of the number of hours past noon.

The populations do not necessarily grow exponentially.

(a) Compare the populations at noon.

(b) Suppose Colony A has 50,000 bacteria at 3:00pm. When does Colony B have 50,000 bacteria? Explain.

(c) Suppose the population of Colony A takes three hours to grow from 20,000 to 50,000. How long does it take the population of Colony B to grow from 20,000 to 50,000?

(d) Suppose the population of Colony A is increasing at the rate of 10,000 bac/hr at 3pm. What is the growth rate of Colony B at 6pm? Explain.

(e) What is the relative growth rate of Colony A at 3pm? What is the relative growth rate of Colony B at 6pm? Explain.

Question 85 The function

$$P = f(u) = 10e^u, -2 \leq u \leq 5$$

Derivatives of Exponential Functions

expresses the population (in millions of bacteria) of a colony of bacteria in terms of the number of hours since noon.

- (a) What are the units of the input to the exponential function in the above expression for P ?
- (b) At what rate is the population growing when there are 30 million bacteria?
- (c) At what rate is the population growing when there are P million bacteria? Do the units of your answer make sense?
- (d) What is the relative instantaneous growth rate of the population?
- (e) Find a function

$$P = g(t)$$

that expresses the population (in millions of bacteria) of the colony of bacteria in terms of the number of minutes since noon. Include a domain.

- (f) Use common sense to evaluate the derivative

$$\left. \frac{dP}{dt} \right|_{P=30}.$$

Explain your reasoning.

- (g) Use the idea of part (f) to find an expression for the derivative

$$\frac{dP}{dt} = g'(t)$$

at time t minutes past noon.

Question 86 The function

$$P = P_0 e^{kt}, -4 \leq t \leq 5,$$

expresses the population of a colony of bacteria in terms of the number of hours past noon.

- (a) What are the units of the constant P_0 ?
- (b) What are the units of the constant k ?
- (c) Use the ideas of the previous question to find an expression for the growth rate of the population at time t hours past noon. Include units in your answer.
- (d) Find an expression for the relative growth rate at time t hours past noon. Include units in your answer.

Question 87 One of the two functions graphed below is an exponential function. Which one? How do you know?

The Derivative as a Magnification Factor

Exploration 88 It sometimes helps to think of the derivative as a magnification factor that maps a small interval around an input to a function to a corresponding interval around the output.

(a) Use this idea for the function $y = f(x)$ graphed below to approximate the derivatives

$$\left. \frac{d}{dx}(f(x)) \right|_{x=4} \quad \text{and} \quad \left. \frac{dy}{dx} \right|_{x=5}.$$

Desmos link: <https://www.desmos.com/calculator/la4f5ots3r>

Desmos activity available at 151: Magnification Factor 1

Differentiating the Exponential Function e^x

Exploration 89 (a) Use the graph of the function $y = f(x)$ below to approximate the derivatives

$$\left. \frac{dy}{dx} \right|_{y=k} \quad \text{for } k = 1, 2, \dots, 6.$$

Note the above derivatives are evaluated at the outputs of the function f .

(b) What do you notice?

Desmos link: <https://www.desmos.com/calculator/k08dphtuca>

Desmos activity available at 151: Magnification Factor 2

Question 90 The function

$$P = 400e^t, 0 \leq t \leq 2$$

expresses the population (in thousands) of a colony of bacteria in terms of the number of hours past noon.

(a) What are the units of the factor 400?

(b) What are the units of the exponent in the factor e^t ? Be careful.

(c) Find the (instantaneous) growth rate of the population when there are 1,200,000 bacteria.

Derivatives of Exponential Functions

- (d) Find the relative (instantaneous) growth rate when there are 1,200,000 bacteria.
 - (e) Find the relative (instantaneous) growth rate at any time.
 - (f) Approximate the population 30 seconds after there are 1,200,000 bacteria.
-

Question 91 Let $k > 0$ be a constant and let

$$f(x) = ke^x.$$

For $a \in \mathbb{R}$ let point P with coordinates $(a, f(a))$ be a curve on the curve $y = f(x)$. Let Q be the point where the tangent line to the curve intersects the x -axis and let R be the point with coordinates $(a, 0)$

- (a) Find the length of segment \overline{QR} .
 - (b) How is part (a) related to part (e) of the previous question?
-

Circular Interpolation

Circular Interpolation

What is usually called *sinusoidal modeling* is kind of a misnomer. First, it's usually easier to model oscillatory behavior with a cosine function. Second, and more importantly, the name *sinusoidal modeling* fails to convey the main idea of replacing linear interpolation with circular interpolation.

Here's an example.

Example 11. Suppose that over the course of a 24-hour period, from midnight October 29 to midnight October 30, the depth of the water at the Edmonds Pier is a sinusoidal function of time. Suppose also that a high tide of 21 feet occurs at 2:00am and the following low tide of 5 feet occurs at 8:00am.

Explanation. Our aim is to find a function

$$h = f(t), 0 \leq t \leq 24,$$

that expresses the depth of the water (in feet) in terms of the number of hours past midnight, October 29.

Note that to say “sinusoidal function” means that the graph of f is generated by uniform circular motion. But the graphs of the sine and cosine functions are both generated this way, so it is ok to express $f(t)$ in terms of the cosine function, and we will do just that.

Here are the steps.

- (a) First we'll use the information above to sketch by hand a graph of the function f . Label the axes with the appropriate variable names and units. Label the coordinates of two key points on the graph.

Desmos activity available at:

142: Edmonds Pier.

Desmos link: <https://www.desmos.com/calculator/x2kocpkcfm>

- (b) Next we'll compute the mean depth of the water over the 24 hours and the maximum deviation of the depth from this mean. Including units in our computation, the mean depth is

$$h_{avg} = 0.5(21 \text{ ft} + 5 \text{ ft}) = 13 \text{ ft}.$$

Learning outcomes:
Author(s):

And the maximum deviation from the mean is

$$21 \text{ ft} - 13 \text{ ft} = 8 \text{ ft}.$$

Activate the Amplitude folder on Line 5 of the above demonstration to draw the horizontal line showing the average depth. Note this line is labeled with its equation. There is also a vertical line that shows the the maximum deviation from the mean.

- (c) Next we'll use the graph to find the period of oscillation. Since high tide occurs at 2:00am and low tide at 8:00am, the period (the time between successive) high (or low) tides is

$$2(8 \text{ hours} - 2 \text{ hours}) = 12 \text{ hours}$$

Activate the Period folder on Line 11 to show the period on the graph.

- (d) Now we'll compute the rotation rate of a uniform circular motion that generates the sinusoidal variation in the depth of the water. This rotation rate is

$$\omega = \frac{2\pi \text{ radians}}{12 \text{ hours}} = \frac{\pi}{6} \text{ radians/hr.}$$

Activate the Uniform Circular Motion folder on Line 18 to show the period on the graph.

- (e) Next we'll use parts (a)-(d) to find an expression for the function

$$h = f(t), 0 \leq t \leq 24,$$

that gives the depth of the water (in feet) at time t hours past midnight. Use the cosine function. Keeping in mind that high tide occurs at 2:00am, our function is

$$h = f(t) = 13 + 8 \cos\left(\frac{\pi}{6}(t - 2)\right), 0 \leq t \leq 24.$$

- (f) To check that our function is correct, we'll use the given information that the depth of the water is 21 feet at 2:00am and 5 feet at 8:00am.

Substituting $t = 2$ gives the depth at 2:00am (in feet) as

$$\begin{aligned} f(2) &= 13 + 8 \cos\left(\frac{\pi}{6}(2 - 2)\right) \\ &= 13 + 8 \cos(0) \\ &= 13 + 8 \\ &= 21. \end{aligned}$$

Substituting $t = 2$ gives the depth at 2:00am (in feet) as

$$\begin{aligned} f(8) &= 13 + 8 \cos\left(\frac{\pi}{6}(8 - 2)\right) \\ &= 13 + 8 \cos(\pi) \\ &= 13 + 8(-1) \\ &= 5. \end{aligned}$$

These check out.

- (g) Finally, we'll use our function to estimate the depth (in feet) of the water at 5:30am, October 29 to be

$$\begin{aligned} f(5.5) &= 13 + 8 \cos\left(\frac{\pi}{6}(5.5 - 2)\right) \\ &= 13 + 8 \cos(7\pi/12) \\ &\sim 10.92 \end{aligned}$$

- (h) Enter the coordinates of the appropriate point in Line 27 of the Desmos Activity above to check that the depth at 5:30am is reasonable.

Example 12. This is a continuation of the previous example, and we'll start with the function

$$h = f(t) = 13 + 8 \cos\left(\frac{\pi}{6}(t - 2)\right), 0 \leq t \leq 24,$$

that expresses the depth of the water (in feet) at the Edmonds Pier in terms of the number of hours past midnight.

Our question now is to find the rate at which the depth of the water is changing the *second* time the water is 10 feet deep.

One approach would be to first determine when the water is 10 feet deep for the second time. But this is not necessary. We can determine the rate without finding the time. Here's how.

- (a) We'll first take time out of the picture by letting

$$\theta = \frac{\pi}{6}(t - 2)$$

Then we'll let t_0 be the second time when the water is 10 feet deep and define the angle θ_0 as

$$\theta_0 = \frac{\pi}{6}(t_0 - 2)$$

- (b) Our aim is to evaluate the derivative

$$\begin{aligned} \left. \frac{dh}{dt} \right|_{t=t_0} &= -\frac{4\pi}{3} \sin\left(\frac{\pi}{6}(t_0 - 2)\right) \\ &= -\frac{4\pi}{3} \sin(\theta_0), \end{aligned}$$

where I've left the computation of the derivative dh/dt to you.

Note the key point. To evaluate this derivative we need *not* know the value of θ_0 , but only the value of $\sin \theta_0$.

- (c) To determine the value of $\sin \theta_0$, we go back to the height function, which we now write in the form (since $\theta = \pi(t - 2)/6$)

$$h = g(\theta) = 13 + 8 \cos \theta.$$

When the water is 10 feet deep for the second time,

$$10 = 13 + 8 \cos \theta_0$$

and

$$\cos \theta_0 = -3/8.$$

- (d) Now since

$$\cos^2 \theta_0 + \sin^2 \theta_0 = 1,$$

we find that

$$\sin \theta_0 = \pm \frac{\sqrt{55}}{8}.$$

- (e) To choose the correct sign, we go back to the graph.

Desmos activity available at:

142: Edmonds Pier 2.

Desmos link: <https://www.desmos.com/calculator/fs9xy11f0e>

Here we can see that at the point Q (with coordinates $(t_0, f(t_0))$),

$$8 < t_0 < 14.$$

This tells us that

$$\theta_0 = \frac{\pi}{6} (t_0 - 2)$$

is between

$$\frac{\pi}{6} (8 - 2) < \theta_0 < \frac{\pi}{6} (14 - 2)$$

or that

$$\pi < \theta_0 < 2\pi.$$

In other words, the angle θ_0 is in the third or fourth quadrant. This means that $\sin \theta_0 < 0$ so

$$\sin \theta_0 = -\frac{\sqrt{55}}{8}.$$

(f) Finally, we evaluate the derivative

$$\begin{aligned}\left.\frac{dh}{dt}\right|_{t=t_0} &= -\frac{4\pi}{3} \sin(\theta_0) \\ &= \left(-\frac{4\pi}{3}\right) \left(-\frac{\sqrt{55}}{8}\right) \\ &= \frac{\pi\sqrt{55}}{6}\end{aligned}$$

So when the water is 10 feet deep for the second time, the depth is increasing at the rate of

$$\frac{\pi\sqrt{55}}{6} \text{ ft/hr} \sim 3.88 \text{ ft/hr.}$$

Derivatives of Trigonometric Functions

Working with trigonometric functions and their derivatives.

1 Exercises

Question 92 *The function*

$$h = f(t) = 1200 + 500 \cos t + 300 \sin t, \quad 0 \leq t \leq 10,$$

expresses the altitude (in feet) of a balloon in terms of the number of hours past noon.

Desmos link: <https://www.desmos.com/calculator/f1ldi6yrek>

151: Trig 1

- (a) *Use calculus to find the minimum and maximum heights of the balloon between noon and 6pm. Do not use a calculator.*
- (b) *Find the balloon's maximum rate of ascent. No calculator.*

Question 93 *The function*

$$h = f(t) = 800 - 200 \tan t, \quad -1.3 \leq t \leq 1.3,$$

expresses the altitude (in feet) of a balloon in terms of the number of hours past noon.

Desmos link: <https://www.desmos.com/calculator/re6nqofgs0>

151: Trig 2

Find exact answers to the following questions without using a calculator.

- (a) *When is the balloon descending at the rate of 150 ft/hour?*

Learning outcomes:
Author(s):

Derivatives of Trigonometric Functions

- (b) How high is the balloon when it is descending at the rate of 500 ft/hour?
 - (c) Express the balloon's rate of ascent (in feet/hour) in terms of its altitude.
-

Question 94 The function

$$h = f(t) = 11 - 7 \cos t, \quad 0 \leq t \leq 6,$$

expresses the depth of the water (in feet) at the Edmonds Pier in terms of the number of two-hour periods past noon.

Desmos link: <https://www.desmos.com/calculator/4bh7kimi7f>

151: Trig 3

Find exact answers to the following questions without using a calculator.

- (a) Is the depth of the water increasing or decreasing at 2pm? At what rate?
 - (b) At what rate is the depth of the water changing when the water is 6 feet deep?
-

Question 95 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

- (a) Use the figure below to find a function

$$h = f(\theta), \quad 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground in terms of the rotation angle of the wheel (measured in radians since the time you boarded). Use the cosine function, not the sine function.

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

- (b) The wheel stops when you are 100 feet above the ground for the second time. It then starts again and turns through a small angle of $\Delta\theta$ radians before stopping again.

Desmos link: <https://www.desmos.com/calculator/gi8obqmnnav>

151: Ferris Wheel 34

- (i) Zoom in on point P above to approximate the change in your height (in feet) if $\Delta\theta = 0.1$.
- (ii) Use a derivative to approximate the change in your height if $\Delta\theta = 0.1$.
- (iii) Use a derivative to approximate the change Δh in your height (measured in feet) as the wheel turns through the small angle $\Delta\theta$ radians.

2 Visualizing Derivatives

Question 96 You ride a ferris wheel for one revolution and get off. Let

$$h = f(s), 0 \leq s \leq ??,$$

be the function that expresses your height above the ground (measured in feet) in terms of your distance traveled, measured (in feet) along your path from your starting point.

- (a) Choose a radius for the ferris wheel that you think is reasonable and fill in the missing upper bound for the domain of f above.
- (b) Use the demonstration below to sketch by hand a graph of the function

$$r = \frac{dh}{ds} = f'(s).$$

Do not make any computations. Just use the demonstration below. Here are the key points to keep in mind:

- Approximate the derivative dh/ds by the ratio $\Delta h/\Delta s$.
- The length of the red arclength (when the ferris wheel is on the way up) is the input s .
- The length of the purple segment is the output h .
- The lengths of the orange arc and orange segments are Δs .
- The (signed) length of the green segment is Δh .

Explain your reasoning thoroughly. Be sure to include at least the following points:

- The units of the input and output to the derivative
- Scales on the vertical and horizontal axes

- A discussion of how a small change in the input to the function f changes the output at various positions along your ride.
- A discussion of where a small change in the input to f gives the greatest positive change in the output and a consideration of the ratios of these changes
- A discussion of where a small change in the input to f gives the negative change in the output with the greatest magnitude and a consideration of the ratios of these changes
- A discussion of where a small change in the input to f barely changes the output and a consideration of the ratios of these changes

(c) How would your graph of the derivative dh/ds change if you doubled the radius of the ferris wheel? Sketch the new graph.

Desmos link: <https://www.desmos.com/calculator/pxsmo04nmg>

3 The Derivative of the Sine Function

Question 97 Desmos link:

<https://www.desmos.com/calculator/jcmcyrpndw>

Desmos activity available at 151: Derivative of Sine

Question 98 (a) Describe a transformation that takes the graph of the function

$$y = f(\theta) = \sin \theta$$

to the graph of the function

$$y = g(\theta) = \cos \theta.$$

(b) Does that same transformation take the graph of the derivative of the sine function to the graph of the derivative of the cosine function? Do not use any particular facts about these functions to answer this question. Instead, give a general answer that would apply to all pairs of functions similarly related.

(c) Use your answer to part (b) and the fact that

$$\frac{d}{d\theta} (\sin \theta) = \cos \theta$$

to find an expression for the derivative

$$\frac{d}{d\theta}(\cos \theta)$$

of the cosine function.

4 Applications 1

Question 99 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

(a) Find a function

$$h = f(\theta), 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground in terms of the rotation angle of the wheel, measured in radians. Use the cosine function, not the sine function.

(b) The wheel stops when you are 100 feet above the ground and on the way up. It then starts again and turns through a small angle of $\Delta\theta$ radians before stopping again. Use the appropriate linear approximation to estimate the change Δh in your height (measured in feet) as the wheel turned through the angle θ .

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

5 Transformations of the Sine Function

Question 100 (a) Describe a transformation that takes the graph of the function

$$y = f(\theta) = \sin \theta$$

to the graph of the function

$$y = g(\theta) = \sin(2\theta).$$

(b) How does that same transformation affect the slope of a line?

(c) Use your answer to part (b) to find an expression for the derivative

$$g'(\theta) = \frac{d}{d\theta}(\sin(2\theta)).$$

6 Applications 2

Question 101 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You travel at a constant speed of 5 ft/sec as you ride the ferris wheel.

(a) Find a function

$$h = f(t), t \geq 0$$

that expresses your height above the ground in terms of the number of seconds since you got on. Use the cosine function, not the sine function.

(b) Are you ascending or descending the second time you are 90 feet above the ground? At what rate? Use the methods of this class, not vectors, to answer this question.

(c) Find your height when you are descending at the rate of 4.8 feet/sec. Give all possibilities. Do not use a calculator except to do arithmetic.

7 MATH 142

Question 102 The graph below shows the x -coordinate function of a beetle moving around a circle at a constant speed.

Desmos link: <https://www.desmos.com/calculator/qi6c9xbhnw>

142: Edmonds Pier 2

Use the graph to answer the following questions. Be sure to include units.

- Find the x -coordinate of the circle's center.
- Find the radius of the circle.
- Find the period of the motion. This the time it takes the beetle to make one revolution about the center of its circular path.
- Find a time when the beetle's x -coordinate is a maximum.
- Use (a)-(d) to find an expression $x = f(t)$ for the function that expresses the x -coordinate of the beetle (measured in feet) in terms of the number of minutes past noon. Include the domain.
- Check your expression from part (e) by substituting the two times given in the graph.

The Chain Rule

An introduction to the chain rule.

The chain rule tells us how to differentiate the composition of two functions. It says that the derivative of the composition is the product of the derivatives of the two functions.

Here is a more formal statement.

Theorem 1. (The Chain Rule) If $y = f(u)$ is a differentiable function of u and $u = g(t)$ is a differentiable function of t , then the composition

$$y = f(g(t))$$

is a differentiable function of t and

$$\frac{dy}{dt} = \frac{dy}{du} \cdot \frac{du}{dt}.$$

Or paying closer attention to the inputs,

$$\left. \frac{dy}{dt} \right|_{t=t_0} = \left. \frac{dy}{du} \right|_{u=g(t_0)} \cdot \left. \frac{du}{dt} \right|_{t=t_0}.$$

We'll go through some examples to get an understanding of how to use the chain rule and also why it works.

Examples

Example 13. Find an equation of the tangent line to the curve

$$y = f(x) = (2x^3 + 1)^2$$

at the point $(1, 9)$.

Explanation. Let

$$y = (2x^3 + 1)^2$$

and

$$u = 2x^3 + 1.$$

Then

$$y = u^2$$

Learning outcomes:
Author(s):

and

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\
 &= \frac{d}{du}(u^2) \cdot \frac{d}{dx}(2x^3 + 1) \\
 &= 2u(6x^2) \\
 &= 2(2x^3 + 1)(6x^2).
 \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 2(3)(6) = 36,$$

and an equation of the tangent line is

$$y - 9 = 36(x - 1).$$

We can get the same result without the chain rule, by rewriting the original function as

$$y = f(x) = (2x^3 + 1)^2 = 4x^6 + 4x^3 + 1.$$

Then

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx}(4x^6 + 4x^3 + 1) \\
 &= 4 \frac{d}{dx}(x^6) + 4 \frac{d}{dx}(x^3) + \frac{d}{dx}(1) \\
 &= 24x^5 + 12x^2.
 \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 14 + 12 = 36,$$

as before.

Example 14. Find an equation of the tangent line to the curve

$$x^2 + y^2 = 25$$

at the point $P(4, -3)$.

Explanation. Solve the above equation for y in terms of x to get

$$y = \pm \sqrt{25 - x^2}.$$

While this equation does not define y as a function of x , near the point $(4, -3)$, the equation

$$y = -\sqrt{25 - x^2}$$

does define y as a function of x .

Now to find the derivative of the function

$$y = -\sqrt{25 - x^2},$$

let

$$u = 25 - x^2.$$

Then

$$y = \sqrt{u} = -u^{1/2}$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(-u^{1/2}) \cdot \frac{d}{dx}(25 - x^2) \\ &= -\frac{1}{2\sqrt{u}}(-2x) \\ &= \frac{x}{\sqrt{25 - x^2}}. \end{aligned}$$

So the slope of the tangent line to the curve

$$x^2 + y^2 = 25$$

at the point $(4, -3)$ is

$$\left. \frac{dy}{dx} \right|_{x=4} = \left(\frac{x}{\sqrt{25 - x^2}} \right) \Big|_{x=4} = \frac{4}{3}$$

and an equation of the tangent line is

$$y + 3 = \frac{4}{3}(x - 4).$$

To find the slope of the tangent line without appealing to the chain rule, note that the curve

$$x^2 + y^2 = 25$$

is a circle centered at the origin. So the tangent line at $P(4, -3)$ is perpendicular to the radius \overline{OP} from the origin to P . Since \overline{OP} has slope $m_1 = -3/4$, the tangent line at $P(4, -3)$ has slope

$$m_2 = -1/m_1 = 4/3.$$

Question 103 (a) Describe a transformation that takes the circle

$$x^2 + y^2 = 25$$

to the ellipse

$$4x^2 + y^2 = 25. \quad (2)$$

(b) Find the image (call it Q) of the point $P(4, -3)$ under the transformation in part (a).

(c) Use the result of Example 2 to find an equation of the tangent line to the ellipse (2) at Q .

(d) Use the chain rule to first find the slope of the tangent line to the ellipse (2) at Q . Then find an equation of the tangent line.

Question 104 The function

$$P = f(t) = 22 + \frac{t}{2} - \frac{3t^2}{4}, 0 \leq t \leq 3,$$

expresses the price of a stock (in dollars/share) in terms of the number of hours past 9am.

(a) Is the price of the stock increasing or decreasing at 11am? At what rate? At what relative rate?

(b) Is the number of shares you can buy with \$1000 increasing or decreasing at 11am? Approximate the rate numerically.

(c) Compute the exact rate in part (b). Compute also the relative rate of change (with respect to time) in the number of shares you can buy with \$1000 at 11am.

Start your solution to this problem by defining a new function with a new function name and a new dependent variable.

Question 105 At 10am the price of a stock is increasing at the relative rate of $p\%/hr$. Is the number of shares you can buy with \$1000 increasing or decreasing at 10am? At what relative rate?

Answer this question in two ways:

(i) Going back to limits and using the definition of the derivative.

(ii) Using the chain rule.

Either way, start your solution with definitions as in the previous problem.

Question 106 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a 14-foot ladder leans against the trunk, the other rests on the horizontal ground.

Let t be the distance between the top of the ladder and the base of the trunk (measured in feet) and s the distance between the bottom of the ladder and the base of the trunk (also measured in feet).

Geogebra link: <https://www.geogebra.org/m/ctf2bcqz>

Geogebra activity available at 151: Ladder and Tree Part 2

- (a) Use the law of cosines to write an equation relating t and s .
- (b) Use the result of part (a) to find the two possible values of t when $s = 16$.

For the remainder of this problem we'll focus on positions of the ladder when $s \sim 16$ and $t \sim 10$.

- (c) For these positions, complete the square to find a function

$$t = f(s)$$

that expresses the distance (in feet) from the top of the ladder to the base of the trunk in terms of the distance (in feet) from the bottom of the ladder to the trunk's base.

- (d) Drag the slider θ in the worksheet above to approximate the value of the derivative

$$\left. \frac{dt}{ds} \right|_{s=16}.$$

Include units and explain your reasoning.

- (e) Use your function from part (c) to find an expression for the derivative dt/ds . Show every step in computing the derivative as in Examples 1 and 2 above.

- (f) Evaluate the derivative

$$\left. \frac{dt}{ds} \right|_{s=16}$$

and compare the exact value with your estimate from part (d).

- (g) What are the units of the derivative in part (f)? Explain the meaning of the derivative, not by giving a standard response about the rate of change of some quantity with respect to another, but by relating small changes.

- (h) Use your response to part (f) to write an approximation for the change

$$\Delta t = f(s) - f(16)$$

in terms of the change

$$\Delta s = s - 16$$

for values of s near $s = 16$. Use this approximation to estimate the distance between the top of the ladder and the base of the trunk when the bottom of the ladder is 15.6 feet from the base of the trunk. Compare your approximation with the exact distance.

(i) Let $\theta = m\angle GBA$ be the radian measure of the angle the ladder makes with the trunk.

part (i) Find an expression for the derivative $d\theta/ds$.

part (ii) Evaluate the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=16}$$

and interpret its meaning in terms of small changes. Be sure to explain the significance of the derivative's sign as well.

Example 15. (a) Find the slope of the tangent line to the curve

$$y = f(\theta) = 6 \sin \theta$$

at the point with coordinates $P(2\pi/3, 3\sqrt{3})$. We do not need the chain rule for this.

(b) Describe a transformation that takes the graph of $y = f(\theta)$ to the graph of the function

$$y = g(\theta) = 6 \sin(\theta/2).$$

(c) Find the image of the point $P(2\pi/3, 3\sqrt{3})$ under the above transformation.

(d) Use the results of parts (a) - (c) and the demonstration below to guess the slope of the tangent line to the curve $y = g(\theta)$ at the point $Q(4\pi, 3\sqrt{3})$.

Desmos link: <https://www.desmos.com/calculator/mqjxpsqyo5>

Desmos activity available at 151: Chain Rule 1

(e) Use the chain rule to confirm your guess from part (d).

Example 16. Find an equation of the tangent line to the curve

$$y = f(t) = 4 \sin^2 \left(\frac{\pi}{6} t \right)$$

at the point $(2, 3)$.

Explanation. Let

$$y = 4 \sin^2 \left(\frac{\pi}{6} t \right) = 4 \left(\sin \left(\frac{\pi}{6} t \right) \right)^2$$

and

$$u = \sin \left(\frac{\pi}{6} t \right).$$

Then

$$y = 4u^2$$

and

$$\begin{aligned}\frac{dy}{dt} &= \frac{dy}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du}(4u^2) \cdot \frac{d}{dt}\left(\sin\left(\frac{\pi}{6}t\right)\right) \\ &= 8u \cos\left(\frac{\pi}{6}t\right) \frac{d}{dt}\left(\frac{\pi}{6}t\right) \\ &= 8 \sin\left(\frac{\pi}{6}t\right) \cos\left(\frac{\pi}{6}t\right) \frac{\pi}{6}.\end{aligned}$$

Question 107 (a) The third equality above uses the chain rule again. Fill in the missing details of this computation by first letting

$$z = \sin\left(\frac{\pi}{6}y\right)$$

and making an explicit substitution using the variable v in place of u .

(b) Complete the solution by finding an equation of the tangent line to the curve at the point $(2, 3)$.

Question 108 Use the chain rule to compute each of the following derivatives. Show all steps as in Examples 1, 2, and 8 above.

(a) $\frac{d}{dx}\left(\frac{12}{1+x^2}\right)$

(b) $\frac{d}{dt}\left(12(3)^{t/5}\right)$

(b) $\frac{d}{dt}\left(e^{3\sin(4t)}\right)$

(b) $\frac{d}{d\theta}(12 + 5\cos(\theta/4))$

Example 17. The function

$$P = f(t) = 10e^{\frac{1}{4}t}, \quad -4 \leq t \leq 12,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

(a) What are the units of $1/4$ in the function above? How do you know?

(b) Use the chain rule to find an expression for the growth rate of the population at time t hours past noon. What are the units of the growth rate?

- (c) Express the growth rate from part (b) in terms of the population $P = f(t)$ at time t hours past noon.
- (d) What is the relative instantaneous growth rate of the population? Include units.
- (e) Find the relative average growth rate of the population over a one-hour period.
- (f) Describe what happens to the population every hour.

Example 18. (a) Use the chain rule to find the relative instantaneous growth rate of the population function

$$P = 10(2^t), \quad -2 \leq t \leq 5,$$

where P is measured in millions of bacteria and t is the number of hours past noon.

- (b) Describe what happens to the population every hour.
- (c) Estimate the population 2 minutes after there are 30 million bacteria.

Question 109 This problem suggests a way to think about the chain rule geometrically.

You ride a ferris wheel for one revolution and get off. The function

$$h = f(\theta) = , \quad 0 \leq \theta \leq 2\pi,$$

expresses your height (in feet) in terms of the wheel's angle of rotation (measured in radians from the moment you boarded).

The function

$$\theta = g(t), \quad 0 \leq t \leq 44,$$

expresses the rotation angle of the wheel in terms of the number of seconds since you boarded.

Our goal is to use the graphs of the function f and g below (take the times t to be positive, not negative as shown) to approximate your rate of ascent $t = 16$ seconds after boarding. We'll do this in two different ways.

- (a) The first way involves a few steps and uses the graphs of both f and g .
- (i) Approximate the value of the derivative

$$\left. \frac{dh}{d\theta} \right|_{\theta=g(16)}$$

and interpret its meaning. Include units.

- (ii) Approximate the value of the derivative

$$\left. \frac{d\theta}{dt} \right|_{t=16}$$

and interpret its meaning. Include units.

(iii) Use the results of (i) and (ii) and common sense (do not appeal to the chain rule directly) to approximate the value of the derivative

$$\left. \frac{dh}{dt} \right|_{t=16} = h'(16),$$

where $h(t) = f(g(t))$. Explain your logic and include all units in your computation. Interpret the meaning of this derivative.

(b) The second way has just one step and that's to use the graph of $h = f(\theta)$ to approximate your rate of ascent at time $t = 16$ using the fact that the horizontal blue lines are drawn at intervals of $\Delta t = 2$ seconds. Be sure to explain your logic.

Desmos link: <https://www.desmos.com/calculator/fkgfypsowe8>

Desmos activity available at 151: Ferris Wheel 2

Question 110 This is a continuation of the previous question.

Approximate the time between times $t = 5$ and $t = 22$ seconds when you are ascending at the slowest rate. Approximate this rate of ascent.

Question 111 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

(a) Use the geometry of the ferris wheel (see the picture below) to find a function

$$h = f(\theta), \quad 0 \leq \theta \leq 2\pi,$$

that expresses your height (in feet) above the ground in terms of the rotation angle of the wheel, measured in radians. Use the cosine function, not the sine function.

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

(b) Suppose you move at the constant speed of 10 ft/sec as you ride the ferris wheel.

(i) Use the result of part (a) to find a function $h = g(t)$ that expresses your height (in feet) in terms of the number of seconds since you boarded. Include the appropriate domain.

(ii) Find a function $r = h(t)$ that expresses your rate of ascent (in ft/sec) in terms of the number of seconds since you boarded. Include the appropriate

domain. Use vectors to interpret this rate geometrically, in terms of the angle your velocity makes with the horizontal and your speed.

(iii) Write an equation that relates your height h (in feet) and your rate of ascent r (in ft/sec) at any instant. Graph the relation by hand.

(iv) Are you going up or down the second time you are 100 feet above the ground? At what rate?

(v) What is your height when you are descending at the rate of 4 ft/sec? Find all possibilities.

Question 112 The function

$$\theta = f(t), t \geq 0,$$

expresses the radian measure of a ferris wheel's rotation angle in terms of the number of seconds since you boarded. The wheel has radius r feet and its center is b feet above the ground.

(a) Find a function

$$h = g(t), t \geq 0,$$

that expresses your height (in feet) in terms of the number of seconds since you boarded.

(b) Find an expression for your rate of ascent (in ft/sec) at time t seconds after you boarded. Assume g is a differentiable function of t .

(c) Interpret your rate of ascent (part (b)) in terms of your speed at time t and the angle your velocity vector makes with the horizontal. Assume here that $\theta = f(t)$ is an increasing function of t .

Exercises

Exercise 113 Use the facts that

$$\frac{d}{d\theta}(\sin \theta) = \cos \theta$$

and

$$\frac{d}{d\theta}(\cos \theta) = -\sin \theta$$

to compute each of the following derivatives. Show all steps in using the chain rule as in Examples 1 and 2 above.

(a) $\frac{d}{d\theta}(\sec \theta)$

$$(b) \frac{d}{d\theta} (\csc \theta)$$

$$(c) \frac{d}{dt} \left(\sqrt{\sin^2 t + (4 - \cos t)^2} \right)$$

Hint: Simplify the function first.

$$(d) \frac{d}{d\theta} (\sec^2 \theta)$$

$$(e) \frac{d}{dt} \left(\frac{1}{5 + 3 \cos(t/2)} \right)$$

Explanation. (d) Let

$$y = \sec^2 \theta = (\sec \theta)^2$$

and let

$$u = \sec \theta.$$

Then

$$y = u^2$$

and

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du} (u^2) \cdot \frac{d}{d\theta} (\sec \theta) \\ &= 2u(\sec \theta \tan \theta) \\ &= 2(\sec \theta)(\sec \theta \tan \theta). \end{aligned}$$

(e) Let

$$y = \frac{1}{5 + 3 \cos(t/2)}$$

and let

$$u = 5 + 3 \cos(t/2).$$

Then

$$y = u^{-1}$$

and

$$\begin{aligned} \frac{dy}{dt} &= \frac{dy}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du} (u^{-1}) \cdot \frac{d}{dt} (5 + 3 \cos(t/2)) \\ &= -\frac{1}{u^2} \left(-3 \sin \left(\frac{t}{2} \right) \frac{d}{dt} \left(\frac{t}{2} \right) \right) \\ &= \frac{3 \sin \left(\frac{t}{2} \right)}{2(5 + 3 \cos(t/2))^2}. \end{aligned}$$

Exercise 114 You jog around a circular track of radius r feet at the constant speed of v ft/sec. A flagpole lies b feet due east of the track's center.

(a) Use the animation below (and nothing else) to sketch by hand a graph of the function

$$s = f(\theta), \theta \geq 0,$$

that expresses your distance (in feet) to the flagpole in terms of your angle $\theta = \angle FOJ$ of rotation about the track's center, measured in radians from the time you start. Assume you start at the point A on the track nearest the flagpole. Be sure to include variable names, units, and scales on your axes. Explain your reasoning. For this particular graph assume that $r = 40$ and $b = 96$ (be sure to adjust the sliders in the worksheet to have these values).

Desmos link: <https://www.desmos.com/calculator/4pndurvhd>

Worksheet available at 151: Jogger

(b) Using only your graph of the function f from part (a), sketch by hand a graph of the derivative

$$y = f'(\theta) = ds/d\theta$$

when $r = 40$ and $b = 96$. Be sure to include variable names, units, and scales on your axes. Explain your reasoning.

(c) Using only the animation (stop the motion), approximate the value of the derivative

$$\left. \frac{ds}{d\theta} \right|_{\theta=3\pi/2}$$

when $r = 40$ and $b = 96$. Include units. Explain your reasoning.

(d) Use trigonometry to find an expression for the function $s = f(\theta)$ in terms of the parameters r and b (ie. do not assume $r = 40$ and $b = 96$).

(e) Find an expression for the derivative $ds/d\theta$. Show every step in using the chain rule as in Examples 1 and 2. Do not assume $r = 40$ and $b = 96$

(f) Use your result from part (e) to compute the exact value of the derivative

$$\left. \frac{ds}{d\theta} \right|_{\theta=3\pi/2}$$

when $r = 40$ and $b = 96$. Compare this with your estimate from part (c).

(g) What are the units of the derivative in part (f)? Explain the meaning of the derivative, not by giving a standard response about the rate of change of some quantity with respect to another, but by relating small changes.

(h) Use your response to part (f) with $r = 40$ and $b = 96$ to write an approximation for the change

$$\Delta s = f(\theta) - f(3\pi/2)$$

in terms of the change

$$\Delta\theta = \theta - 3\pi/2$$

for values of θ near $\theta = 3\pi/2$.

(i) Find a function

$$s = g(t), t \geq 0,$$

that expresses your distance (in feet) to the flagpole in terms of the number of seconds since you started jogging. Assume you start at the point on the track nearest the flagpole.

(j) Find a function that expresses the rate of change (with respect to time) in your distance to the flagpole in terms of the number of seconds since you began jogging.

(k) Express the rate of change in part (b) in terms of your speed and the angle between the following two vectors:

- the vector that gives your position relative to the flagpole
- the vector that points in the direction of your motion

Exercise 115 High tide of 10.83 feet at Edmonds at 12:28am, May 15, low tide at 6:58am of ?? feet, lowest tide of 1.64 feet at 5:32pm.

Desmos link: <https://www.desmos.com/calculator/zta9tkzzmx>

Worksheet available at 151: Edmonds Tides

Exercise 116 A pendulum of length L feet is $L + 5$ oscillates sinusoidally between angles $-\theta_0$ and θ_0 with period $2\pi\sqrt{L/g}$ seconds, where g is a constant. The angles $\pm\theta_0$ are measured in radians from the downward vertical.

(a) Find a function

$$h = f(\theta), -\theta_0 \leq \theta \leq \theta_0,$$

that expresses the height of the pendulum above its stable equilibrium position (ie. its lowest point) in terms of the angle of rotation. Use the cosine function.

(b) Assume now that the pendulum is released from rest from the displacement angle $\theta = \theta_0$ at time $t = 0$ seconds. Find a function

$$\theta = g(t), t \geq 0,$$

that expresses the displacement angle (in radians) in terms of the number of seconds since the pendulum was released. Use the cosine function.

(c) Find a function

$$h = w(t), t \geq 0$$

that expresses the height (in feet) of the pendulum above equilibrium in terms of the number of seconds since the pendulum was released.

(d) Find an expression for the derivative dh/dt and interpret its meaning.

Exercise 117 The function

$$h = f(s) = 3 + 2 \cos s, 0 \leq s \leq 3,$$

expresses the altitude (in thousand of feet) along a mountain trail in terms of the distance (measured in miles) from the summit. The distance is measured along the trail.

The function

$$s = g(t) = t + \frac{t^2}{4}, 0 \leq t \leq 2,$$

expresses your distance from the summit (in miles) in terms of the number of hours past noon.

The goal of this problem is to compute the rate (with respect to time) at which your altitude is changing at 1pm.

Desmos link: <https://www.desmos.com/calculator/twfsqmmgyb>

Worksheet available at 151: Mountain Road

(a) Find an expression for a function $h = k(t)$ that expresses your altitude (in thousands of feet) in terms of the number of hours past noon. Include a domain.

(b) Evaluate the derivative

$$\left. \frac{ds}{dt} \right|_{t=1}$$

and interpret its meaning.

(c) Evaluate the derivative

$$\left. \frac{dh}{ds} \right|_{s=g(1)}$$

and interpret its meaning.

(d) Use the results of (b) and (c) to compute the rate (with respect to time) at which your altitude is changing at 1pm.

(e) Express the rate of part (d) as a derivative.

(f) Express the derivative

$$\left. \frac{dh}{dt} \right|_{t=1}$$

in terms of the derivatives

$$\left. \frac{ds}{dt} \right|_{t=1}$$

and

$$\left. \frac{dh}{ds} \right|_{s=g(1)}.$$

Exercise 118 A weight is attached to the midpoint of a 10-foot rope.

You hold the ends of the rope 10 feet apart at the same height. You then move the ends directly toward each other, each at the constant speed of 2 ft/sec.

Desmos link: <https://www.desmos.com/calculator/grostrmdln>

Worksheet available at 151: *Weighted Rope*

(a) Find a function

$$v = f(t), \quad 0 \leq t \leq 2.5,$$

that expresses the speed of the weight (in feet/sec) in terms of the number of seconds since you began moving your hands.

(b) Describe how the speed of the weight varies.

(c) Comments?

Exponential Functions, Part 2

More on exponential functions and their derivatives.

Question 119 What type of functions change at

- (a) a constant rate?
- (b) a constant relative rate?

Question 120 The function

$$P = f(t) = 50e^{\frac{t}{10}}, 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past midnight. Graph shown below.

Desmos link: <https://www.desmos.com/calculator/kffjsoepmo>

151: Exp Function 5

- (a) Use the graph of the function $P = f(t)$ above to approximate the growth rate when there are 200 million bacteria.
- (b) Find the relative (instantaneous) growth rate of the population.
- (c) Find the exact growth rate when there are 200 million bacteria. No calculator.
- (d) Find a function $r = g(P)$ that expresses the growth rate (measured in millions of bacteria/hour) in terms of the population (measured in millions). Include a domain.
- (e) What is the population when it is increasing at the rate of 8 million bacteria/hour?

Learning outcomes:
Author(s):

Question 121 The function

$$N = f(t) = \frac{20}{1 + 5e^{-t/5}}, \quad 0 \leq t \leq 30,$$

models the spread of a virus throughout a population. It takes as an input the number of months since January 1st and returns as an output the number of infected individuals (measured in millions). Graph shown below.

Desmos link: <https://www.desmos.com/calculator/ikis451wxu>

151: Logistic Model

- (a) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at the rate of 800,000 people/month.
- (b) Find a function

$$r = g(N)$$
 that expresses the infection rate (in millions of people/month) in terms of the population.
- (c) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at the rate of 750,000 people/month.
- (d) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at its fastest rate. Also approximate this rate.
- (e) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at its fastest rate. Then find this exact rate.

Question 122 Use the chain rule and your knowledge of the derivative $d(e^x)/dx$ to compute each of the following derivatives.

(a)
$$\frac{d}{dt}(100 \cdot 2^t)$$

(b)
$$\frac{d}{dt}(100 \cdot 2^{t/5})$$

Explanation. (a) The key is to realize that

$$2 = e^{\ln 2}.$$

So we can write the function to differentiate as

$$P = 100 \cdot 2^t = 100 (e^{\ln 2})^t = 100e^{t \ln 2}.$$

Now we can make a u -substitution and use the chain rule. We let

$$u = t \ln 2.$$

Then

$$P = 100e^{t \ln 2} = 100e^u$$

and

$$\begin{aligned} \frac{dP}{dt} &= \frac{dP}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du} (100e^u) \cdot \frac{d}{dt} (t \ln 2) \\ &= 100e^u (\ln 2) \\ &= P \ln 2 \\ &= (100 \cdot 2^t) \ln 2 \\ &= (100 \ln 2)(2^t) \end{aligned}$$

Question 123 A colony of bacteria has a population of 20 million at noon and a population of 25 million at 2pm. The population grows exponentially between noon and midnight.

- (a) Describe precisely how the population grows.
- (b) Use your description from part (a) to find a function

$$P = f(t), 0 \leq t \leq 12,$$

that expresses the population (in millions) in terms of the number of hours past noon. Do not use e .

- (c) Find the relative instantaneous growth rate of the population.
- (d) Find a function $r = g(P)$ that expresses the (absolute) growth rate in terms of the population.
- (e) At what rate is the population changing when there are 30 million bacteria?

Explanation. (a) The population increases by

$$\frac{25 \text{ million} - 20 \text{ million}}{20 \text{ million}} = 25\%$$

every two hours.

Better yet, at least for determining the function, is to say that the two-hour growth factor is

$$\frac{25 \text{ million}}{20 \text{ million}} = 1.25.$$

This means that every two hours the population gets multiplied by 1.25.

(b) Since the two-hour growth factor is 1.25, the one-hour growth factor is $1.25^{1/2}$. So

$$P = 20(1.25^{1/2})^t = 20(1.25)^{\frac{t}{2}}, \quad 0 \leq t \leq 12.$$

(c) The key to computing the derivative dP/dt is to write the population function using base e . Since

$$1.25 = e^{\ln 1.25},$$

$$\begin{aligned} P &= 20(1.25)^{\frac{t}{2}} \\ &= 20(e^{\ln 1.25})^{\frac{t}{2}} \\ &= 20e^{t(\ln 1.25)/2} \end{aligned}$$

So the growth rate (in millions of bacteria/hour) is

$$\begin{aligned} \frac{dP}{dt} &= \frac{d}{dt} \left(20(1.25)^{\frac{t}{2}} \right) \\ &= 20 \frac{d}{dt} \left(e^{t(\ln 1.25)/2} \right) \\ &= \left(20e^{t(\ln 1.25)/2} \right) \frac{d}{dt} (t(\ln 1.25)/2) \\ &= P(\ln 1.25)/2. \end{aligned}$$

And the relative growth rate (units are hr^{-1}) is

$$\frac{1}{P} \cdot \frac{dP}{dt} = \frac{1}{2} \ln 1.25.$$

(d) From part (c), the function

$$g(P) = \frac{dP}{dt} = P(\ln 1.25)/2$$

expresses the growth rate (in millions of bacteria/hour) in terms of the population.

- (e) When there are 30 million bacteria, the population is increasing at the rate of

$$g(30) = (30 \text{ million bacteria}) \left(\frac{1}{2} \ln 1.25/\text{hr} \right) = 15 \ln 1.25 \text{ million bacteria/hr.}$$

Question 124 Between ground level and an altitude of 10 km, atmospheric pressure on the planet Krypton is an exponential function of altitude. The pressure is 100 kPa at ground level and 80 kPa at an altitude of 0.75 km.

Desmos link: <https://www.desmos.com/calculator/j5h8kaj8xs>

151: Atmospheric Pressure

- (a) Describe exactly how the pressure decreases.
 (b) Use your description from part (a) to find a function

$$P = f(h), \quad 0 \leq h \leq 10,$$

that expresses the atmospheric pressure (in kPa) in terms of the altitude (in kilometers). Do not use the number e .

- (c) Input your expression for f in Line 2 of the worksheet above.
 (d) Evaluate the derivative

$$\left. \frac{dP}{dh} \right|_{h=5}$$

and interpret its meaning.

- (e) Find a function $r = g(P)$ that expresses the derivative dP/dh in terms of P .
 (f) Use calculus to approximate the altitude at which the pressure is 0.2 kPa less at an altitude 0.1 km higher.

Practice Quiz 2

Practice quiz, Weeks 4-6

Directions:

- (a) Show all work.
- (b) Give brief explanations for each problem. Include these explanations in the flow of the solution.
- (c) Show all units in all computations.
- (d) No calculators.
- (e) Show each step when using the chain, product, and quotient rules and use the Leibniz notation when doing so. Here is an example.

Example 19. Find an expression for the derivative

$$\frac{d}{d\theta} (4 + 5 \cos^3(2\theta)) .$$

Explanation. With $y = 4 + 5 \cos^3(2\theta)$, the derivative $dy/d\theta$ is

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{d}{d\theta} (4 + 5 \cos^3(2\theta)) \\ &= \frac{d}{d\theta} (4) + 5 \frac{d}{d\theta} ((\cos(2\theta))^3) \\ &= 5(3)(\cos(2\theta))^2 \frac{d}{d\theta} (\cos(2\theta)) \\ &= 15(\cos(2\theta))^2 (-\sin(2\theta)) \frac{d}{d\theta} (2\theta) \\ &= -30(\cos(2\theta))^2 \sin(2\theta). \end{aligned}$$

Question 125 Find expressions for each of the following derivatives.

(a)

$$\frac{d}{dt} (4 - t^4 e^{-5t})$$

(b)

$$\frac{d}{d\theta} \left(\frac{12}{4 + 5 \tan(4\theta)} \right)$$

Learning outcomes:
Author(s):

(c)

$$\frac{d}{dw} \left(\sqrt{40 - 5w^2} \right)$$

(d)

$$\frac{d}{dt} \left(2^{t/8} \right)$$

Question 126 (a) Find an equation of the tangent line to the curve

$$y = 4 - t^4 e^{-5t}$$

at the point on the curve with t -coordinate $t = 0$.

(b) Find an equation of the tangent line to the curve

$$y = \frac{12}{4 + 5 \tan(4\theta)}$$

at the point on the curve with θ -coordinate $\theta = 0$.

Question 127 Assume for this question that each month has 30 days and that the number of hours of daylight/day in Seattle is a sinusoidal function of time. Assume also that on June 21, Seattle gets a maximum of 16 hours of daylight/day and that on December 21, Seattle gets a minimum of 8 hours of daylight/day.

(a) Find a function

$$H = f(t), \quad 0 \leq t \leq 12,$$

that expresses the number of daylight hours/day in Seattle in terms of the number of months since June 21. Use the cosine function. Start by sketching a graph. Explain your reasoning.

(b) Evaluate the derivative

$$\left. \frac{dH}{dt} \right|_{t=2}.$$

(c) What are the units of the derivative above?

(d) Explain the meaning of the above derivative.

Question 128 The function

$$T = f(m) = 20 + 70e^{-m/20}, \quad 0 \leq m \leq 60,$$

expresses the temperature (in Celsius degrees) of a cup of coffee in terms of the number of minutes past noon.

- (a) At what rate is the temperature changing at 12:40pm?
- (b) Does the temperature change at a constant relative rate? Justify your assertion.
- (c) Find a function $r = g(T)$ that expresses the rate of change in the temperature (measured in $^{\circ}\text{C}/\text{min}$) in terms of the temperature (measured in Celsius degrees).
- (d) Find the temperature when it is decreasing at the rate of $3^{\circ}\text{C}/\text{min}$.
- (e) Find a function $s = h(T)$ that expresses the relative rate of change in the temperature in terms of the temperature (measured in Celsius degrees).
- (f) Find the temperature when it is decreasing at the rate of $3\%/\text{min}$.

Question 129 Assume for this problem that over the course of a 24-hour period beginning at midnight of July 21, the temperature in Shoreline is a sinusoidal function of time. Assume also that the temperature reaches its minimum of 50°F at 5am and its maximum of 80° at 5pm.

At what rate (with respect to time) is the temperature changing when it is 68°F for the second time? Start by finding a function that expresses the temperature in terms of time. Be sure to define your variables.

Question 130 The function

$$h = f(t) = 3 + 5 \tan\left(\frac{t-9}{12}\right), \quad 0 \leq t \leq 12,$$

expresses the height (in thousands of feet) of a balloon in terms of the number of hours past noon.

Determine the possible height(s) of the balloon when it is ascending at the rate $500 \text{ ft}/\text{hour}$.

Question 131 The function

$$G = f(s) = 5e^{-s/20}, \quad 0 \leq s \leq 40,$$

expresses the number of gallons of gas in a truck in terms of the trip odometer reading (measured in miles).

Find a function

$$m = h(s), \quad 0 \leq s \leq 40,$$

that takes expresses the dashboard reading for the number of miles left to drive before running out of gas in terms of the trip odometer reading (in miles). The dashboard reading assumes the truck burns gas at the current rate (ie. the rate at odometer reading s miles) for the remainder of the trip.

Derivatives of Inverse Functions, Short Version

Derivatives of inverse functions.

1 The Main Idea

The main idea here is simple. That the derivative of the *inverse* of a function is the *reciprocal* of the function's derivative. Or expressed more succinctly, *the derivative of the inverse is the reciprocal of the derivative.*

We don't really need to know much about calculus to understand why.

Question 132 Take for example, the function

$$G = f(s), 0 \leq s \leq 20,$$

that expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route and the graph of the function is shown below.

Desmos link: <https://www.desmos.com/calculator/oltnpzth9>

151: Inverse Function 1

- (a) Use the slider s above to approximate (with units) the derivative

$$\left. \frac{dG}{ds} \right|_{s=2}.$$

Explain the derivative's meaning.

- (b) Use the slider s above to approximate (with units) the derivative

$$\left. \frac{ds}{dG} \right|_{G=f(2)}.$$

Explain the derivative's meaning.

- (c) Write an equation that expresses the relationship between the derivatives in parts (a) and (b). Explain why this relationship holds.

Learning outcomes:
Author(s):

Question 133 This question is a continuation of Question 1. Now we are given an expression

$$G = f(s) = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}, \quad 0 \leq s \leq 20,$$

for the function f .

- (a) Use calculus to find an expression for the derivative dG/ds and to compute the exact value of the derivative

$$\left. \frac{dG}{ds} \right|_{s=2}.$$

- (b) Use the idea of Question 1 to find an expression for the derivative ds/dG and the exact value of the derivative

$$\left. \frac{ds}{dG} \right|_{G=f(2)}.$$

- (c) For a more algebraic way to find an expression for the derivative ds/dG , we'll differentiate both sides of the equation

$$G = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}$$

with respect to G . Keeping in mind that this equation, for values of s between 0 and 20, defines s implicitly as a function of G (why?). Then using the chain rule tells us that

$$\frac{dG}{dG} = \left(\frac{s + 23}{1000} \right) \left(\frac{ds}{dG} \right).$$

And solving for the derivative ds/dG gives

$$\frac{ds}{dG} = \frac{1000}{s + 23}.$$

2 Derivatives of Inverse Trig Functions

The idea of computing the derivative of the inverse of a one-to-one function as illustrated above is simple. But if we require that the derivative of the inverse be expressed in terms of *its independent variable*, the algebra can get a bit tricky.

Example 20. Take for example the function

$$y = g(\theta) = \sin \theta.$$

Now this function is *not* one-to-one and so its inverse is not a function. But we can restrict the domain of g to get a new function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2, \quad (3)$$

that is one-to-one and that has the same range $\{y \mid -1 \leq y \leq 1\}$ as g . The inverse of this new function f is the inverse sine function

$$\theta = f^{-1}(y) = \arcsin(y) = \sin^{-1}(y).$$

The function $\arcsin(y)$ gives the angle between $-\pi/2$ and $\pi/2$ whose sine is equal to y . The key to expressing its derivative

$$\frac{d\theta}{dy} = \frac{d}{dy} (\arcsin y)$$

in terms of y , is to express the derivative of the original function $y = f(\theta)$ in terms of its output $y = \sin \theta$.

Now

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (\sin \theta) = \cos \theta.$$

To express this derivative in terms of $y = \sin \theta$, we use the fact that

$$\cos^2 \theta + \sin^2 \theta = 1.$$

So

$$\frac{dy}{d\theta} = \cos \theta = \pm \sqrt{1 - \sin^2 \theta} = \pm \sqrt{1 - y^2}.$$

To choose between \pm , we must remember that by the definition of the arcsine function (see equation (1)),

$$\pi/2 \leq \theta \leq \pi/2.$$

This tells us that $\cos \theta \geq 0$. So we conclude that

$$\frac{d}{dy} (\arcsin y) = \cos \theta = \sqrt{1 - y^2}.$$

3 Exercises

Exercise 134 Express the derivative of each of the following functions in terms of its output.

Derivatives of Inverse Functions, Short Version

(a)

$$x = \cos \theta, 0 \leq \theta \leq \pi$$

(b)

$$x = \cos \theta, \pi \leq \theta \leq 2\pi$$

(c)

$$y = \tan \theta, -\pi/2 < \theta < \pi/2$$

(d)

$$P = e^{t/5}, t \in \mathbb{R}$$

(e)

$$y = \frac{e^x - e^{-x}}{2}, x \in \mathbb{R}$$

(f)

$$y = \frac{e^x - e^{-x}}{e^x + e^{-x}}, x \in \mathbb{R}$$

(g)

$$y = \frac{e^x + e^{-x}}{2}, x \leq 0$$

(h)

$$G = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}, 0 \leq s \leq 20,$$

Exercise 135 Use the results of Exercise 5 to express the derivative of the inverse of each of the functions in that exercise in terms of its independent variable.

Exercise 136 • Find simplified expressions for each of the following derivatives.

- Do not simplify a function before taking its derivative.
- Show all steps in using the chain rule. Use the Leibniz notation.
- Use Desmos to graph each function (before taking the derivative).
- Explain how you could have found an expression for each derivative directly from the graph of the (original) function.

(a)

$$\frac{d}{dx} (\sin(\arcsin x))$$

Derivatives of Inverse Functions, Short Version

(b)

$$\frac{d}{d\theta} (\arcsin(\sin \theta))$$

(c)

$$\frac{d}{dx} (\arctan x + \arctan(1/x))$$

(d)

$$\frac{d}{dx} \left(\arccos x + \arccos \left(\sqrt{1 - x^2} \right) \right)$$

Derivatives of Inverse Functions

Using the chain rule to compute the derivative of the inverse of a function.

Introduction

We saw at the beginning of the class the relationship between the derivative of a function and the derivative its inverse. Recall the two problems:

- You measure the edge length of a cube to be 2 cm and use this measurement to compute its volume. How is the error in computing the volume related to the error (assumed to be small) in your measurement of the edge length?
- You submerge a cube in water and measure its volume to be 8 cm³. You then use this measurement to compute the edge length of the cube. How is the error in computing the edge length related to the error (assumed to be small) in your measurement of the volume?

The key to the first problem was to differentiate the function

$$V = f(s) = s^3, s > 0$$

that expresses the volume (in cubic cm) of the cube in terms of its edge length (in cm). We used limits and found that

$$\left. \frac{dV}{ds} \right|_{s=2} = 12.$$

Question 137 (a) What are the units of the above derivative? How do you know?

(b) Interpret the meaning of the above derivative in terms of the geometry of the cube.

Question 138 From here, we let

$$\Delta s = s - 2$$

Learning outcomes:
Author(s):

be the error in our measurement (measured in cm) and

$$\Delta V = s^3 - 8$$

the error in the computed volume (measured in cubic cm). Then for small errors $\Delta s \sim 0$,

$$\frac{\Delta V}{\Delta s} \sim \left. \frac{dV}{ds} \right|_{s=2} = 12,$$

and so

$$\Delta V \sim 12\Delta s.$$

For the second problem we used limits to differentiate the function

$$s = f^{-1}(V), \quad V > 0,$$

and found that

$$\left. \frac{ds}{dV} \right|_{V=8} = 1/12.$$

Question 138.1 What are the units of the above derivative? How do you know?

Then for small errors ΔV in our measurement of the volume,

$$\Delta s \sim \frac{1}{12}\Delta V$$

But all this work took us back to the same approximate relation between the errors ΔV and Δs that we had already found in solving the first problem.

Expressed another way, having found that

$$\left. \frac{dV}{ds} \right|_{s=2} = 12,$$

we should have known immediately (without any computations) that

$$\left. \frac{ds}{dV} \right|_{V=8} = 1/12.$$

For the same reason, this same relationship between the derivative of any function and the derivative of its inverse holds. That is, for any differentiable function $y = f(x)$, the derivative dx/dy of the inverse function $x = f^{-1}(y)$ is

$$\begin{aligned} \frac{d}{dy} (f^{-1}(y)) &= \frac{dx}{dy} \\ &= \frac{1}{dy/dx}. \end{aligned}$$

Well almost. We need to take some care in evaluating the above derivatives at the correct inputs.

Supposing that $b = f(a)$, a more precise statement of the relation between the derivative of a function $y = f(x)$ and the derivative dx/dy of its inverse is

$$\begin{aligned}\frac{d}{dy} (f^{-1}(y)) \Big|_{y=b} &= \frac{dx}{dy} \Big|_{y=b} \\ &= \frac{1}{\frac{dy}{dx} \Big|_{x=a}}.\end{aligned}$$

That is, the derivative of the inverse (of a function) is the reciprocal of the derivative (of that function).

Well, not quite. The above relationship is true if

$$\frac{dy}{dx} \Big|_{x=a} \neq 0.$$

This condition also guarantees that the function $y = f(x)$ is one-to-one in a sufficiently small neighborhood near $x = a$ and is therefore invertible in that neighborhood.

Here are some examples.

The Derivative of the Natural Log Function

Example 21. (a) Use the graph of the function $y = f(x)$ below to estimate the value of the derivative

$$\frac{dy}{dx} \Big|_{x=1.1} = \frac{dy}{dx} \Big|_{y=3} \sim 3$$

(b) Use the result of part (a) and the graph below to estimate the value of the derivative

$$\frac{dx}{dy} \Big|_{x=1.1} = \frac{dx}{dy} \Big|_{y=3} \sim 1/3.$$

Desmos link: <https://www.desmos.com/calculator/nshzdh6jpp>

(c) Set $u = 0.6931$ and $n = 90$ in the demonstration above and estimate the value of the derivative

$$\frac{dy}{dx} \Big|_{x=0.6931} = \frac{dy}{dx} \Big|_{y=2} \sim 2$$

(d) Use the result of part (c) and the graph above to estimate the value of the derivative

$$\left. \frac{dx}{dy} \right|_{x=0.6931} = \left. \frac{dx}{dy} \right|_{y=2} \sim 0.5$$

Question 139 Parts (a) and (c) and perhaps a few more derivatives suggest that

$$f(x) = e^x.$$

(e) Use the results of parts (b) and (d) to evaluate the derivatives

$$\left. \frac{dx}{dy} \right|_{y=3} = \left. \frac{d}{dy} (\ln y) \right|_{y=3} = 1/3,$$

$$\left. \frac{dx}{dy} \right|_{y=2} = \left. \frac{d}{dy} (\ln y) \right|_{y=2} = 1/2,$$

and more generally

$$\left. \frac{dx}{dy} \right|_{y=a} = \left. \frac{d}{dy} (\ln y) \right|_{y=a} = 1/a.$$

Question 140 Our conclusion is that

$$\frac{d}{dx} (\ln x) = \frac{1}{x}.$$

Geogebra activity available at [151: Magnification Factor 3](#)

Question 141 Here's an equivalent, but more computational way to show that

$$\frac{d}{dx} (\ln x) = 1/x.$$

The key is to recognize that the chain rule tells us that if $u = g(x)$ is a differentiable function of x , then

$$\frac{d}{dx} (e^u) = e^u \cdot \frac{du}{dx}.$$

Now to compute the derivative above, we know that since the functions $f(x) = \ln x$ and $g(x) = e^x$ are inverses of one another,

$$e^{\ln x} = x.$$

Then differentiate both sides of this equation with respect to x to get

$$\frac{d}{dx} (e^{\ln x}) = \frac{dx}{dx}.$$

And by the chain rule we can rewrite this equation as

$$(e^{\ln x}) \frac{d}{dx} (\ln x) = 1.$$

And since $e^{\ln x} = x$,

$$\frac{d}{dx} (\ln x) = 1/x.$$

Question 142 Find an equation of the tangent line to the curve $y = \ln x$ at the point $(4, \ln 4)$.

$$y - \ln 4 = \frac{1}{4} (x - 4).$$

Question 143 (a) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln(4x)).$$

(b) Compute the same derivative without using the chain rule.

Explanation. Let

$$y = \ln(4x)$$

and

$$u = 4x.$$

Then

$$y = \ln u$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du} (\ln u) \cdot \frac{d}{dx} (4x) \\ &= \frac{1}{u} \\ &= \frac{1}{4x}. \end{aligned}$$

(b) We can compute the derivative

$$\frac{d}{dx} (\ln(4x))$$

without the chain rule as follows:

$$\begin{aligned} \frac{d}{dx} (\ln(4x)) &= \frac{d}{dx} (\ln 4 + \ln x) \\ &= \frac{d}{dx} (\ln 4) + \frac{d}{dx} (\ln x) \\ &= 0 + \frac{1}{x}. \end{aligned}$$

Question 144 (a) Compute the derivative

$$\frac{d}{dx} (\ln(x^2))$$

both with and without the chain rule. Follow the steps exactly as in the previous example when using the chain rule.

(b) What are the domains of the functions $f(x) = \ln x$ and $g(x) = \ln(x^2)$ and their derivatives.

(c) Graph the function $g(x) = \ln(x^2)$ and its derivative on the same coordinate system by hand.

Question 145 (a) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln |x|).$$

Do this by noting that

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

and using the chain rule to compute

$$\frac{d}{dx} (-x).$$

(b) A reflection about the y -axis takes the graph of a function $y = f(x)$ to the graph of a function $y = g(x)$. Describe the transformation that takes the graph of $y = f'(x)$ to the graph of $y = g'(x)$. Explain your reasoning.

Question 146 (a) Find a function

$$T = f(k), \quad k \geq 1,$$

that expresses the time (in years) it takes an investment to grow by a factor of k . Assume the investment grows at a constant relative instantaneous rate of $i\%/yr$. So, for example if $i = 5$, $f(3)$ would be the time it takes an investment to triple at an relative instantaneous growth rate of $5\%/yr$.

Hint: Let B_0 be the initial investment. Then the value of the investment T years later is

$$kB_0 = B_0 e^{\frac{i}{100}T}.$$

Solve this equation for T to find an expression for the function f . You should get that

$$T = f(k) = (100 \ln k)/i$$

(b) Suppose $i = 5$ and evaluate the derivative

$$\left. \frac{dT}{dk} \right|_{k=3}.$$

(c) What are the units of the above derivative? Explain its meaning in terms of small changes.

I'll leave the units up to you. But the meaning is that it takes an investment (increasing at the relative instantaneous rate of $5\%/yr$) about **0.067** years longer to increase by 201% than it does to triple.

(d) Use the result of parts (b) and (c) to estimate how much longer it would take an investment growing at an relative instantaneous rate of $5\%/yr$ to increase by 210% than it would take to triple. Just use simple arithmetic (ie. multiplication), no calculator.

Question 147 The function

$$G = f(v) = 40 - 0.08 \left(\frac{v}{2} - 25 \right)^2, \quad 25 \leq v \leq 65,$$

expresses the gas mileage of a car (in miles/gallon) in terms of its speed (in miles/hr).

(a) Explain the meaning of the derivative

$$\frac{d}{dv} (\ln(f(v))).$$

Include units in your explanation. Also, what are the units of 25 in the above expression? How do you know?

(b) Evaluate the above derivative at $v = 30$ and explain its meaning in terms of small changes.

Question 148 The function

$$v = g(h), \quad 50 \leq h \leq 200,$$

expresses the speed (in ft/sec) of a hawk in terms of its altitude (in feet) during a portion of its flight.

Suppose that $f(150) = 80$ and

$$\left. \frac{dv}{dh} \right|_{h=150} = -2.$$

(a) What are the units of the above derivative? Explain the meaning of the derivative in the context of small changes in this particular scenario. Be specific.

I'll leave the units up to you, but if we assume the hawk is descending, then its speed would *increase* by about 2 ft/sec as it falls from 150 to 149 feet.

(b) Evaluate the derivative

$$\left. \frac{d}{dh} (\ln(g(h))) \right|_{h=150}.$$

Show all work and explain your reasoning.

The key here is to use the chain rule to evaluate the derivative. So with

$$v = g(h)$$

and

$$w = \ln(g(h)),$$

we have

$$w = \ln v.$$

Then by the chain rule,

$$\begin{aligned} \frac{dw}{dh} &= \frac{dw}{dv} \cdot \frac{dv}{dh} \\ &= \left(\frac{1}{v} \right) \left(\frac{dv}{dh} \right), \end{aligned}$$

and continue from here by evaluating the above expression at $h = 150$.

(c) What are the units of the derivative in part (b)? Explain the meaning of the derivative in the context of small changes in this particular scenario. Be specific by answering this question much like part (a) above (appropriately modified).

The Inverse Sine Function

Let

$$\theta = g(y) = \arcsin y = \sin^{-1}(y).$$

- (a) What is the domain of the function g ?
- (b) What is its range?
- (c) Explain the meaning of $\arcsin y$.
- (d) True or false: The inverse sine function is the inverse of the sine function.
- (e) What function is the inverse of the inverse sine function?

Question 149 The graph of the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

is shown below.

Desmos link: <https://www.desmos.com/calculator/lxwoeir1pt>

Worksheet available at 151: Arc Sine

- (a) Use the graph to estimate the derivative

$$\left. \frac{d}{dy} (\arcsin y) \right|_{y=0.8}.$$

Explain your reasoning.

- (b) Use the graph and the slider u to estimate the derivative

$$\left. \frac{d}{dy} (\arcsin y) \right|_{y=0.8}.$$

- (c) Express the derivative of the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

in terms of y .

- (d) Use part (c) and the ideas in parts (a),(b) to find an expression for the derivative

$$\frac{d}{dy} (\arcsin y).$$

- (e) Check your answer to part (d) by evaluating the derivative at $y = 0.6, 0.8$.
 (f) What is the domain of the derivative in part (d)?

Question 150 The top of a 25-foot long ladder slides down a vertical wall at the constant rate of 4 ft/sec.

(a) Find a function

$$\theta = f(t), \quad 0 \leq t \leq 6.25,$$

that expresses the angle the ladder makes with the ground (measured in radians) in terms of the number of seconds since the ladder was in the vertical position.

(b) Find the rotation rate of the ladder when the top of the ladder is

(i) 15 feet above the ground.

(ii) 24 feet above the ground.

(iii) 24.9 feet above the ground.

(c) Solve this problem again by working directly with the sine function, not the arcsine function.

Desmos link: <https://www.desmos.com/calculator/5c4lssovbi>

Worksheet available at 151: Ladder and ArcSine

Question 151 The bottom end of a 25-foot long ladder slides across a horizontal floor at the constant rate of 4 ft/sec as the top end slides down a vertical wall.

(a) Find a function

$$\theta = g(u), \quad 0 \leq u \leq 25,$$

that expresses the angle the ladder makes with the wall (measured in radians) in terms of the distance (in feet) between the wall and the bottom end of the ladder.

Hint: Let A be the bottom end of the ladder, B the top end, and point O the point on the wall closest to A (as in the demonstration below). Use right triangle trigonometry in $\triangle AOB$.

(b) Use the slider u in the animation below to approximate each of the following derivatives. Include units. Note that the tick marks on the radian protractor are spaced at intervals of 0.1 radians.

(i) $\left. \frac{d\theta}{du} \right|_{u=1}$

(ii) $\left. \frac{d\theta}{du} \right|_{u=15}$

(iii) $\left. \frac{d\theta}{du} \right|_{u=24.9}$

- (c) What do the above derivatives tell you?
- (d) Find an expression for the derivative $d\theta/du$. Use your expression to evaluate the three derivatives in part (b) and compare these with your estimates.
- (e) Find a function

$$\theta = f(t), \quad 0 \leq t \leq 6.25,$$

that expresses the angle the ladder makes with the ground (measured in radians) in terms of the number of seconds since the ladder was in the vertical position.

Hint: Express the distance u (in feet) between the bottom end of the ladder and the wall in terms of t . Then substitute this expression for u in your function from part (a).

- (f) Find an expression for the derivative

$$r = \omega(t) = d\theta/dt.$$

Interpret its meaning. Include units.

- (g) Find the rotation rate of the ladder when the bottom end of the ladder is
- (i) 1 foot from the wall.
 - (ii) 15 feet from the wall.
 - (iii) 24.9 feet above the wall.

- (h) Evaluate the limit

$$\lim_{t \rightarrow 6.25} \omega(t)$$

and interpret its meaning.

Desmos link: <https://www.desmos.com/calculator/egolipj5qg>

Worksheet available at 151: Ladder and ArcSine 2

Exercises

Question 152 (a) Simplify the derivative

$$\frac{d}{d\theta} (\arcsin(\sin \theta))$$

- (b) Use the result of part (a) to graph the function

$$y = \arcsin(\sin \theta).$$

Explain your reasoning.

Question 153 Find the measure of the acute angle that the tangent line to the curve

$$y = f(\theta) = \ln |\sec \theta|$$

at the point $(\pi/7, f(\pi/7))$ makes with the x -axis Do not use a calculator.

Question 154 The function

$$q = f(p) = 0.2(2p - 40)^2, \quad 5 \leq p \leq 12,$$

expresses the average number of burgers sold per day at Five Guys in Edmonds in terms of the price (in dollars/burger).

(a) Evaluate the derivative

$$\left. \frac{d}{dp} (\ln(f(p))) \right|_{p=7.5}$$

(b) What are the units of the derivative above? Explain its meaning.

Question 155 The bottom end of a 25-foot ladder lies 24 feet from the base of a vertical wall. Use the appropriate linear approximation to estimate the angle through which the ladder rotates when the bottom end is pulled an additional 0.1 feet away from the wall along a horizontal floor.

Solve this problem twice, first using an inverse trig function and again without an inverse trig function.

Start by defining your variables, with units.

Use a calculator if need be, but only for arithmetic and not to evaluate any trigonometric functions.

Compare your estimate with the actual angle of rotation.

Question 156 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a 14-foot ladder leans against the trunk, the other rests on the horizontal ground. The bottom end of the ladder is pulled away from the trunk at the constant speed of 4 ft/sec. At what rate is the ladder rotating when the bottom and tops ends are respectively 16 and 10 feet from the base of the trunk?

Hint: Use the law of sines.

Desmos link: <https://www.desmos.com/calculator/rpms2jqfpm>

Desmos activity available at 151: *Tree and Ladder*

Question 157 (a) The animation below shows water draining from a tank. Play the animation and sketch by hand a graph of the function $V = f(t)$ that expresses the depth of the water as a function of time. Explain your reasoning. Label the axes with units and the appropriate variable names.

Access Desmos interactives through the online version of this text at

Desmos link: <https://www.desmos.com/calculator/pdghky6tie>

(b) Torricelli's law says that the rate, say in cm^3/sec , at which water drains out of a small hole in the bottom of a tank is proportional to the square root of the depth of the water. So if $V = f(h)$ is a function that expresses the volume (in cm^3) of water in the tank in terms of the depth (in feet) and $h = g(t)$ is a function that expresses the depth of the water (in feet) in terms of number of seconds past noon, then

$$\frac{dV}{dt} = -k\sqrt{h}$$

for some positive constant k .

For a cylindrical tank of radius r cm,

$$V = f(h) = \pi r^2 h$$

and

$$\frac{dV}{dh} = \pi r^2.$$

So by the chain rule

$$\begin{aligned} \frac{dV}{dt} &= \frac{dV}{dh} \cdot \frac{dh}{dt} \\ &= \pi r^2 \frac{dh}{dt}. \end{aligned}$$

So for the cylindrical tank we can write Torricelli's law as

$$\frac{dV}{dt} = \pi r^2 \frac{dh}{dt} = -k\sqrt{h}.$$

Or equivalently as

$$\frac{dh}{dt} = -\frac{k}{\pi r^2} \sqrt{h} = -k_2 \sqrt{h},$$

where

$$k_2 = \frac{k}{\pi r^2}$$

is a positive constant.

(c) Which of the following functions might express the depth of water in a cylindrical tank (in terms of time) as the water drains out of a small hole in the bottom of the tank? Justify your reasoning.

(i) $h = g(t) = 2(5 - t)^3, 0 \leq t \leq 5$

(ii) $h = g(t) = 2(5 - t)^2, 0 \leq t \leq 5$

Solution:

The key idea is to express the derivative dh/dt as a function of the depth h of the water.

(i) If $h = 2(5 - t)^3$, then

$$\begin{aligned} \frac{dh}{dt} &= \frac{d}{dt} (2(5 - t)^3) \\ &= 2(3)(5 - t)^2 \cdot \frac{d}{dt} (5 - t) \\ &= -6(5 - t)^2. \end{aligned}$$

Now to express dh/dt in terms of h , solve the equation

$$h = 2(5 - t)^3$$

for t to get

$$t = 5 - \left(\frac{h}{2}\right)^{1/3}.$$

Then, substitute this expression for t into the derivative dh/dt :

$$\begin{aligned} \frac{dh}{dt} &= -6(5 - t)^2 \\ &= -6 \left(\frac{h}{2}\right)^{2/3} \\ &= -\left(\frac{6}{2^{2/3}}\right) h^{2/3}. \end{aligned}$$

This tells us that the rate of change in the depth of the water is not proportional to the square root of the water's depth as Torricelli's law requires. So the function

$$h = 2(5 - t)^3, 0 \leq t \leq 5$$

is not a possible depth function for water draining from a cylindrical tank.

Free Response: Give a similar analysis for the depth function of part (ii).

Question 158 This is a continuation of the previous problem.

Now we pour water into a cylindrical tank at a constant rate, while at the same time water leaks out through a small hole in the bottom of the tank. We'll suppose that the tank starts with some initial volume of water.

Free Response: (a) What do you think happens to the water level in the tank initially?

(b) What do you think happens to the water level in the long run?

To model this situation, we need to modify Torricelli's law. For this, let's suppose that we pour water into the cylindrical tank (of radius r) at the constant rate of $k_3 \text{ cm}^3/\text{sec}$. Then with the same notation as before,

$$\frac{dV}{dt} = k_3 - k\sqrt{h}.$$

But since

$$\begin{aligned}\frac{dV}{dt} &= \frac{dV}{dh} \cdot \frac{dh}{dt} \\ &= \pi r^2 \cdot \frac{dh}{dt},\end{aligned}$$

the above modification of Torricelli's law becomes

$$\begin{aligned}\frac{dh}{dt} &= \frac{k_3}{\pi r^2} - \frac{k}{\pi r^2} \sqrt{h} \\ &= k_1 - k_2 \sqrt{h},\end{aligned}$$

where

$$k_1 = \frac{k_3}{\pi r^2}$$

and

$$k_2 = \frac{k}{\pi r^2}$$

are positive constants.

Free Response: (a) What are the units of k_1 and k_2 ? How do you know?

(b) Explain the meaning of k_1 .

The equation

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}$$

expresses the rate of change in the water's depth as a function of the depth. It is called a differential equation and you will learn a little about how to solve equations like this next quarter.

For this particular differential equation, it is not possible to express the depth of the water explicitly as a function of time. But assuming that the depth of the water is h_0 cm at time $t = 0$, it turns out that the function

$$t = g^{-1}(h) = -\frac{2}{k_2} \left(\sqrt{h} - \sqrt{h_0} + \frac{k_1}{k_2} \ln \left| \frac{k_1 - k_2\sqrt{h}}{k_1 - k_2\sqrt{h_0}} \right| \right), \quad t \geq 0,$$

expresses the time (in seconds) in terms of the depth of the water (in cm). We can check that this is indeed correct as follows:

- (a) Show algebraically that the depth of the water at time $t = 0$ is $h = h_0$.
- (b) Use the above expression for $t = g^{-1}(h)$ to compute and then simplify the derivative dt/dh .
- (c) Use the result of part (b) to show that

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}.$$

Desmos link: <https://www.desmos.com/calculator/c78kv7wifv>

Worksheet available at 151: *Draining Cylinder 2*

- (d) Experiment with the sliders above and summarize your observations about how the graph of the function $h = g(t)$ changes depending on the initial depth of the water and the constants k_1 and k_2 .
- (e) Express the equilibrium depth in terms of k_1 and k_2 . Check that your expression has the correct units. The equilibrium depth is the depth at which the water level remains constant. It is also the depth which the water level approaches (independent of the initial depth).
- (f) What happens to the equilibrium depth when k_1 increases (and k_2 is held constant)? When k_2 changes and k_1 is held constant?

The Quotient Rule

An introduction to the quotient rule.

Discussion Questions

Question A: (a) At 9am on May 29, the balance in an account is increasing at the relative rate of 8%/yr. At the same time, the price of a stock is increasing at the rate of 5%/yr. Is the number of shares you can buy with the balance in the account increasing or decreasing at this time? At what relative rate? No computations. Just explain what you think.

(b) How would your answer to part (a) change if instead the stock price were decreasing at the rate of 5%/yr?

Question B: Let $B = f(t)$ and $P = g(t)$ be functions that respectively express the balance (in dollars) in an account and the price (in dollars/share) of a stock in terms of the number of years past 9am on May 29. Let $S = h(t)$ be the number of shares of the stock you can buy with the balance in the account at time t years past 9am on May 29.

(a) Interpret the meanings of the following derivatives. Include units.

(i) $\frac{d}{dt}(\ln(f(t))) = \frac{d}{dt}(\ln B)$

(ii) $\frac{d}{dt}(\ln(g(t))) = \frac{d}{dt}(\ln P)$

(iii) $\frac{d}{dt}(\ln(h(t))) = \frac{d}{dt}(\ln S)$

(b) Express the derivative

$$\frac{d}{dt}(\ln S) = \frac{d}{dt} \left(\ln \left(\frac{B}{P} \right) \right)$$

in terms of the derivatives

$$\frac{d}{dt}(\ln B)$$

and

$$\frac{d}{dt}(\ln P).$$

Learning outcomes:
Author(s):

The Relative Quotient Rule

Had we never learned about relative changes and relative rates of change, the following questions would have almost forced these ideas upon us.

Question 159 (a) Suppose over a six-month period, the national debt of a small country increases from \$4 billion to \$4.16 billion while the population increases from 20 million to 21 million. Does the per-capita (ie. per-person) share of the national debt increase or decrease during this period?

(b) At noon on July 1, the national debt of a small country with a population of 20 million is \$4 billion. At that same instant the population is increasing at the rate of 1 million people/yr while the national debt is increasing at the rate of \$0.16 billion/yr. Is the per-capita share of the national debt increasing or decreasing at this instant? At what rate?

Explanation. (a) The question is really to determine which is greater,

$$c_1 = \frac{4 \times 10^9 \text{ dollars}}{2 \times 10^7 \text{ person}}$$

or

$$c_2 = \frac{4.16 \times 10^9 \text{ dollars}}{2.1 \times 10^7 \text{ person}} = \frac{4 \times 10^9 \times 1.04 \text{ dollars}}{2 \times 10^7 \times 1.05 \text{ person}} = \left(\frac{1.04}{1.05} \right) c_1.$$

And because the total debt increased by 4% while the population increased by 5%, the per-capita share of the national debt decreased during the six-month period.

But the per-capita share of the debt did *not* decrease by 1%, but rather by about (to the nearest thousandth of a percent) **0.952%**.

(b) For the instantaneous rate, let

$$Q = f(t)$$

be the total national debt in billions of dollars at time t years past noon on July 1 and

$$P = g(t)$$

the population in millions at time t . Also, let

$$C = h(t) = \frac{f(t)}{g(t)}$$

be the per-capita share of the debt in thousands of dollars/person at time t years past noon on July 1. We wish to find the value of the derivative

$$\left. \frac{dC}{dt} \right|_{t=0}.$$

Part (a) suggests we express the relative instantaneous rate of change

$$\frac{1}{C} \cdot \frac{dC}{dt} \Big|_{t=0}$$

in terms of the relative rates

$$\frac{1}{Q} \cdot \frac{dQ}{dt} \Big|_{t=0} = 0.04$$

and

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=0} = 0.05$$

Perhaps the easiest way to do this, although not entirely complete, is to assume that the population and national debt each grow exponentially. In that case, the relative growth rates of each are constant and

$$Q = f(t) = 4e^{0.04t}$$

and

$$P = g(t) = 20e^{0.05t}.$$

This tells us that

$$C = \frac{f(t)}{g(t)} = 0.2e^{-0.01t}$$

and

$$\frac{1}{C} \cdot \frac{dC}{dt} \Big|_{t=0} = -0.01$$

So at noon on July 1, we suspect that the per-capita share of the debt is decreasing at the relative rate of 1%/yr and at the absolute rate of \$2/person/yr.

To be sure this is true even if the relative rates of change in the population and total debt are *not* constant, we can be more general and compute the relative instantaneous rate of change in the per-capita share of the national debt as

$$\begin{aligned} \frac{1}{C} \cdot \frac{dC}{dt} &= \frac{d}{dt} \left(\ln \left(\frac{Q}{P} \right) \right) \\ &= \frac{d}{dt} (\ln Q - \ln P) \\ &= \frac{d}{dt} (\ln Q) - \frac{d}{dt} (\ln P) \\ &= \frac{1}{Q} \cdot \frac{dQ}{dt} - \frac{1}{P} \cdot \frac{dP}{dt}. \end{aligned}$$

Theorem 2. (a) (The Relative Quotient Rule) If $Q = f(t)$ and $P = g(t)$ are differentiable functions of t , then if $g(t) \neq 0$,

$$C = \frac{f(t)}{g(t)} = \frac{Q}{P}$$

is a differentiable function of t and

$$\frac{1}{C} \cdot \frac{dC}{dt} = \frac{1}{Q} \cdot \frac{dQ}{dt} - \frac{1}{P} \cdot \frac{dP}{dt}.$$

The relative rate of change in a quotient of two functions is equal to the difference in the relative rates of change in the functions.

(b) (The Quotient Rule) With the same hypotheses (and obtained by multiplying both side of the previous equation by $C = Q/P$),

$$\frac{d}{dt} \left(\frac{Q}{P} \right) = \frac{1}{P} \cdot \frac{dQ}{dt} - \frac{Q}{P^2} \cdot \frac{dP}{dt}.$$

The Tangent Fuction

Question 160 Compute the derivative

$$\frac{d}{d\theta} (\tan \theta)$$

Explanation. We compute this derivative from scratch by letting

$$y = \tan \theta = \frac{\sin \theta}{\cos \theta}.$$

Then

$$\ln |y| = \ln |\sin \theta| - \ln |\cos \theta|$$

and

$$\frac{d}{d\theta} (\ln |y|) = \frac{d}{d\theta} (\ln |\sin \theta| - \ln |\cos \theta|).$$

So

$$\begin{aligned} \frac{1}{y} \cdot \frac{dy}{d\theta} &= \frac{1}{\sin \theta} \cdot \frac{d}{d\theta} (\sin \theta) - \frac{1}{\cos \theta} \cdot \frac{d}{d\theta} (\cos \theta) \\ &= \cot \theta + \tan \theta. \end{aligned}$$

Then mulitplying both sides by $y = \tan \theta$ gives

$$\frac{d}{d\theta} (\tan \theta) = 1 + \tan^2 \theta.$$

Question 161 (a) Use the graph of the function

$$y = f(\theta) = \tan(2\theta)$$

below to estimate the y -coordinates of all points on the curve where the tangent lines are parallel to the lines

(i) $6x - y = 12$.

(ii) $6x + y = 12$.

(b) Find the exact y -coordinates without using a calculator.

Desmos link: <https://www.desmos.com/calculator/obz6ghw3ej>

Desmos activity available at 151: *Tangent Graph*

Question 162 You stand 50 feet from the base of a tree and measure the angle of elevation to the top of the tree with an error of at most $\pm 2^\circ$. You then compute the height of the tree above eye level to be 100 feet.

Desmos link: <https://www.desmos.com/calculator/yjyghsoeog>

Desmos activity available at 151: *Angle of Elevation 1*

(a) Use the demonstration above to approximate your error in computing the tree's height.

(b) Use the appropriate linear approximation to estimate your error in computing the tree's height. Compare this with your estimate from part (a). Do not use a calculator. Do this as follows:

- Find a function

$$h = f(\theta), \quad 0 < \theta < \pi/2,$$

that expresses the computed height of the tree above eye level (measured in feet) in terms of the measured angle of elevation (in radians). Draw a picture to help with your explanation.

This function is

$$h = f(\theta) = 50 \tan \theta, \quad 0 < \theta < \pi/2.$$

- Next find an expression for the derivative $dh/d\theta$ and evaluate the derivative

$$\left. \frac{dh}{d\theta} \right|_{h=100}.$$

- Now we'll take the exact height of the tree above eye level to be 100 feet and let

$$\Delta h = h - 100$$

be the error in the computed height and

$$\Delta \theta = \theta - \arctan(2)$$

be the error in the measured angle.

Then if $\Delta \theta \sim 0$,

$$\left. \frac{dh}{d\theta} \right|_{h=100} \sim \frac{\Delta h}{\Delta \theta}$$

and so

$$\Delta h \sim 250 (\Delta \theta).$$

- I'll let you continue from here.

The Inverse Tangent Function

Question 163 (a) Explain the meaning of the function

$$\theta = f(y) = \arctan y.$$

In particular, what does it take as an input and what does it return as an output? Include the domain and range of the function as part of your explanation.

(b) Express the derivative

$$\frac{d}{d\theta} (\tan \theta) = 1 + \tan^2 \theta$$

of the function

$$y = \tan \theta$$

in terms of y .

(c) Use part (b) to find an expression for the derivative

$$\frac{d\theta}{dy} = \frac{d}{dy} (\arctan y) = \frac{d}{dy} (\tan^{-1} y).$$

(d) Evaluate the derivatives

$$\left. \frac{d}{d\theta} (\tan \theta) \right|_{\theta=\pi/4}$$

and

$$\frac{d}{dx} (\arctan x) \Big|_{x=1}.$$

Comments?

Question 164 You stand 60 feet from the base of a tree and measure the angle of elevation to the top of an 85-foot tall tree. You then move 4 feet closer to the tree and measure the same angle. Assume your eyes are five feet above the ground.

Desmos link: <https://www.desmos.com/calculator/qqgvq3noah>

Desmos activity available at 151: Tree 3

(a) Drag the slider u in the demonstration above to estimate the change in the angle of elevation after you move 4 feet closer to the tree. Note that the angles of elevation are $\angle TEB$ and $\angle TFB$ and that $\angle ETF$ measures their difference. Consecutive tick marks on the protractor are spaced at intervals of 0.01 radians.

(b) Use derivatives to approximate the change in the angle of elevation and compare your approximation with your estimate in part (a) and with the actual change. Use a calculator with only addition, subtraction, multiplication, and division.

Go about this by first finding a function

$$\theta = f(s), s > 0,$$

that expresses the angle of elevation to the top of the tree (measured in radians) in terms of your distance from the tree (measured in feet). Use the arctangent function in your expression. Do not use the inverse cotangent function.

The function is

$$\theta = f(s) = \arctan(80/s), s > 0.$$

Then continue in a manner similar to Question 4.

Two Motions

Question 165 (a) Play the slider s in the demonstration below to show the motion of a beetle crawling along the y -axis as it leaves behind tracks spaced at equal time intervals.

(b) Use the animation to sketch (by hand) a graph of the function

$$s = f(t)$$

that expresses the position (in this case the y -coordinate) of the beetle as a function of time. Label the axes with the appropriate variable names and units. Then activate the folder in Line 2 to see how you did.

(c) Use the animation to sketch (by hand) a graph of the function

$$v = g(t)$$

that expresses the beetle's velocity (in this case the rate of change, with respect to time, in the beetle's y -coordinate). Label the axes with the appropriate variable names and units. Then activate the folder in Line 7 to see how you did.

Desmos link: <https://www.desmos.com/calculator/srotstrdzm>

Desmos activity available at 151: *Tangent Motion*

Question 166 Repeat parts (a)-(c) of the previous question for the motion below.

Desmos link: <https://www.desmos.com/calculator/h6vq21lfql>

Desmos activity available at 151: *ArcTangent Motion*

Exercises

Question 167 (a) Use the website below to compute an accurate estimate of the current rate (in dollars/year) at which the U.S. national debt is changing. Explain your method.

National Debt Clock

(b) Use the website below to compute an accurate estimate (in people/yr) at which the U.S. population is currently increasing.

Population Clock

(c) Use your estimates above and the current national debt and U.S. population to compute

(i) the current rate of change in the per-capita share of the national debt.

(ii) the current relative rate of change in the per-capita share of the national debt.

Include units in every number in each step of your computations.

Question 168 The function

$$C = f(t) = At^4e^{-kt}, t \geq 0,$$

expresses the concentration of a drug (measured in mg/L) in the bloodstream in terms of the number of hours since the drug was injected. Here A and k are positive constants.

Desmos link: <https://www.desmos.com/calculator/lvsxu3wa9a>

Desmos activity available at 151: [Drug Concentration](#)

- (a) What are the units of the constant A ? How do you know?
- (b) What are the units of k ? How do you know?
- (c) Use the graph of $C = f(t)$ above to sketch by hand a graph of the derivative $r = dC/dt$. Be sure to label the axes with the appropriate variable names and units. Do not use technology. Explain your reasoning.
- (d) Use calculus and algebra to find an expression in terms of k for the time when the concentration is a maximum. Work in general. Do not use a specific value of k . Check your work by following the directions, Lines 4 and 5, in the desmos demonstration. Label also the coordinates of the corresponding point on your hand-drawn graph of the derivative.
- (e) Find expressions (in terms of k) for the times when the concentration is increasing and decreasing at the maximum rates. Check that your expressions have the correct units. Work in general. Do not use a specific value of k . Check your work by following the directions, Lines 6-9, in the desmos demonstration. Label also the coordinates of the corresponding points on your hand-drawn graph of the derivative.
- (f) Find an expression (in terms of k) for the relative rate at which the concentration is changing t hours after the injection. Check that your expression has the correct units. Work in general. Do not use a specific value of k .
- (g) Suppose the concentration is at its maximum five hours after injection and determine when the concentration is increasing at the rate of 50%/hr. Determine also when the concentration is decreasing at the rate 50%/hr.
- (h) Observations?

Question 169 The function

$$P = f(t) = 5 - 3t + t^2, 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

The Quotient Rule

- (a) Use the graphs of the function $P = f(t)$ and the function $r = f'(t)/f(t)$ to estimate when the stock price is increasing at the greatest relative rate.
- (b) Use algebra to find the exact time when the stock price is increasing at the greatest relative rate.

Hint: What is the value of the derivative dr/dt at this time? But start by finding an expression for the instantaneous relative rate of change in the stock price.

Desmos link: <https://www.desmos.com/calculator/xuupp3srqv>

Desmos activity available at 151: Stock Price 4

Question 170 You jog once around a circular track of radius r meters at the constant speed of v m/sec. A flagpole lies b meters due east of the track's center.

- (a) Find a function

$$s = f(t), 0 \leq t \leq 2\pi r/v,$$

that expresses your distance (in meters) to the flagpole in terms of the time (measured in seconds) since you started running. Assume you start at the point A on the track nearest the flagpole. Explain your reasoning. Work with the general parameters r , v , and b , not with any specific values for these parameters.

- (b) Find an expression for the time when your distance to the flagpole is increasing at the greatest rate. Try to give a geometric interpretation of your position at this time.

Desmos link: <https://www.desmos.com/calculator/bxofhvfbs>

Demonstration available at Math 151: Jogger 3

Implicit Differentiation

An introduction to implicit differentiation.

Discussion Questions

Question 171 Which of the following equations define y implicitly as a function of x in a sufficiently small neighborhood of the given point? Supplement your reasoning with a graph of each relation.

- (a) $x^2 + y^2 = 25$ near the point $(4, -3)$
- (b) $x^2 + y^2 = 25$ near the point $(0, -5)$
- (c) $x^2 + y^2 = 25$ near the point $(-5, 0)$
- (d) $x^2 - xy + y^2 = 1$ near the point $(1, 1)$.
- (e) $x^2 - xy + y^2 = 3$ near the point $(1, 2)$.
- (f) $x^2 - xy + y^2 = 3$ near the point $(2, 1)$.

Introduction to Implicit Differentiation

Question 172 Let $p = f(t)$ be a differentiable function of t . Find expressions for each of the following derivatives, first supposing that

$$p = f(t) = t^3 + 1,$$

and then more generally, not assuming any particular expression for the function f .

- (a) $\frac{d}{dp} (p^5)$
- (b) $\frac{d}{dt} (p^5)$
- (c) $\frac{d}{dt} (e^{2p})$
- (d) $\frac{d}{dt} (t^4 \sin(p))$

Learning outcomes:
Author(s):

Question 173 (a) Find equations of the tangent lines to the ellipse

$$x^2 - xy + 2y^2 = 9$$

at the points where the ellipse intersects the x -axis.

(b) Find equations of all horizontal tangent lines to the ellipse.

(c) Find equations of all vertical tangent lines.

Desmos link: <https://www.desmos.com/calculator/ox3wvfitqk>

Desmos activity available at 151: *Implicit Ellipse*

Question 174 (a) Find an equation of the tangent line to the curve

$$4xe^y - 3y \cos(2xy) = 0$$

at the origin.

(b) Enter your equation in Line 5 of the worksheet below to check your answer.

Desmos link: <https://www.desmos.com/calculator/1qprxg226m>

Desmos activity available at 151: *Implicit 1*

Astroids

Question 175 (a) The coordinate axes cut out a segment from the tangent line to the curve

$$x^{2/3} + y^{2/3} = 5$$

at the point $P(8, 1)$. Find the length of that segment. Do this by using implicit differentiation to help find an equation of the tangent line at P and go from there. But first show that P lies on the curve.

(b) As a bug crawls around the above curve and passes the point $P(8, 1)$, it is moving away from the x -axis at the rate of 4 cm/sec. Is the bug moving toward or away from the y -axis at this time? At what rate?

(c) Find the slope of the tangent line in part (a) without using implicit differentiation.

Question 176 (a) Use implicit differentiation to show that segments cut by the coordinate axes from the tangent lines to the astroid

$$x^{2/3} + y^{2/3} = a^{2/3}$$

all have the same length. Here $a > 0$ is a constant.

(b) Prove the same result by using trigonometric functions to parameterize the astroid.

Desmos link: <https://www.desmos.com/calculator/vrythrvjuc>

Desmos activity available at 151: Astroid

Ellipses and Related Curves

Question 177 (a) Use implicit differentiation to find an equation of the tangent line to the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

at the point P with coordinates (x_0, y_0) . Here a, b are positive constants.

(b) Input your equation from part (a) on Line 3 of the desmos worksheet below. Drag point P to check your equation is correct.

(c) Solve part (c) again without calculus by considering a composition of transformations that takes the circle

$$x^2 + y^2 = 1$$

to the ellipse in part (a).

Desmos link: <https://www.desmos.com/calculator/t1e9v7ncpv>

Desmos activity available at 151: Tangents to Ellipse

Question 178 An ellipse through the point $P(0, b)$ has focal points F_1 at the origin and F_2 at the point $A(a, 0)$.

(a) Use the definition of an ellipse as the set of points whose distances to the foci have a constant sum to find an equation of the ellipse.

(b) Use implicit differentiation to find the slope of the tangent line to the ellipse at P .

- (c) Find an equation of the line normal to the ellipse at Q .
- (d) Find the coordinates of the point Q where the normal line intersects the x -axis.
- (e) Express the ratio $F_1Q : QF_2$ in terms of a and b . Interpret the ratio geometrically.

The ratio is

$$\frac{F_1Q}{QF_2} = \frac{b}{\sqrt{a^2 + b^2}}.$$

- (f) Check your work in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/xqywotxxf9>

Desmos activity available at 151: Normal to an Ellipse

Question 179 Let F_1 and F_2 be respectively the origin and the point with coordinates $(a, 0)$. The curve \mathcal{C} passes through the point $P(0, b)$. The curve is defined by the property that the sum of the distance from a point Q of \mathcal{C} to F_1 and k times its distance to F_2 is a constant.

- (a) Find an equation of the curve \mathcal{C} .
- (b) Find an equation of the tangent line to \mathcal{C} at P .
- (c) Find an equation of the line normal to \mathcal{C} at P .
- (d) Find the coordinates of the point Q where the normal line intersects the x -axis.
- (e) Express the ratio $F_1Q : QF_2$ in terms of a and b . Interpret the ratio geometrically.

The ratio is

$$\frac{F_1Q}{QF_2} = \frac{bk}{\sqrt{a^2 + b^2}}.$$

- (f) Check your work in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/uz7w5uh0v1>

Desmos activity available at 151: Generalized Ellipse

The Sliding Ladder

Question 180 You walk along a straight path, moving either directly toward or directly away from a tree at the end of the path.

Let $s = f(t)$ be a function that expresses your distance to the tree in terms of the number of minutes past noon and suppose that

$$\left. \frac{ds}{dt} \right|_{t=5} = -240.$$

- (a) Interpret the meaning of the above derivative.
- (b) What is your speed at 12:05pm?
- (c) Find an expression for your speed at time t minutes past noon.

Question 181 A tree leans precariously with its trunk making an angle of ϕ radians with the ground. One end of a ladder with length L feet leans against the trunk, the other rests on the horizontal ground. You slide the bottom end of the ladder along the ground at random.

- (a) Let s and t be the respective distances (measured in feet) from the base of the trunk to the bottom and top ends of the ladder. Find an equation that relates these distances. Use the parameters L and ϕ , not their particular values in the demonstration below.
- (b) Let u be time measured in seconds and suppose that s and t are (unknown) functions of u . Express the derivative dt/du in terms of the derivative ds/du . Interpret the meanings of these derivatives.
- (c) Switch s and t in your equation from part (b). What do you notice? Explain why.
- (d) Use part (b) to express the speed of the top end of the ladder in terms of the speed of its bottom end.
- (e) Suppose $\phi = \pi/3$ and that at some instant the bottom end of the ladder is three times as far from the trunk's base as the top end. What can you say about the speeds of the ends of the ladder at this instant?
- (f) Still supposing $\phi = \pi/3$, find all possible angles the ladder makes with the ground when the top end of the ladder is moving four times as fast as the bottom end. Characterize each angle in one of two ways:
 - (i) One end of the ladder is moving toward the trunk's base and the other end away from the base.
 - (ii) Both ends of the ladder are either moving toward or away from the trunk's base.

Give exact angles. Then use a calculator to approximate their radian measures to the nearest hundredth.

(g) Use the animation below to check that your answers to parts (e) and (f) are reasonable. The graph is of the relation

Desmos link: <https://www.desmos.com/calculator/58jqjo0inl>

Desmos activity available at 151: *Tree and Ladder*

Question 182 A tree leans precariously with its trunk making an angle of ϕ radians with the ground. One end of a ladder with length L feet leans against the trunk, the other rests on the horizontal ground. You slide the bottom end of the ladder along the ground at random.

Suppose at some instant the ladder makes an angle of β_0 radians with the trunk and that the bottom end is s_0 feet from the trunk's base. Suppose also that the ladder is rotating at the rate of ω_0 rad/sec at this same moment.

- (a) Find an expression for rate of change in the distance from the bottom end of the ladder to the base of the trunk at this time.
- (b) Find an expression for rate of change in the distance from the top end of the ladder to the base of the trunk at this time.
- (c) Find expressions for the speeds of the ends of the ladder at this time.

Hint: Use the law of sines in $\triangle BOT$ in the demonstration from the previous question.

Question 183 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere.

Geogebra link: <https://www.geogebra.org/m/vnhrutwu>

Geogebra activity available at 151: *Declination of Sun 2*

(a) Use the graph of the function f below (at latitude $\phi \sim 1.1$) to sketch a graph of the derivative

$$\frac{dh}{d\delta} = f'(\delta).$$

(b) Suppose the latitude ϕ is held constant and find an expression for the derivative

$$\frac{dh}{d\delta} = \frac{d}{d\delta} (-\tan \phi \tan \delta).$$

(c) What are the units of the derivative in part (b)?

(d) Input your expression for the derivative in Line 4 of the worksheet below (follow the directions there). Then vary the slider ϕ to see how the function f and its derivative vary with latitude. Summarize your observations.

(e) Find an expression for the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0} = \left. \frac{d}{d\delta} (-\tan \phi \tan \delta) \right|_{\delta=0}$$

in terms of the latitude ϕ .

(f) Evaluate the derivative in part (e) at a latitude of $\phi = \pi/4$. Interpret its meaning in terms of small changes.

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at [151: Length of Day 1](#)

(g) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

Find an expression for a function

$$\delta = k(t), \quad t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

(h) Find a function

$$r = g(\phi), \quad -\pi/2 < \phi < \pi,$$

that expresses the rate of change in the number of minutes of daylight per day (measured in (minutes of daylight/day)/day) on the spring equinox in terms of the latitude ϕ . Input this function in Line 1 of the demonstration below.

(i) Use the result of part (f) to approximate the rate in (hours of daylight/-day)/day at which the number of daylight hours per day is changing at a latitude of $\phi = \pi/4$ radians on the spring equinox and on the fall equinox.

(j) Evaluate the rates from part (i) in Fairbanks, Alaska, latitude 64.8°N .

Desmos link: <https://www.desmos.com/calculator/nf8n5uphhl>

Desmos activity available at 151: *Length of Day 2*

Waves

Question 184 The function

$$y = f(x, t) = a \sin(kx - \omega t), t \geq 0, \quad (4)$$

describes a wave on a string. The functions expresses the displacement (in meters) of a point on the string in terms of the position x (in meters) of the point and time t , measured in seconds since the motion began.

(a) Experiment with the sliders k , ω , in the demonstration below, playing the slider u (u is just another name for t). Summarize your observations. In particular, be sure to turn off Line 1 to be better able to see the motion of the individual points of the string.

(b) What are the units of k and ω ? How do you know?

(c) Find an expression for the wavelength λ in terms of k , ω .

(d) Find an expression for the period of oscillation T in terms of k , ω .

(e) Hold y constant and differentiate each side of equation (4) with respect to t to find an expression for the speed of the wave. Turn on the graph in Line 9. Explain the logic behind the computation.

Desmos link: <https://www.desmos.com/calculator/9xmkg9hwi>

Desmos activity available at 151: *Traveling Wave 1*

The Ladder and the Tree

Implicit differentiation.

Question 185 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground. We analyze how a small change in the distance between the bottom of the ladder and the base of the trunk changes the distance between the top of the ladder and the base of the trunk.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/u8bxskxyhl>

151: Ladder and Tree 20

We'll let u be the distance OT between the top of the ladder and the base of the trunk (measured in feet) and s the distance OB between the bottom of the ladder and the base of the trunk (also measured in feet).

We'll focus on the particular position of the ladder when the bottom end is 5 feet from the trunk's base and the top end is 8 feet from the base as illustrated above.

- (a) Use the slider s approximate the derivative

$$\left. \frac{ds}{du} \right|_{(s,u)=(5,8)}.$$

Be sure to include units.

- (b) Find the length of the ladder
- (c) Find an equation that relates s and u . Then activate the Folder in Line 20. Explain the significance of the curve and the slope of the tangent line to the curve at P .
- (d) Find an expression for the derivative ds/du .
- (e) Use your expression for ds/du to evaluate the derivative

$$\left. \frac{ds}{du} \right|_{(s,u)=(5,8)}.$$

Learning outcomes:
Author(s):

- (f) Interpret the meaning of the derivative using the language of small changes.
- (g) Use the slider s to approximate the distance $s = OB$ when moving the top of the ladder a small distance away from the base of the tree makes the bottom end move twice as far toward the base. Then use calculus and algebra to approximate that distance.
- (h) Activate the folder in Line 31 of the worksheet above. Let θ be the angle (measured in radians) the ladder makes with the ground as marked. Take $\theta = 2\pi/3$ when the ladder lies along the tree.
- (i) Use the slider s to approximate the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=5}.$$

Include units.

- (ii) Write an equation relating θ and s .
- (iii) Find an expression for the derivative $ds/d\theta$ and use it to compute the exact value of the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=5}.$$

- (iv) Interpret the meaning of the above derivative using the language of small changes.
- (v) Find the exact value of the derivative

$$\left. \frac{d\theta}{du} \right|_{s=5}.$$

The Ladder and the Tree, Part 2

Implicit differentiation.

Question 186 The bottom end of a seven-foot ladder slides across a horizontal floor as its top end slides down a vertical wall.

- (a) Suppose the bottom of the ladder is sliding toward the wall at a speed of 5 ft/sec when the bottom is 3 feet from the wall. At what rate is the ladder rotating at this instant?
- (b) Drag the slider s in Line 1 of the worksheet below to approximate the angle the ladder makes with the ground when its bottom end is moving twice as fast as its top end.
- (c) Find the exact angle in part (b).

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/mlulonghoa>

151: Ladder and Tree 23

Question 187 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground.

At the moment the bottom end of the ladder is 8 feet from the tree's base (in the position shown below), its bottom end is moving toward the tree's base at a speed of 5 ft/sec. At this same moment the top end of the ladder is 3 feet from the tree's base.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/oftz4vb9qj>

151: Ladder and Tree 22

Learning outcomes:
Author(s):

- (a) Is the ladder rotating clockwise or counterclockwise at this moment?
- (b) At what rate? Do not assume that the bottom end moves at a constant speed. Solve this problem twice. First, with implicit differentiation and then again without. Do not use a calculator.

Question 188 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a seven-foot ladder leans against the trunk, the other rests on the horizontal ground.

- (a) Find all possible angles the ladder makes with the ground when its bottom end is moving twice as fast as its top end.
- (b) Drag the slider ϕ_2 in Line 2 of the worksheet below to check your answer.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/oftz4vb9qj>

151: Ladder and Tree 22

Question 189 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/bscbpblji1>

151: Ladder and Tree 21

- (a) Find a function

$$v_B = f(\theta), 0 \leq \theta \leq 2\pi/3,$$

that expresses the speed of the ladder's bottom end in terms of the angle $\theta = \angle OTP$ the ladder makes with the tree and the ladder's rotation rate ω (measured in rad/sec). Do not assume the rotation rate is constant.

- (b) Check that your expression for the speed has the correct units.

The Ladder and the Tree, Part 2

- (c) Similarly find a function

$$v_T = g(\theta), 0 \leq \theta \leq 2\pi/3,$$

that expresses the speed of the ladder's top end in terms of θ and ω .

- (d) Play the slider in Line 2 of the worksheet above. Then use the animation to sketch by hand graphs of the speed functions in parts (a) and (c) assuming the ladder rotates at a constant rate.
- (e) Activate the speed functions folder in Line 41 to check your graphs.
- (f) Find the ratio v_B/v_T of the speeds when the bottom end of the ladder is five feet from the tree's base.
- (g) What angle does the ladder make with the tree when the ladder's bottom end is moving twice as fast as its top end? Find all possibilities.
- (h) Answer part (g) when the tree is perpendicular to the ground instead.
-

More Related Rates Problems

Related rates.

Directions:

- Start by defining all variables, each in a complete sentence with units. Be sure to precisely define the variable for time as well.
- Answer each problem with a concluding sentence.

1 Kite Flying

Question 190 A kite drifts due east as it maintains a constant altitude. Play the slider u in Line 1 of the worksheet below to watch the motion.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/qdrzdt4erc>

151: Kite

- Which do you think is greater, the rate at which the string is being let out or the speed of the kite? No computations, just go with your intuition.
- Express the speed of the kite (not assumed constant) in terms of the rate at which the string is being let out (not assumed constant) and the angle θ the string makes with the horizontal.
- What can you say if $\theta = \pi/3$?
- Was your intuition correct?

Learning outcomes:
Author(s):

2 A Crawling Beetle

Question 191 As a beetle crawling in the xy -plane passes the point $(-4, -3)$ (coordinates in centimeters) its distance from the x -axis is increasing at the rate of 2 cm/sec and its distance from the origin is increasing at the rate of 3 cm/sec.

Is the beetle rotating clockwise or counterclockwise at this instant? At what rate?

3 A Mechanical Motion

Question 192 As a rod OP of length R meters rotates about its endpoint O , the other endpoint P drags along a second rod PA of length L meters as illustrated below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/r20xwnok5r>

151: Engine

- Express the speed of A in terms of R , L , the angle θ shown above, and the rotation rate ω (measured in radians/sec) of rod OP . Do not assume ω is constant. Work in general, not with the specific values of L and R in the worksheet.
- Check that your expression in part (a) has the correct units.
- Express the signed rotation rate of PA in terms of the same parameters. Measure the rate to be positive when PA rotates counterclockwise. Do not assume ω is constant.
- Check that your expression in part (c) has the correct units.

4 Tracking a Helicopter

Sensors on the ground four hundred feet apart track a helicopter. At the instant shown below, the sensors at A and B are rotating counterclockwise at the respective rates of 2 rad/min and 3 rad/min. And the marked angles at A and B have respective measures $\angle A = \pi/6$ and $\angle B = \pi/3$.

Is the helicopter ascending or descending at this instant? At what rate?

Desmos link: <https://www.desmos.com/calculator/xl8t3toppg>

151: Tracking a Helicopter

5 Distance to the Horizon

Question 193 On a clear day with an unobstructed view (like you might have at the beach or in a hot air balloon), the distance to the horizon is limited by the curvature of the earth as illustrated in the demonstration below. This distance is the length of the (red) arc AT on the earth's surface.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

Desmos activity available at

151:Distance to Horizon 1

Now suppose at some instant you are h miles above the earth and descending at the rate of v miles/hour.

Find an expression for the rate (in miles/hour) at which your distance to the horizon (the length of the red arc AT above) is changing at this instant. Your expression should be in terms of h , v , and the radius R of the earth (measured in miles). Do not assume v is constant.

6 Jar Lid and Rubber Band

Question 194 You wrap a rubber band around a circular jar lid and pull the band tight as illustrated below.

Desmos link: <https://www.desmos.com/calculator/jbku3wrtdq>

151: String and Jar Lid

Suppose at some instant you are pulling the point P of the rubber band directly away from the center of the lid at a speed of v inches/sec. Express the rate (in inches/sec) at which the length of the rubber band is changing at this instant in terms of v and the marked angle θ between the straight segments of the band at P .

Hint: Express the length of the rubber band in terms of θ and the distance h (measured in inches) between P and the center of the lid. Keep in mind the band has a curved section in addition to the straight parts.

7 Disco Dancing

Question 195 A spotlight in a circular dance hall of radius r meters is located b meters from the center of the hall.

At some instant the light is rotating at the rate of ω rad/sec.

Desmos link: <https://www.desmos.com/calculator/m2o267u9ur>

151: Disco Dancing

- (a) Find an expression for the speed of the light beam as it moves along the wall at this instant. Your expression should be in terms of ω , r , b , and the angle θ marked below.
 - (b) Check that your expression in part (a) has the correct units.
-

The Natural Log Function

Log functions and their derivatives.

1 The Derivative of the Natural Log Function

Question 196 (a) *Explain the meaning of*

$$\ln 5 = \log_e 5.$$

(b) *Simplify the function*

$$f(x) = e^{\ln x}.$$

Include the appropriate domain.

(c) *Use the result of part (b) to find an expression for the derivative*

$$\frac{d}{dx}(\ln x).$$

Explanation. To compute the derivative in part (c), we know that since the functions $f(x) = \ln x$ and $g(x) = e^x$ are inverses of one another,

$$e^{\ln x} = x.$$

Then differentiate both sides of this equation with respect to x to get

$$\frac{d}{dx}(e^{\ln x}) = \frac{dx}{dx}.$$

And by the chain rule we can rewrite this equation as

$$(e^{\ln x}) \frac{d}{dx}(\ln x) = 1.$$

And since $e^{\ln x} = x$,

$$\frac{d}{dx}(\ln x) = 1/x.$$

Learning outcomes:
Author(s):

Question 197 Find simplified expressions for each of the following derivatives. Include an appropriate domain for each.

(a)

$$\frac{d}{dx} (\ln(x/2))$$

(b)

$$\frac{d}{dx} (\ln(x^2))$$

Question 198 Find an equation of the tangent line to the curve

$$y = 3 \ln \left(\frac{x^2 + 1}{x^3 + 1} \right)$$

at the point on the curve with x -coordinate $x = 1$.

Question 199 (a) Graph the function

$$y = f(x) = \ln|x|$$

by hand.

(b) Use the graph above to sketch (by hand) a graph of the derivative

$$y = \frac{d}{dx} (\ln|x|).$$

(c) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln|x|).$$

Do this by noting that

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

and using the chain rule to compute

$$\frac{d}{dx} (-x).$$

- (d) A reflection about the y -axis takes the graph of a function $y = f(x)$ to the graph of a function $y = g(x)$. Describe the transformation that takes the graph of $y = f'(x)$ to the graph of $y = g'(x)$. Explain your reasoning.

Question 200 Find an equation of the tangent line to the curve

$$y = 3 \ln \left| \frac{x^2 + 1}{x^3 - 1} \right|$$

at the point on the curve with x -coordinate $x = -1$.

Question 201 Let \mathcal{L} be the tangent line to the curve

$$y = \ln |\sec \theta|, \quad -\pi/2 < \theta < \pi/2,$$

at the point on the curve with $\theta = \pi/7$.

Find the acute angle \mathcal{L} makes with the x -axis.

2 Relative Rates of Change

Question 202 The function

$$G = f(v) = 40 - 0.08 \left(\frac{v}{2} - 25 \right)^2, \quad 25 \leq v \leq 65,$$

expresses the gas mileage of a car (in miles/gallon) in terms of its speed (in miles/hr).

- (a) Explain the meaning of the derivative

$$\frac{d}{dv} (\ln(f(v))).$$

Include units in your explanation. Also, what are the units of 25 in the above expression? How do you know?

- (b) Evaluate the above derivative at $v = 30$ and explain its meaning in terms of small changes.

Question 203 At a certain instant the population of a country is decreasing at the rate of 3%/yr. At this same instant the country's national debt is increasing at the rate of 5%/yr.

The Natural Log Function

- (a) *At what relative rate is the per capita (ie. per person) share of the national debt changing at this time? Hint: Use the idea of the previous question.*
 - (b) *If the per capita share of the national debt is \$200,000/person at this time, at what (absolute) rate is the per-capita share of the debt changing at this instant?*
-

Derivatives and the Shapes of Curves

First and second derivatives and their implications.

Optimization

Optimization.

Question 204 Determine the minimum and maximum values of the function

$$y = f(x) = x^3 - 9x, -3 \leq x \leq 0.$$

Question 205 A vertical wall b feet high runs parallel to a tall building. The wall is a feet from the building. A ladder reaching from the ground to the building rests on the top of the wall as shown below.

Desmos link: <https://www.desmos.com/calculator/4ak46ub8ay>

151: Shortest Ladder

- (a) Find a function that express the length of the ladder (measured in feet) in terms of the angle the ladder makes with the ground. Include an appropriate domain.
- (b) Use part (a) to express the length of the shortest such ladder in terms of a and b . Justify your assertion.
- (c) Check that your expression has the correct units.

Work in general and not with the particular values of a and b in the worksheet above.

Question 206 The bottom and top edges of a painting are respectively a and b feet above eye level.

Desmos link: <https://www.desmos.com/calculator/dkqndsegod>

151: Viewing Angle

Learning outcomes:
Author(s):

Optimization

- (a) *Find a function that expresses the viewing angle in terms of your distance from the painting. Include an appropriate domain.*
 - (b) *How far from the painting should you stand to maximize the viewing angle marked above? Justify your assertion.*
 - (c) *Check that your expression has the correct units.*
-

Hours of Daylight

Modeling the number of hours of daylight.

Question 207 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere. The graph of the function $h = f(\delta)$ for latitude $\phi \sim 1.1$ is shown below.

Don't be confused by the angle δ (delta). It's just the lower case Greek letter for Δ .

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at 151: Length of Day 1

- (a) Use the graph to approximate the number of hours of daylight/day at a latitude of $\phi = 1.1$ radians on March 1, when the sun is about 8° below the plane of the equator. What about at our latitude? At the equator?

- (b) Evaluate the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0}.$$

- (c) What are the units of the derivative above? Interpret its meaning.
- (d) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

- (i) Find an expression for a function

$$\delta = k(t), \quad t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

Learning outcomes:
Author(s):

- (ii) Evaluate the derivative

$$\left. \frac{d}{dt} (f(k(t))) \right|_{t=0} = \left. \frac{dh}{dt} \right|_{t=0}.$$

- (iii) What are the units of the derivative above? Interpret its meaning.

Question 208 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere.

Geogebra link: <https://www.geogebra.org/m/vnhrutwu>

Geogebra activity available at 151: Declination of Sun 2

- (a) Use the graph of the function f below (at latitude $\phi \sim 1.1$) to sketch a graph of the derivative

$$\frac{dh}{d\delta} = f'(\delta).$$

- (b) Suppose the latitude ϕ is held constant and find an expression for the derivative

$$\frac{dh}{d\delta}$$

- (c) What are the units of the derivative in part (b)?
- (d) Input your expression for the derivative in Line 4 of the worksheet below (follow the directions there). Then vary the slider ϕ to see how the function f and its derivative vary with latitude. Summarize your observations.
- (e) Find an expression for the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0}$$

in terms of the latitude ϕ .

- (f) Evaluate the derivative in part (e) at a latitude of $\phi = \pi/4$. Interpret its meaning in terms of small changes.

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at 151: Length of Day 1

- (g) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

Find an expression for a function

$$\delta = k(t), t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

- (h) Find a function

$$r = g(\phi), -\pi/2 < \phi < \pi,$$

that expresses the rate of change in the number of hours of daylight per day (measured in (hours of daylight/day)/day)) on the spring equinox in terms of the latitude ϕ . Input this function in Line 1 of the demonstration below.

- (i) On the spring equinox, all latitudes receive 12 hours of daylight/day. Use the result of part (h) to approximate how many extra minutes of daylight we would get on the following day living in

(i) Shoreline, latitude $47.75^\circ N$

(ii) Fairbanks, Alaska, latitude $64.8^\circ N$

Desmos link: <https://www.desmos.com/calculator/nf8n5uphhl>

Desmos activity available at 151: Length of Day 2

Motion

An introduction to motion.

Question 209 Play the slider u in the activity below to see the motion of a balloon. Use the animation to sketch graphs of

- (a) the altitude of the balloon as a function of time.
- (b) the balloon's rate of ascent as a function of time.
- (c) Activate the folders in Lines 13 and 18 to see how you did.

Desmos link: <https://www.desmos.com/calculator/amv52b9ljt>

Desmos activity available at 151: Balloon

Question 210 The function

$$h = f(t) = 10 - \frac{1}{2}t^2e^{-t/5}, \quad 0 \leq t \leq 40,$$

expresses the altitude of a balloon (in thousands of feet) in terms of the number of hours since noon on August 31, 2023.

Use the graph of this function in Question 1 to first approximate answers to the following questions. Then use calculus and algebra to determine the exact times.

- (a) When is the balloon at its minimum height? At its maximum height?
- (b) When is the balloon ascending at the fastest rate? Descending at the fastest rate?

To compute these times you will end up solving a quadratic equation that may be written in the form

$$t^2 - 20t + 50 = 0.$$

So the balloon is ascending at its fastest rate at time (give the exact times and then approximations to the nearest hundredth of an hour)

$$t = 10 + \sqrt{50} \sim 17.07$$

Learning outcomes:
Author(s):

hours past noon and descending at its fastest rate at time

$$t = 10 - \sqrt{50} \sim 2.93$$

hours past noon.



Review of Differentiation

Derivative Review.

Examples

Example 22. Find an equation of the tangent line to the curve

$$y = f(x) = (2x^3 + 1)^2$$

at the point $(1, 9)$.

Explanation. Let

$$y = (2x^3 + 1)^2$$

and

$$u = 2x^3 + 1.$$

Then

$$y = u^2$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(u^2) \cdot \frac{d}{dx}(2x^3 + 1) \\ &= 2u(6x^2) \\ &= 2(2x^3 + 1)(6x^2). \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 2(3)(6) = 36,$$

and an equation of the tangent line is

$$y - 9 = 36(x - 1).$$

Example 23. Find an expression for the derivative

$$\frac{d}{d\theta}(\theta \cos(5\theta)).$$

Learning outcomes:
Author(s):

Explanation. We use the product rule first to get

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)).\end{aligned}$$

Now we use the chain rule to compute

$$\frac{d}{d\theta}(\cos(5\theta)).$$

For this we hide the composition by letting

$$y = \cos(5\theta)$$

and

$$u = 5\theta.$$

Then

$$y = \cos u$$

and by the chain rule

$$\begin{aligned}\frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du}(\cos u) \cdot \frac{d}{d\theta}(5\theta) \\ &= (-\sin u)(5) \\ &= -5 \sin(5\theta).\end{aligned}$$

The final result is that

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) - 5\theta \sin(5\theta).\end{aligned}$$

Here's a shorter version of the same solution.

We use the product rule first to get

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)).\end{aligned}$$

Then we use the chain rule to differentiate $\cos(5\theta)$, giving

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta(-\sin(5\theta)) \frac{d}{d\theta}(5\theta) \\ &= \cos(5\theta) - 5\theta \sin(5\theta).\end{aligned}$$

Exercises

Directions: Follow the method of Example 1 *exactly* for each of the following problems.

Question 211 Find expressions for each of the following derivatives. Simplify your expressions for the derivatives. Do not simplify the function being differentiated.

Part 1:

- (a) $\frac{d}{dw}(\arctan w + \arctan(1/w))$
 (b) $\frac{d}{dy}(\arcsin(\sqrt{1-y^2}))$

Part 2:

Any comments or observations on the derivatives above?

Question 212 *Desmos link:*

<https://www.desmos.com/calculator/zjqsdhvvz4p>

Desmos activity available at 151: Building Temperature

Question 213 Water is poured into a cylindrical tank at a constant rate. At the same time, water flows out of a small hole in the bottom of the tank. The tank is empty at noon.

The function

$$t = f(h) = -\frac{2}{k_2} \left(\sqrt{h} + \frac{k_1}{k_2} \ln \left| \frac{k_1 - k_2 \sqrt{h}}{k_1} \right| \right), \quad t \geq 0,$$

expresses the time (measured in minutes past noon) in terms of the depth (measured in cm) of water in the tank. Here k_1, k_2 are positive constants.

- (a) Use the above function to verify that the tank is empty at noon.
- (b) Find a simplified expression for the derivative dt/dh .
- (c) Use the result of part (b) to show that

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}.$$

- (d) What are the units of k_1, k_2 ? How do you know?
- (e) For much more on this problem, see Questions 23 and 24 of the chapter Derivatives of Inverse Functions.

Question 214 The function

$$s = f(t) = Ae^{-k_1 t} \cos(k_2 t), \quad t \geq 0,$$

expresses the displacement (in meters) from equilibrium of an oscillating mass on a spring in terms of the number of seconds since the mass was released from rest.

- (a) What are the units of the constants A, k_1 , and k_2 ? Explain how you know.
- (b) Find an expression for the velocity ds/dt of the mass.
- (c) Find an expression for the acceleration

$$\frac{d}{dt} \left(\frac{ds}{dt} \right) = \frac{d^2 s}{dt^2}$$

of the mass.

- (d) Use algebra to show that

$$\frac{d^2 s}{dt^2} = - \left(2k_1 \frac{ds}{dt} + (k_1^2 + k_2^2)s \right).$$

Desmos link: <https://www.desmos.com/calculator/ygikqgj7af>

Desmos activity available at 151: Damped Harmonic Oscillator

Question 215 (a) Find a function

$$s = f(h), \quad h \geq 0,$$

that expresses the distance (in miles) to the horizon in terms of your altitude (in miles). We'll suppose the earth to be a perfect sphere of radius $R = 4000$ miles. The distance to the horizon is the arclength AT below, measured along the surface of the earth (you can think of this distance as the radius of the spherical disk visible to us). Our height is the distance AP .

The function is

$$s = f(h) = 4000 \arccos \left(\frac{4000}{4000 + h} \right), \quad h \geq 0.$$

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

(b) Use the graph of the function $s = f(h)$ show below to sketch a graph of the derivative

$$y = \frac{ds}{dh} = f'(h).$$

by hand. What is the domain of the derivative? Note that the curve $s = f(h)$ has a horizontal asymptote. What is an equation of this asymptote?

Desmos link: <https://www.desmos.com/calculator/nof5l3mtfy>

Then check your sketch by activating the folder in Line 1 of the worksheet below.

Desmos link: <https://www.desmos.com/calculator/pdbjfao316>

(c) Find an expression for the derivative

$$\frac{ds}{dh} = \frac{d}{dh} \left(\arccos \left(\frac{R}{R + h} \right) \right).$$

Hint: Start by using the graphs of the functions $y = \arcsin x$ and $y = \arccos(x)$ below and your knowledge of the derivative $d/dx(\arcsin x)$ to find an expression for $d/dx(\arccos x)$. Or equivalently, recognize that

$$\arcsin x + \arccos x = \pi/2.$$

Desmos link: <https://www.desmos.com/calculator/zuq9rf1j4d>

(d) With $R = 4000$, evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=16}.$$

(e) Interpret the meaning of the derivative in part (d) in terms of specific small changes.

(f) You take a ride on Blue Origin and in two minutes are boosted straight up to an altitude of 32 miles. Suppose that the function

$$h = g(t) = \begin{cases} 10t^2 + 6t^3, & 0 \leq t \leq 1 \\ 32 - 10(2-t)^2 - 16(2-t)^3, & 1 < t \leq 2, \end{cases}$$

expresses your altitude (in miles) in terms of the number of minutes since launch for the first two minutes of your flight.

(i) At what rate (with respect to time) is your distance to the horizon changing when you are one minute into the flight? Use the graph of the function $s = f(g(t))$ shown below to first approximate this rate.

(ii) At what rate (with respect to time) is your distance to the horizon changing at the start of the flight? Use the graph of the function $s = f(g(t))$ shown below to first approximate this rate. Are you surprised given your graph in part (b)?

Desmos link: <https://www.desmos.com/calculator/jchwjpxugf>

Review Problems

Some problems for review.

Question 216 This question is about how the temperature inside a building changes in response to changes in the outdoor temperature. We assume the building has no internal heating or cooling system.

We'll suppose that the function

$$f(t) = M - B \cos\left(\frac{\pi}{12}t\right), \quad t \geq 0,$$

expresses the outdoor temperature (in Fahrenheit degrees) in terms of the number of hours past 4am.

Newton's law of cooling models the rate at which the indoor temperature is changing at any time. It says that this rate of change is proportional to the difference in the indoor and outdoor temperatures. So if the function

$$T = g(t) \quad t \geq 0,$$

expresses the outdoor temperature (in Fahrenheit degrees) in terms of the number of hours past 4am, Newton's law says that

$$\frac{dT}{dt} = k(f(t) - g(t)) \tag{5}$$

for some constant k .

- (a) What are the units of k ? How do you know?
- (b) Is k positive or negative? How do you know?
- (c) Experiment with the sliders in the demonstration below. Summarize your observations.

Desmos link: <https://www.desmos.com/calculator/oag9lhvgo5>

Desmos activity available at 151: *Building Temperature*

Next quarter you will learn how to use the above equation to determine the indoor temperature at any time given the temperature at some specific time.

Learning outcomes:

Author(s):

For now, we'll just claim that the indoor temperature is given by the function

$$\begin{aligned}T &= g(h) \\&= M + Ce^{-kt} - \frac{B}{1 + (\frac{\pi}{12k})^2} \left(\cos\left(\frac{\pi}{12}t\right) + \frac{\pi}{12k} \sin\left(\frac{\pi}{12}t\right) \right) \\&= M + Ce^{-kt} - \frac{B}{\sqrt{1 + (\frac{\pi}{12k})^2}} \cos\left(\frac{\pi}{12}t - \phi\right),\end{aligned}$$

where

$$\phi = \arctan\left(\frac{\pi}{12k}\right).$$

Hanging Chains

Comparing a catenary with a weighted chain.

Question 217 *Desmos link:*

<https://www.desmos.com/calculator/cifsqaas5j>

Desmos activity available at 151: Weighted Chain

Question 218 *Desmos link:*

<https://www.desmos.com/calculator/sv70z21j2j>

Desmos activity available at 151: Catenary

Learning outcomes:
Author(s):

Electron in a Crossed Field

Electron.

Crossed Fields and the Hodograph

The force on an a charge q with velocity \mathbf{v} in a crossed magnetic/electric field is given by

$$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}).$$

Our problem is to parameterize the motion of a charged particle when the fields are uniform and mutually orthogonal. We suppose the magnetic and electric fields point in the respective directions \mathbf{k} and \mathbf{j} , and that the charge has velocity \mathbf{v}_0 perpendicular to \mathbf{k} at time $t = 0$.

Then the charge, assumed to be positive, has acceleration

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = k_1\mathbf{j} + k_2\mathbf{v} \times \mathbf{k},$$

for some constants $k_1, k_2 \geq 0$, with respective units m/sec^2 and sec^{-1} .

This is a differential equation that is easy enough to solve algebraically. But our aim here is to take a geometric approach that uses some key ideas from Calculus 3.

In sketching motion we typically draw the path and perhaps draw a few position vectors at equally-spaced time intervals. These vectors allow us to approximate a few velocity vectors, or at least their relative lengths. But instead of drawing the velocity vectors tangent to the path, it is usually more useful to draw them with their tails pinned at a common point. The curve traced by these tips of these pinned vectors is called the *hodograph* of a motion.

For starters, the hodograph gives us a way to visualize the motion's acceleration as it is tangent to the hodograph. But to get some idea of the acceleration's magnitude, we need to consider not only the direction and magnitude of the velocity vector but also its rotation rate.

For a uniform circular motion that rotates around a circle of radius r meters at a constant rate of ω rad/sec, for example, we *see* from the hodograph (a circle of radius $v = \omega r$ that the acceleration vector points directly toward the center of the path and rotates at the same rate as both the position and velocity vectors.

Learning outcomes:
Author(s):

So without any derivatives, we get the acceleration's magnitude to be

$$|\mathbf{a}| = \omega|\mathbf{v}| = \omega^2 r.$$

Turning off the Electric Field

Returning to our charged particle, with the electric field turned off ($k_1 = 0$), the acceleration and velocity vectors are perpendicular, so the charge moves with constant speed $|\mathbf{v}_0| = v_0$. But then because \mathbf{v} is perpendicular to \mathbf{k} ,

$$|\mathbf{a}| = k_2|v| = k_2 v_0$$

is also constant.

So we are looking for a plane motion having constant speed and an acceleration with constant magnitude. One choice would be uniform circular motion, where

$$|\mathbf{a}| = v_0 \omega,$$

where $\omega = d\theta/dt$ is the (constant) rotation rate of the pinned velocity vector. Then $k_2 = \omega$ and the trajectory has radius

$$r = \frac{v_0}{\omega} = \frac{v_0}{k_2}.$$

To see that there are no other possible motions we could probably appeal to some uniqueness theorem of differential equations. But for a more geometric approach, consider what we know.

- (a) that the charge moves with constant speed and
- (b) that the acceleration vector rotates at a constant rate.

Question 219 What do these conditions imply about the trajectory?

Hint: Think about curvature.

Explanation. Click the arrow to the lower right for the solution.

Let θ be the angle from $\mathbf{i} = \langle 1, 0, 0 \rangle$ to \mathbf{v} . Then because \mathbf{v} and \mathbf{a} are orthogonal, these vectors rotate at the same rate $\omega = d\theta/dt$. And because the speed is constant, the path's radius of curvature

$$r = \frac{|v|}{\left| \frac{d\theta}{dt} \right|} = \frac{v_0}{k_2}$$

is also constant. And because the path is a plane curve, it is a circle with this radius.

1 Newton's Law of Cooling

Exploration 220 *Desmos link:*

<https://www.desmos.com/calculator/yebumxuwms>

Electron 11

Question 221 *The function*

$$P = 40e^{-t/20}, \quad 0 \leq t \leq 15,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

(a) *Find a function*

$$r = g(P)$$

that expresses the growth rate of the population (in millions of bacteria/hour) in terms of the population. Include a domain.

(b) *Use the result of part (a) to determine the population when it is decreasing at the rate of 1,820,000 bacteria/hour. End with a concluding sentence. No credit for other methods.*

Short Quizzes Math 151

Short Quizzes Math 151.

Question 222 The function

$$h = f(v), \quad 80 \leq v \leq 120,$$

expresses the height of a helicopter (measured in feet) in terms of its speed (measured in ft/sec).

Suppose

$$\left. \frac{dh}{dv} \right|_{v=100} = -20.$$

- (a) What are the units of the above derivative?
- (b) Explain the meaning of the derivative using the language of small changes.

Question 223 The function

$$P = f(t) = 40e^{-t/20}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

- (a) Find a function

$$r = g(P),$$

that expresses the growth rate (in millions of bacteria/hour) of the population in terms of the population (in millions of bacteria).

- (b) Use the result of part (a) to determine the population when it is decreasing at the rate of 1,820,000 bacteria/hour.
 - (i) No credit for other methods.
 - (ii) End with a concluding sentence.

Learning outcomes:
Author(s):

Question 224 Find an equation of the tangent line to the curve

$$x^3 \sin(5y) + 4e^{xy} = 4x$$

at the point $(1, 0)$.

- (a) Show all work.
- (b) Use the Leibniz notation for derivatives and their evaluations.
- (c) Include a brief explanation with your work.
- (d) End with a concluding sentence.



Short Quizzes Math 142

Short Quizzes Math 142

Question 225 Let Q be the point on the circle of radius 40 meters that is 90 meters from the point $A(40, 0)$ (coordinates measured in meters). The distance is measured clockwise around the circle from A to Q .

Between 12:06pm and 1:00pm a beetle crawls counterclockwise around this circle at a constant speed of 5 meters/min, passing the point Q at 12:43pm.

- (a) Find an expression for the function

$$\theta = f(t), \quad 6 \leq t \leq 60,$$

that expresses the polar angle (measured in radians) of the beetle (more precisely of the vector giving the beetle's position relative to the origin) in terms of the number of minutes past noon being sure to do the following:

- (i) Explain your reasoning. This means to include a brief description of what you are computing for each of your computations.
- (ii) Include units for each number in each computation.
- (iii) Include a graph (drawn by hand) of the function $\theta = f(t)$ to help with your explanation.
- (iv) Include a sketch (drawn by hand) of the circle on a coordinate system with appropriately labeled points and arclengths. Label the axes with the appropriate variable names and units.

- (b) Find functions

$$x = g_1(t), \quad 6 \leq t \leq 60,$$

and

$$y = g_2(t), \quad 6 \leq t \leq 60,$$

that express the coordinates (in meters) of the beetle in terms of the number of minutes past noon. Include a brief explanation.

- (c) Input the correct coordinate functions on Lines 2 and 3 of the worksheet below. Then play the slider u (another name for t , the number of minutes past noon) in Line 1 to see if your functions are correct.

Desmos link: <https://www.desmos.com/calculator/lkrunhfgxi>

Learning outcomes:
Author(s):

142: Short Quiz 1

- (i) *Explain how you know that your functions are correct.*
 - (ii) *Include at least two screenshots of the animation at different times to help with your explanation.*
-

Question 226 *Solve the equation*

$$9 - 3 \sin \theta = 11.$$

- (a) *Explain your reasoning thoroughly, in complete sentences.*
 - (b) *Include a picture as in class to help with your explanation.*
 - (c) *End with a concluding sentence that expresses the solution as a set.*
-

Sinusoidal Modeling, Math 142

Implicit differentiation.

Question 227 Suppose that during a five-day period, beginning at midnight Monday morning, the depth of the water at the Edmond's Pier is a sinusoidal function of time.

Suppose also that a low tide of 7 feet occurs at 1:00am Monday morning and that the following high tide of 23 feet occurs at 6:45am that same morning.

- (a) Sketch by hand a graph of the function

$$h = f(t)$$

that expresses the depth of the water (in feet) as a function of the number of hours since midnight, Monday morning.

- (b) Find an expression for the above function. Include a domain.
 (c) Find all times during the week when the water is 10 feet deep. Give exact times. Do not use a calculator.
 (d) Approximate the clock times (to the nearest minute) on Friday when the water is 10 feet deep.

Question 228 Assume for this question that each month has 30 days and that the number of hours of daylight/day in Seattle is a sinusoidal function of time. Assume also that on June 21, Seattle gets a maximum of 16 hours of daylight/day and that on December 21, Seattle gets a minimum of 8 hours of daylight/day.

- (a) Find a function

$$H = f(t), 0 \leq t \leq 12,$$

that expresses the number of daylight hours/day in Seattle in terms of the number of months since June 21. Use the cosine function. Start by sketching a graph. Explain your reasoning.

- (b) Use your function to determine the number of hours of daylight/day that Seattle gets on March 1.

Learning outcomes:
 Author(s):

- (c) *About how many more minutes of daylight/day do we get tomorrow than today?*
 - (d) *Use your function to determine the day(s) of the year when Seattle gets 14 hours of daylight/day.*
-

Simple Harmonic Motion

Simple harmonic motion.

Suppose for this problem that the earth is a ball with uniform density of radius 4000 miles. Now imagine drilling a straight tunnel through the earth from the North Pole to the South Pole.

A rock dropped from rest at the north pole falling through the tunnel would then oscillate in simple harmonic motion between the poles and return to the north pole every 84 minutes. This means we can think of the rock as being dragged along by a point moving around the earth at constant speed as illustrated below.

Desmos link: <https://www.desmos.com/calculator/ij8dqowgza>

Desmos activity available at 142: Simple Harmonic Motion

- (a) Activate the *Protractor* folder in Line 15 to see the protractor, where consecutive tick marks subtend equal angles of $\pi/100$ radians about the earth's center. Use the protractor but *not* any trigonometry to answer the following questions. *Hint:* Think proportionately.
 - (i) Estimate the distance of the rock from the South Pole at time $t = 16.8$ minutes after the rock is released.
 - (ii) Estimate the first two times when the rock is 1000 miles from the South Pole.
- (b) Sketch by hand one period of the graph of the function

$$s = f(t) \quad t \geq 0,$$

that expresses the distance of the rock (measured in thousands of miles) from the South Pole in terms of the number of minutes since the rock was released. Assume the rock is dropped at midnight on July 1, 2085.

- (c) Find an expression for the function $s = f(t)$. Start by expressing the distance s in terms of the polar angle θ marked above.
- (d) Use your function to determine the exact distance between the rock and the South Pole at time $t = 16.8$ minutes past noon. Then use a calculator to approximate this distance to the nearest mile and compare it with your estimate in part (i).

Learning outcomes:
Author(s):

Simple Harmonic Motion

- (e) Use your function to find all (exact) times when the rock is 1000 miles from the South Pole. Do this by first finding all (exact) polar angles when the rock is 1000 miles from the South Pole.
- (f) Find the (exact) first two times when the rock is 1000 miles from the South Pole. Then use a calculator to estimate these clock times to the nearest minute
- (g) Find all (exact) times between 8pm and 12pm, July 1, 2085 when the rock is 1000 miles from the South Pole. Then use a calculator to estimate these clock times to the nearest minute