
Calculus 1

May 9, 2026

Contents

Announcements, HW, Math 151 Hybrid	13
1 Homework Assignments	13
2 Announcements	13
Exam 1	14
Stereographic Projection	20
1 Introduction to Differential Calculus	20
2 Stereographic Projection	20
An Introduction to Differentiable Functions, Part 1	24
Introduction, Part 2	27
A Ferris Wheel	34
Lambert's Equal-Area Map, Part 1	37
1 Lambert's Equal-Area Map	37
2 The Mercator Map	38
3 Loxodromes	38
Shadow of a Building, Part 1	40
Looking Down on the Moon	44
Small Changes	47
1 Odometer Readings	47
2 A Projectile	50
3 The Falling Ladder, Part 1	51
4 The Falling Ladder, Part 2	52
5 Riding a Ferris Wheel	54
The Derivative as a Scaling Factor	55
Stereographic Projection	58
1 Introduction to Differential Calculus	58
2 Stereographic Projection	58
Stretching Factors	62
1 Rubber Band Calculus	62
2 Non-Linear Stretching Functions	63
3 A Hanging Slinky	66

4	Gas Consumption	69
5	A Falling Slinky	70
6	Atmospheric Pressure	70
	Average Rates of Change	71
	A Balloon's Rate of Ascent	79
	Rates of Change, Gas Mileage	81
1	Gas Mileage	81
	Practice Quiz 1, Approximating Scaling Factors Numerically	83
	The Rythm of a Function	85
	Differentiable Functions	87
1	Differentiability	87
2	Using Algebra to Compute Derivatives	90
3	Applications	94
	Differentiable Functions, Part 2	95
1	Using Limits to Compute Derivatives	95
	Limits and Derivatives, Part 1	98
1	Limits and Speed	98
2	The Cubing Function, Part 1	99
3	The Cubing Funtion in Context	100
4	The Cube Root Function in Context	100
5	Limits and Gas Mileage	101
	Limits and Derivatives, Part 2, Trigonometric Functions	103
1	The Sine Function	103
2	A Building's Shadow	105
	Practice with Limits and Derivatives	106
1	More Practice with Rational Functions	106
2	The Squaring Function and Its Inverse	110
3	The Cubing Function and its Inverse	113
4	A Square Root Function	113
5	More on the Squaring Function	113
	Limits	114
1	Limits and Tangent Lines	114

2	Limits and Gas Mileage	114
3	Limits, Gas Mileage and Speed	115
4	Limits, Speed and Altitude	116
5	Limits and Gas Mileage	117
6	Limits and Purchasing Power	117
	Free Fall	118
1	Speed vs. Velocity	118
2	A Rock in Freefall	119
3	Takeaways	120
	The Weighted Chain	123
	Purchasing Power	125
1	Purchasing Power	125
2	Weight in Space	129
3	A Rock in Free Fall	130
4	Distance to the Horizon	131
5	Intensity of Sound	132
	Practice Quiz 1	133
1	Part 1	133
	Introduction to Motion	136
	The Derivative	137
1	Limits, Speed and Altitude	142
	Derivatives of Polynomials	144
1	Differentiating Polynomials	144
2	Stock Price	149
3	Homework Solution	160
	Derivatives of Polynomials CW	161
1	The Power Rule	161
2	Stock Price	162
	Derivatives of Polynomials Worksheet	166
	Normals to a Parabola	168
	Gas Consumption	170
	The Quotient and Product Rules	173

1	Relative Rates of Change	173
2	Exercises	174
	Test 1	177
	Derivatives of Exponential Functions	181
1	Exercises	192
	The Chain Rule	194
1	Introduction	194
2	Distance to the Horizon	196
3	Visualizing the Chain Rule	198
4	Computing with the Chain Rule	199
	The Chain Rule and Polynomials	215
1	Practice Questions	215
2	Discussion Questions	219
	The Chain Rule CW1	221
1	Trails	221
2	Purchasing Power	223
3	An Oscillating Ferris Wheel	224
	The Chain Rule and the Price of a Stock	225
	The Doppler Effect: Stretching Time	229
	On Trails	232
1	Trails	232
2	Trails Described with Functions	235
3	Parametrically Defined Trails	235
	Differentiating Trigonometric Functions using Limits	236
1	The Motion of a Balloon	236
2	An Important Limit	237
3	Variations on the Limit	238
4	Sine Proof 1	239
5	Sine Proof 2	240
6	The Derivative of the Cosine Function	241
7	Exercises	241
	Derivatives of Trigonometric Functions	244

1	Introduction	244
2	Velocity	244
3	Derivatives of the Sine and Cosine Functions	245
4	Riding a Ferris Wheel	246
5	Simple Harmonic Motion	247
6	Two Optimization Problems	248
7	Exercises	249
8	Visualizing Derivatives	250
9	The Derivative of the Sine Function	251
10	Applications 1	252
11	Transformations of the Sine Function	253
12	Applications 2	253
13	MATH 142	253
	Trigonometry Tangent Line	255
	The Sliding Ladder, Part 1	257
	The Tangent Function and its Inverse	261
1	The Tangent Function	261
2	Exercises	262
	The Tangent Function CW1	264
1	The Tangent Function	264
2	Exercises	266
	Circular Interpolation	268
1	Three Problems	268
2	A Quicker Approach: Circular Interpolation	273
	Review of Sine and Cosine	275
1	Derivatives	275
2	Transformations	275
3	Simple Harmonic Motion	276
4	Ferris Wheels	277
5	Flagpole and Track	278
	Exponential Functions, Part 2	280
1	Introduction	280

2	Relative Growth Rates	283
3	The Number e	284
4	An Example	285
5	Exercises	286
6	Conceptual Questions	291
	Exponential Functions, Short Version	292
1	Introduction	292
2	Relative Growth Rates	294
3	The Number e	295
4	An Example	296
5	Exercises	298
6	Conceptual Questions	303
	Practice Quiz 2	304
	Derivatives of Inverse Functions, Shortest Version	308
1	The Main Idea	308
2	Theorem	309
3	Examples	309
4	Exercises	311
	Derivatives of Inverse Functions, Short Version	313
1	The Main Idea	313
2	Derivatives of Inverse Power Functions	314
3	Derivative of the Arc Sine Function	317
4	Derivative of the Arc Cosine Function	319
5	Exercises	320
	Derivatives of Inverse Functions	322
	Looking Down on the Moon, Part 2	338
	Inverse Trig CW	341
	The Quotient Rule	344
	Moving Averages and their Rates of Change	357
	Implicit Differentiation	358
	The Ladder and the Tree	366
	Statuary Hall and Implicit Differentiation	368

The Ladder and the Tree, Part 2	370
More Related Rates Problems	373
1 Shadow of a Building	373
2 Kite Flying	374
3 A Crawling Beetle	374
4 A Mechanical Motion	375
5 Tracking a Helicopter	375
6 Distance to the Horizon	376
7 Jar Lid and Rubber Band	377
8 Disco Dancing	377
The Natural Log Function	378
1 The Derivative of the Natural Log Function	378
2 Relative Rates of Change	380
Derivatives and the Shapes of Curves	382
1 A Surge Function	382
2 Changes and Rates of Change, Relative Changes and Relative Rates of Change	383
Optimization	386
Measuring Distances on the Earth	389
1 Trigonometry: Along a Circle of Latitude	389
2 Calculus: Along a Circle of Latitude	390
Disco Dancing, an Optimization Problem	391
Clock Hands	392
Mars and the Outer Planets	395
Hours of Daylight	397
Motion	400
Linear Approximation	402
1 The Derivative as a Scaling Factor	402
2 Interpreting the Derivative	403
2.1 Speed	403
3 Weight and Distance	404
4 Speed and Height	405
5 Distance to the Horizon	406

6	Gas Mileage	406
	Linear Approximation, Part 2	408
1	Review of the Natural Log Fuction	408
2	First Derivatives	408
3	Second Derivatives	410
	Review of Differentiation	411
	Final, Math 151	417
1	Directions	417
2	Examples of How to Show Work	418
	Review Problems	420
	Hanging Chains	422
	Electron in a Crossed Field	423
1	Newton's Law of Cooling	425
	Short Quizzes Math 151	426
	Short Quizzes Math 142	428
	Sinusoidal Modeling, Math 142	430
	Simple Harmonic Motion, Law of Cosines, Math 142	432
	Quiz Solutions	434
1	Quiz 5A	434
	Introduction to Integration, Part 1	436
	Introduction to Integration, Part 2	440
1	Review	440
2	Summation Notation and Technology	440
	Addition	444
1	Two Balloon Problems	444
2	Time, Distance, and Mileage Problems	445
3	On Trails	448
4	A Rollercoaster Problem	449
5	The Mercator Map	451
	Integrating Power Functions	453
1	Two Balloon Problems	453
	Riemann Sums	454

Geometric Integration	458
The Fundamental Theorem of Calculus, Part 2	461
The Fundamental Theorem of Calculus, Part 1	465
1 Gas Consumption	467
2 On Trails	468
Substitution and Parametrically-defined Curves	472
1 The Basics	472
2 A Geometric Interpretation	474
The Limits to Growth	477
Area	479
1 Parabolas and Power Functions, Part 1	479
2 Parabolas, Part 2	481
3 Which Area is Greatest? Least?	484
4 Trigonometric Functions	484
5 The Rectangular Elastica	485
Volumes of Solids in Three-Dimensions	487
1 Square Cross Sections	487
2 The Hoof of Archimedes	490
3 Archimedes' Bicylinder	491
4 Triangles and Cones	491
The Hoof of Archimedes	493
Volumes, Part 2	496
1 Paraboloid of Revolution	496
2 Napkin Ring	498
The Mylar Balloon	499
1 The Rectangular Elastica	499
2 Happy Birthday!	501
Which Tanks Empty Faster?	504
1 Preliminaries	504
2 Thoughts on Differential Equation in Math 152	504
3 Which Tanks Drain Faster?	505
4 A Hemispherical Tank	506

Parametrically-defined curves and Area	507
1 The Basics	507
2 The Area of a Circle	508
Parametrically-defined Curves, CW	510
1 The Basics	510
2 An Application: Gas Consumption	512
3 Exercises	513
Integration by Parts	516
1 The Basics	516
Arclength of Curves in the Plane	517
1 The Ideas	517
2 Circles	518
3 Parameterizing an Astroid by Arclength	519
4 Trail Profiles	520
The Cycloidal Pendulum	521
1 The Cycloid	521
2 The Cycloidal Track	522
152 Review	526
Loxodromes and Equiangular Spirals	536
1 Loxodromes	536
Circles, Spheres, Area, Volume	537
1 The Area of a Circle	537
2 The Hoof of Archimedes	537
3 Archimedes' Bicylinder	538
4 Tori	538
5 The Sphere	538
Hyperbolic Trigonometry	539
1 The Hyperbolic Cosine and Sine Functions	539
2 Hanging Chains	540
3 The Hyperbolic Tangent Function	540
4 Inverse Functions	541
Review of the Chain Rule	542

1	Example	542
2	Exercises	543
	Trails and Hanging Chains, Part 1	545
1	Angles of Inclination	545
2	A Climber's Trail	546
3	A Hanging Chain	548
	The Mercator Map	550
1	Scaling Factors and Elastic Bands	550
2	Lambert's Equal-Area Map	552
3	The Mercator Map	553
4	Loxodromes	553
	Riemann Sums with Geometric Partitions	555
1	Integrating $1/x$	555
2	Computing	556
	Quiz Solutions Fall 2025	557
1	Quiz1B	557
2	Quiz 2A	558
3	Quiz 2B	558
4	Quiz 3A	560
5	Quiz 3B	561
6	Quiz 4A	561
7	Quiz 4B	561
8	Quiz 5A	563
9	Quiz 5B	564
10	Quiz 6A	565
11	Quiz 6B	567
12	Quiz8A	569
13	Quiz 8B	570

Announcements, HW, Math 151 Hybrid

The tangent function and its inverse, classwork.

1 Homework Assignments

2 Announcements

Exam 1

Solution key.

Question 1 Find an equation of the tangent line to the curve

$$y = \frac{48}{\sqrt{x}} - 3x + 2$$

at the point on the curve with x -coordinate $x = 4$.

Explanation. The tangent line to the curve

$$y = f(x) = \frac{48}{\sqrt{x}} - 3x + 2$$

at the point $(x, f(x))$ has slope

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \left(\frac{48}{\sqrt{x}} - 3x + 2 \right) \\ &= \frac{d}{dx} (48x^{-1/2}) - \frac{d}{dx} (3x) + \frac{d}{dx} (2) \\ &= 48 \frac{d}{dx} (x^{-1/2}) - 3 + 0 \\ &= 48 \left(-\frac{1}{2} x^{-3/2} \right) - 3 \\ &= -24x^{-3/2} - 3. \end{aligned}$$

So the tangent line to the curve $y = f(x)$ at the point $(4, f(4))$ has slope

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{x=4} &= \left(-24x^{-3/2} - 3 \right) \Big|_{x=4} \\ &= -6. \end{aligned}$$

To find the y -coordinate of the point of tangency, let $x = 4$. Then

$$y = f(4) = 14.$$

So the tangent line to the curve $y = f(x)$ at the point $(4, 14)$ has slope -6 and equation

$$y - 14 = -6(x - 4).$$

Learning outcomes:
Author(s):

Question 2 The function

$$W = f(r) = \frac{2000}{r^2}, r \geq 4,$$

expresses the weight of an astronaut (measured in pounds) in terms of her distance from the center of the earth (measured in thousands of miles).

- (a) Find a fully-simplified expression for the average rate of change in the astronaut's weight with respect to her distance from the earth's center between distances $r = b$ and $r = c$ thousands of miles from the center. Assume $b, c \geq 4$ and that $b \neq c$. Work vertically, one equal sign per line. Do not split the page into multiple columns.
- (b) Use your expression from part (a) and the definition of the derivative (no credit for other methods) to find a fully-simplified expression for the derivative

$$\left. \frac{dW}{dr} \right|_{r=a}.$$

No credit for other methods.

Explanation. (a) The average rate of change is

$$\begin{aligned} \frac{\Delta W}{\Delta r} &= \frac{f(c) - f(b)}{c - b} \\ &= \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \\ &= \left(\frac{2000}{c - b} \right) \left(\frac{b^2 - c^2}{c^2 b^2} \right) \\ &= \left(\frac{2000}{c - b} \right) \cdot \frac{(b + c)(b - c)}{c^2 b^2} \\ &= \frac{-2000(b + c)}{c^2 b^2}. \end{aligned}$$

(b) The derivative is

$$\begin{aligned} \left. \frac{dW}{dr} \right|_{r=a} &= \lim_{\Delta r \rightarrow 0} \frac{\Delta W}{\Delta r} \\ &= \lim_{\Delta c \rightarrow a} \frac{-2000(a + c)}{c^2 a^2} \\ &= -\frac{2000(a + a)}{a^2 a^2} \\ &= -\frac{4000}{a^3}. \end{aligned}$$

Question 3 Use precalculus and arithmetic, but not calculus, to give numerical evidence that suggests an answer to this question:

A rocket orbits a planet of radius 2000km at an altitude of 400 km above the surface. Is there a scaling factor that converts (by multiplication) a small change in orbit's radius to an accurate approximation of the corresponding change in radius of the portion of the planet's surface visible to an astronaut in the rocket? The radius (orange) is measured along the planet's surface.

- (a) Start by writing a complete sentence that defines a function expressing the radius of the visible portion in terms of the altitude above the surface. Include units in your definition. Use meaningful variable names, not x and y . Then find an expression for the function. Annotate the picture to help with your explanation.
- (b) Then gather your evidence in an appropriate table with at least six rows. If the data suggest the existence of a scaling factor, explain why and approximate the scaling factor to the nearest tenth. Include units. If the data suggest otherwise, explain why.

Explanation. Let

$$r = f(h), h \geq 0,$$

be the function that expresses the spherical radius (in km) of surface visible to the astronaut in terms of her altitude above the surface (in km).

Let θ be the radian measure of $\angle OPT$ between the astronaut's line of sight to the horizon and the downward vertical pointing from the astronaut to the planet's center.

Then

$$\cos \theta = \frac{2000}{2000 + h}.$$

And since $0 \leq \theta < \pi/2$,

$$\theta = \arccos \left(\frac{2000}{2000 + h} \right).$$

So the radius of the visible portion (measured in km) is

$$\begin{aligned} r &= 2000\theta \\ &= 2000 \arccos \left(\frac{2000}{2000 + h} \right). \end{aligned}$$

h (km)	$r = f(h)$ km	$\Delta h = h - 400$ km	$\Delta s = f(h) - f(400)$ (km)	$\Delta s / \Delta h$ (km/km)
397	1167.59	-3	-3.78	1.26
398	1168.85	-2	-2.52	1.26
399	1170.11	-1	-1.26	1.26
400	1171.37	0	0	
401	1172.63	1	1.26	1.26
402	1173.88	2	2.51	1.26
403	1175.13	3	3.76	1.26

The data suggest the scaling factor is approximately 1.26 km/km. It is dimensionless.

Question 4 Between speeds of 100 km/hour and 150 km/hour, the function

$$v = f(G), \quad 4 \leq G \leq 17,$$

expresses the speed of a car (in km/hr) in terms of its gas mileage of a car (in km/liter).

Suppose $f(5) = 140$.

- (a) Which of the following is more likely to be correct? Explain your reasoning.

$$\left. \frac{dv}{dG} \right|_{G=5} = 0.8$$

or

$$\left. \frac{dv}{dG} \right|_{G=5} = -0.8$$

- (b) (1 point) What are the units of the correct derivative above?
- (c) (2 points) Explain the meaning of the derivative as a scaling factor. Write your explanation in complete sentences without any mathematical notation.

Explanation. (a) At such high speeds, an increase in the speed would most likely decrease the gas mileage. We would therefore expect the derivative to be negative. So the second choice is most likely correct.

- (b) The units of the derivative dv/dG are

$$\frac{\text{km/hr}}{\text{km/liter}}.$$

- (c) The derivative is a scaling factor that converts (by multiplication) a small change in the mileage (measured in km/liter from 5 km/liter) to an accurate approximation of the corresponding change in the speed (measured in km/hr).

More specifically, if the gas mileage increases from 5 km/liter to 5.1 km/liter, the speed would decrease by approximately 0.08 km/hr (ie. from 140 km/hr to 139.92 km/hr).

Question 5 The function $T = f(m)$, $0 \leq t \leq 32$, expresses the temperature (in Celsius degrees) of a beaker of water in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/ifsu6n6qt0>

Access this activity online at 151: Normals to Parabola 2

Use a straightedge to approximate the time(s) when the temperature is changing at a rate equal to its average rate of change between 12:00pm and 12:32pm. Explain your reasoning thoroughly in complete sentences. Draw on the graph above to help with your explanation. End your explanation with a concluding sentence that captures both the question and your response.

The parabola below has equation

$$y = \frac{x^2}{2p},$$

for some constant $p \neq 0$. Point A has coordinates $(a, a^2/(2p))$, and point B has coordinates $(b, b^2/(2p))$. Assume $a \neq b$.

Desmos link: <https://www.desmos.com/calculator/oczzppq2n4>

Access this activity online at 151: Exam 1 Figure 1S

Explanation. First draw the secant line through the points $(0, f(0))$ and $(32, f(32))$. The slope of this line is the average rate of change in the temperature with respect to time between 12:00pm and 12:32pm.

Second, holding your ruler parallel to the secant line, slide it so that it becomes tangent to the graph. There are three possibilities.

Now since parallel lines have the same slope and the slope of the tangent line to the curve measures the instantaneous rate of the change of temperature with respect to time, the m -coordinates of the three points of tangency give the desired times. It looks like these coordinates are approximately $m \sim 4, 17, 29$.

Exam 1

So the temperature of the water is changing at a rate equal to its average rate of change between 12:00pmm and 12:32pm at approximately 12:04pm, 12:17pm, or 12:29pm.

Stereographic Projection

Maps of the earth, stereographic projection.

1 Introduction to Differential Calculus

Differential has the same root as difference and this class, *differential calculus*, is about calculating differences. Put another way, it is really about subtraction.

More specifically, the main question of differential calculus is how to relate changes in the output of a function to small changes in its inputs. For *differentiable* functions, we can accurately approximate the change in the output by scaling a small change in the input. The *scaling factor* is the *derivative*.

You've seen scaling factors before. Look at a map of a city, for example, and you'll see a scale, maybe 3 km/cm. But for a larger map, like one of the earth, there is no global scaling factor like this. Instead, the scaling factor varies from place to place and is usually a function of latitude. Even at a specific location, there is often not one, but infinitely many scaling factors, each depending on direction.

But for a *conformal* map (from the Latin *conformalis*, *of the same shape*) the scaling factor does not depend on direction. These maps preserve angles. The most well-know is probably the Mercator map. But we'll start with another.

2 Stereographic Projection

The worksheet below shows a stereogrphic projection of the earth between latitudes $-\pi/6$ and $\pi/2$, projected from the south pole.

Desmos link: <https://www.desmos.com/calculator/wy9zofmbe>

151: Stereographic Projection World

The worksheet here shows how the map is created. With the south pole S at the top of the spherical earth, a point P on the surface is mapped to the point P' in the plane of the equator lying on the line $\overrightarrow{SP}$.

- (a) Drag the sliders θ_1 (the longitude of P) and ϕ_1 (the latitude of P) to see

Learning outcomes:
Author(s):

Stereographic Projection

how the projection varies with latitude and longitude. Summarize your observations here.

Free Response:

Desmos3D link: <https://www.desmos.com/3d/lnhm28tzak>

152: Stereographic Projection

- (b) Stereographic projection maps circles on the sphere to circles in the plane. Experiment with the sliders θ_1 (Line 2), ϕ_1 (Line 4), r_1 (Line 6), t_1 (Line 8), and d (Line 10). Summarize your observations here.

Free Response:

Desmos3D link: <https://www.desmos.com/3d/dukkzdikzg>

152: Stereographic Projection 2

Question 6 Our aim is to see how the scale of the map varies from point to point.

- (a) First, suppose on the map below, the scale along the equator (red) is 100 km/cm. This means each centimeter on the map, measured not as a straight line distance but along an arc of the equator, represents 100 km along the earth's equator.
- (i) Use the number line below and simple arithmetic (no trigonometry) to approximate the scale at the south pole. Note the circles of latitude on the map are plotted equal intervals of latitude on the sphere.
- (ii) Use the number line below and simple arithmetic (no trigonometry) to approximate the scale at points on the orange circle of latitude in the direction due north.

Explain your method here and state a clear conclusion.

Free Response:

Desmos link: <https://www.desmos.com/calculator/wy9zofmbe>

151: Stereographic Projection World

- (b) To compute the actual scale in the direction along a meridian (ie. due north or due south) at a given distance from the center of the map (the south pole). We'll start by defining two parameters. The first is R , the radius of the earth in kilometers. The second is the radius of the equator on the map, measured in centimeters. We'll call this a .

- (c) Our goal now is to find an expression for a function

$$r = f(s), 0 \leq s \leq \pi R,$$

that takes as an input the actual distance of a point from the north pole (measured in km) and returns as an output the distance of its image from the center of the map (measured in cm).

The worksheet below shows a cross-section of the earth through the north and south poles. The point P is s km from the north pole.

Desmos link: <https://www.desmos.com/calculator/ncskufxsk3>

151: Stereo Proj 2D2

- (d) First express the distance $r_* = OQ$, measured in km, from point Q (in the plane of the equator) to the earth's center in terms of s and R .

$$r_* = R \tan\left(\frac{s}{2R}\right).$$

- (e) Next, scale the distance r_* (in km) to get the corresponding distance $r = f(s)$ on the map, measured in cm.

$$r = f(s) = a \tan\left(\frac{s}{2R}\right).$$

- (f) Now find an expression for the derivative dr/ds and interpret it as a scaling factor. Include units.

$$\frac{dr}{ds} = \frac{a}{2R} \sec^2\left(\frac{s}{2R}\right)$$

- (g) Express the derivative dr/ds in terms of r .

$$\frac{dr}{ds} = \frac{r^2 + a^2}{2Ra}$$

- (h) Find an expression for the scale on the map in the direction along the meridian at a point r cm from the map's center.

The scale is

$$\frac{2Ra}{r^2 + a^2} \frac{\text{km}}{\text{cm}}.$$

- (i) Check that your expression in part (h) has the correct units.

Stereographic Projection

- (j) *Use your result from part (h) to find the scale at the origin in the map above, where $R/a = 100$ km/cm.*
- (k) *Use your result from part (h) to find the scale in the direction of the meridians on the orange circle of latitude. Start by using the number line to approximate the ratio r/a .*



An Introduction to Differentiable Functions, Part 1

An introduction to what it means for a function to be differentiable.

Your precalculus classes were about functions, exponential, logarithmic, rational, trigonometric ... , all from a global perspective. Differential calculus is about these same functions, but from a *local* perspective. In one sense, the main idea of this class is to describe how *small* changes to the input of a function change the output. After all, *differential* has the same root as *difference* and *calculus* the same root as *calculate*. So *differential calculus* really means to calculate differences. Put more simply, this course is about subtraction.

One way to understand how small changes in the input of a function affect the output is to zoom in close to a point $(a, f(a))$ on the graph of a function $y = f(x)$. Do this and you'll most likely notice that the graph looks more and more like the graph of a linear function (ie. like a non-vertical line). If so, we say the function $y = f(x)$ is *differentiable* at $x = a$. And the *derivative* of the function, evaluated at $x = a$, written as

$$\left. \frac{dy}{dx} \right|_{x=a},$$

is just the slope of that line or the rate of change of the linear function.

When the independent variable is time, we can interpret the derivative as an instantaneous rate of change.

Example 1. The function

$$T = f(m), 0 \leq m \leq 60,$$

expresses the temperature (in °C) of a cup of coffee in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/fuftb4mq0k>

151: Cooling Coffee

- (a) Zoom in close enough to the point P above to make the graph look like a straight line. Then approximate the slope of this line. Include units. Then interpret the meaning of the slope in part (a) in the context of this scenario.

Learning outcomes:
Author(s):

Free Response:

- (b) Repeat part (a) for other temperatures by dragging the slider T_0 in Line 2 of the worksheet above. Make a table with two columns, temperature and slope. Then look for a relationship between the temperature T and the rate at which the temperature of the coffee is changing. Summarize your observations.

Free Response:

You should have found at the moment the coffee's temperature is 40°C , its temperature is decreasing at the rate of $2^\circ\text{C}/\text{min}$. We write this symbolically as

$$\left. \frac{dT}{dt} \right|_{T=40} = -2^\circ\text{C}/\text{min}.$$

This means that at the moment the temperature of the coffee is 40°C , the temperature is decreasing at the rate of $2^\circ\text{C}/\text{min}$. As long as we do not think too carefully about what exactly we mean by a moment in time, the meaning of the derivative is pretty clear.

But while this interpretation of the derivative as an instantaneous rate of change is often useful, it is fairly limited in scope. Perhaps the biggest problem is that the idea of slope does not extend to higher dimensions.

Example 2. A function $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, like the function

$$(u, v) = f(x, y) = (0.2(x^2 - y^2), 0.4xy),$$

maps the two-dimensional plane to itself. You can get a feel for the global action of this function by dragging the slider u in Line 2 from $u = 0$ to $u = 1$. This transforms the checkerboard of horizontal and vertical lines into two families of intersecting parabolas and gives a global perspective of the function.

Desmos link: <https://www.desmos.com/calculator/59nrhmvvka>

151: Complex Squaring Function 2

By focusing on the shaded square you can get a feeling for the local action of the function near point P . Here f maps this square to a square-like region with a different size and orientation. The image becomes more square-like by shrinking the side length s of the original square (do this by dragging Slider s in Line 4 toward $s = 0.1$).

The derivative of this function at the point P describes how the function acts near P . It can be described by just two numbers. One is a scaling (stretching) factor k that measures the ratio of the sides lengths of the original very small square and its square-like image,

$$k = \lim_{s \rightarrow 0} \frac{\text{side length of square-like image}}{s}.$$

The other number is an angle. It is the angle through which we would rotate the original very small square to make its sides parallel to its square-like image.

Question 7 (a) *Use the animation above to approximate the scaling factor k .*

(b) *Use the animation above to approximate the rotation angle θ .*

Fortunately for us, because our class is about functions $f : \mathbb{R} \rightarrow \mathbb{R}$ that map the real line (or a subset thereof) to itself, rotations do not come into play. So the local behavior of such functions, at least where they are differentiable, can be described by just one number. This number, the derivative, measures the local scaling factor. The factor converts a small change in the input to a function into an approximate change in its output.

Introduction, Part 2

Introduction to small changes and scaling factors.

This course is listed as Calculus I in the catalogue, but it really should be called Differential Calculus. *Differential* has the same root as *difference* and *calculus* the same root as *calculate*. So this class is really about calculating differences, or more simply put it's about subtracting. But it's about subtracting in the context of functions.

Pick a specific input to specific function and you'll likely find that the changes in the function's output are approximately proportional to *small* changes in the input. If so, we say that the function is *differentiable* at that input and we call the proportionality constant the *derivative*.

For example, let's look at the behavior of the function

$$A = f(s) = s^2, s \geq 0,$$

near the input $s = 5$. To emphasize the importance of units, let's define the input s to be the side length of a square measured in feet and the output $f(s)$ to be the area of that square, measured in square feet. The problem before us is to describe a simple relationship between a small change in the side length

$$\Delta s = s - 5$$

of the square and the change

$$\Delta A = f(s) - f(5)$$

in its area.

Question 8 (a) We'll first take a numerical approach and compute some small changes and their ratios. Fill in the missing entries in the table below.

s (ft)	$A = s^2$ (ft ²)	$\Delta s = s - 5$ (ft)	$\Delta A = s^2 - 25$ (ft ²)	$\Delta A / \Delta s$ (ft ² /ft)
4.9	24.01	-0.1	-0.99	9.9
4.99	24.9001	-0.01	-0.0999	9.99
5	25	0	0	—
5.01	25.1001	0.01	0.1001	10.01
5.1	26.01	0.1	1.01	10.1

Learning outcomes:
Author(s):

(b) The data in the table above suggests an approximate proportional relationship between ΔA and Δs . We can guess the scaling factor from the fifth column. As $s \rightarrow 5$ (as s approaches 5), it looks like the ratio $\Delta A/\Delta s$ approaches some number, what we call the constant of proportionality or the scaling factor.

i) What is that number? 10

ii) What are its units?

Free Response:

So for $\Delta s \sim 0$, we suspect that

$$\Delta A \sim 10\Delta s.$$

(c) The scaling factor is called the derivative, in this case of the function $A = s^2$, at the input $s = 5$. We write this as

$$\left. \frac{dA}{ds} \right|_{s=5} = 10.$$

(d) We could have taken an algebraic approach to determine the scaling instead. The idea is to first simplify the quotient $\Delta A/\Delta s$ as

$$\begin{aligned} \frac{\Delta A}{\Delta s} &= \frac{s^2 - 25}{s - 5} \\ &= \frac{(s + 5)(s - 5)}{s - 5} \\ &= s + 5 \text{ if } s \neq 5. \end{aligned}$$

So, for example, if $s = 4.99$, then

$$\frac{\Delta A}{\Delta s} = 4.99 + 5 = 9.99$$

as shown in the last column of the second row of the above table.

The advantage of this algebraic approach is that we can now compute the scaling factor as a limit:

$$\begin{aligned} \left. \frac{dA}{ds} \right|_{s=5} &= \lim_{s \rightarrow 5} \frac{\Delta A}{\Delta s} \\ &= \lim_{s \rightarrow 5} (s + 5) \\ &= 5 + 5 \\ &= 10. \end{aligned}$$

(d) We can also use the graph of the function $A = f(s) = s^2$ to interpret the ratios

$$\frac{\Delta A}{\Delta s} = \frac{f(s) - f(5)}{s - 5}$$

geometrically. Move the slider s in the demonstration below and describe

(i) how the line through the points P and Q is related to the ratio $\Delta A/\Delta s$ show on Line 2,

(ii) what happens to the line PQ as $s \rightarrow 5$, and

(iii) what happens to the line PQ when $s = 5$.

Free Response:

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/vz9ud5txva>

Continuing with the above demonstration,

(i) Open the Code folder in Line 3 and turn off the line PQ in Line 7.

(ii) Write an equation for the line through the point P with slope equal to the scaling factor in the line below and on Line 8 in the desmos worksheet:

$$A = L(s) = 25 + 10(s - 5).$$

(iii) Zoom in close enough to the point P to make the graph of the function $A = f(s)$ look like a line. How do the graph of the function and the graph of the line $A = L(s)$ compare in this close-up view?

Free Response:

(e) **Summary:**

- If we change the side of a square from a length of 5 feet to a length of $s \sim 5$ feet, then the area of the square changes by approximately

$$\Delta A = s^2 - 25 \sim 10\Delta s = 10(s - 5)$$

square feet. The scaling factor 10 has units $ft^2/ft = ft$.

This means that if we increase the side length of a square from 5 feet to 5.1 feet, for example, the area of the square would increase by approximately

$$\Delta A = (10 \text{ ft})(0.1 \text{ ft}) = 1 \text{ ft}^2.$$

- Zoom in close enough to the graph of the function $A = f(s) = s^2$ near the point $P(5, 25)$ and the graph looks like a line with slope equal to the scaling factor.
- We can compute the scaling factor as the limit

$$\left. \frac{dA}{ds} \right|_{s=5} = \lim_{s \rightarrow 5} \frac{f(s) - f(5)}{s - 5}.$$

- Suppose for example, we wanted to approximate the side length s of a square with area 25.06 ft^2 . Then

$$\Delta A = s^2 - 25 = 25.06 - 25 = 0.06.$$

And since

$$\Delta A \sim 10\Delta s = 10(s - 5),$$

$$0.06 \sim 10\Delta s.$$

So

$$\Delta s \sim 0.006$$

and a square with area 25.06 ft^2 has an approximate side length (measured in feet) of

$$s = 5 + \Delta s \sim 5.006.$$

Question 9 On a clear day with an unobstructed view (like you might have at the beach or in a hot air balloon), the distance to the horizon is limited by the curvature of the earth as illustrated in the demonstration below.

In fact, as long as you are not too high above the surface of the earth, the function

$$s = f(h) = 1.22\sqrt{h}, 0 \leq h \leq 20,000,$$

gives a good approximation to the distance to the horizon (the length of the red arc AT below, measured in miles) in terms of your height above the ground (the distance AP below, measured in feet).

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

Desmos activity available at

151:Distance to Horizon 1

Our aim is to approximate the change in the distance to the horizon (in miles) in terms of a small change in height (in feet) from a height of 25 feet.

(i) To start, what are the units of the constant 1.22 above? Explain how you know.

Free Response:

(ii) Go through a similar analysis as in parts (a)-(e) of Example 1, to approximate the change $\Delta s = s - f(25)$ in the distance to the horizon in terms of the change $\Delta h = h - 25$ in your height above the ground. Start by completing the column headings (with units) and the missing entries in the table below.

h (ft)	$s = 1.22\sqrt{h}$ (miles)	$\Delta h = h - 25$ (ft)	$\Delta s = f(h) - f(25)$ (miles)	$\Delta s / \Delta h$ (units?)
4.9^2				
4.99^2				
25		0	0	—
5.01^2				
5.1^2				

Question 10 This question is similar to the last, but suppose instead we are looking down on the earth from the space station or a rocket. Then the approximation to the distance to the horizon from the previous problem will not work.

So our first step is to find a function

$$s = f(h), h \geq 0,$$

that expresses the distance to the horizon (still measured in miles) in terms of our height above the earth's surface, now measured in miles instead of feet. We'll suppose the earth to be a perfect sphere of radius 3960 miles. The distance to the horizon is the arclength AT below, measured along the surface of the earth (you can think of this distance as the radius of the spherical disk visible to us). Our height is the distance AP .

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

(a) Find an expression for the above function.

Hint: Use right triangle $\triangle OTP$ to find an expression for the radian measure of angle $\angle POT$. Then use this angle to find an expression for the arclength AT .

Here are more details.

(i) Enter the two side lengths, measured in feet, in right triangle $\triangle OPT$ below.

$$OT = 3960$$

and

$$OP = h + 3960.$$

(ii) Let θ be the radian measure of $\angle TOP$. Write an equation with a trigonometric function of θ that relates the two lengths in part (i). Use the Math Editor tab to enter the trig function and the angle θ .

$$\cos \theta = \frac{3960}{h + 3960}.$$

(iii) Now solve the equation from part (ii) for θ in terms of h . Then use what you know about measuring arclength along a circle to find an expression for the function f . Use the Math Editor tab to help.

$$s = f(h) = 3960 \arccos\left(\frac{3960}{h + 3960}\right).$$

(b) Now suppose we are 165 miles above the surface of the earth and we wish to approximate how a small change in our altitude changes the distance to the horizon.

To do this, fill in the missing entries in the table below.

h (miles)	$s = f(h)$ (miles)	$\Delta h = h - 165$ (miles)	$\Delta s = f(h) - f(165)$ (miles)	$\Delta s / \Delta h$ (units?)
162				
163				
164				
165				
166				
167				
168				

(c) Do the data above suggest that the quotients $\Delta s / \Delta h$ approach some number as h approaches 165? If so, use the data to approximate that number. If not, explain why not.

(d) Make your own table similar to the one above to get a better approximation, correct to the nearest thousandth, to

$$\lim_{h \rightarrow 165} \frac{f(h) - f(165)}{h - 165}.$$

(e) Use your result from part (d), rounded to the nearest thousandth, to approximate Δs in terms of Δh and enter your result below.

$$\Delta s \sim 3.291\Delta h, \text{ for } \Delta h \sim 0.$$

(f) Explain the meaning of the scaling factor in parts (d) and (e). Be sure to include units in your explanation.

Question 11 The function

$$h = f(v), \quad 80 \leq v \leq 120,$$

expresses the height of a helicopter (measured in feet) in terms of its speed (measured in ft/sec).

Suppose

$$\left. \frac{dh}{dv} \right|_{v=100} = -20.$$

- (a) What are the units of the above derivative?
 - (b) Explain the meaning of the derivative using the language of small changes.
-

A Ferris Wheel

Relating small changes as you ride ferris wheel.

Question 12 This question is about a ferris wheel. The idea is this. After riding a ferris wheel for some time, the wheel gets stuck. Then it turns a through a small angle and gets stuck again. The question is to relate the small change in your height to the small change in the distance you move along your circular path as the wheel turns through this small angle.

- (a) The relationship between these small changes depends upon your position on the ferris wheel. The worksheet below illustrates this. Drag the slider m in Line 2 to see how the relationship between the small change Δh in your height and the small change in the distance Δs along your path changes as your position changes.

Desmos link: <https://www.desmos.com/calculator/vc1xpi5mbq>

151: Ferris Wheel 57

Summarize your observations.

Free Response:

- (b) Now let's focus on the particular position shown below at point P .

Desmos link: <https://www.desmos.com/calculator/9hlj4pkhni>

151: Ferris Wheel 58

Drag the slider n in Line 3 to see how the ratio $\Delta h/\Delta s$ changes as $\Delta s \rightarrow 0$.

Then drag the slider k in Line 4 and again the slider n in Line 3 to make Δs closer to 0. Summarize your observations about how the ratio $\Delta h/\Delta s$ changes as $\Delta s \rightarrow 0$.

Free Response:

- (c) The first step in relating the small changes Δh and Δs is to find a function

$$h = f(s), s \geq 0,$$

Learning outcomes:
Author(s):

A Ferris Wheel

that expresses your height (in feet) above the ground in terms of the distance (in feet) along your path since you boarded the wheel.

For this, we'll assume the wheel has a radius of 30 feet and that its center is 40 feet above the ground.

Desmos link: <https://www.desmos.com/calculator/q6wnvqhnpq>

151: Ferris Wheel 4

Use the figure above to find an expression for the function f . Input your expression in the box below to check your work.

$$h = f(s) = 40 - 30 \cos(s/30).$$

- (d) Now suppose the wheel stops after you have travelled just 15 feet along your path. The wheel then turns through a small angle and stops again. Our aim is to determine if the small change

$$\Delta h = f(s) - f(15)$$

in your height is approximately proportional to the small change

$$\Delta s = s - 15$$

in the distance you move along your path.

Desmos link: <https://www.desmos.com/calculator/uolmkpi5yr>

151: Ferris Wheel 3

- (i) First find a function

$$r = m(v)$$

that expresses the average rate of change in your height with respect to your distance travelled between distances $s = 15$ feet and $s = v$ feet. Enter that expression below with the appropriate domain.

$$r = m(v) = \frac{30 \cos(1/2) - 30 \cos(v/30)}{v - 15}, v \geq 0 \text{ and } v \neq 15.$$

- (ii) Enter your expression for the function $r = m(v)$ in Line 2 of the worksheet below. Use the table and the slider v in Line 4 to approximate the scaling factor that converts a small change

$$\Delta s = s - 15$$

in your distance into an approximation of the change

$$\Delta h = f(s) - f(15)$$

in your height.

Desmos link:

<https://www.desmos.com/calculator/cqprsr778n>

151: Ferris Wheel Table

- (iii) What are the units of the scaling factor?
- (iv) What is the approximate relation between Δh and Δs for small changes Δs ? Enter the scaling factor below, correct to 4 decimal places.

$$\Delta h \sim 0.4794\Delta s \text{ if } \Delta s \sim 0.$$

- (v) Approximate the change in your height if you move an additional 0.04 feet along your circular path from your position $s = 15$ feet.

Lambert's Equal-Area Map, Part 1

Lambert's equal-area map.

1 Lambert's Equal-Area Map

You probably saw the idea of a scaling factor long before taking calculus. Look at a map of Seattle and you'll see a scale, probably near the bottom. It might be 2 miles/inch, or something like that. But you will not see such a scale on a map of the earth for the simple reason that one does not exist. The scale varies from place to place and is usually a function of latitude. Even at a specific location, there is often not one but infinitely many scaling factors that depend on the direction in which you head.

The worksheet below shows a way to create a map of the earth by wrapping a cylinder around its equator. The map sends each point of the earth to its nearest point on the cylinder (ie. straight out, directly away from the earth's axis of rotation). Unwrapping the cylinder gives Lambert's equal-area map, from the earth to the points in a rectangle.

Exploration 13 Drag the slider in Line 6 from $r_1 = 0$ to $r_1 = 1$ to unwrap the cylinder. Drag sliders θ_1, ϕ_1 in Lines 2 and 4 to vary the longitude and latitude of the point on the sphere.

Desmos3D link: <https://www.desmos.com/3d/ehyvvttdoe>

152: Mercator 0

Suppose the earth has radius one. Lambert's map sends the circle of latitude ϕ (the angle between the plane of the equator and the segments from the sphere's center to points on the circle) to a circle on the cylinder with unit radius. So the map stretches the circle of latitude ϕ , with radius $\cos \phi$, by the factor

$$\lambda_p = \frac{1}{\cos \phi}.$$

This is the scaling factor along the circle in the east-west direction at latitude ϕ .

Learning outcomes:
Author(s):

In the north-south direction the scaling factor is also a function of latitude. The key point is that the map sends a point on the circle of latitude ϕ to signed height

$$h = f(\phi) = \sin \phi$$

above the plane of the equator. So the scaling factor at latitude ϕ in the north-south direction is

$$\lambda_m = \frac{dh}{d\phi} = \cos \phi.$$

Drag the slider m in Line 2 of the worksheet below to visualize the scaling factor.

Desmos link: <https://www.desmos.com/calculator/kbryuogdaj>

151: Lambert Map Cosine Function

Desmos link: <https://www.desmos.com/calculator/vwxuwjvxyy>

151: Lambert Map 2

2 The Mercator Map

Watch this video.

[Equal Earth Projection](#)

3 Loxodromes

Rhumb Lines on the Sphere

For a curve on a sphere of radius a that cuts the meridians at a fixed angle α ,

$$d\theta = \frac{\tan \alpha d\phi}{\cos \phi},$$

where ϕ is the latitude and θ the longitude. Taking $\theta = 0$ at the equator where $\phi = 0$,

$$\begin{aligned} \theta &= \tan \alpha \int_0^{\phi^*} \sec \phi^* d\phi^* \\ &= \tan \alpha \operatorname{arctanh}(\sin \phi). \end{aligned}$$

So

$$\sin \phi = \tanh(k\theta),$$

where $k = \cot \alpha$.

Because $-\pi/2 < \phi < \pi/2$,

$$\cos \phi = \frac{1}{\cosh(k\theta)},$$

a parameterization of the rhumb line by longitude is

$$(x, y, z) = \left(\frac{a \cos \theta}{\cosh(k\theta)}, \frac{a \sin \theta}{\cosh(k\theta)}, a \tanh(k\theta) \right).$$

And a parameterization by latitude,

$$(x, y, z) = (a \cos \phi, a \cos \phi, a \sin \phi)$$

Desmos3D link: <https://www.desmos.com/3d/6cqnjsuew7>

152: Mercator 1

Desmos link: <https://www.desmos.com/calculator/rbcbxdrvcp>

Geogebra link: <https://www.geogebra.org/m/mvp9zvge>

152: Mercator

Shadow of a Building, Part 1

Small changes in the length of a building's shadow.

Question 14 This question is about how a small change in the inclination angle of the setting sun changes the length of a building's shadow.

The worksheet below suggests how the length the shadow of a 50-foot tall building changes as the inclination angle of the setting sun changes.

Note, the inclination angle is the angle the sun's rays make with the horizon.

Desmos link: <https://www.desmos.com/calculator/vsnchdgl00>

151: Shadow 1

- (a) Summarize your observations about the relationship between the two changes using the worksheet above. In particular, for what inclination angles does a small change in the angle give a comparatively small change in the shadow's length? A comparatively large change?

Free Response:

- (b) As the sun sets, does its inclination angle increase or decrease? Does the shadow's length increase or decrease?

Free Response:

- (c) What does your response to part (b) tell you about the average rate of change in the length of the building's shadow with respect to the setting sun's inclination angle? Is this rate of change positive or negative? Explain.

Free Response:

- (d) Find a function

$$s = f(\theta), 0 < \theta \leq \pi/2,$$

that expresses the length of the shadow (in feet) in terms of the sun's inclination angle (in radians).

$$s = f(\theta) = 50 \cot \theta, 0 < \theta \leq \pi/2.$$

- (e) Find the exact inclination angle θ_0 when the shadow is 100 feet long.

$$\theta_0 = \arctan(1/2).$$

- (f) Find an expression for the function $r = m(\alpha)$ that expresses average rate of change of the function f between inclination angles θ_0 and α for the angle θ_0 in part (e). Include the domain.

$$r = m(\alpha) = \frac{50(\cot \alpha - 2)}{\alpha - \arctan(1/2)}, \quad 0 \leq \alpha < \pi/2 \text{ and } \alpha \neq \arctan(1/2).$$

- (g) Approximate the value of the scaling factor

$$\left. \frac{ds}{d\theta} \right|_{\theta=\arctan(1/2)} = \lim_{\alpha \rightarrow \arctan(1/2)} \left(\frac{50(\cot \alpha - 2)}{\alpha - \arctan(1/2)} \right)$$

numerically. This factor converts a small change

$$\Delta\theta = \theta - \theta_0 = \theta - \arctan(1/2)$$

in the inclination angle (measured in radians) of the setting sun into an approximation of the change

$$\Delta s = s - 100$$

in the shadow's length (measured in feet).

Approximate the scaling factor as follows:

- (i) First find a rough approximation by looking at the picture above where the tips of the shadows are shown at equal changes

$$\Delta\theta = \frac{\pi}{2n} = \frac{\pi}{68} \text{ (see Line 17 of the worksheet)}$$

in the inclination angle. Be sure to include units. Explain your reasoning.

Free Response:

- (ii) Next find a better approximation by inputting your expression for the function $r = m(\alpha)$ in Line 2 of the worksheet. Then reveal the average rate of change table in Line 3 by clicking the arrow at the left of the line. Summarize your findings.

Free Response:

- (h) Setting aside the particular value $\theta_0 = \arctan(1/2)$ of the inclination angle for the moment, simplify the expression

$$r = m(\alpha) = \frac{50(\cot \alpha - \cot \theta_0)}{\alpha - \theta_0}$$

for the average rate of change function. Do this by expressing the cotangent function in terms of sine and cosine. Then do some algebra and use a trig identity.

$$r = m(\alpha) = \frac{-50 \sin(\alpha - \theta_0)}{(\sin \alpha \sin \theta_0)(\alpha - \theta_0)}.$$

- (i) Use the result of part (h) to find an expression for the scaling factor

$$\left. \frac{ds}{d\theta} \right|_{\theta=\theta_0} = \lim_{\alpha \rightarrow \theta_0} \frac{-50 \sin(\alpha - \theta_0)}{(\sin \alpha \sin \theta_0)(\alpha - \theta_0)}.$$

You'll need to use the fact that

$$\lim_{\phi \rightarrow 0} \frac{\sin \phi}{\phi} = 1.$$

The scaling factor is

$$\begin{aligned} \left. \frac{ds}{d\theta} \right|_{\theta=\theta_0} &= \lim_{\alpha \rightarrow \theta_0} \frac{-50 \sin(\alpha - \theta_0)}{(\sin \alpha \sin \theta_0)(\alpha - \theta_0)} \\ &= -\frac{50}{\sin^2 \theta_0}. \end{aligned}$$

- (j) Use the result of part (i) to find the exact value of the scaling factor when the shadow is 100 feet long. Include units. Compare with your approximation in part (g).
- (k) Approximate the change

$$\Delta s = f(\theta) - 100$$

in the shadow's length (measured in feet) in terms of the small change

$$\Delta \theta = \theta - \arctan(1/2)$$

in the inclination angle (measured in radians).

$$\Delta s \sim -250\Delta\theta.$$

- (1) A student explains the scaling factor

$$\left. \frac{ds}{d\theta} \right|_{s=100} = -250 \frac{\text{ft}}{\text{rad}}$$

by claiming that when the shadow is 100 feet long and the inclination angle of the sun decreases by 1 radian, the shadow's length increases by about 250 feet.

Is this correct? If so, explain why. If not explain why not and correct the statement.

Free Response:

Looking Down on the Moon

Looking at a spherical cap on the lunar surface.

Question 15 This question is about looking at the moon from Artemis II.

The idea is that the astronauts can see only a portion of the moon from any altitude above its surface. We wish to analyze how a small change in altitude changes the radius of the spherical cap in view. The radius is measured on the moon's surface. It is the orange arc shown below. Drag the slider h (altitude, in thousands of miles) below to see how small changes in the altitude affect the radius of the spherical cap.

Desmos3D link: <https://www.desmos.com/3d/paqpozwxm>

151: Spaceship above earth

Here is a two-dimensional cross-section of that picture.

Desmos link: <https://www.desmos.com/calculator/b5c4c7ba17>

151: Distance to Horizon 1

Drag point P_2 toward P_1 in the worksheet below to change the change Δh in the altitude. Keep your eye on the change in the radius Δs of the visible spherical cap and the ratio $\Delta s/\Delta h$ of small changes.

Desmos link: <https://www.desmos.com/calculator/1oj7ofkmda>

151: Distance to Horizon 2

(a) Find a function

$$s = f(h), h \geq 0$$

that expresses the radius of the visible spherical cap (in miles) in terms of the altitude above the surface (in miles). Take the moon to be a perfect sphere of radius 1000 miles.

$$s = f(h) = 1000 \arccos(1000/(1000 + h)), h \geq 0.$$

Click the Hint button above for help.

Learning outcomes:
Author(s):

Looking Down on the Moon

Hint: Use right triangle $\triangle OTP$ to find an expression for the radian measure of angle $\angle POT$. Then use this angle to find an expression for the arclength AT . Here are more details.

- (i) Enter the two side lengths, measured in feet, in right triangle $\triangle OPT$ below.

$$OT = 1000$$

and

$$OP = h + 1000.$$

- (ii) Let θ be the radian measure of $\angle TOP$. Write an equation with a trigonometric function of θ that relates the two lengths in part (i). Use the Math Editor tab to enter the trig function and the angle θ .

$$\cos \theta = \frac{1000}{h + 1000}.$$

- (iii) Now solve the equation from part (ii) for θ in terms of h . Then use what you know about measuring arclength along a circle to find an expression for the function f . Use the Math Editor tab to help.

$$s = f(h) = 1000 \arccos\left(\frac{1000}{h + 1000}\right).$$

- (b) Now suppose we are 250 miles above the lunar surface. Fill in the missing entries in the table below to approximate the scaling factor that converts a small change

$$\Delta h = h - 250$$

in altitude to a good approximation of the change

$$\Delta s = f(h) - f(250)$$

in the radius of the visible spherical cap.

h (miles)	$s = f(h)$ (miles)	$\Delta h = h - 250$ (miles)	$\Delta s = f(h) - f(250)$ (miles)	$\Delta s / \Delta h$ (units?)
247				
248				
249				
250				
251				
252				
253				

- (c) Do the data above suggest that the quotients $\Delta s/\Delta h$ approach some number as h approaches 250? If so, use the data to approximate that number. If not, explain why not.
- (d) Make your own table similar to the one above to get a better approximation, correct to the nearest thousandth, of the scaling factor

$$\lim_{h \rightarrow 250} \frac{f(h) - f(250)}{h - 250}.$$

- (e) Use your result to the nearest thousandth, to approximate Δs in terms of Δh and enter your result below.

$$\Delta s \sim 1.067\Delta h, \text{ for } \Delta h \sim 0.$$

- (f) We now find the exact scaling factor algebraically.

- (i) First find an expression for the function

$$h = f^{-1}(s)$$

that expresses the altitude in terms of the radius of the visible spherical cap, both measured in miles. Include the appropriate domain.

- (ii) Then find a simplified expression for the average rate of change of the function f^{-1} between radii $s = a$ and $s = b$ miles. You should get this:

$$\begin{aligned} \frac{\Delta h}{\Delta s} &= \frac{f^{-1}(b) - f^{-1}(a)}{b - a} \\ &= \left(\frac{1000}{\cos(\frac{a}{1000}) \cos(\frac{b}{1000})} \right) \left(\frac{\cos(\frac{a}{1000}) - \cos(\frac{b}{1000})}{b - a} \right) \\ &= \left(\frac{-1000}{\cos(\frac{a}{1000}) \cos(\frac{b}{1000})} \right) \left(\frac{2 \sin(\frac{a+b}{2000}) \sin(\frac{a-b}{2000})}{b - a} \right) \\ &= \left(\frac{-2000 \sin(\frac{a+b}{2000})}{\cos(\frac{a}{1000}) \cos(\frac{b}{1000})} \right) \left(\frac{\sin(\frac{a-b}{2000})}{b - a} \right) \end{aligned}$$

Small Changes

We explore how small changes to the input of a function change the output.

The main idea of differential calculus is to approximate the change in the output of a function in terms of a small change in the input. For some functions, called *differentiable*, the change in the output is approximately proportional to the (small) change in the input. The proportionality factor is called the derivative. In this chapter we explore this idea.

1 Odometer Readings

Example 3. The graph of the function

$$s = f(t), 0 \leq t \leq 2,$$

that expresses the trip odometer reading (measured in miles) on your car in terms of the number of hours past noon during a two-hour trip is shown below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/iw69lr1bc>

Desmos activity available at

151: Odometer

Our goal is to approximate the car's speed at 12:30pm in three ways:

- (1) geometrically, using the above graph as is.
- (2) geometrically, by zooming in on the above graph.
- (3) arithmetically, using the specific expression for the function f .

(a) Start by using the graph above to describe how the speed of the car varies over the two-hour period. Explain your reasoning. Then play the Slider u in Line 2 and use the animation of the motion to check if your description was accurate. Explain.

Learning outcomes:
Author(s):

Small Changes

(b) Set the slider in Line 6 to $n = 20$. Use *only* the graph (with $u = 2$) to create a table with five columns showing the values of t , s (approximate this), $\Delta t = t - 0.5$, $\Delta s = f(t) - f(0.5)$, and $\Delta s/\Delta t$. Include units in the heading of each column. The table should include five rows, with $t = 0.3, 0.5, \dots 0.7$.

t (hours)	$s = f(t)$ (miles)	$\Delta t = t - 0.5$ (hrs)	$\Delta s = f(t) - f(0.5)$ (miles)	$\Delta s/\Delta t$ (miles/hr)
0.3				
0.4				
0.5				
0.6				
0.7				

(c) Explain the meaning of the fifth column in the table of part (b). What do the entries in this column suggest about the speed of the car at 12:30pm?

(d) Now we'll use the fact that

$$s = f(t) = 48t^2 - 16t^3, 0 \leq t \leq 2,$$

to construct another table like the one in part (b). Do this by finding expressions for

$$\Delta s = f(t) - f(0.5)$$

and

$$r = g(t) = \frac{\Delta s}{\Delta t} = \frac{f(t) - f(0.5)}{t - 0.5},$$

both in terms of t (and *not* Δt). Use these functions to fill in the missing entries in the table below.

t (hours)	$s = f(t)$ (miles)	$\Delta t = t - 0.5$ (hrs)	$\Delta s = f(t) - f(0.5)$ (miles)	$\Delta s/\Delta t$ (miles/hr)
0.49				
0.499				
0.4999				
0.5				
0.5001				
0.501				
0.51				

(e) Does your table from part (d) suggest that the ratios $\Delta s/\Delta t$ approach some number as $t \rightarrow 0.5$? If so, what would be your guess for the exact value of this number? What are its units? What is its meaning?

(f) Activate the folder "Graph of average speed" on Line 9.

(i) Use the graph to check some your entries in the fifth column of your table from part (d). Explain.

- (ii) How is the line PQ , through the fixed point $P(0.5, 10)$ and the variable point $Q(t, s)$ on the graph of the function $r = g(t)$ related to the ratio $\Delta s / \Delta t$?
- (iii) Describe what happens to the line PQ as point Q approaches point P .
- (g) For a quicker way to approximate car's speed at 12:30pm, zoom in sufficiently close to point P in the graph above to make the graph of $s = f(t)$ look like a line. Use the coordinates of point P and a second point in the window far away from P to estimate the car's speed at 12:30pm. Explain your method.
- (h) Summarize your conclusions by comparing your three estimates for the car's speed at 12:15pm. Which estimate do you think is most accurate? Least accurate?

Example 4. This is a continuation of the previous example where we'll algebraically compute the exact speed of the car at 12:30pm, using the odometer function

$$s = f(t) = 48t^2 - 16t^3, 0 \leq t \leq 2.$$

The idea is to first find an algebraic expression for the car's average speed between time t and time $t = 0.5$ hours past noon. Then we'll evaluate the limit of this average speed as $t \rightarrow 0.5$ to find the (instantaneous) speed at 12:30pm.

First we'll find the average speed between time t and time $t = 0.5$.

- (a) Explain in general how to compute a car's average speed over some time interval. What do you need to know? What is the computation? Make up your own specific example.
- (b) Now for our particular odometer function above, the average speed $v_{\text{avg}}(t)$, measured in miles/hour, between time t and time $t = 0.5$ is

$$\begin{aligned} v_{\text{avg}}(t) &= \frac{f(t) - f(0.5)}{t - 0.5} \\ &= \frac{48t^2 - 16t^3 - 10}{t - 0.5} \\ &= \frac{(2t - 1)(-8t^2 + 20t + 10)}{t - 0.5} \\ &= -16t^2 + 40t + 20 \text{ if } t \neq 0.5. \end{aligned}$$

The key step in the computation above is in the third line. How did we know $2t - 1$ was a factor of

$$f(t) - f(0.5) = 48t^2 - 16t^3 - 10?$$

The reason is that $t = 0.5$ is a root of the polynomial $f(t) - f(0.5)$ and therefore $t - 0.5$ is a factor. And so

$$2(t - 0.5) = 2t - 1$$

is also a factor. Then we can use long division to find the quotient.

(c) Show the steps in the long division.

(d) The final step in computing the car's speed v (in miles/hour) at 12:30pm is to evaluate the limit of these average speeds as $t \rightarrow 0.5$. We get

$$\begin{aligned} v &= \lim_{t \rightarrow 0.5} (-16t^2 + 40t + 20) \\ &= 36. \end{aligned}$$

(e) Here's another way to simplify the average speed in part (b). Fill in the missing steps.

$$\begin{aligned} v_{\text{avg}}(t) &= \frac{f(t) - f(0.5)}{t - 0.5} \\ &= \frac{(48t^2 - 16t^3) - (48(0.5)^2 - 16(0.5)^3)}{t - 0.5} \\ &= \frac{(48t^2 - 48(0.5)^2) - (16t^3 - 16(0.5)^3)}{t - 0.5} \\ &= \frac{48(t^2 - (0.5)^2) - 16(t^3 - (0.5)^3)}{t - 0.5} \\ &= \frac{48(t - 0.5)(t + 0.5) - 16(t - 0.5)(t^2 + 0.5t + 0.25)}{t - 0.5} \\ &= 48(t + 0.5) - 16(t^2 + 0.5t + 0.25) \text{ if } t \neq 0.5. \end{aligned}$$

(f) Use the above expression for the average speed function to compute the (instantaneous) speed of the car at 12:30pm by evaluating the appropriate limit.

(g) Sketch *by hand* a graph of the average speed function $y = v_{\text{avg}}(t)$ over the appropriate domain. Be sure also to state this function's domain.

2 A Projectile

Example 5. Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/l4fknr0hpl>

Desmos activity available at

151: Projectile

3 The Falling Ladder, Part 1

Example 6. The top end of a ten-foot ladder leans against a vertical wall and the bottom end rests on the horizontal floor. We analyze how a small change in the distance between the wall and the bottom of the ladder affects the height of the ladder's top above the floor.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/dvyuifyyg4>

Desmos activity available at

151: Ladder 1B

Question 16 (a) The slider s in Line 1 of the demonstration above controls the distance between the wall and the bottom of the ladder, measured in feet. Use the slider s to describe qualitatively how a small change in s changes the height h (also measured in feet) of the ladder's top end above the floor.

(i) Do the small changes have the same or opposite signs?

(ii) At what positions of the ladder does a small change in s result in a comparatively large change in h ?

(b) Now let's focus on the particular position of the ladder where the bottom end is $s = 8$ feet from the wall. For this, turn on the "one position" folder in Line 3.

(i) Drag the slider s close to $s = 8$ and use the coordinates of the endpoints of the ladder to construct a table with five columns showing the values of s , h , $\Delta s = s - 8$, $\Delta h = f(s) - f(8)$, and $\Delta h/\Delta s$. Include units in the heading of each column. The table should include seven rows, with $s = 7.7, 7.8, \dots, 8.2, 8.3$. Here $h = f(s)$ is the function described in part (ii) below.

(ii) Find a function

$$h = f(s), 0 \leq s \leq 10,$$

that expresses the height of the ladder's top end above the ground (in feet) in terms of the distance of its bottom end from the wall (in feet).

(iii) Use your function f to construct another table, like the one in part (i), with $s = 7.9, 7.99, 7.999, 8, 8.0001, 8.01, 8.1$. Do this by finding expressions for

$$\Delta h = f(s) - f(8)$$

and

$$r = g(s) = \frac{\Delta h}{\Delta s} = \frac{f(s) - f(8)}{s - 8},$$

both in terms of s (and not Δs).

(iv) Does your table from part (iii) suggest that the ratios $\Delta h/\Delta s$ approach some number as $s \rightarrow 8$? If so, what would be your guess for the exact value of this limit? What are its units? What is its meaning?

(v) Activate the folder “graph of function” on Line 8. How is the line PQ , through the fixed point $P(8, 6)$ and the variable point $Q(s, h)$ on the graph of the function $h = f(s)$ related to the ratios $\Delta h/\Delta s$?

(vi) Change the bounds for s in Line 1 to run between $s = 7.9$ and $s = 8.1$. Then activate the folder “graph: average rate of change function” on Line 18. Move the slider s and use the graph of the function $r = g(s)$ to check your computations in part (iii).

(vii) Use the result of part (iv) to write an approximation for the change in height

$$\Delta h = h - 8$$

in terms of the change

$$\Delta s = s - 8.$$

(viii) The graph of the function $h = f(s)$ suggests another, geometric way to find the proportionality constant (of part (vii)) that relates Δh to Δs . Explain how.

4 The Falling Ladder, Part 2

Example 7. A tree leans precariously with its trunk making an angle of $\phi = \pi/6$ radians with the ground. One end of a ten-foot ladder leans against the trunk, the other rests on the horizontal ground. We analyze how a small change in the distance between the bottom of the ladder and the base of the trunk changes the distance between the top of the ladder and the base of the trunk.

Access Geogebra interactives through the online version of this text at

Geogebra link: <https://www.geogebra.org/m/qmke5y7x>

We'll let t be the distance between the top of the ladder and the base of the trunk (measured in feet) and s the distance between the bottom of the ladder and the base of the trunk (also measured in feet).

The slider θ in the demonstration above controls the angle that the ladder makes with the ground, but this angle does not come into play in our problem.

(a) Use the slider θ to describe qualitatively how a small change in s (the length of segment GC) changes t (the length of segment GB):

(i) For what positions of the ladder do these small changes have the same signs? Opposite signs?

(ii) For what positions of the ladder does a small change in s result in a comparatively large change in t ?

(b) Now let's focus on the particular position of the ladder when the bottom end C is 16 feet from the trunk's base and the top end D is about 8 feet from the base *as illustrated above*. Our first step is to find a function

$$t = f(s)$$

that expresses t in terms of s for values of s and t near $s = 16$ and $t = 8$ respectively.

To do this, first use the law of cosines to write an equation relating s and t . Then *complete the square* to solve this equation for t in terms of s to find the function $t = f(s)$. Keep in mind that when $s = 16$, we must have $t \sim 8$.

Hint: Here is an outline of the steps to check your work:

(i) Use the law of cosines to relate s and t .

$$s^2 + t^2 - \sqrt{3}st = 100.$$

(ii) Solve the above equation for t in terms of s as follows:

First rewrite the equation with the two terms with t as a factor on the left side and the other two terms on the right.

$$t^2 - \sqrt{3}st = 100 - s^2.$$

Then complete the square by adding the same perfect square to each side.

$$t^2 - \sqrt{3}st + \left(\frac{\sqrt{3}s}{2}\right)^2 = 100 - s^2 + \left(\frac{\sqrt{3}s}{2}\right)^2.$$

Then factor the LHS and simplify the RHS.

$$\left(t - \frac{\sqrt{3}s}{2}\right)^2 = 100 - \frac{s^2}{4}.$$

Next, solve for t in terms of s .

$$t = \frac{\sqrt{3}}{2}s \pm \sqrt{100 - \frac{s^2}{4}}$$

Finally, make the correct choice of $\pm$ to solve for t in terms of s , when s is near 16 and t near 8.

$$t = \frac{\sqrt{3}}{2}s - \sqrt{100 - \frac{s^2}{4}}.$$

(c) Use your function from part (b) to find an expressions for

$$\Delta t = f(s) - f(16)$$

and for the function

$$r = g(s) = \frac{\Delta t}{\Delta s} = \frac{f(s) - f(16)}{s - 16}.$$

Explain what the output of the function g measures. What are its units?

(d) Use the results of part (c) to construct a table with five columns showing the values of s , t , $\Delta s = s - 16$, $\Delta t = f(s) - f(16)$, and $\Delta t/\Delta s$. Include units in the heading of each column. The table should include seven rows, with $s = 15.9, 15.99, 15.999, 16, 16.001, 16.01, 16.1$.

(e) Does your table from part (d) suggest that the ratio $\Delta t/\Delta s$ approaches some number as $s \rightarrow 16$? If so, approximate the value of this number. What are its units?

(f) Check the box “GraphofRelation” in the demonstration above and explain how the line EF is related to part (d).

(g) Use the result of part (e) to write an approximation for the change

$$\Delta t = f(s) - f(16)$$

in terms of the change

$$\Delta s = s - 16$$

for values of s near 16. Use this approximation to estimate the distance between the top of the ladder and the base of the trunk when the bottom of the ladder is 16.4 feet from the trunk’s base. Compare your approximation with the exact distance.

5 Riding a Ferris Wheel

Suppose you ride a ferris wheel

The Derivative as a Scaling Factor

Interpreting the derivative as a scaling factor.

Example 8. This question is about how a small change in the inclination angle of the setting sun changes the length of a building's shadow.

We'll suppose the function

$$s = f(\theta), \quad 0 < \theta \leq \pi/2$$

expresses the length of the shadow of a 50-foot tall building as the sun sets in terms of the sun's inclination angle, measured (in radians) relative to the horizontal.

- (a) One of the following is correct. Which one? Why?

$$\left. \frac{ds}{d\theta} \right|_{s=100} = 250$$

or

$$\left. \frac{ds}{d\theta} \right|_{s=100} = -250$$

- (b) What are the units of the correct derivative above?
 (c) Explain the meaning of the derivative as a scaling factor.

Explanation. (a) As the sun sets the inclination angle decreases and the shadow's length increases. Therefore, the derivative

$$\frac{ds}{d\theta} = \lim_{\Delta\theta \rightarrow 0} \frac{\Delta s}{\Delta\theta}$$

is negative.

- (b) The units of the derivative are ft/radian (ie. the units of s divided by the units of θ).
 (c) The scaling factor is

$$\left. \frac{ds}{d\theta} \right|_{s=100} = -250 \frac{\text{ft}}{\text{rad}} = -2.50 \frac{\text{ft}}{0.01 \text{ rad}}.$$

This means when we multiply a small change in the inclination angle by the scaling factor we get a good approximation to the change in the

Learning outcomes:
 Author(s):

shadow's length. So, for example, when the shadow is 100 feet long and the inclination angle decreases by 0.01 radians the shadow's length increases by approximately 2.5 feet.

Note: It would be incorrect to say that if the inclination angle decreases by 1 radians the shadow's length increases by approximately 250 feet. The problem here would be that 1 radian too large of a change in the angle for the scaling factor to give a good approximation to the change in the shadow's length.

Example 9. The function

$$v = f(h), 0 \leq h \leq 100,$$

expresses the speed (in meters/sec) of a rock dropped from rest near the surface of the planet Krypton in terms of its height above the surface (in meters).

- (a) One of the following is more likely to be correct? Why?

$$\left. \frac{dv}{dh} \right|_{h=20} = 5$$

or

$$\left. \frac{dv}{dh} \right|_{h=20} = -5$$

- (b) What are the units of the correct derivative above?
 (c) Explain the meaning of the derivative as a scaling factor.

Explanation. (a) As the rock falls its speed increases as its height above the surface decreases. The derivative

$$\frac{dv}{dh} = \lim_{\Delta h \rightarrow 0} \frac{\Delta v}{\Delta h}$$

is therefore negative.

- (b) The units of the derivative are

$$\frac{\text{units of } v}{\text{units of } h} = \frac{\text{meters/sec}}{\text{meter}}.$$

- (c) The scaling factor is

$$\left. \frac{dv}{dh} \right|_{h=20} = -5 \frac{\text{meters/sec}}{\text{meter}}.$$

This means when we multiply a small change in the rock's height by the scaling factor we get a good approximation to the change in the rock's speed. So, for example, the rock's speed increases by approximately 5 m/sec as it falls from a height of 20 meters to a height of 19 meters.

The Derivative as a Scaling Factor

Note: In general, you should **not** simplify the units of a derivative. Doing so almost always obscures its meaning. So here, it would not help to write sec^{-1} for the units of the derivative.

It would also be incorrect to say for *every* meter the rock falls from a height of 20 meters, its speed increases by about 5 metes/sec. The problem here is that the word *every* implies the scaling factor is constant and would work for large as well as small changes. This is not correct. In fact, the scaling factor here, like most derivatives, continuously changes.

Stereographic Projection

Maps of the earth, stereographic projection.

1 Introduction to Differential Calculus

Differential has the same root as difference and this class, *differential calculus*, is about calculating differences. Put another way, it is really about subtraction.

More specifically, the main question of differential calculus is how to relate changes in the output of a function to small changes in its inputs. For *differentiable* functions, we can accurately approximate the change in the output by scaling a small change in the input. The *scaling factor* is the *derivative*.

You've seen scaling factors before. Look at a map of a city, for example, and you'll see a scale, maybe 3 km/cm. But for a larger map, like one of the earth, there is no global scaling factor like this. Instead, the scaling factor varies from place to place and is usually a function of latitude. Even at a specific location, there is often not one, but infinitely many scaling factors, each depending on direction.

But for a *conformal* map (from the Latin *conformalis*, *of the same shape*) the scaling factor does not depend on direction. These maps preserve angles. The most well-know is probably the Mercator map. But we'll start with another.

2 Stereographic Projection

The worksheet below shows a stereogrphic projection of the earth between latitudes $-\pi/6$ and $\pi/2$, projected from the south pole.

Desmos link: <https://www.desmos.com/calculator/wy9zofmbe>

151: Stereographic Projection World

The worksheet here shows how the map is created. With the south pole S at the top of the spherical earth, a point P on the surface is mapped to the point P' in the plane of the equator lying on the line $\overrightarrow{SP}$.

- (a) Drag the sliders θ_1 (the longitude of P) and ϕ_1 (the latitude of P) to see

Learning outcomes:
Author(s):

Stereographic Projection

how the projection varies with latitude and longitude. Summarize your observations here.

Free Response:

Desmos3D link: <https://www.desmos.com/3d/lnhm28tzak>

152: Stereographic Projection

- (b) Stereographic projection maps circles on the sphere to circles in the plane. Experiment with the sliders θ_1 (Line 2), ϕ_1 (Line 4), r_1 (Line 6), t_1 (Line 8), and d (Line 10). Summarize your observations here.

Free Response:

Desmos3D link: <https://www.desmos.com/3d/dukkzdikzg>

152: Stereographic Projection 2

Question 17 Our aim is to see how the scale of the map varies from point to point.

- (a) First, suppose on the map below, the scale along the equator (red) is 100 km/cm. This means each centimeter on the map, measured not as a straight line distance but along an arc of the equator, represents 100 km along the earth's equator.
- (i) Use the number line below and simple arithmetic (no trigonometry) to approximate the scale at the south pole. Note the circles of latitude on the map are plotted equal intervals of latitude on the sphere.
- (ii) Use the number line below and simple arithmetic (no trigonometry) to approximate the scale at points on the orange circle of latitude in the direction due north.

Explain your method here and state a clear conclusion.

Free Response:

Desmos link: <https://www.desmos.com/calculator/wy9zofmbe>

151: Stereographic Projection World

- (b) To compute the actual scale in the direction along a meridian (ie. due north or due south) at a given distance from the center of the map (the south pole). We'll start by defining two parameters. The first is R , the radius of the earth in kilometers. The second is the radius of the equator on the map, measured in centimeters. We'll call this a .

- (c) Our goal now is to find an expression for a function

$$r = f(s), 0 \leq s \leq \pi R,$$

that takes as an input the actual distance of a point from the north pole (measured in km) and returns as an output the distance of its image from the center of the map (measured in cm).

The worksheet below shows a cross-section of the earth through the north and south poles. The point P is s km from the north pole.

Desmos link: <https://www.desmos.com/calculator/ncskufxsk3>

151: Stereo Proj 2D2

- (d) First express the distance $r_* = OQ$, measured in km, from point Q (in the plane of the equator) to the earth's center in terms of s and R .

$$r_* = R \tan\left(\frac{s}{2R}\right).$$

- (e) Next, scale the distance r_* (in km) to get the corresponding distance $r = f(s)$ on the map, measured in cm.

$$r = f(s) = a \tan\left(\frac{s}{2R}\right).$$

- (f) Now find an expression for the derivative dr/ds and interpret it as a scaling factor. Include units.

$$\frac{dr}{ds} = \frac{a}{2R} \sec^2\left(\frac{s}{2R}\right)$$

- (g) Express the derivative dr/ds in terms of r .

$$\frac{dr}{ds} = \frac{r^2 + a^2}{2Ra}$$

- (h) Find an expression for the scale on the map in the direction along the meridian at a point r cm from the map's center.

The scale is

$$\frac{2Ra}{r^2 + a^2} \frac{\text{km}}{\text{cm}}.$$

- (i) Check that your expression in part (h) has the correct units.

Stereographic Projection

- (j) *Use your result from part (h) to find the scale at the origin in the map above, where $R/a = 100 \text{ km/cm}$.*
- (k) *Use your result from part (h) to find the scale in the direction of the meridians on the orange circle of latitude. Start by using the number line to approximate the ratio r/a .*



Stretching Factors

Interpreting the derivative as a local stretching/scaling factor.

1 Rubber Band Calculus

Think of a function and the first thing that might come to mind is its graph. But we can also think of a function's domain as an elastic band and the function as acting on that band by stretching or compressing it.

For example, imagine a thin elastic band of length 4 meters running along the horizontal L -axis from $L = 0$ meters to $L = 4$ meters. Now hold the left end fixed at $L = 0$ and stretch the band by moving right end an additional four meters to the right. Then the function

$$H = g(L) = 2L, 0 \leq L \leq 4,$$

describes this stretching action. It takes as an input the distance (in meters) of a point on the band from the origin ($L = 0$) and returns as an output the distance between the origin and the corresponding point on the stretched band. The exploration below shows this stretching action.

Exploration 18 Drag the slider k in Line 2 below to illustrate the stretching action.

Desmos link: <https://www.desmos.com/calculator/qejivz36ui>

151: Rubber Band 1

The *global stretching factor* for a linear function like the one above,

$$H = g(L) = 2L, 0 \leq L \leq 4,$$

is the slope of its graph. We can calculate this factor as an average stretching factor (ie. an average rate of change) between the points $L = a$ meters and $L = b$ meters from the origin. For the function $H = f(L) = 2L$, the stretching

Learning outcomes:
Author(s):

factor is

$$\begin{aligned}\frac{\Delta H}{\Delta L} &= \frac{f(b) - f(a)}{b - a} \\ &= \frac{2(b - a)}{b - a} \\ &= 2,\end{aligned}$$

This tells us that any two points on the stretched band are twice as far apart as they were on the unstretched band.

Question 19 (a) What are the units of the stretching factor?

Free Response:

(b) Find an expression for the inverse function

$$L = f^{-1}(H).$$

Include a domain.

(c) Interpret the inverse function as a deformation of a thin elastic band. What is the global stretching factor for this function?

Free Response:

2 Non-Linear Stretching Functions

Example 10. Here's an example

$$H = f(L) = 10 - \sqrt{100 - L^2}, \quad 0 \leq L \leq 10,$$

of a non-linear stretching function (where L and H are measured in meters as before). Like most functions in this class, it acts like a linear function near most points in its domain.

To stretch the elastic band in the demonstration below, drag the slider u in Line 2 from $u = 0$ to $u = 1$. Then zoom in close enough to the point $H = f(2)$ in the stretched band (highlighted in black) to make the stretching function look linear.

Desmos link: <https://www.desmos.com/calculator/nyd60dbezj>

151: Rubber Band Ladder 5

- (a) Use the close-up view of the stretched band to approximate the stretching factor at the input $L = 2$.
- (b) Drag the slider m in Line 4 to $m = 95$ and repeat part (a) to approximate the stretching factor at the input $L = 9.5$ meters.
- (c) Parts of the elastic band get stretched, others compressed. Identify these.

We can also approximate the stretching factors at different inputs by zooming in near enough to the graph of the function $H = f(L)$ (shown below) to make the graph look like a line.

Desmos link: <https://www.desmos.com/calculator/jfwvbbtrts1>

151: Rubber Band Ladder 6

- (a) Zoom in close enough to the point $P(2, f(2))$ to make the graph above look like a straight line. Then click on the graph to get the coordinates of P and a point close to P . Use these to approximate the stretching factor at $L = 2$.
- (b) Zoom back out and use the graph of the function $H = f(L)$ to determine the stretching factors at $L = 0$ and $L = 10$.

We can also approximate the stretching factors numerically using the expression

$$H = f(L) = 10 - \sqrt{100 - L^2}, \quad 0 \leq L \leq 10,$$

for the stretching function.

For example, at the input $L = 2$, we'll first compute the *average stretching factor* over the interval between lengths $L = 2$ and $L = v$. This average factor is

$$\begin{aligned} k(v) &= \frac{\Delta H}{\Delta L} \\ &= \frac{f(v) - f(2)}{v - 2} \\ &= \frac{10 - \sqrt{100 - v^2} - (10 - \sqrt{96})}{v - 2} \\ &= \frac{\sqrt{100 - v^2} - \sqrt{96}}{v - 2}, \quad v \neq 2. \end{aligned}$$

- (a) The idea to approximate the stretching factor at $L = 2$ is to see if these average stretching factors appear to approach some number as $v \rightarrow 2$ (ie. as v approaches 2). To do this we'll use the Table in Line 3 of the worksheet above. But first you'll need to input the correct expression for the average stretching function in Line 2. Then use the table to approximate the stretching factor at the input $L = 2$.

- (b) Repeat part (a) to approximate the stretching factor at the input $L = 9.5$ and compare this with your earlier estimate.
- (c) Repeat part (a) to approximate the stretching factor at the right end where $L = 10$. Anything different here?
- (d) Use the geometry of the graph of the function $H = f(L)$ to compute the *exact* stretching factors at $L = 2, 9.5, 10$.

Example 11. Here's another example

$$H = f(L) = 4 - 4 \cos(L/2), \quad 0 \leq L \leq 2\pi.$$

of a non-linear stretching function. We still need to talk about a *local* stretching factor, one that varies from point to point.

Exploration 20 Drag the slider u in Line 2 below from $u = 0$ to $u = 1$ to illustrate the stretching action.

Desmos link: <https://www.desmos.com/calculator/hqvyhormhf>

151: Rubber Band Cosine

-
- (a) Use the worksheet above to approximate the local stretching factor at the point $L = \pi \sim 3.1$ meters from the origin on the unstretched band. Explain your reasoning.
 - (b) Drag slider m in Line 4 above to approximate the input(s) at which the local stretching factor is equal to 1. (ie. where the stretched band is neither in tension or in compression). Explain your reasoning.
 - (c) Use the graph of the stretching function below to approximate the local stretching factor at the input $L = \pi$. Start by dragging the slider u in Line 2 below from $u = 0$ to $u = 1$ to illustrate the stretching action and change the graph from $H = L$ to $H = f(L)$. Then zoom in close to the point $P(\pi, 4)$.

Desmos link: <https://www.desmos.com/calculator/aczf35qj>

151: Rubber Band Cosine 3

- (d) Finally, approximate the local stretching factor at $L = \pi$ numerically by finding entering an expression for the average stretching factor $k(v)$ between inputs $L = 2$ and $L = v$ in Line 3 of the worksheet above. Then use the table in Line 4.

3 A Hanging Slinky

Example 12. A hanging slinky stretched under its own weight gives an example of a stretching function.

Desmos link: <https://www.desmos.com/calculator/zqjjgae15j>

151: Slinky Photo

We can model the stretching of an ideal spring stretched under its own weight with the function

$$H = f(L) = L + \frac{g\rho}{2kL_0}L^2, \quad 0 \leq L \leq L_0,$$

where

- L_0 is the length (in meters) of the relaxed (unstretched) spring,
- g is the magnitude of the gravitational acceleration (measured in meter-sec/sec),
- ρ is the linear density (in kg/meter) of the spring,
- k (measured in Newtons/meter) is the spring constant,
- L is the distance of a point on the relaxed spring from the spring's bottom end, and
- $H = f(L)$ is the distance from the corresponding point on the stretched spring to the spring's bottom end.

Free Response: Use the above information to check that the above expression for $H = f(L)$ has the correct units. You will need to know that Newtons (a measure of force) have units $\text{kg} \cdot \text{m}/\text{sec}^2$.

Drag the slider g in Line 5 of the worksheet below from $g = 0$ to $g = g_0 = 0.5\text{m}/\text{sec}^2$ to turn on the gravitational field and stretch the slinky.

Desmos link: <https://www.desmos.com/calculator/vjjibjkdrrz>

151: Slinky 3

We'll now work with the stretching function in the worksheet above,

$$H = f(L) = L + \frac{1}{4}L^2, \quad 0 \leq L \leq 4.$$

- (a) Use the pictures of the springs at the left of the graph to estimate the local stretching factor

$$\left. \frac{dH}{dL} \right|_{L=3}$$

at the point in the relaxed spring $L = 3$ meters from its bottom end.

- (b) Find an expression for the average stretching factor between inputs $L = 3$ and $L = v$ meters. This factor is

$$\begin{aligned} a = m(v) &= \frac{\Delta H}{\Delta L} \\ &= \frac{f(v) - f(3)}{v - 3} \\ &= \frac{v^2 + 4v - 21}{4(v - 3)} \end{aligned}$$

- (c) Enter this expression in Line 1 of the worksheet below. Then
- (i) Use the table in Line 2 to guess the exact value of the local stretching factor at $L = 3$. Compare this with your estimate.
 - (ii) Drag the slider v in Line 11 and use the output in Line 12 ($m(v)$) to guess the exact value of the stretching factor at $L = 3$.
 - (iii) Describe how the line PQ is related to the average stretching factor $a = m(v)$.

Desmos link: <https://www.desmos.com/calculator/zsbupxubm6>

151: Slinky 4

- (d) Now we'll compute the local stretching factor at $L = 3$ algebraically. This factor is the derivative

$$\left. \frac{dH}{dL} \right|_{L=3}$$

of H with respect to L evaluated at $L = 3$. It is equal to the limit, as v approaches 3, of the average stretching factor $m(v)$ between lengths $L = 3$ and $L = v$. Here's the computation with some algebra left for you near the end:

$$\begin{aligned}
 \left. \frac{dH}{dL} \right|_{L=3} &= \lim_{\Delta L \rightarrow 0} \frac{\Delta H}{\Delta L} \\
 &= \lim_{v \rightarrow 3} \frac{f(v) - f(3)}{v - 3} \\
 &= \lim_{v \rightarrow 3} \frac{v^2 + 4v - 21}{4(v - 3)} \\
 &= \lim_{v \rightarrow 3} \frac{(v+7)(v-3)}{4(v-3)} \\
 &= \lim_{v \rightarrow 3} \frac{v+7}{4} \\
 &= \frac{3+7}{4} \\
 &= 2.5.
 \end{aligned}$$

- (e) So our conclusion is that the local stretching factor of the stretching function

$$H = f(L) = L + \frac{1}{4}L^2, \quad 0 \leq L \leq 4,$$

at $L = 3$ is equal to 2.5 meters/meter. This is a dimensionless scaling factor. It means that a small interval of length $\Delta L \sim 0$ meters around the point $L = 3$ on the relaxed spring gets stretched to a small interval approximately 2.5 times as long. Symbolically, we can write this as

$$\Delta H \sim 2.5\Delta L,$$

where

$$\Delta H = f(L) - f(3)$$

and

$$\Delta L = L - 3 \sim 0.$$

- (f) What about the inverse function $L = f^{-1}(H)$ that relaxes the stretched band? What is the local scaling factor

$$\left. \frac{dL}{dH} \right|_{L=3}$$

of the inverse at $L = 3$? Explain your reasoning.

- (g) Use the algebra of limits to find the local stretching factor

$$\left. \frac{dH}{dL} \right|_{L=b}$$

of this same function at the point $L = b$ meters from the bottom end of the relaxed spring by evaluating the limit (as $v \rightarrow b$) of the average stretching factor between $L = v$ and $L = b$. Then check your result by substituting $b = 3$. Check also with the above worksheets at a few other values of b .

4 Gas Consumption

Most of our applications will not be about springs or elastic bands. Nevertheless, we can still think of the derivative as a local stretching factor. But it might be better to talk about a local *scaling factor* instead. The derivative scales (ie. multiplies) a small change in the input to give an approximate corresponding change in the output. Here's an example.

Example 13. The function

$$G = f(s) = \frac{1}{2000}(s - 100)^2, \quad 0 \leq s \leq 80,$$

expresses the number of gallons of gas in a car in terms of the trip odometer reading in miles. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/wdnmaszvqb>

151: Gas Consumption A

- What are the units of the factor $1/2000$ in the function f ? How do you know?
- Find the average rate (in gal/mile) at which the car burned gas during the trip. Then find the average gas mileage (in miles/gal) over the entire trip.
- Use the graph above to approximate the rate (in gal/mile) at which the car burns gas when the odometer reads 60 miles. Then approximate the gas mileage (in miles/gal) at this odometer reading.
- Enter the correct expression in Line 5 of the worksheet for the average rate at which the car burns gas between odometer readings $s = v$ and $s = 60$ miles. Then use the table to guess the exact rate at which the car burns gas at the odometer reading $s = 60$ miles.
- Use the algebra of limits to compute the exact rate (in gal/mile) at which the car burns gas when the odometer reads 60 miles.
- Approximate the change

$$\Delta G = f(v) - f(60)$$

in the number of gallons of gas in the tank from odometer reading $s = 60$ miles to odometer reading $s = v$ miles in terms of the change

$$\Delta v = v - 60$$

in the odometer reading. Assume $\Delta v \sim 0$.

- Use the algebra of limits as above to compute the exact rate (in gal/mile) at which the car burns gas at the odometer reading $s = u$ miles.

5 A Falling Slinky

Falling Slinky

6 Atmospheric Pressure

Atmospheric Pressure

Average Rates of Change

Graphing average rate of change functions.

With time as the independent variable, the derivative is most naturally interpreted as an instantaneous rate of change. Here's an example.

Question 21 Due to a late-season frost in eastern Washington, the price of Cosmic Crisp apples is rising precipitously.

The function

$$N = f(t), 0 \leq t \leq 8,$$

expresses the number of pounds of apples you can buy with \$36 in terms of the number of hours past 9am on April 23, 2005.

The graph of the function $N = f(t)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/njanpkrqex>

151: Apples Drought

- (a) Let $b \in [0, 8]$ be a constant. Interpret the meaning of the function

$$m(v) = \frac{f(v) - f(b)}{v - b}$$

in the context of this scenario. Include units. State the domain of the function as well.

- (b) Use the graph of the function $N = f(t)$ above and the slider v on Line 4 to sketch a rough graph of the function $r = m(v)$ when $b = 5$. Label the axes with the appropriate variable names and units.
- (c) Activate the folder (avg. rate of change function) in Line 11 to check your graph of the function $r = m(v)$.
- (d) Click on the appropriate points of the curve $r = m(v)$ to find the average rate of change (with respect to time) in the number of pounds of apples you can buy with \$36 between
- (i) 11:00 am and 2:00 pm

Learning outcomes:
Author(s):

- (ii) 1:00 pm and 2:00 pm
 Be sure to include units. Note that because of the different scale on the vertical axis for the function $r = m(t)$, you will need to divide the r -coordinates by 10.
- (e) Zoom in close enough near the missing point on the curve $r = m(v)$ to approximate the rate of change at 2pm in the number of pounds of apples you can buy with \$36 at 2:00pm.
- (f) Open the Table in Line 1 by clicking the right arrow at the left of the line. Use it to guess the rate of change (with respect to time) in the number of pounds of apples you can buy with \$36 at 2:00pm.
- (g) Now suppose between 9am and 5pm the price of apples increases at a constant rate. Suppose also that the price is \$2/lb at 9am and \$4/lb at 2pm.

- (i) Find an expression for the function $N = f(t)$.

Hint: First find a function

$$P = g(t), 0 \leq t \leq 8,$$

that expresses the price (in dollars/pound) in terms of the number of hours past 9am.

- (ii) Find an expression for the function $r = m(v)$. Include the appropriate domain.

Explanation. Working with the function

$$\begin{aligned} N &= f(t) \\ &= \frac{36}{2 + \frac{2}{5}t} \\ &= \frac{90}{5 + t}, 0 \leq t \leq 8, \end{aligned}$$

we have

$$\begin{aligned} m(v) &= \frac{f(v) - f(5)}{v - 5} \\ &= \frac{1}{v - 5} \left(\frac{90}{5 + v} - 9 \right) \\ &= \frac{1}{v - 5} \left(\frac{45 - 9v}{v + 5} \right) \\ &= \frac{-9}{v + 5} \text{ if } v \neq 5. \end{aligned}$$

- (iii) Use your expression for $m(v)$ and the algebra of limits to compute the rate of change (with respect to time) at 2pm in the number of pounds of apples you can buy with \$36.

Explanation. The question is asking us to compute the derivative

$$\left. \frac{dN}{dt} \right|_{t=5} = \lim_{v \rightarrow 5} \frac{f(v) - f(5)}{v - 5}.$$

Using the previous part, we get

$$\begin{aligned} \left. \frac{dN}{dt} \right|_{t=5} &= \lim_{v \rightarrow 5} \frac{f(v) - f(5)}{v - 5} \\ &= \lim_{v \rightarrow 5} \frac{-9}{v + 5} \\ &= \frac{-9}{5 + 5} \\ &= -0.9. \end{aligned}$$

Conclusion: At 2pm we can buy 9 pounds of apples with \$36. And at 2pm the number of pounds of apples we can buy with \$36 is decreasing at the rate of 0.9 lb/hour.

- (iv) Compare at 2pm the relative rate of change in the number of pounds of apples you can buy with \$36 and the relative rate of change in the price.

Explanation. To compute the *relative* rate of change in the price at 2pm, we divide the absolute rate of change, $0.40(\$/\text{lb})/\text{hr}$, by the price $P = f(5) = \$4/\text{lb}$ at that time. This tells us the price is increasing at the relative rate of

$$\frac{1}{P} \cdot \left. \frac{dP}{dt} \right|_{t=5} = \frac{0.40(\$/\text{lb})/\text{hr}}{4(\$/\text{lb})} = 10\%/\text{hr}$$

at 2pm.

At the same time, the number of pounds of apples we can buy with \$36 is changing at the relative rate of

$$\frac{1}{N} \cdot \left. \frac{dN}{dt} \right|_{t=5} = \frac{-0.9 \text{ lb/hr}}{9 \text{ lb}} = -10\%/\text{hr}.$$

So at 2pm the number of pounds of apples we can buy with \$36 is decreasing at the relative rate of 10%/hr.

Free Response: Notice anything?

Question 22 This question is a continuation of Question 1. We'll work with the same function

$$N = f(t) = \frac{36}{2 + \frac{2}{5}t}, \quad 0 \leq t \leq 8$$

and the expression

$$\frac{\Delta N}{\Delta t} = \frac{f(v) - f(b)}{v - b}$$

for the average rate of change of N (the number of pounds of apples we can buy with \$36) with respect to time over the time interval between $t = b$ and $t = v$ hours past 9am.

- (a) Use the algebra of limits and the average rate of change function above to compute the derivative

$$\left. \frac{dN}{dt} \right|_{t=b}.$$

- (b) Use the result of part (a) to compare at 9am the relative rates of change in the price and the number of pounds of apples we can buy with \$36.
- (c) Express the derivative from part (a) in terms of N and

$$P = 2 + \frac{2}{5}t.$$

Use this equation to relate the relative rates of change in the price and the number of pounds of apples we can buy with \$36 at any instant.

Question 23 We can often get more insight into a problem by taking away the numbers. Here we'll generalize the first two questions by supposing the (differentiable) function

$$P = g(t), 0 \leq t \leq 8,$$

expresses the price (in dollars/pound) of Cosmic Crisp apples in terms of the number of hours past 9am. And we'll let

$$N = f(t), 0 \leq t \leq 8,$$

be the function that expresses the number of pounds of apples we can buy with \$100 dollars in terms of the number of hours past 9am.

- (a) Find an expression for the function $N = f(t)$ in terms of the function $g(t)$.
- (b) Use your expression from part (a) to find an expression (in terms of the function g) for the average rate of change

$$\frac{\Delta N}{\Delta t} = \frac{f(v) - f(b)}{v - b}.$$

- (c) Use your expression from part (b) to find an expression for the derivative

$$\left. \frac{dN}{dt} \right|_{t=b}$$

in terms of the function $P = g(t)$ and its derivative dP/dt .

- (d) Express the derivative from part (c) in terms of N , P , and dP/dt .
- (e) Use your equation from part (c) to compare the relative rates of change in the price and the number of pounds of apples we can buy with \$100 at any instant.

Question 24 The function

$$G = f(s) = \frac{1}{2000}(s - 100)^2, \quad 0 \leq s \leq 80,$$

expresses the number of gallons of gas in a car in terms of the trip odometer reading in miles. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/igubjr4nw1>

151: Gas Consumption B

- (a) What are the units of the factor $1/2000$ in the function f ? How do you know?
- (b) Find the average gas mileage (in mile/gal) between odometer readings $s = 20$ and $s = 60$ miles.
- (c) Zoom in on the appropriate point in the graph above to approximate the gas mileage (in miles/gal) at the instant the odometer reads 20 miles.
- (d) Find an expression for the average gas mileage between odometer readings $s = b$ and $s = v$ miles. Simplify this expression.
- (e) Use the previous part to find an expression for the gas mileage at the instant the odometer reads b miles.
- (f) What is the gas mileage at the instant the odometer reads $s = 20$ miles.
- (g) At what odometer reading is the car getting 20 miles/gal?

Question 25 The function

$$g = f(v), \quad 10 \leq v \leq 55,$$

expresses the rate (in gal/hr) at which a car burns gas in terms of its speed (in miles/hour).

The graph of the function $g = f(v)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/3ubcc1x8kh>

151: Rate of Gas Consumption

- (a) Let $b \in [10, 55]$ be a constant. Interpret the meaning of the function

$$m(w) = \frac{f(w) - f(b)}{w - b}$$

in the context of this scenario. Include units. State the domain of the function as well.

- (b) Use the graph of the function $g = f(v)$ above and the slider w on Line 4 to sketch a rough graph of the function $r = m(w)$ when $b = 20$. Label the axes with the appropriate variable names and units.
- (c) Activate the folder (avg. rate of change function) in Line 12 to check your graph of the function $r = m(v)$.
- (d) Click on the appropriate points of the curve $r = m(v)$ to find the average rate of change (with respect to speed) in the fuel consumption rate between speeds of

(i) 15 miles/hour and 20 miles/hour

(ii) 20 miles/hour and 30 miles/hour

Be sure to include units. Note that because of the different scale on the vertical axis for the function $r = m(v)$, you will need to divide the r -coordinates by 10.

- (e) Zoom in close enough near the missing point on the curve $r = m(v)$ to approximate the rate of change (with respect to speed) in the rate of fuel consumption at a speed of 20 miles/hr.
- (f) Open the Table in Line 1 by clicking the right arrow at the left of the line. Use it to guess the rate of change (with respect to speed) in the rate of fuel consumption at a speed of 20 miles/hr.
- (g) Now suppose between speeds of 10 miles/hour and 55 miles/hour that the car's gas mileage is a linear function of its speed. Suppose also that the car gets 25 miles/gal at a speed of 20 miles/hour and 35 miles/gal at a speed of 40 miles/hour.
- (i) Find an expression $G = k(v)$ for the function that expresses the car's gas mileage (in miles/gal) in terms of its speed (in miles/hr).
- (ii) Find an expression for the function $g = f(v)$ that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hour). Note: This is not a linear function.

- (iii) Find an expression for the function $r = m(v)$. Include the appropriate domain.
- (iv) Use your expression for $m(v)$ and the algebra of limits to compute the rate of change (with respect to speed) in the rate of fuel consumption at a speed of 20 miles/hr.
- (v) Use the language of small changes to interpret the meaning of the derivative

$$\left. \frac{dg}{dv} \right|_{v=20}.$$

- (vi) Simplify the units of the derivative above. What do these units suggest about its meaning? Is this correct? Why or why not?

Question 26 The function

$$h = f(t), 0 \leq t \leq 2.2,$$

expresses the height of a balloon (in hundreds of feet) in terms of the number of minutes past noon.

The graph of the function $h = f(t)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/yd4xm6x6ub>

151: Balloon

- (a) Let $b \in [0, 2.2]$ be a constant. Interpret the meaning of the function

$$m(v) = \frac{f(v) - f(b)}{v - b}$$

in the context of this scenario. Include units. State the domain of the function as well.

- (b) Use the graph of the function $h = f(t)$ above and the slider u to sketch a rough graph of the function $r = m(v)$ when $b = 2$. Label the axes with the appropriate variable names and units.
- (c) Activate the folder (avg. rate of change function) in Line 11 to check your graph of the function $r = m(v)$.
- (d) Click on the appropriate points of the curve $r = m(v)$ to find the balloon's average rate of ascent between
 - (i) 12:00pm and 12:02pm
 - (ii) 12:01pm and 12:02pm

- (e) Zoom in close enough near the missing point on the curve $r = m(v)$ to approximate the balloon's rate of ascent at 12:02pm.
- (f) Open the Table in Line 1 by clicking the right arrow at the left of the line. Use it to guess the instantaneous rate of ascent at 12:02pm.
- (g) Now suppose

$$f(t) = -2t^3 + 7t^2 - 8t + 8, \quad 0 \leq t \leq 2.2.$$

- (i) Find a simplified expression for the function $r = m(v)$. Include the appropriate domain.
- (ii) Use your simplified expression to compute the balloon's rate of ascent at 12:02pm.

Question 27 (a) Drag the slider u from $u = 0$ to $u = 1$ in Line 2 below. Then use the graph/animation of the stretching function

$$H = f(L) = 0.1 + 0.5L + 0.1(L - 1)^3, \quad 0 \leq L \leq 3,$$

to approximate the local stretching factor at $L = 2$. Include units.

Desmos link: <https://www.desmos.com/calculator/boubpczsne>

151: Rubber Band 12

- (b) Use the graph of the same stretching function $H = f(L)$ show again below to graph (by hand) the function

$$r = m(v)$$

that gives the average stretching factor of the portion of the band between lengths $L = v$ and $L = 2$ meters. Label the axes with the appropriate variable names and units. Then activate the average rate of change folder in Line 11 to see how you did.

Desmos link: <https://www.desmos.com/calculator/vss7ofbwii>

151: Rubber Band 12

- (c) Use the graph of the average rate of change function and the Table in Line 1 to guess the local stretching factor at $L = 2$.
- (d) Use the algebra of limits to compute the local stretching factor at $L = 2$.

A Balloon's Rate of Ascent

Graphing an average rate of change function and using limits.

Question 28 The function

$$h = f(t), 0 \leq t \leq 2.2,$$

expresses the height of a balloon (in hundreds of feet) in terms of the number of minutes past noon.

The graph of the function $h = f(t)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/yd4xm6x6ub>

151: Balloon

- (a) Let $b \in [0, 2.2]$ be a constant. Interpret the meaning of the function

$$m(v) = \frac{f(v) - f(b)}{v - b}$$

in the context of this scenario. Include units. State the domain of the function as well.

- (b) Use the graph of the function $h = f(t)$ above and the slider u to sketch a rough graph of the function $r = m(v)$ when $b = 2$. Label the axes with the appropriate variable names and units.
- (c) Activate the folder (avg. rate of change function) in Line 11 to check your graph of the function $r = m(v)$.
- (d) Click on the appropriate points of the curve $r = m(v)$ to find the balloon's average rate of ascent between
- (i) 12:00pm and 12:02pm
 - (ii) 12:01pm and 12:02pm
- (e) Zoom in close enough near the missing point on the curve $r = m(v)$ to approximate the balloon's rate of ascent at 12:02pm.
- (f) Open the Table in Line 1 by clicking the right arrow at the left of the line. Use it to guess the instantaneous rate of ascent at 12:02pm.

Learning outcomes:
Author(s):

(g) Now suppose

$$f(t) = -2t^3 + 7t^2 - 8t + 8, \quad 0 \leq t \leq 2.2.$$

- (i) Find a simplified expression for the function $r = m(v)$. Include the appropriate domain.
- (ii) Use your simplified expression and limits to compute the balloon's rate of ascent at 12:02pm.
- (iii) Assuming a constant rate of ascent, find the best possible approximation to the change

$$\Delta h = f(t) - f(2)$$

in the balloon's height (in hundreds of feet) over the time interval

$$\Delta t = t - 2$$

from 12:02pm to time t minutes past noon.

- (iv) Use the result of part (iii) to approximate when the balloon is 380 feet high. Explain your reasoning.
- (v) Use the graph of the function $h = f(t)$ to determine if your estimate in part (iii) is before or after the actual time. Explain your reasoning.

Rates of Change, Gas Mileage

Limits in context.

1 Gas Mileage

Example 14. The function

$$G = f(s) = \frac{2}{5} + \frac{1}{5000} (40(s+2)^2 - (s+2)^3), \quad 0 \leq s \leq 23,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/xzknfkw3>

Desmos activity available at 151: [Gas as a Function of Distance](#)

- Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
- Find your average gas mileage (in miles/gallon) over the interval $s \in [8, 18]$.
- Use the graph to approximate your gas mileage at the moment you are 18 miles from home. Do this by zooming in on the appropriate point.
- Find an expression for the function $r = h(s)$, that gives your average gas mileage between distances s miles and 18 miles from home. Include the correct domain.
- Use the algebra of limits to determine your exact gas mileage at the moment you are 18 miles from home.
- Use the result of part (e) to approximate the change in the volume of gas

$$\Delta G = f(s) - f(18)$$

in terms of the change

$$\Delta s = s - 18$$

in your distance from home for values of s near 18 miles. What are the units of the scaling factor?

Learning outcomes:
Author(s):

Rates of Change, Gas Mileage

- (g) Use the result of part (f) to approximate your distance from home when there are 1.9 gallons of gas in your tank.

Practice Quiz 1, Approximating Scaling Factors Numerically

Practice Quiz 1.

Directions:

- Give thorough explanations in complete sentences to each of the following questions. Include pictures for Questions 1 and 2 to help with your explanation.
- You will need a scientific calculator, but do not use any other technology.

Question 29 Use precalculus and arithmetic, but no calculus, to give numerical evidence that suggests an answer to the question below:

In the late afternoon, a 100-foot tall building casts a 300-foot long shadow on level ground. Is there a scaling factor that converts (by multiplication) a small change in the inclination angle of the setting sun to an accurate approximation of the corresponding change in the shadow's length?

- Start by writing a complete sentence that defines a function expressing the length of the shadow in terms of the inclination angle. Include units in your definition. Use meaningful variables, not x and y . Then find an expression for the function.*
- If your evidence suggests that such a scaling factor exists, explain why and approximate the scaling factor to the nearest hundredth. Include units. Otherwise, if your evidence suggests that no such scaling factor exists, explain why.*

Question 30 Use precalculus and arithmetic, but no calculus, to give numerical evidence that suggests an answer to the question below:

A rocket orbits a planet of radius 2000km at an altitude of 400 km above the surface. Is there a scaling factor that converts (by multiplication) a small change

Learning outcomes:
Author(s):

Practice Quiz 1, Approximating Scaling Factors Numerically

in orbit's radius to an accurate approximation of the corresponding change in radius of the portion of the planet's surface visible to an astronaut in the rocket? The radius is measured along the planet's surface as shown below.

Desmos3D link: <https://www.desmos.com/3d/paqpozwxm>

Desmos activity available at 151: *Spaceship above earth*

- (a) Start by writing a complete sentence that defines the appropriate function. Include units in your definition. Use meaningful variables, not x and y . Then find an expression for the function.
- (b) If your evidence suggests that such a scaling factor exists, explain why and approximate the scaling factor to the nearest hundredth. Include units. Otherwise, if your evidence suggests that no such scaling factor exists, explain why.

Question 31 Use precalculus and arithmetic, but no calculus, to give numerical evidence that suggests an answer to the question below:

The atmospheric pressure is 3 atm at an altitude of 5 km above the surface of the planet Krypton. The pressure is 2 atm at an altitude of 12km. Assume the pressure is an exponential function of altitude.

A weather balloon is 5km above the surface. Is there a scaling factor that converts (by multiplication) a small change in the balloon's altitude to an accurate approximation of the corresponding change in atmospheric pressure?

- (a) Start by writing a complete sentence that defines the appropriate function. Include units in your definition. Use meaningful variables, not x and y . Then find an expression for the function. Include a graph of the function.
- (b) If your evidence suggests that such a scaling factor exists, explain why and approximate the scaling factor to the nearest hundredth. Include units. Otherwise, if your evidence suggests that no such scaling factor exists, explain why.

The Rhythm of a Function

Listening to the beat of a function.

Have you ever sat by the train tracks listening to the trains roll by? If so, and you're near a joint in a rather old track, you'll hear a regular beat of the train as it passes by. The video below (especially recommended for engineers), explains how modern technology gets around these types of joints.

Expansion Joints

Exploration 32 *But we can use this idea, the old-fashioned clickity-clack rhythm of a train, to listen to the beat of a function.*

We'll imagine shrinking our train to a point that moves upward along a straight track (the s -axis in the demonstration below). And we'll start listening to our train at time $t = 0$ seconds as it passes the point $s = 0$. For the demonstration below, the joints are spaced just 0.1 meters apart.

Desmos link: <https://www.desmos.com/calculator/qjsfuvhsv9>

151: Sound Squaring Function 3

Follow the directions in the exploration above and then do the following.

- (a) Describe how the train's speed varies.
- (b) Sketch a rough graph of the function $v = f(t)$, $0 \leq t \leq 1$, that expresses the train's speed (in meters/sec) as function of the number of seconds since the train passed the point $y = 0$. Label the axes with the correct variable names and units.
- (c) Use your graph from part (b) to sketch a graph of the function $s = g(t)$, that expresses the train's distance (in meters) from the point $s = 0$ as a function of time (measured in seconds as in part (b)).
- (d) Finally, use your graphs from part (b) and (c) to sketch a graph of the function $v = h(s)$, that expresses the train's speed (in meters/sec) in terms of its distance (in meters) from $s = 0$.

Desmos link: <https://www.desmos.com/calculator/xaajoakuug>

Learning outcomes:
Author(s):

151: Sound Squaring Function 5B

Exploration 33 *Desmos link:*

<https://www.desmos.com/calculator/xyfteumpre>

151: Sound Function Cube

Exploration 34 *Desmos link:*

<https://www.desmos.com/calculator/uwym1ow6zq>

151: Sound Function Exp2

Exploration 35 *Can you hear an inflection point?*

Desmos link: <https://www.desmos.com/calculator/oojiur3a93>

151: Sound Function Oscillating

Exploration 36 *Desmos link:*

<https://www.desmos.com/calculator/xwj3qnoqia>

151: Sound Function Root

Differentiable Functions

An introduction to what it means for a function to be differentiable.

1 Differentiability

Zoom in closer and closer near a point $(a, f(a))$ on the graph of a common function f and you'll most likely notice that the graph looks more and more like the graph of a linear function (ie. like a non-vertical line). If so, we say the function $y = f(x)$ is *differentiable* at $x = a$. And the *derivative* of the function, evaluated at $x = a$, written as

$$\left. \frac{dy}{dx} \right|_{x=a},$$

is just the slope of that line or the rate of change of the linear function.

When the independent variable is time, we can interpret the derivative as an instantaneous rate of change.

Example 15. The graph of the function

$$T = f(m), \quad 0 \leq m \leq 60,$$

expressing the temperature (in °C) of a cup of coffee in terms of the number of minutes past noon is shown below.

Desmos link: <https://www.desmos.com/calculator/fuftb4mq0k>

151: Cooling Coffee

- Zoom in close enough to the point P above to make the graph look like a straight line. Then approximate the slope of this line. Include units.
- Use the result of part (a) to interpret the value and the meaning of the derivative

$$\left. \frac{dT}{dm} \right|_{T=40}.$$

- Repeat parts (a) and (b) for other temperatures by dragging the slider T_0 in Line 2 of the worksheet above. Then see if you can find a relationship between the temperature T and the rate dT/dm at which the temperature of the coffee is changing.

Learning outcomes:
Author(s):

We'll get back to applications a bit later. The next few examples focus on the geometric meaning of differentiability.

Exploration 37 Use the graph of the function $y = f(x)$ below to answer the following questions.

Desmos link: <https://www.desmos.com/calculator/ojdj4r3r9v>

151: Diff0

- (a) Zoom in toward the point A to determine if the function $y = f(x)$ is differentiable at $x = 1$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=1} = \left. \frac{d}{dx} (f(x)) \right|_{x=1}.$$

The derivative is

$$\left. \frac{dy}{dx} \right|_{x=1} = 3.$$

- (b) Zoom in toward the point B to determine if the function $y = f(x)$ is differentiable at $x = 0$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0} = \left. \frac{d}{dx} (f(x)) \right|_{x=0}.$$

The derivative is

$$\left. \frac{dy}{dx} \right|_{x=0} = -2.$$

Exploration 38 Use the graph of the function

$$y = f(x) = |x|$$

below to evaluate the following derivatives if possible. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/us2fruzbra>

151: Diff1

- (a)

$$\left. \frac{d(|x|)}{dx} \right|_{x=2} = 1$$

(b)

$$\left. \frac{d(|x|)}{dx} \right|_{x=0} =$$

(c)

$$\left. \frac{d(|x|)}{dx} \right|_{x=-3} = -1$$

Exploration 39 Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0$. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/ov8qt938ot>

151: Diff2

Exploration 40 (a) Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0.2$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0.2} = \left. \frac{d}{dx} (f(x)) \right|_{x=0.2}.$$

Explain your reasoning.

(b) Use the graph of the function $y = f(x)$ below to determine if the function is differentiable at $x = 0$. If so, approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=0} = \left. \frac{d}{dx} (f(x)) \right|_{x=0}.$$

Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/tvwtbx9hco>

151: Not Differentiable

Exploration 41 Use the graph of $y = f(x)$ below to determine whether each of the following derivatives are negative, positive, or zero. Explain your reasoning.

(a)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=0.5}$$

(b)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=1}$$

(c)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=1.5}$$

(d)

$$\left. \frac{d}{dx} (f(x)) \right|_{x=2.29}$$

Desmos link: <https://www.desmos.com/calculator/ks2yui6ofs>

151:Diff 7

Question 42 (a) Summarize your understanding of the main ideas of this section.

(b) What questions do you have about this section?

Free Response:

2 Using Algebra to Compute Derivatives

Exploration 43 The aim of this problem is to use numerical methods to approximate and algebra to evaluate the derivative

$$\left. \frac{d}{dx} (x^3) \right|_{x=1}.$$

(a) First use the graph of the function $y = f(x) = x^3$ below to approximate or guess the value of the above derivative by zooming in on the appropriate point.

$$\left. \frac{d}{dx} (x^3) \right|_{x=1} = 3.$$

Desmos link: <https://www.desmos.com/calculator/rto22qzlvn>

151: Cubing Function

- (b) The idea to compute the derivative algebraically is this: Fix the point $P(1,1)$ on the graph of $y = f(x) = x^3$. Then choose a variable point Q on the graph, different from P , with coordinates (v, v^3) . When Q is sufficiently close to P , the line PQ approximates the curve $y = x^3$ near $P(1,1)$ and the slope of this line approximates the derivative above.
- (i) Our first step is to find an expression for the slope of line PQ as a function of v . The slope is

$$\begin{aligned} m(v) &= \frac{\Delta y}{\Delta v} \\ &= \frac{f(v) - f(1)}{v - 1} \\ &= \frac{v^3 - 1}{v - 1} \end{aligned}$$

all assuming $v \neq 1$.

- (ii) Now we'll use the slope (or average rate of change) function $m(x)$ to create a table of slopes for the lines PQ . Reveal the contents of the Table folder in Line 1 of the worksheet above by clicking the Right Arrow just to the left of "Table".
- Add a few entries to the table to get better approximations to the above derivative.
 - The slopes $m(v)$ should appear should appear to approach some number as v approaches 1. What is that number?
 - This suggests that

$$\left. \frac{d}{dx} (x^3) \right|_{x=1} = \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} = 3.$$

- (iii) Another approach to approximating the derivative is to graph the slope function

$$m(v) = \frac{v^3 - 1}{v - 1}, v \neq 1.$$

Activate the Average Rate of Change folder on Line 11 by clicking the camera icon to the left of the line to see the graph of this function. Activate also the Table folder in Line 1.

- Drag the slider v on Line 4. Describe the relationship between the graph of the slope function and the line PQ .
- There is a hole in the graph of the function $y = m(x)$. Where is it? Why is it there?

(iv) To verify the numerical and graphical evidence that

$$\frac{d}{dx}(x^3) \Big|_{x=1} = \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} = 3,$$

we'll use algebra to evaluate the above limit.

The idea is to factor the numerator $v^3 - 1$. Since

$$(v^3 - 1) \Big|_{v=1} = 1^3 - 1 = 0,$$

we know that $v - 1$ is a factor of $v^3 - 1$. To simplify the quotient

$$\frac{v^3 - 1}{v - 1}, \quad v \neq 1,$$

we could use long division or factor the difference of two cubes:

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2).$$

Either way, the result is that

$$v^3 - 1 = (v - 1)(v^2 + v + 1).$$

So

$$\frac{v^3 - 1}{v - 1} = v^2 + v + 1, \quad v \neq 1.$$

Putting this all together, we get

$$\begin{aligned} \frac{d}{dx}(x^3) \Big|_{x=1} &= \lim_{v \rightarrow 1} \frac{v^3 - 1}{v - 1} \\ &= \lim_{v \rightarrow 1} \frac{(v - 1)(v^2 + v + 1)}{v - 1} \\ &= \lim_{v \rightarrow 1} (v^2 + v + 1) \\ &= 1^2 + 1 + 1 \\ &= 3. \end{aligned}$$

Exploration 44 Repeat all parts of Exploration 2 for the following derivatives.

(a)

$$\frac{d}{dx}(x^3) \Big|_{x=2} = 12.$$

(b)

$$\frac{d}{dx} (x^3) \Big|_{x=a} = 3a^2.$$

Use the demonstration below to check your work for part (b) by dragging the sliders v (Line 4) and a (Line 16). Then activate the Derivative Function folder on Line 17.

Desmos link: <https://www.desmos.com/calculator/ewy7jqij6s>

151: Cubing Function 2

Keep following the method of Exploration 2 to evaluate the following derivatives.

(c)

$$\frac{d}{dx} \left(\frac{1}{x} \right) \Big|_{x=1} = -1.$$

(d)

$$\frac{d}{dx} \left(\frac{1}{x} \right) \Big|_{x=a} = -1/a^2.$$

(e)

$$\frac{d}{dx} \left(\frac{1}{x^2} \right) \Big|_{x=3} = -2/27.$$

(f)

$$\frac{d}{dx} \left(\frac{1}{x^2} \right) \Big|_{x=a} = -2/a^3.$$

(g)

$$\frac{d}{dx} \left(\frac{1}{1+x^2} \right) \Big|_{x=1} = -\frac{1}{2}.$$

(h)

$$\frac{d}{dx} \left(\frac{1}{1+x^2} \right) \Big|_{x=a} = -\frac{2a}{(1+a^2)^2}.$$

3 Applications

Question 45 The function

$$h = f(t), 0 \leq t \leq 2.2,$$

expresses the height of a balloon (in hundreds of feet) in terms of the number of minutes past noon.

The graph of the function $h = f(t)$ is shown below.

Desmos link: <https://www.desmos.com/calculator/yd4xm6x6ub>

151: Balloon

- (a) Find an expression for the function $r = m(v)$ that gives the balloon's average rate of ascent (measured in hundreds of feet per minute) between time v minutes past noon and 12:02 pm. Include also the function's domain.
- (b) How is the average rate of ascent function in part (a) related to the line PQ in the demonstration above?
- (c) Use the graph of the function $h = f(t)$ above to sketch a rough graph of the function $r = m(v)$.
- (d) Activate the folder (avg. rate of change function) in Line 11 to check your graph of the function $r = m(v)$.
- (e) Open the Table in Line 1 by clicking the right arrow at the left of the line.
 - (i) What is the balloon's average rate of ascent between 12:02:00 pm and 12:02:06 pm?
 - (ii) Use the graph of the function $h = f(t)$ above to estimate the balloon's rate of ascent at 12:02pm by zooming in on the appropriate point.
 - (iii) What does the table suggest about the balloon's rate of ascent at 12:02pm? Explain.

(f) Now suppose

$$f(t) = -2t^3 + 7t^2 - 8t + 8.$$

- (i) Find a simplified expression for the function $r = m(v)$. Include the appropriate domain.
- (ii) Use your simplified expression to compute the balloon's rate of ascent at 12:02pm.

Differentiable Functions, Part 2

Using limits to compute the derivative of a function at a general input.

1 Using Limits to Compute Derivatives

Exploration 46 In part (b) of Exploration 7 from the previous chapter where we computed

$$\left. \frac{d}{dx} (x^3) \right|_{x=1}.$$

Now we'll use limits to evaluate the derivative

$$\left. \frac{d}{dx} (x^3) \right|_{x=a}.$$

In other words, we'll compute the derivative of the function $f(x) = x^3$ at a general input $x = a$. Think about zooming in on the graph of $y = x^3$ sufficiently close to the point (a, a^3) so that the graph looks like a straight line. We'll compute the slope of that line.

The algebra here is nearly identical to what we did earlier. You should compare the two computations.

Desmos link: <https://www.desmos.com/calculator/8eiffwbgt5>

151: Cubing Function 3

The idea to compute the derivative algebraically is this: Fix the point $P(a, a^3)$ on the graph of $y = f(x) = x^3$. Then choose a variable point Q on the graph, different from P , with coordinates (v, v^3) . When Q is sufficiently close to P , the line PQ approximates the curve $y = x^3$ near $P(a, a)$ and the slope of this line is the derivative.

- (a) Our first step is to find an expression for the slope of line PQ as a function

Learning outcomes:
Author(s):

of v . The slope is

$$\begin{aligned} m(v) &= \frac{\Delta y}{\Delta v} \\ &= \frac{f(v) - f(a)}{v - a} \\ &= \frac{v^3 - a^3}{v - a} \end{aligned}$$

all assuming $v \neq a$.

- (b) The next step is to factor the numerator $v^3 - a^3$. Since

$$(v^3 - a^3) \Big|_{v=a} = a^3 - a^3 = 0,$$

we know that $v - a$ is a factor of $v^3 - a^3$. To simplify the quotient

$$\frac{v^3 - a^3}{v - a}, \quad v \neq a,$$

we could use long division or factor the difference of two cubes:

$$A^3 - B^3 = (A - B)(A^2 + AB + B^2).$$

Either way, the result is that

$$v^3 - a^3 = (v - a)(v^2 + va + a^2).$$

So

$$\frac{v^3 - a^3}{v - a} = v^2 + va + a^2, \quad v \neq a.$$

Putting this all together, we get

$$\begin{aligned} \frac{d}{dx}(x^3) \Big|_{x=a} &= \lim_{v \rightarrow a} \frac{v^3 - a^3}{v - a} \\ &= \lim_{v \rightarrow a} \frac{(v - a)(v^2 + va + a^2)}{v - a} \\ &= \lim_{v \rightarrow a} (v^2 + va + a^2) \\ &= a^2 + a^2 + a^2 \\ &= 3a^2. \end{aligned}$$

- (c) More simply put, we just write the derivative

$$\frac{d}{dx}(x^3) = 3x^2$$

as a function of x .

(d) We can check our result using the desmos worksheet above. Do this as follows:

- (i) Input the correct expression for the derivative $d(x^3)/dx$ on Line 17.
- (ii) For a differentiable function $f(x)$ and values of h near zero, we can approximate the derivative $d(f(x))/dx$ as the slope

$$\frac{d}{dx}(f(x)) = \frac{f(x+h) - f(x-h)}{2h}$$

of the line through the points $(x-h, f(x-h))$ and $(x+h, f(x+h))$. Now activate the folder Approximation to the derivative function on Line 19. Then drag the slider h on Line 21 to get a better approximation to the derivative.

Free Response: Describe what happens to the above approximation as $h \rightarrow 0$.

Exercise 47 (a) Follow the method of Exploration 1 for the function $f(x) = x^4$ to compute the derivative

$$\left. \frac{d}{dx}(x^4) \right|_{x=a}.$$

- (b) Modify the desmos worksheet below for the function $f(x) = x^4$.

151: Cubing Function 3

Then to check your work, do part (d) of Exploration 1 for $f(x) = x^4$. Describe also what happens to the approximation in part (d) as $h \rightarrow 0$. Include a screenshot to help with your description.

Exercise 48 Repeat all parts of Exercise for the function $f(x) = 1/x^4$.

Exercise 49 Repeat all parts of Exercise for the function $f(x) = 1 + x^2$.

Exercise 50 Repeat all parts of Exercise for the function $f(x) = 1/(1 + x^2)$.

Limits and Derivatives, Part 1

Limits in context.

1 Limits and Speed

Example 16. A rock is dropped from rest near the surface of the moon. The function

$$s = f(t) = 0.8t^2, \quad 0 \leq t \leq 10,$$

expresses the distance (in meters) the rock falls in terms of the number of seconds since it was dropped.

- Find a function $v = g(t)$ that gives the average speed of the rock (in meters/sec) between times $t = 3$ and $t = b$ seconds. Simplify the function. State the domain of g .
- Graph the function $v = g(t)$ by hand. Label the axes with the appropriate variable names and units.
- How is the rock's average speed between times $t = 3$ and $t = b$ related to the line AB in the worksheet below?

Desmos link: <https://www.desmos.com/calculator/tcti9t3iub>

Desmos activity available at 151: Moon Rock 1

- Drag point B close to point A and approximate the rock's speed 3 seconds after being dropped. Do this by approximating the coordinates of a second point P on the line AB far from A . Then use the points A and P to approximate the slope of line AB .
- Use the result of part (a) and the algebra of limits to find the exact speed of the rock three seconds after being dropped.
- Find an expression for the rock's average speed between times $t = a$ and $t = b$.
- Use part (f) and the algebra of limits to find the speed of the rock at time $t = a$ seconds after being dropped.

Learning outcomes:

Author(s):

- (h) Compare this rock's speed at time $t = a$ with the rock's average speed during the first a seconds of its fall.
- (i) Interpret the two speeds in part (h) geometrically on the graph below.

Desmos link: <https://www.desmos.com/calculator/v42lwmlag>

Desmos activity available at 151: Moon Rock 2

2 The Cubing Function, Part 1

Example 17. This question is about the function

$$y = f(x) = x^3, -2 \leq x \leq 2.$$

Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/bb5oac9pik>

Desmos activity available at 151: Cubing Function 1

- (a) Zoom in on the graph near the point $(1, 1)$ to approximate the derivative

$$\left. \frac{dy}{dx} \right|_{x=1} = \left. \frac{d}{dx} (x^3) \right|_{x=1}.$$

As you zoom in closer to the point $(1, 1)$ the graph looks more and more like a straight line. The derivative is the slope of this line.

- (b) Graph by hand the function $r = m(b)$ that gives the average rate of change of the function $y = f(x) = x^3$ between $x = 1$ and $x = b$. Drag point B in the worksheet above to help.
- (c) Find a simplified expression for the function $r = m(b)$. Include a domain.
- (d) Input the expression for the function $r = m(x)$ in Line 1 of the worksheet. Then activate and open the Average Rate of Change folder in Line 2. Use the table to guess the value of the derivative in part (a).
- (e) Use the algebra of limits and your expression for the function $r = m(b)$ to compute the derivative in part (a).
- (f) Find an equation of the tangent line to the curve $y = x^3$ at the point $(1, 1)$.
- (g) Find the coordinates of the points where the tangent line in part (f) intersects the curve $y = x^3$.

3 The Cubing Function in Context

Example 18. (a) Find a function

$$V = f(s), s \geq 0,$$

that expresses the volume of a cube (measured in cubic cm) in terms of its edge length (measured in cm).

(b) Find a simplified expression for the average rate of the function $f(s)$ between edge lengths $s = a$ and $s = b$.

(c) Use the result of part (b) and the algebra of limits to evaluate the derivative

$$\left. \frac{dV}{ds} \right|_{s=a}.$$

(d) What are the units of the derivative in part (b)? Explain how you know.

(e) Find the scaling factor that converts a small change

$$\Delta s = s - 2$$

in the cube's edge length (in cm) to a good approximation of the corresponding change

$$\Delta V = f(s) - f(2)$$

in the cube's volume.

(f) What are the units of the scaling factor?

(g) Approximate ΔV in terms of Δs for small changes Δs .

4 The Cube Root Function in Context

Example 19. (a) Find a function

$$s = g(V), V \geq 0,$$

that expresses the edge length of a cube (measured in cm) in terms of its volume (measured in cubic centimeters).

(b) Find an expression for the average rate of the function $g(V)$ between volumes $V = V_1$ and $V = V_2$.

$$\frac{\Delta s}{\Delta V} =$$

- (c) Use the algebra of limits to evaluate the derivative

$$\left. \frac{ds}{dV} \right|_{V=V_1}.$$

Do this by first letting $s_1 = V_1^{1/3}$ and $s_2 = V^{1/3}$ and rewriting the average rate of change in part (b) in terms of s_1 and s_2 .

$$\frac{\Delta s}{\Delta V} =$$

- (d) What are the units of the derivative in part (b)? Explain how you know.
 (e) Find the scaling factor that converts a small change

$$\Delta V = V - 8$$

in the cube's volume (in cubic cm) to a good approximation of the corresponding change

$$\Delta s = g(V) - g(8)$$

in the cube's edge length.

- (f) What are the units of the scaling factor?
 (g) Approximate Δs in terms of ΔV for small changes ΔV .

5 Limits and Gas Mileage

Example 20. The function

$$G = f(s) = \frac{2}{5} + \frac{1}{5000}(40(s+2)^2 - (s+2)^3), \quad 0 \leq s \leq 23,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/xzkknfpkw3>

Desmos activity available at 151: Gas as a Function of Distance

- (a) Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
 (b) Find your average gas mileage (in miles/gallon) between distances $s = 8$ and $s = 18$ miles from home.

- (c) Use the graph to approximate your gas mileage at the moment you are 8 miles from home. Do this by zooming in on the appropriate point.
- (d) Find a simplified expression for your average gas mileage (in miles/gal) between distances $s = 8$ and $s = b$ miles from home.

Hint: Do *not* simplify the expression

$$f(8) = \frac{2}{5} + \frac{1}{5000}(40(10)^2 - (10)^3)$$

for $f(8)$. Instead, leave as is and factor the difference of two squares

$$A^2 - B^2 = (A + B)(A - B)$$

and the difference of two cubes

$$A^3 - B^3 = (A - B)(A^2 + AB + B^2).$$

- (e) Use your expression from part (d) and the algebra of limits to determine your exact gas mileage at the moment you are 8 miles from home.
- (f) Find a simplified expression for your average gas mileage (in miles/gal) between distances $s = a$ and $s = b$ miles from home. Use this expression and the algebra of limits to find an expression for your gas mileage at the moment you are a miles from home.

Limits and Derivatives, Part 2, Trigonometric Functions

Derivatives of some trig functions.

1 The Sine Function

Example 21. (a) Zoom in near the origin on the graph of the function

$$y = f(\theta) = \sin \theta, \quad -2\pi \leq \theta \leq 2\pi.$$

below to approximate the value of the derivative

$$\left. \frac{dy}{d\theta} \right|_{\theta=0} = \left. \frac{d}{d\theta} (\sin \theta) \right|_{\theta=0}.$$

Geometrically, the derivative is the slope of the tangent line to the curve $y = \sin \theta$ at the origin.

Desmos link: <https://www.desmos.com/calculator/tordzgkr1za>

Desmos activity available at 151: [Sine Graph Basic](#)

- (b) Find a simplified expression for the function $m = r(\theta_1)$ that gives the average rate of change of the function $f(\theta)$ over the interval between $\theta = 0$ and $\theta = \theta_1$. State the domain of this function.
- (c) Sketch by hand a graph of the function $m = r(\theta_1)$. Drag point B in the graph above to help.
- (d) Input your expression for the function m in Line 1 of the worksheet. Then open and activate the average rate of change folder in Line 2. Describe what you see? What does this suggest about the value of the derivative

$$\left. \frac{d}{d\theta} (\sin \theta) \right|_{\theta=0} = \lim_{\theta_0 \rightarrow 0} \frac{\sin \theta_0}{\theta_0}?$$

Example 22. Let

$$g(\theta) = \sin \left(\frac{1}{3} \theta \right), \quad -6\pi \leq \theta \leq 6\pi.$$

Learning outcomes:
Author(s):

- (a) Start by graphing this function by hand. Describe the transformation that takes the graph of $y = f(\theta) = \sin \theta$ to the graph of

$$y = g(\theta) = f\left(\frac{1}{3}\theta\right).$$

- (b) Use part (a) to guess the value of the derivative

$$\frac{d}{d\theta} \left(\sin \left(\frac{1}{3}\theta \right) \right) \Big|_{\theta=0}.$$

Explain your thinking.

- (c) Find a simplified expression for the function $m = r(\theta_1)$ that gives the average rate of change of the function $g(\theta)$ over the interval between $\theta = 0$ and $\theta = \theta_1$. State the domain of this function.

- (d) Use the fact that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

and the algebra of limits to evaluate the derivative in part (b).

Example 23. Repeat parts (a), (b), and (d) of Example 1 for the derivative

$$\frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\pi/6}.$$

Use the graph of the function $y = \sin \theta$ below to help.

Desmos link: <https://www.desmos.com/calculator/bf2ic534ik>

Desmos activity available at 151: Sine Graph Basic 2

Example 24. Let

$$f(\theta) = \sin \theta.$$

- (a) Find an expression for the average rate of change of the function $f(\theta)$ between angles $\theta = \theta_1$ and $\theta = \theta_2$.
- (b) Use part (a), the algebra of limits, and the fact that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

to find an expression for the derivative

$$\frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\theta_1}.$$

You'll need the identity

$$\sin B - \sin A = 2 \sin \left(\frac{1}{2} (B + A) \right) \cos \left(\frac{1}{2} (B - A) \right).$$

- (c) Use the result of part (c) to evaluate the derivative

$$\frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\pi/6}.$$

Compare the exact value with your approximation from Example 3.

2 A Building's Shadow

Example 25. The function

$$s = f(\theta) = 50 \cot \theta, \quad 0 < \theta \leq \pi/2,$$

expresses the length of the shadow of a 50-foot tall building in terms of the sun's inclination angle.

- (a) Find an expression for the average rate of change of the function $f(\theta)$ between angles $\theta = \theta_1$ and $\theta = \theta_2$.
 (b) Use part (a), the algebra of limits, and the fact that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

to find an expression for the derivative

$$\frac{ds}{d\theta} \Big|_{\theta=\theta_1} = \frac{d}{d\theta} (50 \cot \theta) \Big|_{\theta=\theta_1}.$$

You'll need the identity

$$\sin(B - A) = \sin B \cos A - \cos B \sin A.$$

Express the derivative in part (b) in terms of the length

$$s = f(\theta_1)$$

of the shadow.

Use part (c) to find the exact value of the derivative

$$\frac{ds}{d\theta} \Big|_{s=100}.$$

Compare with your work in the chapter

[Shadow of a Building, Part 1](#)

Practice with Limits and Derivatives

Practice with limits and derivatives.

1 More Practice with Rational Functions

Question 51 *The function*

$$W = f(r) = \frac{2000}{r^2}, r \geq 4,$$

expresses the weight of an astronaut (measured in pounds) in terms of her distance from the center of the earth (measured in thousands of miles).

- (a) *Find a fully-simplified expression for the average rate of change in the astronaut's weight with respect to her distance from the earth's center between distances $r = b$ and $r = c$ thousands of miles from the center. Assume $b, c \geq 4$ and that $b \neq c$.*
- (b) *Use your expression from part (a) directly and the definition of the derivative to find an expression (fully simplified) for the derivative*

$$\left. \frac{dW}{dr} \right|_{r=b}.$$

- (c) *Use part (b) to evaluate the derivative*

$$\left. \frac{dW}{dr} \right|_{r=5}.$$

- (d) *What are the units of the derivative above?*
- (e) *Explain the meaning of the derivative in part (c) using the language of scaling factors and small changes.*

Learning outcomes:
Author(s):

Explanation. (a) The average rate of change is

$$\begin{aligned}
 \frac{\Delta W}{\Delta r} &= \frac{f(c) - f(b)}{c - b} \\
 &= \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \\
 &= \left(\frac{2000}{c - b} \right) \left(\frac{b^2 - c^2}{c^2 b^2} \right) \\
 &= \left(\frac{2000}{c - b} \right) \cdot \frac{(b + c)(b - c)}{c^2 b^2} \\
 &= \frac{-2000(b + c)}{c^2 b^2}.
 \end{aligned}$$

(b) The derivative is

$$\begin{aligned}
 \left. \frac{dW}{dr} \right|_{r=b} &= \lim_{\Delta r \rightarrow 0} \frac{\Delta W}{\Delta r} \\
 &= \lim_{\Delta c \rightarrow b} \frac{-2000(b + c)}{c^2 b^2} \\
 &= -\frac{2000(b + b)}{b^2 b^2} \\
 &= -\frac{4000}{b^3}.
 \end{aligned}$$

(c) The value of the derivative is

$$\begin{aligned}
 \left. \frac{dW}{dr} \right|_{r=2} &= \left(-\frac{4000}{r^3} \right) \Big|_{r=5} \\
 &= -32.
 \end{aligned}$$

(d) The units of the derivative are pounds/(thousands of miles).

(e) The derivative

$$\left. \frac{dW}{dr} \right|_{r=2} = -32 \frac{\text{lbs}}{\text{thousands of miles}}.$$

is a scaling factor that converts a small change

$$\Delta r = r - 5$$

in the astronaut's distance (in thousands of miles) from the center of the earth into an approximation for the small change

$$\Delta W = W = f(5) = W - 80$$

in her weight. This means that for $\Delta r \sim 0$

$$\Delta W \sim -32\Delta r.$$

So if the astronaut's distance to the earth's center increased from $r = 5$ miles to $r = 5.1$ miles, her weight would change by approximately

$$\begin{aligned}\Delta W &\sim \left(-32 \frac{\text{lbs}}{\text{thousands of miles}}\right) (0.1 \text{ thousands of miles}) \\ &= -3.2 \text{ lbs.}\end{aligned}$$

Question 52 (a) Find a function

$$G = f(P), \quad P > 0,$$

that expresses the number of gallons of gas you can buy with \$100 in terms of the price (measured in dollars/gallon).

- (b) Sketch a graph of the function f by hand.
- (c) Find a fully-simplified expression for the average rate of change in the number of gallons of gas you can buy with \$100 with respect to the price (measured in dollars/gallon) between prices $P = a$ and $P = b$, both measured in dollars/gallon.
- (d) Use your expression from part (c) directly and the definition of the derivative to find an expression (fully simplified) for the derivative

$$\left. \frac{dG}{dP} \right|_{P=a}.$$

- (e) What are the units of the derivative? Do not simplify the units.
- (f) Evaluate the derivative

$$\left. \frac{dG}{dP} \right|_{P=5}.$$

- (g) Explain the meaning of the derivative in part (f) using the language of scaling factors and small changes.

Question 53 Between speeds of 20 miles/hour and 70 miles/hour the gas mileage of a car (in miles/gallon) is a quadratic function $G = f(v)$ of its speed (in miles/hr). The graph of the function f is shown below.

Desmos link: <https://www.desmos.com/calculator/6bbnti8zm5>

151: Gas Mileage Quadratic Function

- (a) Find an expression for the function $G = f(v)$.
- (b) Find a fully-simplified expression for the average rate of change in the gas mileage (in miles/gal) with respect to speed (measured in miles/gallon) between speeds $v = a$ and $v = b$, both measured in miles/hour.
- (c) Use your expression from part (b) directly and the definition of the derivative to find an expression (fully simplified) for the derivative

$$\left. \frac{dG}{dv} \right|_{v=a}.$$

- (d) What are the units of the derivative? Do not simplify the units.
- (e) Evaluate the derivative

$$\left. \frac{dG}{dv} \right|_{v=30}.$$

- (f) Explain the meaning of the derivative in part (e) using the language of scaling factors and small changes.

Question 54 The function

$$g = f(v) = \frac{1}{2}v + 10, \quad 10 \leq v \leq 55,$$

expresses the gas mileage (in miles/gal) of a car in terms of its speed.

- (a) At what rate (in gal/hr) does the car burn gas at a speed of 40 miles/hour?
- (b) Find a function

$$r = g(v), \quad 10 \leq v \leq 55,$$

that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed.

Answer: The function is

$$r = g(v) = \frac{v}{\frac{1}{2}v + 15}, \quad 10 \leq v \leq 55.$$

- (c) Find an expression for the average rate of change in the rate at which the car burns gas with respect to its speed between speeds of $v = w$ and $v = b$ miles/hour. Then use this expression and the algebra of limits to find an expression for the derivative

$$\left. \frac{dr}{dv} \right|_{v=b}.$$

- (d) Evaluate the derivative

$$\left. \frac{dr}{dv} \right|_{v=40}.$$

- (e) What are the units of the derivative above.? Do not simplify the units.
- (f) Explain the meaning of the derivative in part (d) using the language of scaling factors and small changes.

2 The Squaring Function and Its Inverse

Question 55 The function

$$A = f(s) = s^2, \quad s \geq 0,$$

expresses the area of a square (measured in square feet) in terms of its side length (in feet).

- (a) Find an expression for the average rate of change in the area of a square with respect to its side length between side lengths $s = b$ and $s = w$ feet.
- (b) Use your expression from part (a) and the algebra of limits to find an expression for the derivative

$$\left. \frac{dA}{ds} \right|_{s=b}.$$

- (c) Use the result of part (b) to evaluate the derivative

$$\left. \frac{dA}{ds} \right|_{s=5}.$$

- (d) What are the units of the derivative in part (c).
- (e) Explain the meaning of the derivative (c) in terms scaling factors and small changes.

Explanation. You should have found that

$$\left. \frac{dA}{ds} \right|_{s=5} = 10 \text{ ft}^2/\text{ft}.$$

This is a scaling factor that converts a small change

$$\Delta s = s - 5 \sim 0,$$

in the side length to a good approximation of the change

$$\Delta A = f(s) - f(5) = f(s) - 25$$

in the area. That is, for $\Delta s \sim 0$,

$$\frac{\Delta A}{\Delta s} \sim \left. \frac{dA}{ds} \right|_{s=5} = 10 \text{ ft}^2/\text{ft},$$

and

$$\Delta A \sim 10\Delta s.$$

For example, if

$$\Delta s = 0.1,$$

then

$$\Delta A \sim (10 \text{ ft})(0.1 \text{ ft}) = 1 \text{ ft}^2.$$

Compare this approximation with the actual change

$$\Delta A = f(5.1) - f(5) = 1.01 \text{ ft}^2$$

in the area. Pretty good.

Question 56 *The function*

$$s = g(A) = \sqrt{A}, \quad A \geq 0,$$

expresses the side length (in feet) of a square in terms of its area (in square feet).

(a) *Use the algebra of limits to find an expression for the derivative*

$$\left. \frac{ds}{dA} \right|_{A=a}.$$

(b) *Evaluate the derivative*

$$\left. \frac{ds}{dA} \right|_{A=25}.$$

What are its units? Explain its meaning in terms of small changes.

Explanation. The standard approach goes something like this.

$$\begin{aligned}
 \left. \frac{ds}{dA} \right|_{A=a} &= \lim_{w \rightarrow a} \frac{g(w) - g(a)}{w - a} \\
 &= \lim_{w \rightarrow a} \frac{\sqrt{w} - \sqrt{a}}{w - a} \\
 &= \lim_{w \rightarrow a} \frac{\sqrt{w} - \sqrt{a}}{(\sqrt{w} - \sqrt{a})(\sqrt{w} + \sqrt{a})} \\
 &= \lim_{w \rightarrow a} \frac{1}{\sqrt{w} + \sqrt{a}} \\
 &= \frac{1}{2\sqrt{a}}.
 \end{aligned}$$

This approach obscures a key point. Namely, that the function g is the *inverse* of the function f in Question 1. We can exploit this by instead rewriting the equation

$$s = \sqrt{A} = g(A)$$

as

$$A = s^2 = f(s).$$

Now we'll find an expression for the derivative

$$\left. \frac{ds}{dA} \right|_{s=b}$$

in terms of the side length $s = b$. This gives

$$\begin{aligned}
 \left. \frac{ds}{dA} \right|_{s=b} &= \lim_{w \rightarrow b} \frac{w - b}{f(w) - f(b)} \\
 &= \lim_{w \rightarrow b} \frac{w - b}{w^2 - b^2} \\
 &= \lim_{w \rightarrow b} \frac{1}{w + b} \\
 &= \frac{1}{2b}.
 \end{aligned}$$

Letting $a = b^2$ be the area, we have $b = \sqrt{a}$, and

$$\left. \frac{ds}{dA} \right|_{A=a} = \frac{1}{2\sqrt{A}}.$$

This makes it clear that the derivative of the inverse of a function is the *reciprocal* of the function's derivative.

(c) Use the result of part (b) to evaluate the derivative

$$\left. \frac{dA}{ds} \right|_{s=5}.$$

- (d) What are the units of the derivative in part (c).
- (e) Explain the meaning of the derivative (c) in terms of small changes.

3 The Cubing Function and its Inverse

Question 57 Repeat the two questions of the previous section with the function

$$V = f(s) = s^3, \quad s \geq 0,$$

that expresses the volume of a cube in terms of its side length.

4 A Square Root Function

Question 58 A rock dropped from a height of 100 feet falls to the surface of the planet Krypton without air resistance.

The function

$$v = f(h) = 6\sqrt{100 - h}, \quad 0 \leq h \leq 100,$$

expresses the rock's speed (in ft/sec) in terms of its height (in feet).

- (a) Find an expression for the average rate of change in the rock's speed with respect to its height between heights $h = a$ and $h = b$ feet.
- (b) Use your expression from part (a) directly and the definition of the derivative to find an expression (fully simplified) for the derivative

$$\left. \frac{dv}{dh} \right|_{h=b}.$$

- (c) Use part (b) to evaluate the derivative

$$\left. \frac{dv}{dh} \right|_{h=36}.$$

- (d) What are the units of the derivative above?
- (e) Explain the meaning of the derivative in part (c) using the language of small changes.

5 More on the Squaring Function

Check back later.

Limits

Limits in context.

1 Limits and Tangent Lines

Example 26. Let

$$g(x) = \frac{x^2 - 9}{3x - 9}.$$

(a) Evaluate each of the following expressions.

(i) $g(7)$

(ii) $\lim_{x \rightarrow 7} g(x)$

(iii) $g(3)$

(iv) $\lim_{x \rightarrow 3} g(x)$

(b) Simplify and then graph the function $g(x)$.

(c) Interpret the expressions in part (a) geometrically by considering the graph of the function $f(x) = x^2/3$ as in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/u0uvuchnrk>

Desmos activity available at 151: Parabola Basic

2 Limits and Gas Mielage

Example 27. The function

$$G = f(s) = \frac{2}{5} + \frac{1}{5000}(40(s+2)^2 - (s+2)^3), \quad 0 \leq s \leq 23,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/xzknfkw3>

Learning outcomes:
Author(s):

Desmos activity available at 151: Gas as a Function of Distance

- (a) Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
- (b) Find your average gas mileage (in miles/gallon) over the interval $s \in [8, 18]$.
- (c) Use the graph to approximate your gas mileage at the moment you are 18 miles from home. Do this by zooming in on the appropriate point.
- (d) Use the algebra of limits to determine your exact gas mileage at the moment you are 18 miles from home.
- (e) Use the result of part (d) to approximate the change in the volume of gas

$$\Delta G = f(s) - f(18)$$

in terms of the change

$$\Delta s = s - 18$$

in your distance from home for values of s near 18 miles. What are the units of the proportionality constant?

- (f) Use the result of part (e) to approximate your distance from home when there are 1.9 gallons of gas in your tank.

3 Limits, Gas Mileage and Speed

Example 28. Suppose that between speeds of 30 miles/hour and 70 miles/hour the gas mileage of a car is a quadratic function of its speed. Suppose also that the car gets a maximum of 42 miles/gal at a speed of 50 miles/hour and that the car gets 38 miles/gallon at a speed of 40 miles/hour.

- (a) Find an expression for the function

$$G = f(v), \quad 30 \leq v \leq 70,$$

that gives the gas mileage (in miles/gal) in terms of the speed (in miles/hour).

- (b) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the car's speed at 60 miles/hour gives an approximately proportional change in its gas mileage.
- (c) Use the results of part (b) to approximate the proportionality constant. What are its units?
- (d) Use the algebra of limits to find the exact value of the proportionality constant.
- (e) Explain the meaning of the proportionality constant.
- (f) Approximate the change

$$\Delta G = g - f(60)$$

in gas mileage in terms of a small change

$$\Delta v = v - 60$$

in the car's speed.

- (g) Use part (f) to approximate the speed at which the car gets 36 miles/gallon.
- (h) Would you expect your approximation in part (g) to be greater or less than the actual speed? Explain your reasoning with a graph.
- (i) Simplify the units of the proportionality constant. What might these units suggest about a way to interpret the constant?
- (j) At what rate (in gal/hr) does the car burn gas at a speed of 60 miles/hour?
- (k) How is the rate in part (j) related to the proportionality constant?

4 Limits, Speed and Altitude

Question 59 A rock dropped from a height of 100 feet falls to the surface of Planet Krypton without air resistance.

(a) By considering only the physical situation and without doing any computations, sketch a graph of the function

$$v = g(h), 0 \leq h \leq 100$$

that expresses the rock's speed (in ft/sec) in terms of its height (in feet).

(b) Use the results from part (a) to choose a reasonable expression for the function g from the list below.

Multiple Choice:

- (a) $g(t) = 100 - 9t^2, 0 \leq t \leq 10/3$
- (b) $g(h) = 100 - 9h^2, 0 \leq h \leq 100$
- (c) $g(h) = 0.005(100 - h)^2, 0 \leq h \leq 100$
- (d) $g(h) = 6\sqrt{100 - h}, 0 \leq h \leq 100$ ✓

(c) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the rock's height from 64 feet gives an approximately proportional change in its speed.

(d) Use the results of part (c) to approximate the proportionality constant. What are its units?

(e) Use the algebra of limits to find the exact value of the proportionality constant.

(f) Explain the meaning of the proportionality constant.

(g) Approximate the change

$$\Delta v = v - g(64)$$

in the rock's speed in terms of a small change

$$\Delta h = h - 64$$

in its height.

(h) Use part (g) to approximate the rock's speed at a height of 63 feet.

(i) Would you expect your approximation in part (h) to be greater or less than the actual speed? Explain your reasoning with a graph.

(j) Simplify the units of the proportionality constant. Does this simplification help to understand or obscure the meaning of the proportionality constant?

5 Limits and Gas Mileage

6 Limits and Purchasing Power

Free Fall

Rocks in free fall.

1 Speed vs. Velocity

In colloquial English, speed and velocity are typically used interchangeably. A baseball announcer might say, for example, that a pitcher has good velocity, when he really means the pitcher is throwing hard (fast).

But we need to be precise and for us *velocity* is a vector. It is the rate of change of position (also a vector) with respect to time. As such, we can talk about *average velocity* over a time interval and *instantaneous velocity* at a moment in time. For example, at some instant we might be driving due south at a speed of 50 miles/hour and this would be a description of our instantaneous velocity. Or over a three-hour period we might have driven 120 miles and ended up at a point due south of where we started. Then we could describe our average velocity over this three-hour period as a displacement (ie. change in position) due south at an average speed of 40 miles/hour.

In colloquial English *acceleration* is also used incorrectly. An advertisement might claim a car can accelerate from rest to a speed of 60 miles/hour in 4 seconds. But acceleration, being the rate of change of velocity with respect to time, is also a vector. So instead, we might say that a car is speeding up at a constant rate of 22 (ft/sec)/sec. This implies, for example, that every second the speed of the car increases by 22 ft/sec. It would *not* be correct to say that the car's acceleration is 22 (ft/sec)/sec. Acceleration, like velocity, needs a direction.

It also obscures meaning if we say that the speed of a car is increasing at the rate of 22 ft/sec². What exactly do we mean by a square second. Better to say what we really mean, that the speed is increasing at the rate of 22 (ft/sec)/sec.

Velocity plays only a very small part in our class, and this activity is all about speed. Velocity is defined as a rate of change (of position with respect to time), but what about speed? To think of speed this way we need a *cumulative distance function*. The odometer of your car is one such function. It records the total number of miles the car has been driven since its birth. The trip odometer also keeps track of cumulative distance. In either case, the average rate of change of the odometer reading with respect to time measures average speed (distance travelled / time) over an interval of time. The limit of these average rates of

Learning outcomes:
Author(s):

change, as the time interval shrinks to zero, measures the (instantaneous) speed at any moment. This is the reading on the speedometer.

For this class, motion is generally restricted to a line, and the cumulative distance function measures distance along this line from some select point. Later on, you'll learn about *arclength*, distance measured along a curve. Typically the parameter s is used for this distance function. Don't confuse this with speed. In this activity we drop a rock and measure the cumulative distance function $s = f(t)$ straight down, from the rock's initial position.

It's a consequence of Newton's second law and his law of universal gravitation, that in the absence of air resistance the speed of a rock dropped near the surface of a planet increases at a constant rate. On the earth, this rate is about 32 (ft/sec)/sec (or about 9.8 (m/s)/s). On the moon, the rate is about one-sixth these values. On Mars, the *magnitude* of the gravitational acceleration is approximately 3.73 (m/s)/s.

2 A Rock in Freefall

Suppose we drop a rock from rest near the surface of a planet without an atmosphere (or very near the surface of the earth). Our focus here will be in addressing two questions. Don't try to answer these now. Wait until the end of the activity.

- Question 60** (a) How does the speed at which the rock hits the surface compare with its average speed during the time it takes to hit the surface?
- (b) Through approximately what fraction of its initial height does the rock fall during the last 1/20 of the time it takes to hit the surface? Or during the last similarly small fraction of the falling time?
- (c) What is the exact fraction?

Question 61 Play the slider u in Line 2 of the desmos worksheet below to watch two motions. One shows a rock falling to the surface of a planet when it is dropped from rest near the surface. The other is a constant-speed motion.

Desmos link: <https://www.desmos.com/calculator/dmlrxahkld>

151: Free Fall 2

- (a) Which motion is which?

- (b) What does the animation suggest about the speed of the rock as it hits the surface compared with its average speed during the time it takes to hit the ground?
- (c) What does the animation suggest about the fraction of its initial height the rock falls through during the last $1/20$ of the time it takes to hit the surface? Give a quick estimate.
- (d) Now suppose the rock is dropped from a height of H meters and that it takes T seconds to hit the surface. Then, as we'll verify another day, the function expressing the rock's distance from its starting point (in meters) in terms of the number of seconds since it was dropped is of the form

$$s = f(t) = kt^2$$

for some positive constant k .

- (i) What are the units of the constant k ? How do you know?
- (ii) Express the constant k in terms of H and T .
- (iii) Find a fully-simplified expression for the average speed of the rock over the time interval $b \leq t \leq w$. Assume $b < w$.
- (iv) Use the algebra of limits to find an expression for the (instantaneous) speed of the rock b seconds after being dropped. Check that your expression has the correct units.
- (v) Use the result of part (iv) to express in terms of H and T the speed of the rock (in metes/sec) as it hits the surface. Check that your expression has the correct units. Hint: Evaluate the derivative above at the appropriate time $t = b$.
- (vi) Compare the speed of the rock as it hits the ground with its average speed during the time it falls.
- (vii) Use the result of part (vi) to approximate the fraction of its initial height through which the rock fall during the last $1/20$ of the time it takes to hit the surface
- (viii) Through what exact fraction of its initial height the rock fall through during the last $1/20$ of the time it takes to hit the surface? Compare this with your earlier estimate.

3 Takeaways

In the last problem you showed that the derivative of the function

$$y = f(x) = kx^2$$

is

$$\frac{dy}{dx} = \frac{d}{dx}(x^2) = 2kx.$$

But in some contexts, like for the falling rock, it is more insightful to write to write this derivative as

$$\begin{aligned}\frac{dy}{dx} &= 2kx \\ &= 2\left(\frac{2kx^2}{x}\right), x \neq 0, \\ &= 2\left(\frac{y}{x}\right), x \neq 0.\end{aligned}$$

Geometrically, this says that the slope ($2ka$) of the tangent line to the parabola

$$y = f(x) = kx^2, k \neq 0,$$

at the point $P(a, ka^2)$ is *twice* the slope (ka) of the line OP through the origin and P .

Desmos link: <https://www.desmos.com/calculator/aamjtxiqgi>

151: Squaring Function

In the context of a rock dropped from rest, this means that the rock's speed at any time t is twice its average speed during the time interval from the moment of release to time t .

This way of writing the derivative has other implications as well.

For example, in the context of the function

$$A = s^2, s > 0,$$

that expresses the area of a square of its side length, we have for $\Delta s \sim 0$,

$$\frac{\Delta A}{\Delta s} \sim \frac{dA}{ds} = \frac{2A}{s}.$$

So that for $\Delta s \sim 0$,

$$\frac{\Delta A}{A} \sim 2\left(\frac{\Delta s}{s}\right).$$

In other words, the relative change in the area of a square (or the relative error in computing the area) is approximately twice the relative change in its side length (or twice the relative error in measuring its side length), at least for *small* changes in (or errors in measuring) the side length.

We don't really need calculus to understand this. A picture does just as well.

Desmos link: <https://www.desmos.com/calculator/h2fm6mm8ua>

151: Square Error

Question 62 *Explain exactly what the picture suggests and how it does so.*

Free Response:

For more on this idea, see the chapter about *Elasticity*.

The Weighted Chain

Lifting a suspended weight.

Question 63 Attach a small weight to the center of a twenty-inch string, hold the ends of the string together and then pull your hands apart. Play the slide *a* in Line 2 of the animation below to watch the motion.

Desmos link: <https://www.desmos.com/calculator/jmqscra2if>

Desmos activity available at 151: [Weighted Chain 22](#)

What we're most interested in here is how moving your hands a bit farther apart changes the height of the weight. You might, for example, move each hand 1 cm outward. What happens to the weight? Perhaps it moves upward about half as far, only 0.5 cm. Or perhaps about twice as far. The ratio of these two small distances depends upon the distance between your hands. And while we're not literally stretching an elastic band here, the effect is the same. We can talk about the local scaling factor, the one that converts a small outward movement of your hands to an approximate change in height. This local scaling factor, in some configurations less than one, in others greater than one, is the derivative. Our aim here is to see how this factor depends on the distance between your hands, or really half that distance.

- (a) Play the animation. Then sketch by hand a graph of the function

$$h = f(s), 0 \leq s \leq 10$$

that expresses the distance (in inches) of the weight from its starting point in term of half the distance (in inches) between your hands (ie. the distance *MR* above). Activate the folder in Line 15 to see how you did.

- (b) Use the graph of the height function $h = f(s)$ to sketch by hand a rough graph of its derivative $r = dh/ds = f'(s)$. Be sure to label the axes with the appropriate variable names and units. Keep in mind the following when drawing your graph:
- (i) For what values of s is the scaling factor less than one?
 - (ii) For what value(s) of s is the scaling factor equal to one?
 - (iii) Greater than one?
 - (iv) Equal to zero?

Learning outcomes:
Author(s):

Then activate the folder in Line 20 to see how you did.

- (c) Find an expression for the function $h = f(s)$.
- (d) Use the geometry of the curve $h = f(s)$ to evaluate the local scaling factor

$$\left. \frac{dh}{ds} \right|_{s=b}.$$

- (e) Now we'll go about computing this scaling factor algebraically:
 - (i) Find an expression for the average rate of change of h with respect to s between $s = b$ and $s = w$.
 - (ii) Use the algebra of limits to find an expression for the derivative

$$\left. \frac{dh}{ds} \right|_{s=b}.$$

- (f) Drag the slider a in Line 2 to $a = 9.5$ and use the animation to approximate the derivative

$$\left. \frac{dh}{ds} \right|_{s=9.5}$$

- (g) Compute the exact value of the above derivative. Include units. Compare this with your approximation.
- (h) Explain the meaning of the above derivative in terms of small changes.

Question 64 The last question had nothing to do with time. But suppose now, for example, we move our hands outward at a constant speed of 4 inches/sec. Then the speed of the weight would depend on the half-distance between our hands. How?

We'll work more generally, and suppose we have a function

$$s = g(t), \quad 0 \leq t \leq T_0,$$

that expresses the distance MR (ie. half the distance between our hands), measured in inches, in terms of the number of seconds since we began moving our hands.

Desmos link: <https://www.desmos.com/calculator/x0e7h7sobg>

Desmos activity available at 151: Weighted Chain 23

Purchasing Power

How a small change in price affects the quantity of an item you can buy.

1 Purchasing Power

Question 65 This problem investigates how a small change in the price of apples would affect the number of pounds we can buy with \$10.

- (a) We'll start by finding a function

$$n = f(p), p > 0,$$

that expresses the number of pounds of apples we can buy with \$10 in terms of the price (measured in dollars/pound).

The function is

$$n = f(p) = 10/p, p > 0.$$

- (b) Would you expect the derivative dn/dp to be positive, negative, or zero? Explain.
- (c) Use limits to find an expression for the derivative

$$\left. \frac{dn}{dp} \right|_{p=a}.$$

The derivative is (click the Hint tab above for help)

Hint:

$$\begin{aligned} \left. \frac{dn}{dp} \right|_{p=a} &= \lim_{p \rightarrow a} \frac{1}{p-a} (f(p) - f(a)) \\ &= \lim_{p \rightarrow a} \frac{1}{p-a} \left(\frac{10}{p} - \frac{10}{a} \right) \\ &= \lim_{p \rightarrow a} \frac{10}{p-a} \left(\frac{a-p}{pa} \right) \\ &= \lim_{p \rightarrow a} \frac{-10}{pa} \\ &= \frac{-10}{a^2}. \end{aligned}$$

Learning outcomes:
Author(s):

- (d) Evaluate the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

- (i) What are the units of the above derivative?
 (ii) What do you get by “simplifying” the units? Explain how simplifying the units of this derivative gives you insight into its meaning.
- (e) Interpret the meaning of the above derivative. Click the arrow to the lower right for help.

For a way to interpret the meaning of the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

we can ask ourselves what happens to the number of pounds of apples we can buy with \$10 if the price changes by a small amount from \$2/lb. For this, let Δp be a small change in price from \$2/lb (measured in dollars/lb) and let Δn be the corresponding change in the number of pounds we can buy with \$10.

For $\Delta p \sim 0$, we have

$$-2.5 = \left. \frac{dn}{dp} \right|_{p=2} \sim \frac{\Delta n}{\Delta p}$$

and (type “Delta” for Δ)

$$\Delta n \sim -2.5 \Delta p.$$

This tells us that if the price increases from \$2/lb to say \$2.10/lb, then

$$\Delta n \sim \left(-2.5 \frac{\text{lb}}{\$/\text{lb}} \right) \left(0.10 \frac{\$}{\text{lb}} \right) = -0.25 \text{ lbs.}$$

So if the price increases from \$2/lb to say \$2.10/lb, we can buy about 0.25 fewer pounds of apples with ten dollars.

- (f) Next we’ll use the worksheet below to visualize the approximate change

$$\begin{aligned} \Delta n &\sim \left(\left. \frac{dn}{dp} \right|_{p=2} \right) \Delta p \\ &= \left(-2.5 \frac{\text{lb}}{\$/\text{lb}} \right) \left(0.10 \frac{\$}{\text{lb}} \right) \\ &= -0.25 \text{ lbs.} \end{aligned}$$

in the number of pounds of apples we can buy with \$10 as the price increases from \$2/lb to \$2.10/lb.

Desmos link: <https://www.desmos.com/calculator/qw7wis1q0c>

151: Apples

- (i) To get started, find an equation (in point-slope form) of the tangent line to the curve $n = f(p)$ at the point P with coordinates $(2, 5)$. The tangent line has a slope equal to the derivative

$$\left. \frac{dn}{dp} \right|_{p=2}.$$

So its equation is

$$n = 5 + -2.5(p - 2).$$

- (ii) Enter your equation of the tangent line on Line 17 of the desmos worksheet.
- (iii) Activate the folders tangent line and linear approximation on Lines 18 and 22.
- (iv) Explain why the difference in the n -coordinates of points P and R' (ie. n -coordinate of R' minus n -coordinate of P) is equal to our approximation of Δn above when Q as coordinates $(2.10, f(2.10))$.
- (v) Drag Slider v on Line 2 to make Q approach P . What so you think happens to the ratio of Δn to our approximation of Δn as $v \rightarrow 2$?
- (g) We can get a better understanding of the changes

$$\Delta p = 2.1 - 2 = 0.10 \text{ dollars/pound}$$

and

$$\Delta n = f(2.1) - f(2) \sim 0.25 \text{ pounds}$$

by thinking about relative instead of absolute change.

An increase in price from \$2/lb to \$2.10/lb is a relative change of

$$\frac{\Delta p}{p} = \frac{\$0.1/\text{lb}}{\$2/\text{lb}} = 0.05 = 5\%.$$

This causes a relative change in the number of pounds we can buy of approximately

$$\frac{\Delta n}{n} \sim \frac{-0.25\text{lbs}}{5\text{lbs}} = -0.05 = -5\%.$$

- (h) We can get a the same result relating the two relative changes of the previous question by working in general.

Suppose we increase the price of apples by $Q\%$, where $Q \sim 0$. What can we say about the relative change in the number of pounds of apples we can buy with \$10?

Well, if the change in price Δp is near zero and Δn is the corresponding change in the number of pounds we can buy, then

$$\frac{\Delta n}{\Delta p} \sim \frac{dn}{dp} = -10/p^2.$$

So

$$\Delta n \sim \left(\frac{-10}{p^2} \right) \Delta p.$$

Dividing both sides by $n = 10/p$ tells us that

$$\frac{\Delta n}{n} \sim \left(\frac{-10}{p^2} \right) \left(\frac{\Delta p}{\frac{10}{p}} \right) = -\frac{\Delta p}{p} = -Q\%.$$

So if we increase the price by $Q\% \sim 0$, then the number of apples we can buy decreases by approximately that same $Q\%$.

- (i) We can think about this relationship between the relative changes geometrically. To do this, remove your equation of the tangent line on Line 17 and activate the folder tangent line in the worksheet above. Then drag slider a on Line 4 to move point P . What do you notice about the ratio $PB : PA$ of the distances PA and PB as P moves? More on this later.

Question 66 Due to a printing error, the graph of the function

$$n = f(p), \quad p > 0,$$

expressing the number of pounds of an item we can buy with \$100 in terms of the price (measured in \$/lb) is not shown below. All we can see is a point A on the graph.

Desmos link: <https://www.desmos.com/calculator/y5nqs8fkvj>

151: Printers Error 1

The problem is to draw the tangent line to the graph at A without sketching the graph or doing any kind of computation. Click the arrow below for a hint.

The key idea is to relate the slope of the tangent line at A to the slope of the line OA through the origin and A .

We know that

$$n = f(p) = \frac{100}{p}, \quad p > 0,$$

and

$$\frac{dn}{dp} = -100/p^2.$$

Now write this derivative in terms of both n and p ,

$$\frac{dn}{dp} = -\frac{100}{p^2} = -\frac{n}{p}.$$

Now relate this to the slope of line OA and draw the tangent line at A .

2 Weight in Space

Question 67 The weight of an object is the gravitational force that the earth exerts on the object's mass and varies with the object's height above the surface.

The function

$$W = f(h) = \frac{k}{(h + 4)^2}, \quad h \geq 0,$$

expresses the weight (in pounds) of an object in terms of its height (in thousands of miles) above the surface of the earth. We'll suppose the object weighs 200 pounds on the surface.

Desmos link: <https://www.desmos.com/calculator/zklsucctjp>

151: Weight in Space

- Find the value of the constant k . What are its units?
- Find the average rate of change of W with respect to h over the interval between heights of h and v thousands of miles.
- Use part (b) to find an expression for the derivative dW/dh .
- Use the graph of the function $W = f(h)$ above to approximate the derivative

$$\left. \frac{dW}{dh} \right|_{h=4}.$$

Include units

- Evaluate the derivative

$$\left. \frac{dW}{dh} \right|_{h=4}$$

using your expression for dW/dh and compare this with your estimate.

- What are the units of the derivative above? Interpret its meaning.

- (i) At approximately what altitude does your weight decrease by 4 pounds when your altitude increases by 200 miles? Use your expression for the derivative dW/dh to help.
- (ii) Compute the exact change in your weight over the interval you found in part (i).

3 A Rock in Free Fall

Question 68 A rock is dropped from a height of 100 meters on the planet Krypton.

The function

$$v = f(h), 0 \leq h \leq 100,$$

expresses the speed (in meters/sec) of the rock in terms of its height (in meters) above the surface.

Desmos link: <https://www.desmos.com/calculator/hcqb5nxc18>

151: Free Fall Speed and Height

- (a) Would you expect the derivative

$$\left. \frac{dv}{dh} \right|_{h=36}$$

to be positive or negative? Why?

- (b) Suppose

$$v = f(h) = 4\sqrt{100 - h}, 0 \leq h \leq 100$$

and find an expression for the average rate of change of the rock's speed (measured in meters/sec) with respect to its height (in meters) between heights of v meters and a meters.

- (c) Use your expression from part (b) to find an expression for the derivative

$$\left. dv/dh \right|_{h=a}.$$

- (d) Use the graph above to approximate the derivative

$$\left. dv/dh \right|_{h=64}.$$

Include units.

- (e) Evaluate the derivative

$$dv/dh \Big|_{h=64}$$

and compare the exact value with your estimate.

- (f) What are the units of the derivative above? What is its meaning?
- (g) “Simplify the units of the derivative”. What insight does this give you into its meaning?
- (h) Use part (e) to approximate the rock’s speed at a height of 61 feet. Compare your estimate with the actual speed.
- (i) Use part (c) to approximate the interval over which the rock falls two meters as its speed increases by 0.5m/s.

4 Distance to the Horizon

Question 69 The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 10,000$$

expresses the distance to the horizon (measured in miles) in terms of your altitude (measured in feet).

- (a) Find an expression for the average rate of change in the distance to the horizon with respect to altitude between altitudes h feet and v feet.
- (b) Use your expression from part (a) to find an expression for the derivative ds/dh .
- (c) Evaluate the derivative

$$\frac{ds}{dh} \Big|_{h=25}.$$

- (d) What are the units of the derivative above? Interpret its meaning.
- (e) Use the result of part (c) to approximate the distance to the horizon at an altitude of 24 feet. Then compare this approximation with the actual distance.
- (f) Approximate the relative change in the distance to the horizon in terms of a small relative change in altitude.
- (g) Use your result from part (b) to approximate the altitude at which moving 10 feet higher increase the distance to the horizon by 0.5 miles.

5 Intensity of Sound

Question 70 The intensity of sound is measured in Watts per square meter and is a function of the distance from the source.

Suppose for a jet taking off, this function is

$$I = f(r) = \frac{100}{r^2}, \quad r \geq 10,$$

where r is the distance (in meters) from the jet.

- (a) Find an expression for the average rate of change of the sound intensity (in Watts/m²) with respect to the distance from the source (in meters) between distances r and a meters.
- (b) Use your expression from part (a) to find an expression for the derivative

$$dI/dr \Big|_{r=a}$$

- (c) Evaluate the derivative

$$dI/dr \Big|_{r=20}.$$

- (d) What are the units of the derivative above? What is its meaning?
- (e) Approximate the relative change in the sound intensity in terms of a small relative change in the distance to the source.



Practice Quiz 1

First practice quiz, Weeks 1-2

Directions:

- (a) Show all work.
- (b) Give brief explanations for each problem. Include these explanations in the flow of the solution.
- (c) Show all units in all computations.

1 Part 1

Question 71 Explain what it means intuitively for a function to be differentiable at some input in terms of the graph of that function.

Question 72 The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 10,000,$$

expresses the distance to the horizon (measured in miles) in terms of your altitude (measured in feet).

- (a) Find an expression for the average rate of change in the distance to the horizon with respect to altitude between altitudes b feet and v feet.
- (b) Use your expression from part (a) to find an expression for the derivative

$$\left. \frac{ds}{dh} \right|_{h=b}.$$

- (c) Evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=25}.$$

- (d) What are the units of the derivative above? Interpret its meaning.

Learning outcomes:
Author(s):

- (e) Use the result of part (c) to approximate the distance to the horizon at an altitude of 24 feet. Then compare this approximation with the actual distance.
- (f) Approximate the relative change in the distance to the horizon in terms of a small relative change in altitude.
- (g) Use your result from part (b) to approximate the altitude at which moving 10 feet higher increase the distance to the horizon by 0.5 miles.

Question 73 The intensity of sound is measured in Watts/(square meter) and is a function

$$I = f(r) = \frac{100}{r^2}, \quad r \geq 0.5,$$

of the distance from the source.

Suppose r is measured in meters and that the intensity of the sound emitted by a vacuum cleaner is 10^{-4} watts/ m^2 at a distance of 0.5 meters.

- (a) Compute the value of the constant k . What are its units?
- (b) Find an expression for the average rate of change of the sound intensity (in Watts/ m^2) with respect to the distance from the source (in meters) between distances b and r meters.
- (c) Use your expression from part (b) to find an expression for the derivative

$$\left. \frac{dI}{dr} \right|_{r=b}.$$

- (d) Evaluate the derivative

$$\left. \frac{dI}{dr} \right|_{r=2}.$$

- (e) What are the units of the derivative above? What is its meaning?
- (f) Approximate the relative change in the sound intensity in terms of a small relative change in the distance to the source.

Question 74 Between speeds of 70 miles/hr and 84 miles/hr, the gas mileage of a car (in miles/gal) is a one-to-one function $G = f(v)$ of its speed (in miles/hour). The car gets 10 miles/gal at a speed of 80 miles/hour.

- (a) Which of the following is more likely to be true? Explain your reasoning.

(i) $\left. \frac{dG}{dv} \right|_{v=80} = 0.25$ or

(ii) $\left. \frac{dG}{dv} \right|_{v=80} = -0.25$

(b) What are the units of the correct derivative above? Explain its meaning.

(c) Assuming the correct choice in part (b), evaluate the derivative

$$\left. \frac{dv}{dG} \right|_{G=10}.$$

(d) Simplify the units of the derivative in part (c). What does this tell you about its meaning?

(e) At what rate (in gal/hour) does the car burn gas at a speed of 80 miles/hour?

Question 75 Between speeds of 55 miles/hour and 70 miles/hour the gas mileage of a car is a linear function of its speed. The car gets 40 miles/gal at a speed of 55 miles/hour and 30 miles/gal at a speed of 70 miles/hour.

(a) Find a function

$$r = f(v), \quad 55 \leq v \leq 70,$$

that express the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hour). Note: This function is not linear

(b) Find an expression for the average rate of change of r with respect to v between speeds of b miles/hour and w miles/hour.

(c) Use your expression from part (b) to find an expression for the derivative

$$\left. \frac{dr}{dv} \right|_{v=b}.$$

(d) Evaluate the derivative

$$\left. \frac{dr}{dv} \right|_{v=60}.$$

(e) What are the units of the derivative above? Interpret its meaning.

Introduction to Motion

An introduction to motion

Exploration 76 *Desmos link:*

<https://www.desmos.com/calculator/bk1z9cwbhb>

151: Sound Squaring Function

Learning outcomes:
Author(s):

The Derivative

Computing derivatives with limits.

Example 29. Suppose

$$y = f(x) = x^2$$

and let's use limits to evaluate

$$f'(3) = \left. \frac{dy}{dx} \right|_{x=3} = \left. \frac{d(x^2)}{dx} \right|_{x=3}.$$

We have

$$\begin{aligned} f'(3) &= \left. \frac{d(x^2)}{dx} \right|_{x=3} = \lim_{v \rightarrow 3} \frac{f(v) - f(3)}{v - 3} \\ &= \lim_{v \rightarrow 3} \frac{v^2 - 9}{v - 3} \\ &= \lim_{v \rightarrow 3} \frac{(v - 3)(v + 3)}{v - 3} \\ &= \lim_{v \rightarrow 3} (v + 3) \\ &= (3 + 3) \\ &= 6. \end{aligned}$$

Next let's do almost the same thing and compute

$$f'(x) = \frac{dy}{dx}$$

for the function

$$y = f(x) = x^2$$

by replacing 3 in the above computation with x .

Learning outcomes:
Author(s):

We get

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d(x^2)}{dx} = \lim_{v \rightarrow x} \frac{f(v) - f(x)}{v - x} \\
 &= \lim_{v \rightarrow x} \frac{v^2 - x^2}{v - x} \\
 &= \lim_{v \rightarrow x} \frac{(v - x)(v + x)}{v - x} \\
 &= \lim_{v \rightarrow x} (v + x) \\
 &= (x + x) \\
 &= 2x.
 \end{aligned}$$

Just as a check, when $x = 3$,

$$\left. \frac{dy}{dx} \right|_{x=3} = (2x) \Big|_{x=3} = 2(3) = 6.$$

Question 77 Use the result of the previous example to solve each of the following problems. Do not use limits.

- (a) Find an equation of the tangent line to the parabola $y = x^2$ at the point $(-4, 16)$.
- (b) Find an equation of the tangent line to the parabola perpendicular to the tangent line in part (a).
- (c) Find the coordinates of the point where the lines in parts (a) and (b) intersect.
- (d) Let $\mathcal{L}$ be the line through the points of tangency of the lines in parts (a) and (b). Find the coordinates of the point where $\mathcal{L}$ intersects the y -axis.
- (e) Repeat parts (a)-(d) above for the tangent line to the parabola $y = x^2$ at the general point (a, a^2) . What do you notice?

Question 78 (a) Use the method of Example 1 for the function

$$y = g(x) = 1/x^2$$

to compute

$$g'(3) = \left. \frac{dy}{dx} \right|_{x=3} = \left. \frac{d(1/x^2)}{dx} \right|_{x=3}$$

and

$$g'(x) = \frac{dy}{dx}.$$

(b) Use the result of part (a) to find an equation of the tangent line to the curve $y = 1/x^2$ at the point $(3, 1/9)$.

Question 79 (a) Use numerical methods to estimate the slope of the tangent line to the curve

$$y = f(x) = x^3$$

at the point $(2, 8)$. Include enough data to suggest a progression toward a limit.

(b) Use the algebra of limits to find the exact slope of the tangent line in part (a).

(c) Use algebra to find the coordinates of the all pointw where the tangent line in part (a) intersects the curve $y = x^3$.

(d) Suppose you measure the edge length of a cube to be 2cm and then use this measurement to compute the volume of the cube. Use the result of part (b) to approximate your error in computing the volume in terms of your error in measuring the edge length. Assume the latter error is small.

Then compare your exact error in computing the volume with your approximation for some specific edge length near 2cm. You should start this problem defining a function with meaningful variable names (do not use x and y).

Question 80 (a) Find a function

$$s = g(V), V \geq 0$$

that expresses the edge length (measured in cm) of a cube in terms of its volume (measured in cubic centimeters).

(b) Use the algebra of limits to evaluate the derivative

$$g'(V_0) = \left. \frac{ds}{dV} \right|_{V=V_0}.$$

(c) What are the units of the derivative in part (b)? Explain how you know.

(d) Suppose you submerge the cube in water and measure its volume to be 8 cm^3 . You then use this measurement to compute the edge length of the cube. Use the result of part (b) to approximate your error in computing the edge length in terms of your error in measuring the volume. Assume the latter error is small.

Start this problem by defining the errors Δs and ΔV in terms of the actual edge length s of the cube.

(e) Use the result of part (d) to approximate your error in computing the edge length of the cube if the cube's actual volume is 7.7 cm^3 .

(f) Explain how this question is related to Question 4.

Question 81 Suppose that between speeds of 60 miles/hr and 72 miles/hr, the gas mileage of a car is a linear function of its speed. Suppose also that the car gets 36 miles/gallon at a speed of 60 miles/hour and 32 miles/gallon at a speed of 72 miles/hour.

(a) Find a function

$$r = f(v), 60 \leq v \leq 72,$$

that expresses the rate (measured in gal/hr) at which the car burns gas in terms of its speed (measured in miles/hour). Explain your reasoning. This function is not linear.

Hint:

$$r = f(v) = \frac{3v}{168 - v}, 60 \leq v \leq 72.$$

(b) Use numerical methods to estimate the value of the derivative

$$f'(63) = \left. \frac{dr}{dv} \right|_{v=63}.$$

Make a table that shows enough data to suggest a progression toward a limit. Include units in all column headings.

(c) Use the algebra of limits to find an expression for the derivative

$$f'(v) = \frac{dr}{dv}.$$

Then use this expression to find the exact value of the derivative in part (b).

(d) What are the units of the derivative in part (b)? Explain its meaning.

(e) Use the result of part (c) to approximate the change

$$\Delta r = f(v) - f(63)$$

in the rate at which your car burns gas in terms of the change

$$\Delta v = v - 63$$

of the car's speed. Assume $\Delta v \sim 0$.

Question 82 Suppose that between speeds of 30 miles/hour and 70 miles/hour the gas mileage of a car is a quadratic function of its speed. Suppose also that

the car gets a maximum of 42 miles/gal at a speed of 50 miles/hour and 34 miles/gallon at a speed of 30 miles/hour.

(a) Find a function

$$r = h(v), \quad 30 \leq v \leq 70,$$

that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hour).

Hint: (i) At what rate does the car burn gas at a speed of 50 miles/hour? 25/21 gal/mile

(ii) Find a function that expresses the gas mileage G (measured in miles/gallon) in terms of the speed (measured in miles/hr).

$$G = 42 - 0.02(v - 50)^2, \quad 30 \leq v \leq 70.$$

(ii) The rate (in gal/hr) at which the car burns gas as a function of its speed (in miles/hr) is

$$r = h(v) = \frac{v}{42 - 0.02(v - 50)^2}, \quad 30 \leq v \leq 70.$$

(b) Use the algebra of limits to evaluate the derivative

$$h'(40) = \left. \frac{dr}{dv} \right|_{v=40}.$$

(c) What are the units of the above derivative? How do you know?

(d) Express the meaning of the derivative in the context of small changes.

Question 83 At 10:00am on April 18, the wholesale price of Cosmic Crisp apples is \$2.00/lb and is decreasing at the rate of \$0.10/lb/hour.

Use the algebra of limits to determine the rate (in pounds/hour) at which the number of pounds of apples a store can purchase with \$1000 is changing at this time.

Start this question by defining a function that expresses the number of pounds of apples the store can buy with \$1000 in terms of the price (in \$/lb). Choose meaningful variable names (not x and y). Do not assume the price is decreasing at a constant rate.

1 Limits, Speed and Altitude

Question 84 A rock dropped from a height of 100 feet falls to the surface of Planet Krypton without air resistance.

(a) By considering only the physical situation and without doing any computations, sketch a graph of the function

$$v = g(h), 0 \leq h \leq 100$$

that expresses the rock's speed (in ft/sec) in terms of its height (in feet). Explain your reasoning.

(b) Use the results from part (a) to choose a reasonable expression for the function g from the list below.

Multiple Choice:

- (a) $g(t) = 100 - 9t^2, 0 \leq t \leq 10/3$
- (b) $g(h) = 100 - 9h^2, 0 \leq h \leq 100$
- (c) $g(h) = 0.005(100 - h)^2, 0 \leq h \leq 100$
- (d) $g(h) = 6\sqrt{100 - h}, 0 \leq h \leq 100$ ✓

(c) Give numerical and graphical evidence that either supports or refutes the claim that a small change in the rocks height from 64 feet gives an approximately proportional change in its speed. Then approximate the proportionality constant. What are its units?

(d) Use the algebra of limits to find an expression for the derivative

$$g'(h) = \frac{dv}{dh}.$$

Then use this expression to find the exact value of the proportionality constant in part (c).

(e) Explain the meaning of the proportionality constant.

(f) Approximate the change

$$\Delta v = v - g(64)$$

in the rock's speed in terms of a small change

$$\Delta h = h - 64$$

in its height.

- (g) Use part (g) to approximate the rock's speed at a height of 63 feet.
 - (h) Would you expect your approximation in part (h) to be greater or less than the actual speed? Explain your reasoning with a graph.
 - (i) Simplify the units of the proportionality constant. Does this simplification help to understand or obscure the meaning of the proportionality constant?
-

Derivatives of Polynomials

Working with polynomials and their derivatives.

1 Differentiating Polynomials

Question 85 Let $f(x) = x^5$.

(a) Use the algebra of limits to find an expression for the derivative

$$\frac{d}{dx}(f(x)) = \frac{d}{dx}(x^5).$$

Explanation. The average rate of change of the function $y = f(x) = x^5$ between $x = a$ and $x = b$ ($b \neq a$) is

$$\begin{aligned} \frac{\Delta y}{\Delta x} &= \frac{f(b) - f(a)}{b - a} \\ &= \frac{b^5 - a^5}{b - a} \\ &= \frac{(b - a)(b^4 + b^3a + b^2a^2 + ba^3 + a^4)}{b - a} \\ &= b^4 + b^3a + b^2a^2 + ba^3 + a^4 \text{ since } b \neq a. \end{aligned}$$

So

$$\begin{aligned} \left. \frac{d}{dx}(x^5) \right|_{x=a} &= \lim_{b \rightarrow a} \frac{f(b) - f(a)}{b - a} \\ &= \lim_{b \rightarrow a} (b^4 + b^3a + b^2a^2 + ba^3 + a^4) \\ &= a^4 + a^3a + a^2a^2 + aa^3 + a^4 \\ &= 5a^4 \end{aligned}$$

and

$$\frac{d}{dx}(x^5) = 5x^4.$$

(b) Use the result of part (a) to find an expression for the derivative

$$\frac{d}{dx}(f^{-1}(x)) = \frac{d}{dx}(x^{1/5}).$$

Use set-builder notation to state the domains of $f^{-1}(x)$ and its derivative.

Learning outcomes:
Author(s):

Explanation. Let

$$y = f^{-1}(x) = x^{1/5}.$$

Then the slope of the line between the points $(a, a^{1/5})$ and $(b, b^{1/5})$, $a \neq b$, on the graph of the function $y = x^{1/5}$ is

$$\frac{\Delta y}{\Delta x} = \frac{b^{1/5} - a^{1/5}}{b - a}.$$

To make this average rate of change look more like the one in part (a), let $a_1 = a^{1/5}$ and $b_1 = b^{1/5}$. Then $a = a_1^5$ and $b = b_1^5$. So

$$\begin{aligned}\frac{\Delta y}{\Delta x} &= \frac{b^{1/5} - a^{1/5}}{b - a} \\ &= \frac{b_1 - a_1}{b_1^5 - a_1^5}.\end{aligned}$$

Now as $b \rightarrow a$, $b_1 \rightarrow a_1$, so

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{x=a} &= \lim_{b \rightarrow a} \frac{b^{1/5} - a^{1/5}}{b - a} \\ &= \lim_{b_1 \rightarrow a_1} \frac{b_1 - a_1}{b_1^5 - a_1^5} \\ &= \frac{1}{5a_1^4}\end{aligned}$$

where the last equality follows from our work in part (a). To express this derivative in terms of a , recall that $a_1 = a^{1/5}$. So

$$a_1^4 = \left(a^{1/5}\right)^4 = a^{4/5}$$

and

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{x=a} &= \frac{1}{5a_1^4} \\ &= \frac{1}{5a^{4/5}} \\ &= \frac{1}{5}a^{-4/5}.\end{aligned}$$

Our conclusion is that

$$\frac{d}{dx} \left(x^{1/5} \right) = \frac{1}{5} x^{-4/5}.$$

The domain of the function $f(x) = x^{1/5}$ is the set of all real numbers, but the domain of its derivative is the set

$$\{x \in \mathbb{R} | x \neq 0\}.$$

Explain geometrically why the derivative is undefined at $x = 0$. Include a graph of the function $y = x^{1/5}$ to help with your explanation.

Free Response:

- (c) Use the result of part (a) and the algebra of limits to find an expression for the derivative

$$\frac{d}{dx} \left(\frac{1}{f(x)} \right) = \frac{d}{dx} (x^{-5}).$$

Explanation.

$$\begin{aligned} \frac{d}{dx} (x^{-5}) \Big|_{x=a} &= \lim_{b \rightarrow a} \frac{b^{-5} - a^{-5}}{b - a} \\ &= \lim_{b \rightarrow a} \frac{a^5 - b^5}{b^5 a^5 (b - a)} \\ &= - \lim_{b \rightarrow a} \frac{1}{b^5 a^5} \lim_{b \rightarrow a} \frac{b^5 - a^5}{b - a} \\ &= - \left(\frac{1}{a^{10}} \right) (5a^4) \\ &= - \frac{5}{a^6} \\ &= 5a^{-6}. \end{aligned}$$

Our conclusion is that

$$\frac{d}{dx} (x^{-5}) = 5x^{-6}.$$

- (d) Use the results of parts (a)-(c) to find equations of the tangent lines to the three curves $y = f(x)$, $y = f^{-1}(x)$, and $y = 1/f(x)$ at the point $(1, 1)$. Graph the curves and their tangent line in Desmos to check your work.

Explanation. The slope of the tangent line to the curve

$$y = x^5$$

at the point $(1, 1)$ is

$$\begin{aligned} \frac{dy}{dx} \Big|_{x=1} &= \frac{d}{dx} (x^5) \Big|_{x=1} \\ &= 5x^4 \Big|_{x=1} \\ &= 5. \end{aligned}$$

So an equation of the tangent line to the curve $y = x^5$ at the point $(1, 1)$ is

$$y - 1 = 5(x - 1).$$

I'll leave the other two equations and the graphs for you.

Question 86 (a) Answer the question at the end of the Explanation to Question 1(b).

(b) Complete the last part of Question 1.

(c) Repeat all parts of Question 1 for the function

$$y = f(x) = x^7.$$

(d) Make a conjecture about the derivative

$$\frac{d}{dx}(f(x)) = \frac{d}{dx}(x^n)$$

of the function $f(x) = x^n$.

(e) What do you get for the derivative when $n = 0$? When $n = 1$? Are these results correct? Explain.

Question 87 Due to a printing error, the graph of the function $y = f(x) = 234x^5$ is missing below. All we see is a point A on the graph of the function.

Desmos link: <https://www.desmos.com/calculator/pjyqjtdcxm>

151: Printers Error 2

Without sketching a graph, drag the slider m to draw the tangent line to the curve $y = 234x^5$. Explain your reasoning.

Question 88 (a) Suppose one giraffe is always twice as tall as another. What can you say about their growth rates at any instant?

(b) Suppose one giraffe is always two feet taller than another. What can you say about their growth rates at any instant?

(c) What do parts (a) and (b) suggest about how to compute the derivatives

$$\frac{d}{dx}(f(x) + b),$$

$$\frac{d}{dx}(af(x)),$$

and

$$\frac{d}{dx}(af(x) + b)$$

for constants $a, b \in \mathbb{R}$?

(d) Make up your own scenario that suggests how to compute the derivative

$$\frac{d}{dx}(f(x) + g(x))$$

of the sum of two functions.

Question 89 (a) Find an equation of the tangent line to the curve $y = 36/x$ at the point $P(6, 6)$. Then explain how you could have solved this problem without using calculus.

(b) Find an equation of the tangent line to the curve

$$y = -x^3 + 4x^2 - 3x + 1$$

at the point $(2, 3)$. Graph the curve and its tangent line on Desmos.

Explanation. Since

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(-x^3 + 4x^2 - 3x + 1) \\ &= \frac{d}{dx}(-x^3) + \frac{d}{dx}(4x^2) - \frac{d}{dx}(3x) + \frac{d}{dx}(1) \\ &= -\frac{d}{dx}(x^3) + 4\frac{d}{dx}(x^2) - 3\frac{d}{dx}(x) + 0 \\ &= -3x^2 + 4(2x) - 3(1) \\ &= -3x^2 + 8x - 3, \end{aligned}$$

the slope of the tangent line to the curve at $(2, -3)$ is

$$\left. \frac{dy}{dx} \right|_{x=2} = (-3x^2 + 8x - 3) \Big|_{x=2} = 1.$$

So an equation of the tangent line to the curve

$$y = -x^3 + 4x^2 - 3x + 1$$

at the point $(2, 3)$ is

$$y - 2 = x + 3.$$

(c) The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 30000,$$

expresses the distance to the horizon (in miles) in terms of your altitude (in feet).

(i) Evaluate $f(25)$ and interpret its meaning.

- (ii) Evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=25}.$$

Include units.

- (iii) Use the language of small changes to interpret the meaning of the derivative above.

2 Stock Price

Question 90 The function

$$P = f(t) = 0.1t^3 - t^2 + t + 20, \quad 0 \leq t \leq 8,$$

expresses the price (in dollars/share) of a stock in terms of the number of hours past 9am. The graph of the function is shown below.

Desmos link: <https://www.desmos.com/calculator/gpqicgxvk3>

151: Stock Price

- (a) Is the price increasing or decreasing at 10am? Use the graph to estimate this rate as follows:
 - (i) First drag the slider u (another name for t) on Line 5 to the appropriate value.
 - (ii) Then drag the slider m (the slope of the orange line) so that the orange line looks parallel to the (red) tangent line to the curve $P = f(t)$.
- (b) Use the function to compute the exact rate at which the price is changing at 10am. Include units.
- (c) At what relative rate is the price changing at 10am? Include units.
- (d) At what rate is the number of shares you can buy with \$1000 changing at 10am? Do not use the quotient rule or the chain rule (we have not learned these yet). Use limits instead, but work in general (ie. not with this specific price function). See the Explanation below for help.
- (e) Use the sliders m and b to control the line $P = b + mt$ and approximate when the price is decreasing at the rate of $(\$1.8/\text{share})/\text{hr}$. Then use calculus and algebra to compute the exact time(s).

- (f) Use calculus and technology to approximate the time(s) when the price is decreasing at the relative rate of 10%/hour.
- (g) Use the graph to approximate the minimum and maximum prices of the stock during the eight-hour period. Then use calculus and algebra to compute these exact prices. Hint: What is the value of the derivative dP/dt when the price is a maximum/minimum?
- (h) Use the slider u (another name for the variable t) to approximate the time(s) when the price is decreasing at the fastest rate. Then use calculus to compute the exact time(s).

Hint: It helps to graph the function

$$dP/dt = 0.03t^2 - 2t + 1, 0 \leq t \leq 8,$$

Then think about how you would use calculus to compute the exact time when the price is decreasing at the fastest rate. This is similar to the previous question.

- (i) Use the graph above to approximate the end of the time interval beginning at 9am over which the price decreases at the greatest average rate. Do this by activating the folder secant line on Line 6 (click the circle on the left) and dragging the slider u .
- (j) Use calculus and algebra to find the end of the time interval beginning at 9am over which the price decreases at the greatest average rate.

Hint: Think about the function

$$m(t) = \frac{f(t) - f(0)}{t - 0}, 0 < t \leq 8,$$

that expresses the average rate of change of the price (with respect to time) over the time interval between 9am and time t hours past 9am. Then use this function and the ideas of parts (g) and (h).

Explanation. (d) Here's the solution to part (d).

- Start by finding a function

$$S = g(t), 0 \leq t \leq 8,$$

that expresses the number of shares we can buy with \$1,000 in terms of the number of hours past 9am.

This function is

$$S = g(t) = \frac{1000}{f(t)}, 0 \leq t \leq 8$$

- Next we'll use limits to find an expression for the derivative

$$\left. \frac{dS}{dt} \right|_{t=a}$$

as follows.

$$\begin{aligned} \left. \frac{dS}{dt} \right|_{t=a} &= \lim_{v \rightarrow a} \frac{g(v) - g(a)}{v - a} \\ &= \lim_{v \rightarrow a} \left(\frac{1000}{v - a} \right) \left(\frac{1}{f(v)} - \frac{1}{f(a)} \right) \\ &= \lim_{v \rightarrow a} \left(\frac{1000}{v - a} \right) \left(\frac{f(a) - f(v)}{f(v)f(a)} \right) \\ &= \lim_{v \rightarrow a} \left(\frac{-1000}{f(v)f(a)} \right) \left(\frac{f(v) - f(a)}{v - a} \right) \\ &= \lim_{v \rightarrow a} \frac{-1000}{f(v)f(a)} \cdot \lim_{v \rightarrow a} \frac{f(v) - f(a)}{v - a} \end{aligned}$$

Let's pause here for a moment. The last equality above follows from the fact that both limits in the product exist. The first limit is easy enough to evaluate. The second limit exists because *it is equal to the derivative*

$$\left. \frac{dP}{dt} \right|_{t=a}.$$

That's the key point. Now we know that

$$\begin{aligned} \left. \frac{dS}{dt} \right|_{t=a} &= \lim_{v \rightarrow a} \frac{-1000}{f(v)f(a)} \cdot \lim_{v \rightarrow a} \frac{f(v) - f(a)}{v - a} \\ &= \left(\frac{-1000}{(f(a))^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right) \end{aligned}$$

This is enough to compute the rate at which the number of shares we can buy with \$1000 is changing at 10am.

But we can get a better understanding of this by writing the above expression in a more meaningful way. Remember that $f(a) = P$ is the price at time t . So

$$\begin{aligned} \left. \frac{dS}{dt} \right|_{t=a} &= - \left(\frac{1000}{(f(a))^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right) \\ &= - \left(\frac{1000}{P^2} \right) \left(\left. \frac{dP}{dt} \right|_{t=a} \right) \\ &= - \left(\frac{1000}{P} \right) \left(\frac{1}{P} \cdot \left. \frac{dP}{dt} \right|_{t=a} \right) \\ &= -S \left(\frac{1}{P} \cdot \left. \frac{dP}{dt} \right|_{t=a} \right), \end{aligned}$$

where $S = 1000/P$ is the number of shares we can buy with \$1,000 at a price of P dollars/share.

Now remember that the product

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=a}$$

is the *relative* rate of change in the price (with respect to time) at time $t = a$ hours past 9am.

From part (c) we know at 10am this relative rate is equal to

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=1} = -\frac{0.7}{20.1} \text{hr}^{-1} \sim -3.4826\%/\text{hour}$$

So at 10am we can buy

$$S = \frac{\$1000}{\$20.1/\text{share}} \sim 47.7512 \text{ shares}$$

with \$1,000. And the number of shares we can buy with is increasing at the rate of approximately

$$(47.7512 \text{ shares}) (-3.4826\%/\text{hour}) \sim 1.733 \text{ shares}/\text{hour}.$$

at 10am.

Thinking about Parabolas

Question 91 (a) Find an equation of the tangent line to the parabola $y = x^2$ at the point $(-3, 9)$.

(b) Find an equation of the tangent line to the parabola perpendicular to the tangent line in part (a).

(c) Find the coordinates of the point where the lines in parts (a) and (b) intersect.

(d) Let $\mathcal{L}$ be the line through the points of tangency of the lines in parts (a) and (b). Find the coordinates of the point where $\mathcal{L}$ intersects the y -axis.

(e) Repeat parts (a)-(d) above for the tangent line to the parabola $y = x^2$ at the general point (b, b^2) . What do you notice? Enter your work in the Desmos activity below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/qe7mgnu5sv>

Question 92 Let

$$y = f(x) = ax^2$$

where $a \in \mathbb{R}$ is a constant, and the variables x, y are measured in meters.

- (a) What are the units of the constant a ? How do you know?
- (b) Answer part (e) of the previous question for this function. Modify the desmos activity in the previous question to check your work.

Question 93 In the absense of air resistance, a rock released from rest near the surface of the earth falls $s = 16t^2$ feet during the first t seconds of its fall.

Compare the speed of the rock when it hits the ground with the average speed of the rock during the entire time interval of its fall.

Question 94 (a) The demonstration below shows two normal lines to a parabola and their point of intersection P . What do you think happens to P as point B approaches A ? Answer this question without dragging the slider b .

Free Response:

- (b) Now drag the slider b near $a = 2$ and observe what happens to point P . Were you correct? Were you surprised?

Free Response:

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/ybaivhc2t1>

Access this activity online at 151: Normals to Parabola

- (c) The parabola has equation $y = x^2/4$, point A has coordinates $(2, 1)$, and point B has coordinates $(b, b^2/4)$.
 - (i) Find an equation of the normal lines to the parabola at A and B .

- (ii) Use algebra to find an equation of the point P where the normal lines intersect.
- (iii) The point P approaches some point Q as B approaches A . Use the algebra of limits to find the coordinates of Q .
- (iv) Repeat parts (i)-(iii), replacing the point $A(2, 1)$ with the point $A(a, a^2)$.

Applications

Question 95 Play the slider u in the animation below to watch the motion of a balloon as it leaves behind track's at equal time intervals. Ignore the balloon's rightward jogs. Their purpose is to prevent the tracks from overlapping.

Desmos link: <https://www.desmos.com/calculator/h91txxjcmi>

151: Balloon Motion 7

- (a) Use the animation to sketch by hand the graph of the function

$$h = f(t), \quad -1 \leq t \leq 1.8,$$

that expresses the balloon's height (in thousands of feet) in terms of the number of hours past noon.

- (b) Use the animation or your graph from part (a) to sketch by hand the graph of the function

$$r = g(t), \quad -1 \leq t \leq 1.8,$$

that expresses the balloon's rate of ascent (in thousands of feet/hour in terms of the number of hours past noon.

- (c) Use your graph(s) or the animation to approximate when the balloon descends at the fastest rate.

Question 96 The function

$$h = f(t) = 5 - 2t - \frac{t^2}{4} + t^3, \quad -1 \leq t \leq 1.8,$$

expresses the height (in thousands of feet) of a balloon in terms of the number of hours past noon.

- (a) Find the balloon's average rate of ascent between 11:00am and 11:30am.
- (b) Is the balloon rising or falling at 1:00pm? At what rate? Use the graph of the function $h = f(t)$ below to approximate the rate. Then compute the exact rate.
- (c) When is the balloon descending at the rate of 1000 ft/hour? Use the sliders m and b below to approximate the time(s). Then compute the exact time(s).
- (d) Use the graph below to approximate when the balloon is descending at the fastest rate. Approximate this rate from the graph. Then compute the exact time and rate.
- (e) Use the graph below to approximate when the balloon is at its lowest point. Then compute the exact time.
- (f) Use the graph below to approximate when the balloon is at its highest point between 11am and 1:36pm. Then compute the exact time.
- (g) Use algebra to find all half-hour time intervals during which the balloon descends at an average rate of 500 ft/hour.

Desmos link: <https://www.desmos.com/calculator/bvtukd0v1c>

Access this activity online at 151: Height of Balloon

Question 97 The function

$$T = f(m) = \frac{m^3}{3} - 3m^2 + m + 35, \quad 0 \leq m \leq 8,$$

expresses the temperature (in Celsius degrees) of a beaker of water in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/jrzsv06pwt>

Desmos activity available at 151: Beaker Temperature

- (a) Use the sliders above to approximate the time(s) when the temperature is decreasing at the rate of $5^\circ\text{C}/\text{min}$.
- (b) Use calculus and algebra to find the exact times in part (a). Do not rely on the quadratic formula. Complete the square instead.
- (c) When is the temperature a minimum? Use the graph to approximate the time. Then use calculus and algebra to find the exact time.

Question 98 The function

$$P = f(t) = 8 + 4t - t^2, 0 \leq t \leq 5,$$

expresses the price (in dollars/share) of a stock in terms of the number of hours past 9am. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/jrwe6t0x41>

Desmos activity available at 151: *Stock Price Short*

- Is the price increasing or decreasing at 10am? At what relative rate?
- Use the slider above to approximate the time(s) when the price is decreasing at the relative rate of 40%/hour.
- Use calculus and algebra to find the exact time(s) in part (b). Do not rely on the quadratic formula. Complete the square instead.

Question 99 The function

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), 3 \leq s \leq 28,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/cphmgntm7>

Desmos activity available at 151: *Gas as a Function of Distance 2c*

- Use the graph of the function f shown above to determine if you are driving toward or away from home. Explain your reasoning.
- Zoom in on the graph to approximate your gas mileage at the moment you are 20 miles from home. Show a screenshot to help explain how you got your approximation. Then compute the exact gas mileage.
- Use the sliders m and b_1 in the graph to approximate your distance from home at the moment your car gets 30 miles/gallon. Show a screenshot to help explain how you got your approximation. Then compute the exact distances.
- Use the sliders m and b_1 in the graph to approximate an interval beginning or ending when you are 20 miles from home over which your average gas mileage is equal to you gas mileage at the moment you are 20 miles from home. Show a screenshot to help explain how you got your approximation. Then compute the exact interval.

(e) Sensors on your car measure both the (instantaneous) gas mileage and the number of gallons of gas in your tank at each instant. A computer then uses these measurements to estimate the number of additional miles you can drive before running out of gas. Use this idea to find a function

$$m = g(s), \quad 3 \leq s \leq 28,$$

that expresses the number of miles you can drive before running out of gas (assuming your gas mileage remains constant for the remainder of your trip) in terms of your distance from home. Explain your reasoning.

Question 100 (a) Make up your own quadratic function

$$v = f(G) = aG^2 + bG + c,$$

with a, b , and c all not equal to zero, that expresses the speed of a car (measured in miles/hour) in terms of its gas mileage (measured in miles/gallon). Be sure to include a domain. Explain why you think your function is reasonable.

(b) Compute the derivative dv/dG and evaluate it at a specific gas mileage. Include units.

(c) Evaluate the derivative dv/dG at a specific gas mileage and its meaning. Include units in your explanation.

(d) Use your function from part (a) to find a function

$$r = h(G)$$

that expresses the rate (in gal/hr) at which the car burns gas in terms of its gas mileage (in miles/gal). Explain your logic thoroughly.

(e) Evaluate $h(G)$ at the same gas mileage, say G_0 , you used in part (c). Compare the units of $h(G_0)$ and the derivative $dv/dG \Big|_{G=G_0}$. Are these two numbers related? What does this tell you about simplifying the units of a derivative?

(f) Use the ideas of this chapter (ie. derivatives of polynomials, and nothing beyond) to find an expression for the derivative dr/dG .

(g) What are the units of the derivative dr/dG ?

(h) Evaluate the derivative dr/dG at a specific gas mileage and explain its meaning. Include units in your explanation.

(i) Make up and answer your own question about the derivative dr/dG at a specific gas mileage.

A Few More Problems

Question 101 The function

$$h = f(t) = kt^3, t \geq 0,$$

expresses the height of a balloon (in thousands of feet) in terms of the number of hours past noon. Here $k > 0$ is a positive constant.

- (a) What are the units of the constant k ? Explain how you know.
- (b) Find the balloon's rate of ascent (measured in thousands of ft/hr) at time $t = 3$ hours past noon. Then find its average rate of ascent (measured in thousands of ft/hr) between noon and 3pm. How are these rates related? Note that both will be expressed in terms of k .
- (c) Find a function

$$r = g(a)$$

that expresses the balloon's rate of ascent (measured in thousands of ft/hr) at time $t = u$ hours past noon in terms of its average rate of ascent (measured in thousands of ft/hr) over the time interval $t \in [0, u]$. Assume $u > 0$.

- (d) Interpret your result from part (b) geometrically on the graph of the function f . Follow the directions on Lines 4, 6, and 8 in the demonstration below to help explain your interpretation.
- (e) Which of the following expresses the balloon's rate of ascent at time t hours past noon in terms of t and $h = f(t)$?

Multiple Choice:

- (a) $\frac{h}{t}$
- (b) $\frac{h}{3t}$
- (c) $\frac{3h}{t}$ ✓

Desmos link: <https://www.desmos.com/calculator/0byjcy77yw>

Desmos activity available at 151: Subtangent 1

Question 102 The function

$$v = f(G)$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon) for speeds between 55 miles/hour and 70 miles/hour.

Suppose $f(30) = 60$.

(a) Which of the following is more likely to be true?

$$\left. \frac{dv}{dG} \right|_{G=30} = 4$$

or

$$\left. \frac{dv}{dG} \right|_{G=30} = -4?$$

Explain your reasoning.

(b) What are the units of the above derivative? Do not simplify the units and do not write “per” in place of “/”.

(c) Explain the meaning of the derivative in part (a). It is not enough to say “the rate of change of something with respect to something else.” Remember this class is all about small changes and your explanation should be about an approximate relationship between small changes in this setting.

(d) Simplify the units of the derivative. What does this suggest about its meaning?

(e) At what rate does the car burn gas (in gal/hour) at a speed of 60 miles/hour?

(f) What does this problem suggest about simplifying the units of a derivative?

Question 103 The function

$$v = f(G), 20 \leq G \leq 40,$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon). Use the graph of the function f to find approximate answers to the following questions. Change the position and slope of line AB by dragging either the line or the points A or B . Change the position of the tangent line by dragging the slider G .

(a) Label the axes with the appropriate variable names and units.

(b) At what speed does the car burn gas as the fastest rate?

Hint: One of questions (b), (c) is related to the tangent lines to the curve, the other is related to the lines through the origin and the points of the curve.

(c) At what speed does increasing the speed by 0.1 miles/hour result in the greatest change in the gas mileage? Approximate that change.

Geogebra link: <https://www.geogebra.org/m/vjdf6x6z>

Geogebra activity available at 151: Gas Mileage

3 Homework Solution

Desmos link: <https://www.desmos.com/calculator/jku58rp2ve>

Derivatives of Polynomials CW

Working with polynomials and their derivatives.

1 The Power Rule

Question 104 Let

$$y = f(x) = kx^6,$$

where k is a constant and the variables x, y are measured in meters.

- (a) What are the units of k ? How do you know?
- (b) Find an expression for the average rate of change of the function $f(x)$ between $x = a$ and $x = b$, $a \neq b$.
- (c) Use part (b) and the algebra of limits to find an expression for the derivative

$$\left. \frac{dy}{dx} \right|_{x=a} = \frac{d}{dx} (kx^6) \Big|_{x=a}.$$

- (d) Check that your expression for the derivative has the correct units as follows:

- (i) Use the units x and y to find the units of the derivative

$$\left. \frac{dy}{dx} \right|_{x=a} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{b \rightarrow a} \frac{f(b) - f(a)}{b - a}$$

without using your expression for the derivative from part (c).

- (ii) Then use your expression for the derivative from part (c) and the units of k to find the units of the derivative

$$\left. \frac{dy}{dx} \right|_{x=a} = 6ka^5.$$

- (iii) Check that the units in parts (i) and (ii) are identical.
- (e) Due to a printing error, the graph of the function $y = f(x) = 234x^6$ is missing below. All we see is a point A on the graph.

Desmos link: <https://www.desmos.com/calculator/qcolims10c>

Learning outcomes:
Author(s):

151: Printers Error 3

Without sketching a graph, drag the slider m so that the blue line is tangent to the curve $y = 234x^6$ at A . Explain your reasoning.

- (f) Make a conjecture about the derivative

$$\frac{d}{dx}(kx^n)$$

for constants n and k .

Question 105

The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 30000,$$

expresses the distance to the horizon (in miles) in terms of your altitude (in feet).

- (a) Evaluate $f(25)$ and interpret its meaning.
 (b) Evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=25}.$$

Include units.

- (c) Use the language of small changes to interpret the above derivative as a scaling factor.

2 Stock Price

Question 106 The function

$$P = f(t) = 0.1t^3 - t^2 + t + 20, \quad 0 \leq t \leq 8,$$

expresses the price (in dollars/share) of a stock in terms of the number of hours past 9am. The graph of the function is shown below.

Desmos link: <https://www.desmos.com/calculator/gpqicgxvk3>

151: Stock Price

- (a) What are the units of the coefficient 0.1 in the leading term $0.1t^3$? Explain how you know.
- (b) Is the price increasing or decreasing at 10am? Use the graph to estimate this rate as follows:
 - (i) First drag point P to the appropriate position.
 - (ii) Then drag the slider m (the slope of the orange line) so that the orange line looks parallel to the (red) tangent line to the curve $P = f(t)$.
- (c) Use the function to compute the exact rate at which the price is changing at 10am. Include units.
- (d) Drag sliders m and b (Lines 2 and 3) to approximate the time(s) when the price is decreasing at the rate of $(\$1.8/\text{share})/\text{hr}$.
- (e) Use the function to find the exact time(s) when the price is decreasing at the rate of $(\$1.8/\text{share})/\text{hr}$.
- (f) Find all three-hour time intervals during which the price decreases at an average rate of $(\$2/\text{share})/\text{hr}$. Start by defining an unknown in a complete sentence, with units.
- (g) Use the graph to approximate the minimum and maximum prices of the stock during the eight-hour period. Then use calculus and algebra to compute these exact prices. Hint: What is the value of the derivative dP/dt when the price is a maximum/minimum?

Question 107 The function

$$G = f(s), \quad 4 \leq s \leq 24,$$

expresses the number of gallons of gas in your car in terms of the trip odometer reading (in miles).

- (a) Find an expression for your average gas mileage (measured in miles/gal) between odometer readings $s = a$ and $s = b$. Assume $4 \leq a, b \leq 24$ and $a \neq b$.
 Note: Gas mileage is positive and has the units stated, miles/gallon.
- (b) Use part (a) to write an expression with a limit for your (instantaneous) gas mileage at the odometer reading $s = a$ miles, $4 \leq a \leq 24$.
- (c) Use part (b) to write an expression with a derivative for your (instantaneous) gas mileage at the odometer reading $s = a$ miles, $4 \leq a \leq 24$.

Question 108

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), \quad 4 \leq s \leq 24,$$

expresses the number of gallons of gas in your car in terms of the trip odometer reading (in miles).

Desmos link: <https://www.desmos.com/calculator/ny3z9ixyr1>

Desmos activity available at 151: *Gas as a Function of Distance 2c*

- (a) Zoom in on the graph to approximate your (instantaneous) gas mileage when the odometer reading is 20 miles. Show a screenshot to help with your explanation.
- (b) Compute the exact (instantaneous) gas mileage when the odometer reading is 20 miles.
- (c) Use the sliders m and b_1 in the graph to approximate the odometer reading(s) when your car gets 30 miles/gallon. Show a screenshot to help with your explanation.
- (d) Compute the exact odometer reading(s) when your car gets 30 miles/gal.
- (e) Use the sliders m and b_1 in the graph to approximate an interval beginning or ending at odometer reading $s = 20$ miles over which your average gas mileage is equal to your gas mileage at the moment the odometer reads 20 miles. Show a screenshot to help explain how you got your approximation. Then compute the exact interval.
- (f) Sensors on your car measure both the (instantaneous) gas mileage and the number of gallons of gas in your tank at each instant. A computer then uses these measurements to estimate the number of additional miles you can drive before running out of gas.
 - (i) Use the worksheet below to approximate the computer reading for the number of miles you can drive before running out of gas when the odometer reads 20 miles.

Desmos link:

<https://www.desmos.com/calculator/ulfojucqoc>

Desmos activity available at 151: *Gas as a Function of Distance 9*

(ii) Find a function

$$m = w(s), \quad 4 \leq s \leq 24,$$

that expresses the number of miles you can drive before running out of gas (assuming your gas mileage remains constant for the remainder of your trip) in terms of the odometer reading (in miles). Explain your reasoning.

Derivatives of Polynomials Worksheet

Working with polynomials and their derivatives.

Question 109 Find an equation of the tangent line to the curve

$$y = \frac{12}{x^2} - 3x + 4$$

at the point $(2, 1)$. Show all steps.

Question 110 The function $h = f(t)$, $-1 \leq t \leq 1.8$, expresses the altitude (in thousands of feet) of a balloon in terms of the number of hours past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/8pgzf0orms>

151: Height of Balloon 54

- (a) Use the sliders in Lines 2 and 3 above to approximate the balloon's rate of ascent at 12:30pm.
- (b) Use the sliders in Lines 2 and 3 above to approximate the time(s) when the balloon is ascending at the rate of 4,000 ft/hr.
- (c) Use the sliders in Lines 2 and 3 above to approximate the time(s) when the balloon's rate of ascent is equal to its average rate of ascent between 11:00am and 1:00pm.
- (d) Suppose

$$h = f(t) = 5 - 2t - \frac{t^2}{4} + t^3, \quad -1 \leq t \leq 1.8,$$

and find the exact answers to parts (a)-(c).

Question 111 The function

$$h = g(t) = h_0 - \frac{1}{2}gt^2$$

Learning outcomes:
Author(s):

expresses the height of a rock (in meters) dropped from rest from a height of h_0 meters above the lunar surface in terms of the number of seconds since it was released. Here g is a constant.

- (a) Find an expression for the rock's speed (in m/sec) in terms of the number of seconds since it was dropped.
- (b) Express the speed of the rock when it hits the surface in terms of its average speed during the time it takes to reach the surface.

Question 112 The function $G = f(s)$, $4 \leq s \leq 28$, expresses the number of gallons of gas in your car in terms of the trip odometer reading (measured in miles). Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/nnoxloqoro>

151: Gas as a Function of Distance 99 2c

- (a) Drag point P above to approximate your (instantaneous) gas mileage when the odometer reading is 10 miles. Gas mileage is measured in miles/gal and is positive.
- (b) A dashboard display shows the number of miles you have left to drive assuming your gas mileage remains equal to the current gas mileage for the remainder of your trip. Drag point P to approximate this reading when the odometer reads 15 miles. **No computations.**
- (c) Now suppose

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), \quad 4 \leq s \leq 28.$$

- (i) Find the exact answers to parts (a) and (b).
- (ii) Find a function

$$r = w(s), \quad 4 \leq s \leq 28.$$

that expresses the dashboard display for number of miles you have left to drive in terms of the odometer reading.

Normals to a Parabola

Normals to a parabola.

Question 113 Access Desmos interactives through the online version of this text at

The parabola below has equation

$$y = \frac{x^2}{2p},$$

for some constant $p \neq 0$. Point A has coordinates $(a, a^2/(2p))$, and point B has coordinates $(b, b^2/(2p))$. Assume $a \neq b$.

Desmos link: <https://www.desmos.com/calculator/ifsu6n6qt0>

Access this activity online at 151: Normals to Parabola 2

- Find an equation of the normal lines to the parabola at A and B .
- Use algebra to find an equation of the point P where the normal lines intersect.
- The point P approaches some point Q as B approaches A . Use the algebra of limits to find the coordinates of Q .

Explanation. An equation of the normal line to the parabola at $A(a, a^2/(2p))$ is

$$y = \frac{a^2}{2p} - \frac{p}{a}(x - a)$$

and an equation of the normal at $B(b, b^2/(2p))$ is

$$y = \frac{b^2}{2p} - \frac{p}{b}(x - b).$$

To find the coordinates of the point where the normals intersect, we need to solve the system

$$\begin{cases} y = \frac{a^2}{2p} - \frac{p}{a}(x - a) \\ y = \frac{b^2}{2p} - \frac{p}{b}(x - b) \end{cases}$$

Learning outcomes:
Author(s):

Substituting gives

$$\frac{a^2}{2p} - \frac{p}{a}(x - a) = \frac{b^2}{2p} - \frac{p}{b}(x - b).$$

After some algebra

$$\frac{1}{2p}(a - b)(a + b) = \frac{p}{ab}(b - a)x$$

and

$$x = -\frac{ab}{2p^2}(a + b).$$

So as $B \rightarrow A$, the x -coordinate of the intersection point approaches the x -coordinate

$$x = -\lim_{b \rightarrow a} \frac{ab}{2p^2}(a + b) = -\frac{a^3}{p^2}.$$

of Q .

Since Q lies on the normal through A , the y -coordinate of Q is

$$\begin{aligned} y &= \frac{a^2}{2p} - \frac{p}{a} \left(-\frac{a^3}{p^2} - a \right) \\ &= \frac{a^2}{2p} + \frac{a^2}{p} + p \\ &= \frac{3a^2}{2p} + p. \end{aligned}$$

So Q has coordinates

$$\left(-\frac{a^3}{p^2}, \frac{3a^2}{2p} + p \right).$$

Question 114 The lines normal to the parabola

$$y = \frac{x^2}{2p}$$

are tangent to another curve as illustrated below.

- Find a parameterization of that curve.
- Find a Cartesian equation of the curve.

Desmos link: <https://www.desmos.com/calculator/xjly7skb9p>

Access this activity online at 151: Normals to Parabola 3

Gas Consumption

Gas consumption and derivatives.

Question 115 (a) Use calculus to find an expression for the x -coordinate of the vertex of the parabola

$$y = ax^2 + bx + c, a \neq 0.$$

(b) Find an equation of the parabola's axis of symmetry.

Question 116 The function

$$\begin{aligned} v &= f(G) \\ &= \frac{1}{20}G^2 - 2G + 40, \quad 20 \leq G \leq 40, \end{aligned}$$

graphed below expresses the speed (in miles/hr) of a car in terms of its gas mileage (in miles/gal).

Desmos link: <https://www.desmos.com/calculator/lkxc2lrvz>

151: Gas Mileage 34

- Use the graph to determine the rate (in gal/hr) at which the car burns gas at a speed of 30 miles/hour.
- Drag the slider u in Line 2 to approximate the speeds between 20 miles/hour and 40 miles/hour at which the car burns gas at the maximum and minimum rates (measured in gal/hr). Explain your reasoning.
- Use the slider u to sketch by hand a rough graph of the function $r = h(G)$ that expresses the rate (in gal/hr) at which the car burns gas in terms of its gas mileage (in miles/gal).
- Use calculus to determine the exact speeds in part (b).
- Explain how you could justify your answers to part (d) without relying on the graph of the function $v = f(G)$.

Learning outcomes:
Author(s):

Question 117 The function

$$\begin{aligned} G &= f(v) \\ &= -\frac{v^2}{20} + 5v - 90, \quad 35 \leq v \leq 70, \end{aligned}$$

graphed below expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

Desmos link: <https://www.desmos.com/calculator/fapdhcqml>

151: Burning Gas

- Use the graph to determine the rate (in gal/hr) at which the car burns gas at a speed of 50 miles/hour.
- Drag the slider v in Line 1 to approximate the speeds between 35 miles/hour and 70 miles/hour at which the car burns gas at the maximum and minimum rates (measured in gal/hr). Explain your reasoning.
- Use calculus to determine the exact speeds in part (b). Find a way that avoids using the quotient rule.

Question 118 The function

$$G = f(s) = \frac{11}{5} + \frac{1}{5000} (s^3 - 50s^2 + 300s), \quad 3 \leq s \leq 28,$$

expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route.

Desmos link: <https://www.desmos.com/calculator/pb8v4t3cxg>

Desmos activity available at 151: Gas as a Function of Distance 33

- Use the graph above to determine if you are driving toward or away from home. Explain your reasoning.
- Sensors on your car measure both the (instantaneous) gas mileage and the number of gallons of gas in your tank at each instant. A computer then uses these measurements to estimate the number of additional miles you can drive before running out of gas.

- (i) Use the graph and the slider s_0 above to approximate the reading for the number of additional miles you can drive when you are 10 miles from home and 20 miles from home. Explain your reasoning.

- (ii) Find a function

$$m = g(s), 3 \leq s \leq 28,$$

that expresses the number of miles you can drive before running out of gas (assuming your gas mileage remains constant for the remainder of your trip) in terms of your distance from home. Explain your reasoning.

- (iii) Use your function g to find the reading for the number of additional miles you can drive when you are 10 miles from home. When you are 20 miles from home. Compare these with your estimates above.

The Quotient and Product Rules

Quotient and product rules.

Here are examples of how to use the Leibniz notation correctly in computing the derivative of a product and quotient of functions.

Example 30. Find expressions for the derivatives

(a)

$$\frac{d}{dx} ((5x^3 - 2)(4 - 6x))$$

(b)

$$\frac{d}{dt} \left(\frac{5t^3 - 2}{4 - 6t} \right)$$

Explanation. (a)

$$\begin{aligned} \frac{d}{dx} ((5x^3 - 2)(4 - 6x)) &= (4 - 6x) \frac{d}{dx} (5x^3 - 2) + (5x^3 - 2) \frac{d}{dx} (4 - 6x) \\ &= (4 - 6x)15x^2 + (5x^3 - 2)(-6). \end{aligned}$$

(b)

$$\begin{aligned} \frac{d}{dt} \left(\frac{5t^3 - 2}{4 - 6t} \right) &= \frac{1}{(4 - 6t)^2} \left((4 - 6t) \frac{d}{dt} (5t^3 - 2) - (5t^3 - 2) \frac{d}{dt} (4 - 6t) \right) \\ &= \frac{(4 - 6t)15t^2 - (5t^3 - 2)(-6)}{(4 - 6t)^2}. \end{aligned}$$

1 Relative Rates of Change

Suppose $P = f(t)$ and $Q = g(t)$ are differentiable functions of t and $g(t) \neq 0$. Then the quotient and product rules are better written in the forms

$$\frac{d}{dt} \left(\frac{P}{Q} \right) = \frac{P}{Q} \left(\frac{1}{P} \cdot \frac{dP}{dt} - \frac{1}{Q} \cdot \frac{dQ}{dt} \right) \quad (1)$$

and

$$\frac{d}{dt} (PQ) = PQ \left(\frac{1}{P} \cdot \frac{dP}{dt} + \frac{1}{Q} \cdot \frac{dQ}{dt} \right).$$

Learning outcomes:
Author(s):

Question 119 (a) Verify the statements above.

(b) What do they say about the relative rate of change in the quotient of two functions? In their product?

2 Exercises

Exercise 120 Between speeds of 45 miles/hr and 75 miles/hr, the function

$$G = f(v), 45 \leq v \leq 75,$$

expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

Suppose $f(50) = 25$ and

$$\left. \frac{dG}{dv} \right|_{v=50} = 0.8.$$

- (a) What are the units of the derivative above? What is the meaning? Explain in terms of small changes.
- (b) What are the simplified units of the derivative above? What insight do these units give you about the derivative's meaning?
- (c) Approximate the gas mileage at a speed of 48 miles/hour.
- (d) Let

$$r = g(v), 45 \leq v \leq 75,$$

be the function that expresses the rate (in gal/hr) at which the car burns gas in terms of its speed (in miles/hr).

- (i) Use Equation (1) above to evaluate the derivative

$$\left. \frac{dr}{dv} \right|_{v=50}.$$

Include units for all numbers.

- (ii) What are the units of the derivative above? What is the meaning? Explain in terms of small changes.
 - (iii) What are the simplified units of the derivative above? What insight do these units give you about the derivative's meaning?
 - (iv) Approximate the rate (in gal/hr) at which the car burns gas at a speed of 48 miles/hr.
-

Exercise 121 At 9:00am on February 23, 2023, the price of oil was decreasing at the relative rate of 2%/hour. At what relative rate was the number of gallons of oil you could buy with \$100,000 changing at that time? Use calculus to justify your assertion.

Exercise 122 The function

$$P = f(t) = 5 - 3t + t^2, 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) Use the graphs of the function $P = f(t)$ and the function $r = f'(t)/f(t)$ to estimate when the stock price is increasing at the greatest relative rate.
- (b) Use algebra to find the exact time when the stock price is increasing at the greatest relative rate.

Hint: What is the value of the derivative dr/dt at this time? But start by finding an expression for the instantaneous relative rate of change in the stock price.

Desmos link: <https://www.desmos.com/calculator/xuupp3srqv>

Desmos activity available at 151: Stock Price 4

Exercise 123 An h -foot tall giraffe walks directly toward a spotlight on the ground as the light casts its shadow on a vertical wall as illustrated below. The wall is b feet from the light.

Desmos link: <https://www.desmos.com/calculator/2eiyjjpu9n>

Desmos activity available at 151: Spotlight

Suppose at a certain instant the giraffe is w feet from the spotlight and is walking at a speed of v ft/sec.

- (a) Is the length of the giraffe's shadow increasing or decreasing at this instant?
- (b) At what rate?
- (c) Check that your answer in part (b) has the correct units.

- (d) Assume v is constant and describe what happens to the rate in part (b) as the giraffe approaches the spotlight. How is your conclusion revealed in the animation?
-

Exercise 124 (a) On January 1, 2024 the national debt of a country was decreasing at the rate of 3%/yr and the population was increasing at the rate of 2%/yr. Was the per-capita (ie. per person) share of the national debt increasing or decreasing at this time? At what relative rate?

- (b) During the year 2024 the national debt of a country decreased 3% and the population increased 2%. Did the per-capita share of the national debt increase or decrease during the year? By what percent?

- (c) Compare the two questions above and their answers.
-

Test 1

Test 1.

- (a) (6 points) The function

$$W = f(r) = \frac{2000}{r^2}, r \geq 4,$$

expresses the weight of an astronaut (measured in pounds) in terms of her distance from the center of the earth (measured in thousands of miles).

- (i) (2 points) Find an expression for the average rate of change in the astronaut's weight with respect to her distance from the earth's center between distances $r = b$ and $r = c$ thousands of miles from the center. Assume $b, c \geq 4$ and that $b \neq c$.
- (ii) (4 points) Use your *expression from part (a) directly and the definition of the derivative* to find an expression (fully simplified) for the derivative

$$\left. \frac{dW}{dr} \right|_{r=b}.$$

In particular, do *not* use the power rule. Show all work in its mathematically correct form. Write vertically, one equal sign per line. No need to explain algebra.

Explanation. (i) The average rate of change is

$$\begin{aligned} \frac{\Delta W}{\Delta h} &= \frac{f(c) - f(b)}{c - b} \\ &= \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \end{aligned}$$

- (ii) (Some algebra omitted, left for you to fill in).

$$\begin{aligned} \left. \frac{dW}{dr} \right|_{r=b} &= \lim_{c \rightarrow b} \left(\frac{1}{c - b} \right) \left(\frac{2000}{c^2} - \frac{2000}{b^2} \right) \\ &= \lim_{c \rightarrow b} \frac{-2000(c + b)}{c^2 b^2} \\ &= -\frac{2000(b + b)}{b^2 b^2} \\ &= -\frac{4000}{b^3}. \end{aligned}$$

Learning outcomes:
Author(s):

- (b) (5 points) At 3pm, you can buy 5 pounds of cod with \$100. And at 3pm the number of pounds of cod you can buy with \$100 is decreasing at the rate of 0.4 lbs/hour.

At what rate (with respect to time) is the price of cod changing at 3pm? Explain your reasoning and be sure to end with a concluding sentence.

Explanation. Let $P = f(t)$ be the function that expresses the price of cod (in dollars/pound) in terms of the number of hours since 3pm.

Let $Q = g(t)$ be the function that expresses the number of pounds of cod you can buy with \$100 in terms of the number of hours since 3pm.

Then

$$P = \frac{100}{Q}$$

and the quotient rule gives

$$\begin{aligned} \frac{dP}{dt} &= \frac{1}{Q^2} \left(Q \frac{d}{dt} (100) - 100 \frac{dQ}{dt} \right) \\ &= -\frac{100}{Q^2} \cdot \frac{dQ}{dt} \end{aligned}$$

Since at 3pm the number of pounds we can buy with \$100 is decreasing at the rate of 0.4 lbs/hour, we know that

$$\left. \frac{dQ}{dt} \right|_{t=0} = -0.4 \text{ lb/hr.}$$

And since we can buy $Q = g(0) = 5$ pounds with \$100 at 3pm,

$$\begin{aligned} \left. \frac{dP}{dt} \right|_{t=0} &= - \left(\frac{100}{Q^2} \cdot \frac{dQ}{dt} \right) \Big|_{t=0} \\ &= \left(-\frac{100 \text{ dollars}}{25 \text{ lb}^2} \right) \left(-0.4 \frac{\text{lb}}{\text{hr}} \right) \\ &= (1.6 \text{ dollars/lb})/\text{hr.} \end{aligned}$$

So at 3pm the price of cod is increasing at the rate of (1.6 dollars/lb)/hr.

- (c) (5 points) Between speeds of 60 miles/hour and 80 miles/hour, the function

$$v = f(G), \quad 8 \leq G \leq 20,$$

expresses the speed of a car (in miles/hour) in terms of its gas mileage (in miles/gallon).

Suppose $f(12) = 72$.

- (i) (1 point) Which would be more likely, that

$$\left. \frac{dv}{dG} \right|_{G=12} = 4 \quad \text{or that} \quad \left. \frac{dv}{dG} \right|_{G=12} = -4?$$

Explain your reasoning.

- (ii) (1 point) What are the units of the correct derivative above?
 (iii) (2 points) Explain the meaning of the correct derivative above using the language of *small changes*.
 (iv) (1 point) Simplify the units of the correct derivative above. What do these simplified units tell you about the derivative's meaning?

Explanation. (i) At the high speed of 72 miles/hour it is likely that a small increase in the speed will decrease the gas mileage. So we would expect the derivative to be negative, so

$$\left. \frac{dv}{dG} \right|_{G=12} = -4.$$

- (ii) Since v has units miles/hour and G has units miles/gal, the derivative dv/dG has units

$$\frac{\text{miles/hr}}{\text{miles/gal}}.$$

- (iii) We know that

$$\left. \frac{dv}{dG} \right|_{G=12} = -\frac{0.4 \text{ miles/hr}}{0.1 \text{ miles/gal}} \sim \frac{\Delta v}{\Delta G}$$

if $\Delta G \sim 0$. So increasing the speed by 0.4 mile/hour (from 72 miles/hr to 72.4 miles/hr) would decrease the gas mileage by about 0.1 miles/gal (from 12 miles/gal to about 11.9 miles/gal).

- (iv) The units of the simplified derivative are

$$\frac{\text{miles/hr}}{\text{miles/gal}} = \frac{\text{gal}}{\text{hr}}.$$

Although this suggests the derivative is related to the rate at which the car burns gas, this is *not* true. Our interpretation of the derivative had nothing to do with this rate and in fact the car burns gas at the rate of 6 gal/hour at a speed of 72 miles/hour. So the simplified units are misleading and tell us *nothing* about the derivative's meaning.

- (d) (5 points) The function

$$G = f(s) = \frac{3s + 25}{s + 5}, \quad 0 \leq s \leq 10,$$

expresses the number of gallons of gas in a car in terms of the trip odometer reading (measured in miles).

Sensors on the car measure both the number of gallons of gas in the tank and the current (instantaneous) gas mileage (measured in miles/gallon).

A computer then calculates the number of miles you have left to drive assuming the car's gas mileage remains constant for the remainder of your trip (and equal to the current gas mileage). This number is displayed on the dashboard.

Determine the exact trip odometer reading when the dashboard reading indicates that you have 13 miles left to drive. Explain your reasoning and be sure to end with a concluding sentence.

Explanation. I will not solve this problem, but will give you a hint instead by solving a closely related problem.

Suppose, for example, that we wish to find the dashboard reading (for the number of miles left to drive) at the odometer reading $s = 5$ miles.

First, we would find the number of gallons in the tank at this odometer reading. At this time, the tank has

$$f(5) = \frac{3(5) + 25}{5 + 5} = 4$$

gallons of gas.

Now at $s = 5$, the derivative dG/ds is (left for you to verify)

$$\left. \frac{dG}{ds} \right|_{s=5} = -0.10 \frac{\text{gal}}{\text{mile}}.$$

This means that five miles into our trip the car is burning gas at the rate of 0.10 gallons/mile. So if this rate were to remain constant for the remainder of the trip, we would be able to drive an additional

$$\frac{4 \text{ gallons}}{0.10 \frac{\text{gal}}{\text{mile}}} = 40 \text{ miles}$$

before running out of gas.

Derivatives of Exponential Functions

Working with exponential functions and their derivatives.

Relative Changes and Relative Rates of Change

Relative changes and relative errors are often more meaningful than absolute changes and errors. For example, I might measure the distance from Shoreline's Central Market to the Richmond beach library to be 5 km with an error of at most 0.2 km, while NASA might measure the between the earth and the moon on the first day of spring to be 384,400 km with an error of at most 100 km. The relative error in my measurement is at most

$$\frac{0.2 \text{ km}}{5 \text{ km}} = 0.04 = 4\%,$$

while the relative error in NASA's measurement is at most

$$\frac{100 \text{ km}}{384,400 \text{ km}} \sim 0.00026 = 0.026\%.$$

Relatively speaking, NASA's measurement was about 150 times more accurate than mine.

Question 125 At 10:00am the prices of Stock A and Stock B are both increasing at the rate of (\$2/share)/hour. At 10:00am Stock A sells for \$50/share and Stock B for \$10/share. Compare the relative rates at which the share prices are changing at 10:00am.

Question 126 The function

$$P = f(t) = 5 - 3t + t^2, \quad 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) At what relative rate is the price of the stock changing at 10am?
- (b) When is the share price increasing at a rate of 60%/hr?
- (c) During what time interval is the price of the stock increasing?

Learning outcomes:
Author(s):

(d) During what time interval is the relative rate of change in the price of the stock increasing?

Desmos link: <https://www.desmos.com/calculator/hhkveu6lxp>

Desmos activity available at 151: Stock Price

Question 127 The function

$$P = f(t), 1 \leq t \leq 3.6,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am. Use the graph below to approximate the answers to the following questions without putting a scale on the P axis.

- (a) At what relative rate is the price changing at 11am? At 12:30pm?
- (b) When is the stock price increasing at its maximum relative rate? At its minimum relative rate? Approximate these rates.

Desmos link: <https://www.desmos.com/calculator/jyebaj5jif>

Desmos activity available at 151: Stock Price 2

Question 128 The function

$$W = f(t) = 200 + 4t + 2t^2, 0 \leq t \leq 12,$$

expresses the weight (in pounds) of a baby elephant in terms of its age (in months).

- (a) Find the average rate at which the elephant gained weight between ages 4 and 10 months.
- (b) Find the relative average rate at which the elephant gained weight between ages 4 and 10 months.
- (c) Find the relative instantaneous rate at which the elephant is gaining weight at age 4 month.
- (d) Find the relative instantaneous rate at which the elephant is gaining weight at age 10 months.
- (e) Use the graph below to interpret your answers to parts (b)-(d) geometrically.

Desmos link: <https://www.desmos.com/calculator/2xj6xy7ggo>

Desmos activity available at 151: Elephant

Exponential Growth

Question 129 (a) What does it mean for a population to grow exponentially?
 (b) Is it possible for a population to increase by 20% every year and not grow exponentially?

Question 130 Suppose between noon and 10pm a colony of bacteria grows exponentially. The population is 200,000 at noon and 242,200 at 1pm.

- (a) Describe how the population grows. Keeping Question 2(b) in mind, is your description sufficient?
- (b) How might we find a complete description of the exponential growth?
- (c) Determine the relative average growth rate between noon and 12:30pm. Between 1pm and 1:30pm. Over any half-hour time period. Use the slider u in the graph below to interpret these rates geometrically.
- (d) Approximate the instantaneous relative growth rates in the population at noon, at 1pm, and at 2pm. Modify the definition of $v = u + 1/2$ in the demonstration below and interpret these rates geometrically.
- (e) Use limits to write an expression that gives the instantaneous growth rate at time t hours past noon. What can you conclude?
- (f) Try to answer part (b) again.

Desmos link: <https://www.desmos.com/calculator/wvpsotdhby>

Desmos activity available at 151: [Exp Growth 1](#)

Question 131 The function

$$P = f(t) = 3(2)^t, -2 \leq t \leq 4,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

- (a) Describe how the population grows. Is your description sufficient?

The population *doubles* every *hour*.

- (b) Find an expression for the relative growth rate between time $t = u$ hours past noon and time $t = u + h$ hours past noon. Measure the rate relative to the population at time $t = u$. Is this question asking about an average or an instantaneous relative growth rate?

The relative growth rate is

$$\begin{aligned}\frac{1}{P} \left(\frac{\Delta P}{\Delta t} \right) &= \frac{1}{f(u)} \left(\frac{f(u+h) - f(u)}{h} \right) \\ &= \frac{1}{3(2)^u} \left(\frac{3(2)^{u+h} - 3(2)^u}{h} \right) \\ &= \frac{1}{3(2)^u} \left(\frac{3(2)^u(2^h - 1)}{h} \right) \\ &= \frac{2^h - 1}{h}\end{aligned}$$

- (c) What are the units of the relative average growth rate in part (b)?
- (d) Input your function from part (b) on Line 5 in the worksheet below.
- (e) What do you notice about the distance between points R and S as you drag the slider u below. How is this distance related to the relative average growth rate in part (b)?
- (f) Use limits to write an expression for the relative growth rate at time $t = u$ hours past noon. Simplify this expression as much as possible. What can you conclude about how the relative instantaneous growth rate varies with time?

The relative growth rate is

$$\lim_{h \rightarrow 0} \frac{2^h - 1}{h}.$$

- (g) Interpret your expression from part (f) as the derivative of a specific function evaluated at a specific input. What does this tell you about the relative growth rate of this particular population?

The relative instantaneous growth rate of the population is equal to the derivative of the function

$$f(x) = 2^x$$

evaluated at $x = 0$.

- (h) Use part (f) to numerically approximate the relative (instantaneous) growth rate of the population. Show a table that suggests a progression toward a limit.
- (i) Use a similar method to approximate the relative instantaneous growth rate of the population

$$P = f(t) = 5(3)^t.$$

Desmos link: <https://www.desmos.com/calculator/omjbec2hpu>

Desmos activity available at 151: Exponential Growth 1

Question 132 Parts (h) and (i) of the previous question suggest that there is a number e between 2 and 3 that makes the relative growth rate of the function

$$P = f(t) = P_0 e^t, -3 \leq t \leq 5$$

equal to 100%/hr, where we assume here that t is measured in hours.

- (a) What is the one-hour growth factor for this population?
- (b) Describe what happens to the population every hour.
- (c) At what relative rate is the population increasing at 1:00pm?
- (d) Suppose at 1:00pm the population is 500,000. Approximate the population at 1:03pm and compare your approximation to the actual population at that time.

Exponential Functions with Bases other than e

Question 133 The function

$$P = g(t) = P_0 e^{t/2}, -6 \leq t \leq 10$$

expresses the population (Colony B) of bacteria in terms of the number of hours past noon.

- (a) Describe a transformation that takes the graph of the population function

$$P = f(t) = P_0 e^t, -3 \leq t \leq 5$$

for Colony A (where t is also the number of hours past noon) to the graph of $P = g(t)$.

- (b) Suppose that the population of Colony A is 400,000 at 4:00pm.
 - (i) When is the population of Colony B equal to 400,000?
 - (ii) What are the growth rates of the two populations when they each have respective populations of 400,000 bacteria?
 - (iii) What are the relative growth rates of the two populations when they each have respective populations of 400,000 bacteria?

Question 134 Here's another way to think about differentiating the function

$$P = g(t) = P_0 e^{t/2}, -6 \leq t \leq 10$$

that expresses the population of a colony of bacteria in terms of the number of hours past noon.

We'll let $u = t/2$ be the number of two-hour periods since noon.

- (a) Express the population in terms of u .
- (b) Use what you know about the exponential function base e to express the growth rate of the population in terms of u .
- (c) Use part (b) to find the growth rate of the population at 6pm. Pay careful attention to units.
- (d) Use the idea of part (c) to express the growth rate

$$\frac{dP}{dt} = g'(t)$$

in terms of t .

- (e) Suppose instead that the population grows exponentially and doubles every hour. Find the relative instantaneous growth rate of the population.

Question 135 Between 11am and 8pm, a population of bacteria grows exponentially. The population is 4 million at noon and 5 million at 1pm.

- (a) What is the one-hour growth factor? [1.25](#)
- (b) Describe how the population grows.

The population increases by [25%](#) every hour.

- (c) Use your description from part (b) to find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon. Do not use e in your function. Define meaningful variables and include a domain.

The function

$$P = f(t) = 4(5/4)^t, -1 \leq t \leq 8$$

expresses the population (in millions of bacteria) in terms of the number of hours past noon.

- (d) Use the fact that $k = e^{\ln k}$ for $k > 0$ to express your function from part (c) using an exponential function with base e .

$$P = f(t) = 4e^{t \ln(1.25)}, -1 \leq t \leq 8$$

(e) Use u -substitution and the chain rule (show all steps) along with the fact that

$$\frac{d}{du}(e^u) = e^u$$

to find the relative instantaneous growth rate of the population. Include units in your conclusion.

The relative instantaneous growth rate of the population is

$$\frac{1}{P} \left(\frac{dP}{dt} \right) = \ln(5/4) \sim 22.314\%/hour.$$

(f) At what rate is the population growing when there are 10 million bacteria? Do this the easy way. Note that the question is not asking for a relative rate.

(g) Find the population when the population is increasing at the rate of 3 million bacteria/hour. Do this the easy way.

Question 136 Between 11am and 8pm, a population of bacteria decreases exponentially. The population is 5 million at noon and 4 million at 4pm.

(a) What is the four-hour growth factor? (This is the number you multiply the current population by to get the population four hours later). 0.8

(b) What is the one-hour growth factor? $(4/5)^{1/4}$

(c) Describe how the population decreases.

The population decreases by 20% every 4 hours.

(d) Use your description from part (c) to find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon. Do not use e in your function. Start by define meaningful variables. Include a domain with your function.

The function

$$P = f(t) = 5(4/5)^{t/4}, -1 \leq t \leq 8$$

expresses the population (in millions of bacteria) in terms of the number of hours past noon.

(e) Express your function from part (c) using an exponential function with base e . Then use u -substitution and the chain rule to find the relative instantaneous growth rate of the population. Show all steps in using the chain rule.

The population function is

$$P = f(t) = 5e^{0.25 \ln(0.8)t}, -1 \leq t \leq 8.$$

The relative instantaneous growth rate is

$$\frac{1}{P} \left(\frac{dP}{dt} \right) = 0.25 \ln(0.8) \sim -5.58\%/hour.$$

(f) At what rate is the population decreasing when there are 2 million bacteria? Do this the easy way.

(g) Find the population when the population is decreasing at the rate of 300,000 bacteria/hour. Do this the easy way.

(h) Use the result of part (f) to approximate the population 4 minutes after there are 2 million bacteria.

Relative Rates Again

Question 137 The function

$$P = f(t), 0 \leq t \leq 2,$$

expresses the balance (in dollars) in an account in terms of the number of years since the start of 2022. Suppose

$$\left. \frac{1}{P} \frac{dP}{dt} \right|_{P=5000} = 0.08.$$

- (a) What are the units of the above derivative? How do you know?
- (b) Interpret the meaning of the above derivative.
- (c) Approximate the balance in the account four days after the account has \$5,000. Explain your reasoning.

Question 138 The function

$$q = f(p) = 0.5(p - 18)^2, 6 \leq p \leq 15,$$

expresses the average number of burgers/day sold at Five Guys of Edmonds in terms of the price (in \$/burger).

- (a) At what relative rate does the quantity sold (q) change with respect to the price (p) at a price of \$10/burger?
- (b) What are the units of the above relative rate of change?
- (c) Explain the meaning of the relative rate of change in part (a).

(d) Use the graph of the function $q = f(p)$ and the slider u in the desmos activity below to interpret the relative rate of change in part (a) geometrically. Explain your reasoning.

(e) Use the result of part (a) to approximate the relative change in the average number of burgers sold per day if the Five Guys increases the price from \$10/burger to \$10.25/burger. Explain your reasoning.

(f) Use the result of part (a) to approximate the relative change $\Delta a\%$ in the average number of burgers sold per day in terms of a small relative change $\Delta r\%$ in the price from \$10/burger. Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/ylgk03oaza>

Desmos activity available at 151: Burgers 1

Question 139 The function

$$P = f(t), -2 \leq t \leq 5,$$

expresses the population of a colony (call it Colony A) of bacteria in terms of the number of hours past noon.

The function

$$P = g(t) = f(t/2), -4 \leq t \leq 10,$$

expresses the population of Colony B in terms of the number of hours past noon.

The populations do not necessarily grow exponentially.

(a) Compare the populations at noon.

(b) Suppose Colony A has 50,000 bacteria at 3:00pm. When does Colony B have 50,000 bacteria? Explain.

(c) Suppose the population of Colony A takes three hours to grow from 20,000 to 50,000. How long does it take the population of Colony B to grow from 20,000 to 50,000?

(d) Suppose the population of Colony A is increasing at the rate of 10,000 bac/hr at 3pm. What is the growth rate of Colony B at 6pm? Explain.

(e) What is the relative growth rate of Colony A at 3pm? What is the relative growth rate of Colony B at 6pm? Explain.

Question 140 The function

$$P = f(u) = 10e^u, -2 \leq u \leq 5$$

expresses the population (in millions of bacteria) of a colony of bacteria in terms of the number of hours since noon.

- (a) What are the units of the input to the exponential function in the above expression for P ?
- (b) At what rate is the population growing when there are 30 million bacteria?
- (c) At what rate is the population growing when there are P million bacteria? Do the units of your answer make sense?
- (d) What is the relative instantaneous growth rate of the population?
- (e) Find a function

$$P = g(t)$$

that expresses the population (in millions of bacteria) of the colony of bacteria in terms of the number of minutes since noon. Include a domain.

- (f) Use common sense to evaluate the derivative

$$\left. \frac{dP}{dt} \right|_{P=30}.$$

Explain your reasoning.

- (g) Use the idea of part (f) to find an expression for the derivative

$$\frac{dP}{dt} = g'(t)$$

at time t minutes past noon.

Question 141 The function

$$P = P_0 e^{kt}, -4 \leq t \leq 5,$$

expresses the population of a colony of bacteria in terms of the number of hours past noon.

- (a) What are the units of the constant P_0 ?
- (b) What are the units of the constant k ?
- (c) Use the ideas of the previous question to find an expression for the growth rate of the population at time t hours past noon. Include units in your answer.
- (d) Find an expression for the relative growth rate at time t hours past noon. Include units in your answer.

Question 142 One of the two functions graphed below is an exponential function. Which one? How do you know?

The Derivative as a Magnification Factor

Exploration 143 It sometimes helps to think of the derivative as a magnification factor that maps a small interval around an input to a function to a corresponding interval around the output.

(a) Use this idea for the function $y = f(x)$ graphed below to approximate the derivatives

$$\left. \frac{d}{dx}(f(x)) \right|_{x=4} \quad \text{and} \quad \left. \frac{dy}{dx} \right|_{x=5}.$$

Desmos link: <https://www.desmos.com/calculator/la4f5ots3r>

Desmos activity available at 151: Magnification Factor 1

Differentiating the Exponential Function e^x

Exploration 144 (a) Use the graph of the function $y = f(x)$ below to approximate the derivatives

$$\left. \frac{dy}{dx} \right|_{y=k} \quad \text{for } k = 1, 2, \dots, 6.$$

Note the above derivatives are evaluated at the outputs of the function f .

(b) What do you notice?

Desmos link: <https://www.desmos.com/calculator/k08dphtuca>

Desmos activity available at 151: Magnification Factor 2

Question 145 The function

$$P = 400e^t, 0 \leq t \leq 2$$

expresses the population (in thousands) of a colony of bacteria in terms of the number of hours past noon.

(a) What are the units of the factor 400?

(b) What are the units of the exponent in the factor e^t ? Be careful.

(c) Find the (instantaneous) growth rate of the population when there are 1,200,000 bacteria.

- (d) Find the relative (instantaneous) growth rate when there are 1,200,000 bacteria.
- (e) Find the relative (instantaneous) growth rate at any time.
- (f) Approximate the population 30 seconds after there are 1,200,000 bacteria.

Question 146 Let $k > 0$ be a constant and let

$$f(x) = ke^x.$$

For $a \in \mathbb{R}$ let point P with coordinates $(a, f(a))$ be a curve on the curve $y = f(x)$. Let Q be the point where the tangent line to the curve intersects the x -axis and let R be the point with coordinates $(a, 0)$

- (a) Find the length of segment $\overline{QR}$.
- (b) How is part (a) related to part (e) of the previous question?

1 Exercises

Exercise 147 Find an expression for the derivative

$$\frac{d}{dt}(e^{\cos t}).$$

Explanation. We need to use the chain rule because we are differentiating the composition of functions. So let

$$y = e^{\cos t}$$

and

$$u = \cos t.$$

Then

$$y = e^u$$

and

$$\begin{aligned} \frac{dy}{dt} &= \frac{dy}{du} \cdot \frac{du}{dt} \\ &= \left(\frac{d}{du}(e^u) \right) \left(\frac{d}{dt}(\cos t) \right) \\ &= e^u (-\sin t) \\ &= (-\sin t)e^{\cos t}. \end{aligned}$$

Exercise 148 Find expressions for each of the following derivatives. Follow the previous exercise exactly in using the chain rule.

(a) $\frac{d}{dt} \left(1000e^{\frac{t}{5}} \right)$

(b) $\frac{d}{dt} \left(\frac{10}{3 + 5e^t} \right)$

(c) $\frac{d}{dx} (\sin(e^{4x}))$

(d) $\frac{d}{dt} (1000(2^t))$

Hint: Write 2^t as

$$2^t = (e^{\ln 2})^t = e^{t \ln 2}$$

and use the chain rule.

The Chain Rule

An introduction to the chain rule.

1 Introduction

The chain rule tells us how to differentiate the composition of functions. Computationally, it is the *most* important part of this course. Conceptually, it is the most intuitive. In fact, even though you might not realize it, you already know the chain rule. And you've known it for a long time, well before you ever heard of a derivative.

Here's an example. Suppose you're ascending in a hot air balloon at the constant rate of 200 ft/min. Suppose also that the air temperature decreases at the constant rate of $0.003^{\circ}\text{C}/\text{ft}$. The question is at what rate, with respect to time, is the temperature of your surrounding environment changing.

It should be intuitive that we need only multiply these rates. So the temperature is changing at the constant rate of

$$\left(-0.003 \frac{^{\circ}\text{C}}{\text{ft}}\right) \left(200 \frac{\text{ft}}{\text{min}}\right) = -0.6 \frac{^{\circ}\text{C}}{\text{min}}.$$

Here we have two linear functions, one

$$h = g(t)$$

that expresses our altitude (in feet) as a function of time (say in minutes past 1pm), the other

$$C = f(h)$$

that expresses the temperature (in Celsius degrees) in terms of altitude (in feet). And the composition

$$C = f(g(t))$$

expresses the temperature in terms of time. Since the height and temperature functions h and f have respective constant rates of change $\Delta h/\Delta t$ and $\Delta C/\Delta h$, the composition has constant rate of change

$$\frac{\Delta C}{\Delta t} = \frac{\Delta C}{\Delta h} \cdot \frac{\Delta h}{\Delta t}.$$

Learning outcomes:

Author(s):

And this is what the chain rule says, that the derivative of the composition of functions is the product of their derivatives.

Well, not quite. We need to be more precise because things are more subtle when the rates of change are not constant. Suppose for example, we know only that at precisely 2pm our balloon is rising at the rate of 200 ft/min. Suppose also at 2pm we are 2500 feet above the ground and that at this altitude the temperature decreases at the rate of $0.003^{\circ}\text{C}/\text{ft}$. What can we say about the rate, with respect to time, at which the temperature of our surrounding environment is changing at 2pm?

Well, there's nothing else we can really do except multiply the two rates and end up with almost the same conclusion, that at 2pm the temperature of our surrounding environment is decreasing at the rate of $0.6^{\circ}\text{C}/\text{min}$. But this reasoning rests on the subtle assumption that the temperature ($C = f(h)$) and altitude ($h = g(t)$) functions are *differentiable* at their respective inputs $h = 2500$ and $t = 60$. What this means intuitively is that these functions act sufficiently like linear functions near their respective inputs to make the above computation remain valid.

It all boils down to what we mean by these rates of change. The idea that at 2pm our balloon is ascending at the rate of 200 ft/min seems intuitive, but is it? What do we really mean by a moment in time? Instead of addressing this question directly, we can think of the ascent rate as a *scaling factor* that converts a small interval of time

$$\Delta t = t - 60 \sim 0$$

to an approximation of the actual change in height

$$\Delta h = f(t) - f(60) = f(t) - 2500$$

during that interval. We might say, for example, that our altitude increases by about 200 feet from 2:00pm to 2:01pm. But because our rate of ascent might change considerably during time interval, this approximation might not be very accurate. It would be better to say that in the 1/100th of a minute following 2pm, our altitude increases by about two feet.

More generally, we could approximate the change in height (in feet) over the time interval Δt minutes beginning at 2pm by the product

$$\Delta h \sim \left(\frac{dh}{dt} \Big|_{t=60} \right) \Delta t = 200\Delta t.$$

If we assume the altitude function is *differentiable* at 2pm (which would seem almost certain) then this is a good approximation for small time intervals $\Delta t \sim 0$ in the sense that the error

$$\Delta h - 200\Delta t$$

in the approximation approaches zero *faster* than Δt . That is, we should expect that

$$\lim_{\Delta t \rightarrow 0} \frac{\Delta h - 200\Delta t}{\Delta t} = 0.$$

And this is equivalent to saying that the altitude function is differentiable at 2pm and that

$$\left. \frac{dh}{dt} \right|_{t=60} = \lim_{\Delta t \rightarrow 0} \frac{\Delta h}{\Delta t} = 200 \text{ ft/min.}$$

Similarly, the rate of change of temperature with respect to altitude,

$$\left. \frac{dC}{dh} \right|_{h=2500} = -0.003^\circ\text{C/ft},$$

is best understood in terms of small changes. Over a small change in altitude of $\Delta h = h - 2500$ feet, the change in temperature

$$\Delta C = g(h) - g(2500)$$

is well-approximated by the product

$$\Delta C \sim \left(\left. \frac{dC}{dh} \right|_{h=2500} \right) \Delta h = -0.003\Delta h.$$

By this we mean that

$$\left. \frac{dC}{dh} \right|_{h=2500} = \lim_{\Delta h \rightarrow 0} \frac{\Delta C}{\Delta h} = -0.003^\circ\text{C/ft}.$$

With these assumptions we can compute the rate of change in temperature with respect to time at 2pm as

$$\begin{aligned} \left. \frac{dC}{dt} \right|_{t=60} &= \lim_{t \rightarrow 0} \frac{\Delta C}{\Delta t} \\ &= \lim_{t \rightarrow 0} \frac{\Delta C}{\Delta h} \cdot \frac{\Delta h}{\Delta t} \\ &= \lim_{h \rightarrow 0} \frac{\Delta C}{\Delta h} \cdot \lim_{\Delta t \rightarrow 0} \frac{\Delta h}{\Delta t} \\ &= \left. \frac{dC}{dh} \right|_{h=2500} \cdot \left. \frac{dh}{dt} \right|_{t=60} \\ &= \left(-0.003 \frac{^\circ\text{C}}{\text{ft}} \right) \left(200 \frac{\text{ft}}{\text{min}} \right) \\ &= -0.6 \frac{^\circ\text{C}}{\text{min}}. \end{aligned}$$

2 Distance to the Horizon

Here's an example of a question more typical of what you see in this class. It looks more complicated, but it is really just like the last example.

Example 31. The function

$$h = g(t) = -4t^2 + 16t + 100, \quad -2 \leq t \leq 5$$

expresses the altitude (in feet) of a balloon in terms of the number of seconds past 2pm.

The function

$$s = f(h) = 1.22\sqrt{h}, \quad 0 \leq h \leq 30000,$$

expresses the distance to the horizon (in miles) in terms of altitude (in feet).

Our problem is to determine the rate at which the distance to the horizon (limited by the curvature of the earth) is changing (with respect to time) for the occupants of the balloon at 2pm.

Explanation. To determine this rate, just as in the previous question, we need only multiply two rates of change:

- (a) the balloon's rate of ascent at 2pm and
- (b) the rate of change in the distance to the horizon with respect to altitude, evaluated at the height of the balloon at 2pm.

These rates are both derivatives. The rate of ascent at 2pm is the derivative

$$\begin{aligned} \left. \frac{dh}{dt} \right|_{t=0} &= \left. \frac{d}{dt} (-4t^2 + 16t + 100) \right|_{t=0} \\ &= \left. (-8t + 16) \right|_{t=0} \\ &= 16 \text{ ft/sec.} \end{aligned}$$

For the second rate of change, in the distance to the horizon with respect to altitude, we first compute the derivative

$$\frac{ds}{dh} = \frac{d}{dh} (1.22\sqrt{h}) = \frac{0.61}{\sqrt{h}} \text{ miles/ft}$$

and then evaluate it at the altitude

$$h = g(0) = 100 \text{ feet.}$$

We get

$$\left. \frac{ds}{dh} \right|_{h=100} = \left. \frac{0.61}{\sqrt{h}} \right|_{h=100} = 0.061 \text{ miles/ft.}$$

Then the rate of change in the distance to the horizon for the occupants of the

balloon at 2pm is

$$\begin{aligned}\frac{ds}{dt}\Big|_{t=0} &= \left(\frac{ds}{dh}\Big|_{h=g(0)}\right) \left(\frac{dh}{dt}\Big|_{t=0}\right) \\ &= \left(0.061 \frac{\text{miles}}{\text{ft}}\right) \left(16 \frac{\text{ft}}{\text{sec}}\right) \\ &= 0.976 \text{ miles/sec.}\end{aligned}$$

Note: What we really did here was to evaluate the derivative of the composition

$$s = f(g(h)) = 1.22\sqrt{-4t^2 + 16t + 100},$$

expressing the distance to the horizon in terms of time, at time $t = 0$.

3 Visualizing the Chain Rule

The image of stretching an elastic band gives us a way to visualize the chain rule.

Suppose we have two stretching functions

$$H = g(L), \quad 0 \leq L \leq 10$$

and

$$s = f(H),$$

where the inputs and outputs are all measured in meters.

We wish to approximate the local stretching factor of the composition

$$s = f(g(L)), \quad 0 \leq L \leq 10$$

at the points $L = 9.5$ meters from the origin on the relaxed band.

Let's think first about what this means. Imagine a thin elastic band running along the x -axis, here between $L = 0$ and $L = 10$ meters. First we stretch the band with the function g . This action sends the points L meters from the origin on the relaxed band to the points $H = g(L)$ meters on the stretched band.

- (a) Watch this first stretching action by sliding the Slider u in Line 2 of the worksheet below from $u = 0$ to $u = 1$.

Desmos link: <https://www.desmos.com/calculator/8whlpa61hn>

Worksheet available at [151: Rubber Band Composition](#)

Next we stretch the stretched band with function f . This action takes the points $H = g(L)$ meters from the origin on the stretched band to the points $s = f(g(L))$ meters from the origin.

- (b) Watch the second stretch by sliding the Slider u_2 on Line 4 from $u_2 = 0$ to $u_2 = 1$.
- (c) Our goal now is to approximate the local stretching factor

$$\left. \frac{ds}{dL} \right|_{L=9.5}$$

of the composition $f \circ g$ at the points $L = 9.5$ meters from the origin in the relaxed band.

Here are two approaches:

- (i) Use the animation to approximate the local stretching factors

$$\left. \frac{dH}{dL} \right|_{L=9.5} \quad \text{and} \quad \left. \frac{ds}{dH} \right|_{H=g(9.5)}.$$

Then use these factors to approximate the local stretching factor of the composition $f \circ g$ at the points $L = 9.5$ meters from the origin in the relaxed band. Explain your reasoning in detail. Include screen shots to help with your explanation. Include units in all computations.

Free Response:

- (ii) Use the animation to approximate the local stretching factor

$$\left. \frac{ds}{dL} \right|_{L=9.5}$$

directly. Explain your reasoning in detail. Include screen shots to help with your explanation. Include units in all computations.

Free Response:

- (iii) Compare the results of the two methods.

Free Response:

4 Computing with the Chain Rule

The chain rule tells us how to differentiate the composition of functions. It says that the derivative of the composition is the product of the derivatives.

Here is a more formal statement.

Theorem 1. (The Chain Rule) If $y = f(u)$ is a differentiable function of u and $u = g(t)$ is a differentiable function of t , then the composition

$$y = f(g(t))$$

is a differentiable function of t and

$$\frac{dy}{dt} = \frac{dy}{du} \cdot \frac{du}{dt}.$$

Or paying closer attention to the inputs,

$$\left. \frac{dy}{dt} \right|_{t=t_0} = \left. \frac{dy}{du} \right|_{u=g(t_0)} \cdot \left. \frac{du}{dt} \right|_{t=t_0}.$$

We'll go through some examples to get an understanding of how to use the chain rule and also why it works.

Examples

Example 32. Find an equation of the tangent line to the curve

$$y = f(x) = (2x^3 + 1)^2$$

at the point $(1, 9)$.

Explanation. Let

$$y = (2x^3 + 1)^2$$

and

$$u = 2x^3 + 1.$$

Then

$$y = u^2$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(u^2) \cdot \frac{d}{dx}(2x^3 + 1) \\ &= 2u(6x^2) \\ &= 2(2x^3 + 1)(6x^2). \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 2(3)(6) = 36,$$

and an equation of the tangent line is

$$y - 9 = 36(x - 1).$$

We can get the same result without the chain rule, by rewriting the original function as

$$y = f(x) = (2x^3 + 1)^2 = 4x^6 + 4x^3 + 1.$$

Then

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} (4x^6 + 4x^3 + 1) \\ &= 4 \frac{d}{dx} (x^6) + 4 \frac{d}{dx} (x^3) + \frac{d}{dx} (1) \\ &= 24x^5 + 12x^2.\end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 14 + 12 = 36,$$

as before.

Example 33. Find an equation of the tangent line to the curve

$$x^2 + y^2 = 25$$

at the point $P(4, -3)$.

Explanation. Solve the above equation for y in terms of x to get

$$y = \pm \sqrt{25 - x^2}.$$

While this equation does not define y as a function of x , near the point $(4, -3)$, the equation

$$y = -\sqrt{25 - x^2}$$

does define y as a function of x .

Now to find the derivative of the function

$$y = -\sqrt{25 - x^2},$$

let

$$u = 25 - x^2.$$

Then

$$y = \sqrt{u} = -u^{1/2}$$

and

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(-u^{1/2}) \cdot \frac{d}{dx}(25 - x^2) \\ &= -\frac{1}{2\sqrt{u}}(-2x) \\ &= \frac{x}{\sqrt{25 - x^2}}.\end{aligned}$$

So the slope of the tangent line to the curve

$$x^2 + y^2 = 25$$

at the point $(P(4, -3))$ is

$$\left. \frac{dy}{dx} \right|_{x=4} = \left(\frac{x}{\sqrt{25 - x^2}} \right) \Big|_{x=4} = \frac{4}{3}$$

and an equation of the tangent line is

$$y + 3 = \frac{4}{3}(x - 4).$$

To find the slope of the tangent line without appealing to the chain rule, note that the curve

$$x^2 + y^2 = 25$$

is a circle centered at the origin. So the tangent line at $P(4, -3)$ is perpendicular to the radius $\overline{OP}$ from the origin to P . Since $\overline{OP}$ has slope $m_1 = -3/4$, the tangent line at $P(4, -3)$ has slope

$$m_2 = -1/m_1 = 4/3.$$

Question 149 (a) Describe a transformation that takes the circle

$$x^2 + y^2 = 25$$

to the ellipse

$$4x^2 + y^2 = 25. \quad (2)$$

(b) Find the image (call it Q) of the point $P(4, -3)$ under the transformation in part (a).

(c) Use the result of Example 2 to find an equation of the tangent line to the ellipse (2) at Q .

(d) Use the chain rule to first find the slope of the tangent line to the ellipse (2) at Q . Then find an equation of the tangent line.

Question 150 The function

$$P = f(t) = 22 + \frac{t}{2} - \frac{3t^2}{4}, 0 \leq t \leq 3,$$

expresses the price of a stock (in dollars/share) in terms of the number of hours past 9am.

(a) Is the price of the stock increasing or decreasing at 11am? At what rate? At what relative rate?

(b) Is the number of shares you can buy with \$1000 increasing or decreasing at 11am? Approximate the rate numerically.

(c) Compute the exact rate in part (b). Compute also the relative rate of change (with respect to time) in the number of shares you can buy with \$1000 at 11am.

Start your solution to this problem by defining a new function with a new function name and a new dependent variable.

Question 151 At 10am the price of a stock is increasing at the relative rate of $p\%/hr$. Is the number of shares you can buy with \$1000 increasing or decreasing at 10am? At what relative rate?

Answer this question in two ways:

(i) Going back to limits and using the definition of the derivative.

(ii) Using the chain rule.

Either way, start your solution with definitions as in the previous problem.

Question 152 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a 14-foot ladder leans against the trunk, the other rests on the horizontal ground.

Let t be the distance between the top of the ladder and the base of the trunk (measured in feet) and s the distance between the bottom of the ladder and the base of the trunk (also measured in feet).

Geogebra link: <https://www.geogebra.org/m/ctf2bcqz>

Geogebra activity available at 151: Ladder and Tree Part 2

(a) Use the law of cosines to write an equation relating t and s .

(b) Use the result of part (a) to find the two possible values of t when $s = 16$.

For the remainder of this problem we'll focus on positions of the ladder when $s \sim 16$ and $t \sim 10$.

(c) For these positions, complete the square to find a function

$$t = f(s)$$

that expresses the distance (in feet) from the top of the ladder to the base of the trunk in terms of the distance (in feet) from the bottom of the ladder to the trunk's base.

(d) Drag the slider θ in the worksheet above to approximate the value of the derivative

$$\left. \frac{dt}{ds} \right|_{s=16}.$$

Include units and explain your reasoning.

(e) Use your function from part (c) to find an expression for the derivative dt/ds . Show every step in computing the derivative as in Examples 1 and 2 above.

(f) Evaluate the derivative

$$\left. \frac{dt}{ds} \right|_{s=16}$$

and compare the exact value with your estimate from part (d).

(g) What are the units of the derivative in part (f)? Explain the meaning of the derivative, not by giving a standard response about the rate of change of some quantity with respect to another, but by relating small changes.

(h) Use your response to part (f) to write an approximation for the change

$$\Delta t = f(s) - f(16)$$

in terms of the change

$$\Delta s = s - 16$$

for values of s near $s = 16$. Use this approximation to estimate the distance between the top of the ladder and the base of the trunk when the bottom of the ladder is 15.6 feet from the base of the trunk. Compare your approximation with the exact distance.

(i) Let $\theta = m\angle GBA$ be the radian measure of the angle the ladder makes with the trunk.

part (i) Find an expression for the derivative $d\theta/ds$.

part (ii) Evaluate the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=16}$$

and interpret its meaning in terms of small changes. Be sure to explain the significance of the derivative's sign as well.

Example 34. (a) Find the slope of the tangent line to the curve

$$y = f(\theta) = 6 \sin \theta$$

at the point with coordinates $P(2\pi/3, 3\sqrt{3})$. We do not need the chain rule for this.

(b) Describe a transformation that takes the graph of $y = f(\theta)$ to the graph of the function

$$y = g(\theta) = 6 \sin(\theta/2).$$

(c) Find the image of the point $P(2\pi/3, 3\sqrt{3})$ under the above transformation.

(d) Use the results of parts (a) - (c) and the demonstration below to guess the slope of the tangent line to the curve $y = g(\theta)$ at the point $Q(4\pi, 3\sqrt{3})$.

Desmos link: <https://www.desmos.com/calculator/mqjxpsqyo5>

Desmos activity available at 151: Chain Rule 1

(e) Use the chain rule to confirm your guess from part (d).

Example 35. Find an equation of the tangent line to the curve

$$y = f(t) = 4 \sin^2 \left(\frac{\pi}{6} t \right)$$

at the point $(2, 3)$.

Explanation. Let

$$y = 4 \sin^2 \left(\frac{\pi}{6} t \right) = 4 \left(\sin \left(\frac{\pi}{6} t \right) \right)^2$$

and

$$u = \sin \left(\frac{\pi}{6} t \right).$$

Then

$$y = 4u^2$$

and

$$\begin{aligned} \frac{dy}{dt} &= \frac{dy}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du} (4u^2) \cdot \frac{d}{dt} \left(\sin \left(\frac{\pi}{6} t \right) \right) \\ &= 8u \cos \left(\frac{\pi}{6} t \right) \frac{d}{dt} \left(\frac{\pi}{6} t \right) \\ &= 8 \sin \left(\frac{\pi}{6} t \right) \cos \left(\frac{\pi}{6} t \right) \frac{\pi}{6}. \end{aligned}$$

Question 153 (a) The third equality above uses the chain rule again. Fill in the missing details of this computation by first letting

$$z = \sin\left(\frac{\pi}{6}y\right)$$

and making an explicit substitution using the variable v in place of u .

(b) Complete the solution by finding an equation of the tangent line to the curve at the point $(2, 3)$.

Question 154 Use the chain rule to compute each of the following derivatives. Show all steps as in Examples 1, 2, and 8 above.

(a) $\frac{d}{dx} \left(\frac{12}{1+x^2} \right)$

(b) $\frac{d}{dt} \left(12(3)^{t/5} \right)$

(c) $\frac{d}{dt} \left(e^{3 \sin(4t)} \right)$

(d) $\frac{d}{d\theta} (12 + 5 \cos(\theta/4))$

Example 36. The function

$$P = f(t) = 10e^{\frac{1}{4}t}, \quad -4 \leq t \leq 12,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

- What are the units of $1/4$ in the function above? How do you know?
- Use the chain rule to find an expression for the growth rate of the population at time t hours past noon. What are the units of the growth rate?
- Express the growth rate from part (b) in terms of the population $P = f(t)$ at time t hours past noon.
- What is the relative instantaneous growth rate of the population? Include units.
- Find the relative average growth rate of the population over a one-hour period.
- Describe what happens to the population every hour.

Example 37. (a) Use the chain rule to find the relative instantaneous growth rate of the population function

$$P = 10(2^t), \quad -2 \leq t \leq 5,$$

where P is measured in millions of bacteria and t is the number of hours past noon.

(b) Describe what happens to the population every hour.

(c) Estimate the population 2 minutes after there are 30 million bacteria.

Question 155 *This problem suggests a way to think about the chain rule geometrically.*

You ride a ferris wheel for one revolution and get off. The function

$$h = f(\theta), \quad 0 \leq \theta \leq 2\pi,$$

expresses your height (in feet) in terms of the wheel's angle of rotation (measured in radians from the moment you boarded).

The function

$$\theta = g(t), \quad 0 \leq t \leq 44,$$

expresses the rotation angle of the wheel in terms of the number of seconds since you boarded.

Our goal is to use the graphs of the function f and g below (take the times t to be positive, not negative as shown) to approximate your rate of ascent $t = 16$ seconds after boarding. We'll do this in two different ways.

(a) *The first way involves a few steps and uses the graphs of both f and g .*

(i) *Approximate the value of the derivative*

$$\left. \frac{dh}{d\theta} \right|_{\theta=g(16)}$$

and interpret its meaning. Include units.

(ii) *Approximate the value of the derivative*

$$\left. \frac{d\theta}{dt} \right|_{t=16}$$

and interpret its meaning. Include units.

(iii) *Use the results of (i) and (ii) and common sense (do not appeal to the chain rule directly) to approximate the value of the derivative*

$$\left. \frac{dh}{dt} \right|_{t=16} = h'(16),$$

where $h(t) = f(g(t))$. Explain your logic and include all units in your computation. Interpret the meaning of this derivative.

(b) The second way has just one step and that's to use the graph of $h = f(\theta)$ to approximate your rate of ascent at time $t = 16$ using the fact that the horizontal blue lines are drawn at intervals of $\Delta t = 2$ seconds. Be sure to explain your logic.

Desmos link: <https://www.desmos.com/calculator/fkgfpsowe8>

Desmos activity available at 151: Ferris Wheel 2

Question 156 This is a continuation of the previous question.

Approximate the time between times $t = 5$ and $t = 22$ seconds when you are ascending at the slowest rate. Approximate this rate of ascent.

Question 157 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

(a) Use the geometry of the ferris wheel (see the picture below) to find a function

$$h = f(\theta), 0 \leq \theta \leq 2\pi,$$

that expresses your height (in feet) above the ground in terms of the rotation angle of the wheel, measured in radians. Use the cosine function, not the sine function.

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

(b) Suppose you move at the constant speed of 10 ft/sec as you ride the ferris wheel.

(i) Use the result of part (a) to find a function $h = g(t)$ that expresses your height (in feet) in terms of the number of seconds since you boarded. Include the appropriate domain.

(ii) Find a function $r = h(t)$ that expresses your rate of ascent (in ft/sec) in terms of the number of seconds since you boarded. Include the appropriate domain. Use vectors to interpret this rate geometrically, in terms of the angle your velocity makes with the horizontal and your speed.

(iii) Write an equation that relates your height h (in feet) and your rate of ascent r (in ft/sec) at any instant. Graph the relation by hand.

(iv) Are you going up or down the second time you are 100 feet above the ground? At what rate?

(v) What is your height when you are descending at the rate of 4 ft/sec? Find all possibilities.

Question 158 The function

$$\theta = f(t), t \geq 0,$$

expresses the radian measure of a ferris wheel's rotation angle in terms of the number of seconds since you boarded. The wheel has radius r feet and its center is b feet above the ground.

(a) Find a function

$$h = g(t), t \geq 0,$$

that expresses your height (in feet) in terms of the number of seconds since you boarded.

(b) Find an expression for your rate of ascent (in ft/sec) at time t seconds after you boarded. Assume g is a differentiable function of t .

(c) Interpret your rate of ascent (part (b)) in terms of your speed at time t and the angle your velocity vector makes with the horizontal. Assume here that $\theta = f(t)$ is an increasing function of t .

Exercises

Exercise 159 Use the chain rule and the facts that

$$\frac{d}{d\theta}(\sin \theta) = \cos \theta$$

and

$$\frac{d}{d\theta}(\cos \theta) = -\sin \theta$$

to compute each of the following derivatives. Show all steps in using the chain rule as in the Explanation to Examples 2 and 9 above.

(a) $\frac{d}{dt} \left(12 - 9 \sin \left(\frac{\pi}{12} t \right) \right)$

(b) $\frac{d}{d\theta} (\sec \theta)$

(c) $\frac{d}{d\theta} (\csc \theta)$

$$(d) \frac{d}{d\theta} (\cos^2 \theta)$$

$$(e) \frac{d}{dt} \left(\sqrt{4 + \sin^4 t} \right)$$

$$(f) \frac{d}{d\theta} (\sec^2 \theta)$$

$$(g) \frac{d}{dt} \left(\frac{1}{5 + 3 \cos(t/2)} \right)$$

Explanation. (f) Let

$$y = \sec^2 \theta = (\sec \theta)^2$$

and let

$$u = \sec \theta.$$

Then

$$y = u^2$$

and

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du} (u^2) \cdot \frac{d}{d\theta} (\sec \theta) \\ &= 2u(\sec \theta \tan \theta) \\ &= 2(\sec \theta)(\sec \theta \tan \theta). \end{aligned}$$

(e) Let

$$y = \frac{1}{5 + 3 \cos(t/2)}$$

and let

$$u = 5 + 3 \cos(t/2).$$

Then

$$y = u^{-1}$$

and

$$\begin{aligned} \frac{dy}{dt} &= \frac{dy}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du} (u^{-1}) \cdot \frac{d}{dt} (5 + 3 \cos(t/2)) \\ &= -\frac{1}{u^2} \left(-3 \sin \left(\frac{t}{2} \right) \frac{d}{dt} \left(\frac{t}{2} \right) \right) \\ &= \frac{3 \sin \left(\frac{t}{2} \right)}{2(5 + 3 \cos(t/2))^2}. \end{aligned}$$

Exercise 160 You jog around a circular track of radius r feet at the constant speed of v ft/sec. A flagpole lies b feet due east of the track's center.

(a) Use the animation below (and nothing else) to sketch by hand a graph of the function

$$s = f(\theta), \theta \geq 0,$$

that expresses your distance (in feet) to the flagpole in terms of your angle $\theta = \angle FOJ$ of rotation about the track's center, measured in radians from the time you start. Assume you start at the point A on the track nearest the flagpole. Be sure to include variable names, units, and scales on your axes. Explain your reasoning. For this particular graph assume that $r = 40$ and $b = 96$ (be sure to adjust the sliders in the worksheet to have these values).

Desmos link: <https://www.desmos.com/calculator/4pndurvhd>

Worksheet available at 151: Jogger

(b) Using only your graph of the function f from part (a), sketch by hand a graph of the derivative

$$y = f'(\theta) = ds/d\theta$$

when $r = 40$ and $b = 96$. Be sure to include variable names, units, and scales on your axes. Explain your reasoning.

(c) Using only the animation (stop the motion), approximate the value of the derivative

$$\left. \frac{ds}{d\theta} \right|_{\theta=3\pi/2}$$

when $r = 40$ and $b = 96$. Include units. Explain your reasoning.

(d) Use trigonometry to find an expression for the function $s = f(\theta)$ in terms of the parameters r and b (ie. do not assume $r = 40$ and $b = 96$).

(e) Find an expression for the derivative $ds/d\theta$. Show every step in using the chain rule as in Examples 1 and 2. Do not assume $r = 40$ and $b = 96$

(f) Use your result from part (e) to compute the exact value of the derivative

$$\left. \frac{ds}{d\theta} \right|_{\theta=3\pi/2}$$

when $r = 40$ and $b = 96$. Compare this with your estimate from part (c).

(g) What are the units of the derivative in part (f)? Explain the meaning of the derivative, not by giving a standard response about the rate of change of some quantity with respect to another, but by relating small changes.

(h) Use your response to part (f) with $r = 40$ and $b = 96$ to write an approximation for the change

$$\Delta s = f(\theta) - f(3\pi/2)$$

in terms of the change

$$\Delta\theta = \theta - 3\pi/2$$

for values of θ near $\theta = 3\pi/2$.

(i) Find a function

$$s = g(t), t \geq 0,$$

that expresses your distance (in feet) to the flagpole in terms of the number of seconds since you started jogging. Assume you start at the point on the track nearest the flagpole.

(j) Find a function that expresses the rate of change (with respect to time) in your distance to the flagpole in terms of the number of seconds since you began jogging.

(k) Express the rate of change in part (b) in terms of your speed and the angle between the following two vectors:

- the vector that gives your position relative to the flagpole
- the vector that points in the direction of your motion

Exercise 161 High tide of 10.83 feet at Edmonds at 12:28am, May 15, low tide at 6:58am of ?? feet, lowest tide of 1.64 feet at 5:32pm.

Desmos link: <https://www.desmos.com/calculator/zta9tkzzmx>

Worksheet available at 151: Edmonds Tides

Exercise 162 A pendulum of length L feet is $L + 5$ oscillates sinusoidally between angles $-\theta_0$ and θ_0 with period $2\pi\sqrt{L/g}$ seconds, where g is a constant. The angles $\pm\theta_0$ are measured in radians from the downward vertical.

(a) Find a function

$$h = f(\theta), -\theta_0 \leq \theta \leq \theta_0,$$

that expresses the height of the pendulum above its stable equilibrium position (ie. its lowest point) in terms of the angle of rotation. Use the cosine function.

(b) Assume now that the pendulum is released from rest from the displacement angle $\theta = \theta_0$ at time $t = 0$ seconds. Find a function

$$\theta = g(t), t \geq 0,$$

that expresses the displacement angle (in radians) in terms of the number of seconds since the pendulum was released. Use the cosine function.

(c) Find a function

$$h = w(t), t \geq 0$$

that expresses the height (in feet) of the pendulum above equilibrium in terms of the number of seconds since the pendulum was released.

(d) Find an expression for the derivative dh/dt and interpret its meaning.

Exercise 163 The function

$$h = f(s) = 3 + 2 \cos s, 0 \leq s \leq 3,$$

expresses the altitude (in thousand of feet) along a mountain trail in terms of the distance (measured in miles) from the summit. The distance is measured along the trail.

The function

$$s = g(t) = t + \frac{t^2}{4}, 0 \leq t \leq 2,$$

expresses your distance from the summit (in miles) in terms of the number of hours past noon.

The goal of this problem is to compute the rate (with respect to time) at which your altitude is changing at 1pm.

Desmos link: <https://www.desmos.com/calculator/twfsqmmgyb>

Worksheet available at 151: Mountain Road

(a) Find an expression for a function $h = k(t)$ that expresses your altitude (in thousands of feet) in terms of the number of hours past noon. Include a domain.

(b) Evaluate the derivative

$$\left. \frac{ds}{dt} \right|_{t=1}$$

and interpret its meaning.

(c) Evaluate the derivative

$$\left. \frac{dh}{ds} \right|_{s=g(1)}$$

and interpret its meaning.

(d) Use the results of (b) and (c) to compute the rate (with respect to time) at which your altitude is changing at 1pm.

(e) Express the rate of part (d) as a derivative.

(f) Express the derivative

$$\left. \frac{dh}{dt} \right|_{t=1}$$

in terms of the derivatives

$$\left. \frac{ds}{dt} \right|_{t=1}$$

and

$$\left. \frac{dh}{ds} \right|_{s=g(1)}.$$

Exercise 164 A weight is attached to the midpoint of a 10-foot rope.

You hold the ends of the rope 10 feet apart at the same height. You then move the ends directly toward each other, each at the constant speed of 2 ft/sec.

Desmos link: <https://www.desmos.com/calculator/grostrmdln>

Worksheet available at 151: *Weighted Rope*

(a) Find a function

$$v = f(t), \quad 0 \leq t \leq 2.5,$$

that expresses the speed of the weight (in feet/sec) in terms of the number of seconds since you began moving your hands.

(b) Describe how the speed of the weight varies.

(c) Comments?

The Chain Rule and Polynomials

Problems with polynomials and the chain rule.

1 Practice Questions

Question 165 The function

$$h = f(s), 0 \leq s \leq 3,$$

expresses the altitude (in thousands of feet) of a trail in terms of the distance (in miles) from the trailhead.

The function

$$s = g(t), 0 \leq t \leq 4,$$

expresses your distance from the trailhead (in miles) as you hike along the trail in terms of the number of hours past noon.

The functions f and g are graphed below.

Desmos link: <https://www.desmos.com/calculator/sm5rxml9zi>

Worksheet available at 151: [Hiking Trail](#)

- (a) Use the graphs above to estimate the following. Include units and write each as a derivative using the Leibniz notation.
 - (i) your elevation at 2:30pm.
 - (ii) your hiking speed at 2:30pm.
 - (iii) the steepness of your part of the trail at 2:30pm. Interpret the meaning of this in terms of small changes.
 - (iv) the rate (in feet/hour) at which you are gaining or losing altitude at 2:30pm.

- (b) Now suppose that

$$h = f(s) = -\frac{s^3}{3} + s^2 + s + 2, 0 \leq s \leq 3$$

and

$$s = g(t) = 0.1t^2 + 0.35t, 0 \leq t \leq 4.$$

Learning outcomes:
Author(s):

- (i) Find the exact values of each of the three derivatives in part (a). Use the desmos worksheet to save time by entering $f'(\text{?})$ and $g'(\text{?})$ in a blank line, where you need to determine the inputs to these derivatives.

Question 166 Due to a late-season frost in eastern Washington, the price of Cosmic Crisp apples is rising precipitously. The function

$$P = f(t), 0 \leq t \leq 8,$$

expresses the price (in dollars/pound) in terms of the number of hours past 9am on April 25, 2025.

- (a) Find an expression for the function

$$N = g(P)$$

for the number of pounds of apples you can buy with \$36 in terms of the price (in dollars/pound).

- (b) Evaluate the derivative

$$\left. \frac{dN}{dP} \right|_{P=4}.$$

Include units.

- (c) Interpret the meaning of the above derivative in terms of small changes.
 (d) Find an expression for the derivative dN/dt in terms of P and the derivative dP/dt .
 (e) Suppose

$$P = f(t) = 2 + \frac{2}{5}t, 0 \leq t \leq 8,$$

and determine the rate of change (with respect to time) at 2pm in then number of pounds of apples you can buy with \$36.

- (f) Suppose (for an arbitrary differentiable price function) that at some instant the price is decreasing at the relative rate of 10%/hour. Is the number of pounds of apples you can buy with \$36 increasing or decreasing at this instant? At what relative rate?

Question 167 The bottom end of a ten-foot ladder slides across a horizontal floor as its top end slides along a vertical wall.

- (a) Find a function

$$h = f(s), 0 \leq s \leq 10,$$

that expresses the height of the top end above the floor (in feet) in terms of the distance of the bottom end from the wall (in feet).

- (b) Use the demonstration below to estimate the derivative

$$\left. \frac{dh}{ds} \right|_{s=8}.$$

Include units. Then use calculus to find the exact value.

Desmos link: <https://www.desmos.com/calculator/0r6g0avjbc>

Worksheet available at 151: Ladder and Wall 25

- (c) Interpret the meaning of the above derivative in terms of small changes.
- (d) Suppose when the ladder's bottom end is eight feet from the wall, it is sliding away from the wall at a speed of 5 ft/sec. Is the top end moving up or down the wall at this instant? At what speed?
- (e) Now suppose the function

$$s = g(t) = 10 - \frac{2}{5}(t - 5)^2, 0 \leq t \leq 10,$$

expresses the distance of the ladder's bottom end from the wall (in feet) in terms of the number of seconds past noon.

- (i) Play the slider in Line 2 in the animation below to watch the motion. Then sketch by hand a rough graph of the composition

$$h = f(g(t)), 0 \leq t \leq 10.$$

Label the axes with the appropriate variable names and units. Then activate the folder in Line 43 to see how you did.

Desmos link:

<https://www.desmos.com/calculator/duq8yqxozk>

Worksheet available at 151: Ladder and Tree 25

- (ii) Use the graph of $h = f(g(t))$ to sketch a graph of the function

$$r = \frac{dh}{dt}.$$

Label the axes with the appropriate variable names and units. Then activate the folder in Line 47 to see how you did. Explain the meaning of this function.

- (iii) What comments/questions do you have about the graph of $r = dh/dt$?
- (iv) At time $t = 7$ seconds past noon, is the top end of the ladder moving up or down? At what speed? Use calculus to find the exact speed.

Question 168 The function

$$g = f(v), \quad 20 \leq v \leq 50,$$

expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

The function

$$G = h(w), \quad 32 \leq w \leq 90,$$

expresses the gas mileage (in km/liter) of the same car in terms of its speed (in km/hour).

Assume there are exactly 1.6 km in one mile and exactly 4 liters in one gallon.

Suppose also that

$$\left. \frac{dg}{dv} \right|_{v=40} = 0.3.$$

- (a) What are the units of the derivative above? Explain its meaning in terms of small changes.
- (b) Express the function h in terms of the function f .
- (c) Use common sense to evaluate the derivative

$$\left. \frac{dG}{dw} \right|_{w=64}.$$

- (d) Use common sense to express the derivative dG/dw in terms of dg/dv . Then use the chain rule and check if you get the same result.

Question 169 The function

$$W = f(h) = \frac{2400}{(h+4)^2}, \quad h \geq 0,$$

expresses the weight (in pounds) of an astronaut in terms of her height (in thousands of miles above the surface of the earth).

- (a) What are the units of the constant 2400? How do you know?

- (b) Suppose when her spaceship is 100 miles above the earth's surface, the astronaut is moving directly toward the earth at a speed of 15,000 miles/hour. At what rate (with respect to time) is her weight changing at this instant. Compute this rate as the product of two other rates. Include units for these rates and explain their meanings in terms of small changes.

Question 170 Suppose an oil spill in the Gulf of Mexico retains the shape of a perfect circle as it expands. At some moment the circle's radius is increasing at the relative rate of 5%/hour. At what relative rate is its area changing at this moment?

2 Discussion Questions

Question 171 The function

$$P = f(t), 0 \leq t \leq 7,$$

expresses the price of a stock (in dollars/share) in terms of the number of hours past 9am. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/0yv49behlw>

Worksheet available at 151: *Stock Chain Rule*

- (a) Is the number of shares you can buy with \$200 increasing or decreasing at 3pm? At what rate? Drag points P and Q in the graph to help approximate the rate. Explain your reasoning.
- (b) Suppose

$$P = f(t) = -\frac{1}{2}t^2 + 4t + 4, 0 \leq t \leq 7.$$

- (i) Find the exact rate (with respect to time) at which the number of shares you can buy with \$200 is changing at 3pm.
- (ii) Find the relative rates of change (with respect to time) at 3pm in the stock price and the number of shares you can buy with \$200. Include units. Compare these rates.

Question 172 A rock is dropped from rest near the surface of some planet and allowed to fall until it hits the surface. The function

$$v = f(h), \quad 0 \leq h \leq 50,$$

expresses the speed of the rock (in meters/sec) in terms of its height above the surface (in meters).

- (a) Would you expect the derivative

$$\left. \frac{dv}{dh} \right|_{h=18}$$

to be positive or negative? Explain your reasoning.

- (b) Suppose

$$v = g(h) = \frac{\sqrt{100 - 2h}}{3}, \quad 0 \leq h \leq 50.$$

- (i) Use the graph of the function $y = \sqrt{x}$ and the sliders u and v below to help approximate the derivative in part (a).

Desmos link:

<https://www.desmos.com/calculator/ilb4wptgxe>

Worksheet available at 151: *Stock Chain Rule*

- (ii) Use the expression for the function $v = g(h)$ to find the exact value of the derivative in part (a).
- (iii) Interpret the meaning of the derivative in part (a) in terms of small changes.
- (iv) Use part (iii) to approximate the rock's speed at a height of 17.7 meters.

Question 172.1 Go back to Question 3, part (e) of this chapter and determine all possible angles the ladder makes with the ground when its top end is moving twice as fast as its bottom end.

The Chain Rule CW1

Problems with the chain rule.

1 Trails

Question 173 *The differentiable function*

$$h = f(s), 0 \leq s \leq 3,$$

expresses the altitude (in meters) of a trail in terms of the distance (in km) from the trailhead.

The differentiable function

$$s = g(t), 0 \leq t \leq 4,$$

expresses your distance from the trailhead (in km) as you hike along the trail in terms of the number of hours past noon.

- (a) Suppose three kilometers from the trailhead the scaling factor that converts a small change in the distance from the trailhead (in km) into an accurate approximation of the corresponding change in altitude (in meters) is -250 . Suppose also that $f(3) = 1200$.
 - (i) What are the units of the scaling factor.
 - (ii) Explain the meaning of the scaling factor in terms of specific small changes.
 - (iii) Approximate the change

$$\Delta h = h - 1200$$

in altitude in terms of a small change

$$\Delta s = s - 3$$

in distance from the trailhead.

- (iv) Use the Leibniz evaluation notation to write the scaling factor as a derivative.

Learning outcomes:
Author(s):

- (b) Suppose also that $g(1.5) = 3$ and that

$$\left. \frac{ds}{dt} \right|_{t=1.5} = 2.$$

- (i) What are the units of this derivative?
- (ii) Use everyday English to explain the meaning of the derivative.
- (iii) Approximate the change

$$\Delta s = s - 3$$

in your distance from the trailhead (in km) in terms of a small change

$$\Delta t = t - 1.5$$

in the number of hours past noon.

- (c) Use the results of parts (a) and (b) to approximate your change in altitude (Δh) in terms of a small change in the number of hours past noon (Δt).
- (d) Use everyday English to explain the meaning of the scaling factor in part (c).
- (e) Our conclusion is

$$\left. \frac{dh}{dt} \right|_{t=1.5} = \left. \frac{dh}{ds} \right|_{s=3} \cdot \left. \frac{ds}{dt} \right|_{t=1.5}$$

or more generally that the derivative of the function

$$h = f(g(t)), \quad 0 \leq t \leq 4,$$

expressing your altitude (in km) in terms of the number of hours past noon is

$$\left. \frac{dh}{dt} \right|_{t=a} = \left. \frac{dh}{ds} \right|_{s=g(a)} \cdot \left. \frac{ds}{dt} \right|_{t=a}.$$

Question 174 The function

$$h = f(s) = 2100 - 200 \sin s, \quad -1 \leq s \leq 3.8,$$

expresses the altitude (in meters) of a 4-km trail in terms of the signed distance (in km) from the one-kilometer mark.

The function

$$s = g(t) = -1 + 2t + \frac{t^2}{5}, \quad 0 \leq t \leq 2,$$

expresses your signed distance (in km) from the one-kilometer mark as you hike along the trail in terms of the number of hours past noon. The signed distance is negative between the trailhead and the one-kilometer mark.

- (a) Find your speed at 12:00pm. Include units. Use the Leibniz evaluation notation.
- (b) Find the steepness of the trail beneath your feet at 12:00pm. Include units. Use the Leibniz evaluation notation. Explain the meaning of the steepness as a scaling factor.
- (c) Use the Leibniz evaluation notation to express your rate of ascent at 12:00pm in terms of the derivatives in parts (a) and (b). Then compute this rate of ascent. Include units.
- (d) Find the angle the trail beneath your feet makes with the horizontal at 12:00pm.

2 Purchasing Power

Question 175 (a) Find a function

$$N = w(p), p > 0,$$

that expresses the number of barells of oil you can buy with \$10,000 in terms of the price (in dollars/barell).

(b) Now suppose the function

$$P = h(t) = 100 - 10 \sin t, -5 \leq 5,$$

expresses the price of oil (in dollars/barrel) in terms of the number of hours past noon.

(i) Is the price of oil increasing or decreasing at noon? At what rate? Use the Leibniz evaluation notation for the derivative.

(ii) Evaluate the derivative

$$\left. \frac{dN}{dp} \right|_{t=0}.$$

Interpret its meaning as a scaling factor. Include units.

(iii) Is the number of barrels you can buy with \$10,000 increasing or decreasing at noon? At what rate? Use the Leibniz evaluation notation.

3 An Oscillating Ferris Wheel

Question 176 (a) Your ride a ferris wheel with a radius of 40 feet. The wheel's center is 50 feet above the ground.

- (i) Find a function

$$h = f(\theta), \quad -\pi \leq \theta \leq \pi,$$

that expresses your height above the ground (in feet) in terms of the wheel's signed angle of rotation, measured in radians from the downward-pointing vertical. Use the cosine function.

- (ii) Find an expression for the derivative $dh/d\theta$. Include units. Interpret its meaning as a scaling factor.
- (b) Now suppose the wheel oscillates sinusoidally between rotation angles $-\pi/2$ and $\pi/2$ with a period of 60 seconds and that you are 50 feet above the ground at time $t = 0$ seconds.

Desmos link: <https://www.desmos.com/calculator/ob9jxby6v6>

Worksheet available at 151: Ferris Wheel 89

- (i) Find a function

$$\theta = a(t), \quad t \geq 0,$$

that expresses the signed angle of rotation in terms of time. Use the cosine function. Start by writing a sentence that defines time precisely and sketching a graph of this function. Label the axes with the appropriate variable names and units.

- (ii) Interpret the meaning of the derivative $d\theta/dt$ in everyday English without using the words velocity or speed.
- (c) Find an expression for the derivative dh/dt of the composition

$$h = f(a(t)).$$

Use the Leibniz notation. Explain the meaning of the derivative in everyday English. This problem is not about velocity. Avoid this word.

- (d) Find your rate of ascent ten seconds after you are 50 feet above the ground.
- (e) Find your speed at the bottom of the ferris wheel.

The Chain Rule and the Price of a Stock

A stock problem with the chain rule.

Question 177 The function

$$P = f(t) = 64 - 20 \cos\left(\frac{1}{4}t\right), \quad 8 \leq t \leq 16,$$

expresses the price of a stock (in dollars/share) in terms of the number of hours past 9am.

- (a) What are the units of the constant 64 in the function above? The constant 20? The constant $1/4$? Explain your reasoning.
- (b) Sketch by hand a reasonably accurate graph of the function. Label the axes with the appropriate variable names and units. Include scales on both axes. Label the coordinates of any turning points.
- (c) Find an expression for the derivative dP/dt . Do this by making an explicit u -substitution and showing each step of the chain rule. But do not use u . Use a more meaningful variable instead.
- (d) What are the units of the derivative dP/dt ? Interpret its meaning.
- (e) Find the stock price when it is decreasing at the rate of $(\$3/\text{share})/\text{hr}$.
- (f) Find a function

$$P = g(r)$$

that expresses the stock price (in dollars/share) in terms of the rate of change in the price (measured in $(\text{dollars}/\text{share})/\text{hr}$). Simplify the expression for your function.

- (g) Sketch by hand a graph of the function $P = g(r)$. Label the axes with the appropriate variable names and units.
- (h) Evaluate the derivative

$$\left. \frac{dP}{dr} \right|_{r=-3}.$$

Include units and interpret its meaning as a scaling factor. Include an example of how the scaling factor acts on a specific small change.

Learning outcomes:
Author(s):

- (i) Is the number of shares you can buy with \$1000 increasing or decreasing when the price is decreasing at the rate of (\$4/share)/hour?
- (i) Use numerical methods, no calculus, to approximate the rate of change (with respect to time) in the number of shares you can buy with \$1,000 when the price is decreasing at the rate of (\$4/share)/hour. No calculus.
- (ii) Use calculus to compute the rate in part (i). Quotient rule not allowed.

Explanation. (a) Because P has units dollars/share, so does the constant 64. And because the outputs to the cosine function are dimensionless, the coefficient 20 has these same units.

Because the inputs to the cosine function are measured in radians (ie. are dimensionless) and t is measured in hours, the units of the coefficient $1/4$ are rad/hour or hr^{-1} .

- (b) The function describes an oscillation about the mean price of 64 dollars/share, with a maximum oscillation about the mean of 20 dollars/share and a period of

$$\frac{2\pi \text{rad}}{\frac{1}{4} \text{ rad/hr}} = 8\pi \text{ hrs.}$$

- (c) The derivative (omitting some of the required steps) is

$$\begin{aligned} \frac{dP}{dt} &= \frac{d}{dt} \left(64 - 20 \cos \left(\frac{1}{4}t \right) \right) \\ &= 20 \sin \left(\frac{1}{4}t \right) \frac{d}{dt} \left(\frac{1}{4}t \right) \\ &= 5 \sin \left(\frac{1}{4}t \right). \end{aligned}$$

- (d) The units of dP/dt are (dollars/share)/hour. The derivative measures the rate at which the stock price is changing with respect to time. Multiplying the derivative by a small time interval (in hours) gives a good approximation to the change in price (in dollars/sh) over that time interval.
- (e) Suppose the stock price is decreasing at the rate of 3 (dollars/share)/hr at time $t = t_0$ hours past 9am. Then

$$\begin{aligned} -3 &= \frac{dP}{dt} \Big|_{t=t_0} \\ &= 5 \sin \left(\frac{1}{4}t \right) \Big|_{t=t_0} \\ &= 5 \sin \left(\frac{1}{4}t_0 \right). \end{aligned}$$

So

$$\sin\left(\frac{1}{4}t_0\right) = -\frac{3}{5}$$

and

$$\begin{aligned}\cos\left(\frac{1}{4}t_0\right) &= \pm\sqrt{1 - \sin^2\left(\frac{1}{4}t_0\right)} \\ &= \pm\frac{4}{5}.\end{aligned}$$

To choose the correct sign, we need to remember the domain of the function $P = f(t)$. Since

$$8 \leq t_0 \leq 16,$$

$$2 \leq \frac{t_0}{4} \leq 4.$$

So

$$\frac{\pi}{2} < \frac{t_0}{4} \leq \frac{3\pi}{2}$$

and therefore

$$\cos\left(\frac{1}{4}t_0\right) < 0.$$

So

$$\cos\left(\frac{1}{4}t_0\right) = -\frac{4}{5}.$$

Finally, the stock price at this time is

$$\begin{aligned}f(t_0) &= 64 - 20\cos\left(\frac{1}{4}t_0\right) \\ &= 64 + 16 \\ &= 80.\end{aligned}$$

So when the stock price is decreasing at the rate of 3 (dollars/share)/hour, the stock is selling at a price of 80 dollars/share.

- (f) The solution to this part is much the same as the solution to part (f). We just replace $-3/5$ (dollars/share)/hr with r (dollars/share)/hr).

Omitting the details, we get

$$\cos(t_0) = -\sqrt{1 - \frac{r^2}{25}} = -\frac{1}{5}\sqrt{25 - r^2},$$

where t_0 is the time (measured in hours past noon) when the price is increasing at the rate of r (dollars/share)/hr.

And the stock price at this time is

$$P = f(t_0) = 64 + 4\sqrt{25 - r^2}.$$

The Chain Rule and the Price of a Stock

and our function is

$$P = g(r) = 64 + 4\sqrt{25 - r^2}.$$

(g) The graph is part of a semi-circle opening downward. I'll leave it to you to find the correct domain of this function.

(h)

$$\begin{aligned}\frac{dP}{dr}\Big|_{r=-3} &= \frac{d}{dr} \left(64 + 4\sqrt{25 - r^2} \right) \Big|_{r=-3} \\ &= -\frac{4r}{\sqrt{25 - r^2}} \Big|_{r=-3} \\ &= 3.\end{aligned}$$

The derivative is

$$3 \frac{\text{dollars/share}}{(\text{dollars/share})/\text{hour}}.$$

LEAVE THE UNITS LIKE THIS. DO NOT SIMPLIFY THEM.

So, for example, during the time it takes the rate of change in the stock price to increase from -3 (dollars/share)/hr to -2.90 (dollars/share)/hr, the price of the stock increases by approximately 0.30 dollars/share.

The Doppler Effect: Stretching Time

The Doppler effect.

The Doppler effect, in its simplest form, describes how the pitch of a sound emitted from a moving source changes frequency as heard by a stationary observer. It's difficult for me to think about frequency and pitch, so we'll follow Hermann Bondi (*Relativity and Common Sense*, 1962, pp. 41-49) and imagine instead a moving source that emits pings at frequent, regularly spaced intervals, say at the rate of f pings/second. Then we'll compute the frequency (measured in 1/sec) at which these pings are heard by a stationary observer.

Relativity and Common Sense

Example 38. In this the simplest case, we'll suppose that the source is heading directly toward the observer at the constant speed of v meters/sec. We'll take the speed of sound to be w meters/sec. What determines the frequency heard by the observer is the ratio $\lambda = v/w$ and we'll assume $\lambda < 1$.

We'll also assume the source emits pings at the rate of f_s pings/second. Our problem is to express the frequency f_o of the pings heard by the observer in terms of λ and f_s .

Drag the slider t_1 below to start the motion of the source P . The observer is at the origin O . We assume the source passes harmlessly through the observer and then begins to move directly away. The pings are emitted at the frequency of 1/sec, on the second (the blinking light at A indicates when the pings are emitted). You should be able to approximate the frequency of the pings heard by the observer in the two cases, when the source is moving directly toward the observer and when it is moving directly away.

What are these frequencies?

Free Response:

Desmos link: <https://www.desmos.com/calculator/h2rw1sr8li>

Doppler Stationary Observer 1

We don't need calculus to analyze this problem. Nevertheless, we'll take an approach that will work in the next variation when the source does not move either directly toward or away from the observer.

Suppose the source collides with the observer at noon. The idea is to find a function $T = f(t)$ that takes as an input the time (measured in the number of

Learning outcomes:
Author(s):

seconds past noon, possibly negative) when a ping was emitted and returns as an output the time (measured in the same manner) that the observer hears that ping.

- (a) Find an expression for the function

$$T = f(t), t \in \mathbb{R}.$$

Work in general with speeds v m/sec for the source and w m/sec for the speed of sound. Note the important point that the speed of sound is constant (at a given air temperature) and independent of the motion of the source.

- (b) Compute the derivative dT/dt . What are its units? There are two cases to consider, when $t < 0$ and when $t > 0$. Evaluate the derivative (two cases) for $\lambda = 1/2, 2/3, 9/10$.
- (c) Interpret the meanings of these derivatives. Then write an equation expressing the frequency f_0 of the pings heard by the observer in terms of $\lambda = v/w$ and the frequency f_s of the pings emitted by the source. There are two cases to consider as above.
- (d) Check your results with the demonstration above where $\lambda = 1/2$.

Example 39. This is a continuation of the previous example, where we now suppose the source (suppose it's an airplane) flies at a constant altitude h km and passes directly over the observer at noon.

In the demonstration below, the pings are emitted once/second and the light at A flashes each second. The light at B flashes each time the observer (still at the origin O) hears a ping.

Desmos link: <https://www.desmos.com/calculator/ow5li8h6o6>

Doppler Stationary Observer 2

- (a) Use the demonstration to approximate the frequency of the pings received by the observer when the plane is
- (i) 10 km west (to the left) of O
 - (ii) directly above O
 - (iii) 10 km east of O
- (b) Now find a function

$$T = g(t), t \in \mathbb{R}.$$

as in the first example. This function takes as an input the time (measured in the number of seconds since noon) when a ping was emitted and returns as an output the time (measured the same way) when the observer hears this ping. Work in general, with v km/sec the speed of the source, w km/hour the speed of sound, and h (in km) the altitude of the plane.

The Doppler Effect: Stretching Time

The problem here should be to relate the frequencies at time $t = -0.5$ when $v = 1$, $w = 2$, and $h = 4$. The result should be that

$$f_1 = 0.7f_2.$$

Doppler Effect with Sound:

Desmos link: <https://www.desmos.com/calculator/gxzmjpgkrr>

Doppler Effect Stationary Observer 2 With Sound

On Trails

Hiking bugs.

1 Trails

Question 178 *The differentiable function*

$$h = f(s), \quad 0 \leq s \leq 5,$$

expresses the altitude (in km) along a trail in terms of the distance from the trailhead (measured along the trail in km).

(a) *Suppose*

$$\left. \frac{dh}{ds} \right|_{s=3} = 0.2.$$

What are the units of this derivative? Interpret the meaning of the derivative using the language of small changes.

(b) *What acute angle does the curve $h = f(s)$ make with the s -axis at the point $(3, f(3))$.*

(c) *What acute angle does the trail make with the horizontal at the point on the trail 3 km from the trailhead?*

(d) *Suppose*

$$\left. \frac{dh}{ds} \right|_{s=4} = 1.$$

Explain what this means.

Explanation. (a) The derivative has units 0.2 km/km.

It is a dimensionless scaling factor that converts a small change in the distance to the trailhead from the 3-km mark to an accurate approximation of the corresponding change in altitude (measured in the same units as the distance from the trailhead, whatever those units may be).

For example, if you walk 1 meter farther along the trail from the 3-km mark, your altitude increases by approximately 0.2 meters.

Learning outcomes:
Author(s):

- (b) Let θ be the acute angle (in radians) the curve $h = f(s)$ makes with the s -axis at the point $(3, f(3))$. Then

$$\tan \theta = \left. \frac{dh}{ds} \right|_{s=3} = 0.2.$$

And since $-\pi/2 \leq \theta \leq \pi/2$,

$$\theta = \arctan(0.2).$$

- (c) Let ϕ be the acute angle (in radians) the trail makes with the horizontal at the 3-km mark. From our description in part (a), we know if you walk $\Delta s \sim 0$ meters farther *along the trail* your altitude increases by approximately $0.2\Delta h$ meters. This tells us

$$\sin \phi \sim \frac{\Delta h}{\Delta s} \sim 0.2.$$

Taking a limit as $\Delta s \rightarrow 0$ shows that

$$\sin \phi = \left. \frac{dh}{ds} \right|_{s=3} = 0.2.$$

Since $-\pi/2 \leq \phi \leq \pi/2$, at the 3-km mark the trail slopes upward at an angle of

$$\phi = \arcsin(0.2) \text{ rad}$$

relative to the horizontal.

- (d) This means that the trail is vertical at the 4-km mark. You would be rock-climbing at this point of the trail, not hiking.

Question 179 The function

$$h = f(s) = 2000 + \frac{1800}{\pi} \tan\left(\frac{\pi}{12}s\right), 0 \leq s \leq 4.2,$$

expresses the altitude of a trail (in meters) in terms of the distance from the trailhead (in km).

- (a) What angle does the trail make with the horizontal at the 4-km mark?
- (b) At what altitude does the trail slope upward at a 30° angle relative to the horizontal?

Question 180 The function

$$h = f(s), 0 \leq s \leq 15,$$

expresses the altitude (in meters) of a beetle's path in terms of the distance along the path (in meters) from the start. Its graph is shown below (blue).

The profile of the actual trail (red) is also graphed below.

Point P with coordinates $(s, h) = (s, f(s))$ lies on the graph of the altitude function. Point B is the corresponding point on the trail. It is s km from the trailhead and has altitude $h = f(s)$.

- (a) Use the graphs to explain the following:
- (i) Why and how you would use the inverse tangent function to compute the acute angle the curve $h = f(s)$ makes with the s -axis at P .
 - (ii) Why and how you would use the inverse sine function to compute the acute angle the trail makes with the horizontal at B .

Free Response:

- (b) Suppose

$$h = f(s) = 5 \sin\left(\frac{s}{5}\right), 0 \leq s \leq 15.$$

- (i) Find a function

$$\theta = g(s), 0 \leq s \leq 15$$

that expresses the acute angle the trail makes with the horizontal in terms of the distance from the start of the trail.

- (ii) At what point on the trail does the trail slope downward at an angle of $\arcsin(4/5)$ below the horizontal?
- (iii) Evaluate the derivative $d\theta/ds$ at the point on the trail where it slopes downward at an angle of $\arcsin(4/5)$ below the horizontal. Include units.

Then interpret the meaning of the derivative as a scaling factor in terms of two specific small changes.

Desmos link: <https://www.desmos.com/calculator/lktn53d7ju>

151: Hiking Trail 54C

2 Trails Described with Functions

In this section we model a trail with a curve $h = f(x)$, where x and h are measured in meters. We take the h -coordinate to measure the height (in meters) above sea-level and suppose the positive x -axis points due east.

Question 181 A beetle crawls eastward along the trail $h = f(x)$ at a constant speed of v_0 meters/sec.

At what rate is the beetle gaining altitude if

- (a) $f(x) = x$?
- (b) $f(x) = 2x$?
- (c) $f(x) = mx$?
- (d) Find the limit of the beetle's rate of ascent as $m \rightarrow \infty$ as it crawls on the trail $h = mx$ at the constant speed of v_0 meters/sec.

Question 182 (a) A beetle crawls along the parabola

$$h = 9 - \frac{1}{4}(x - 6)^2, \quad 0 \leq x \leq 12,$$

(coordinates in meters) in the eastward direction at a constant speed of v_0 m/sec. At what rate is it ascending when it is 10 meters above the ground for the second time?

(b) A beetle crawls along the curve

$$h = 4 - 3 \cos\left(\frac{x}{2}\right), \quad 0 \leq x \leq 12,$$

(coordinates in meters) in the eastward direction at a constant speed of v_0 m/sec. At what rate is it ascending when it is 5 meters above the ground for the second time?

3 Parametrically Defined Trails

Differentiating Trigonometric Functions using Limits

Differentiating trig functions from the definition of the derivative.

1 The Motion of a Balloon

Example 40. Play the slider in Line 1 of the worksheet below to watch the motion of a balloon. The balloon leaves behind tracks at equal intervals of time (see Line 4). Drag the slider n in Line 2 to change the time interval.

Desmos link: <https://www.desmos.com/calculator/xsk95weesg>

151: Motion Cosine Derivative

- (a) Sketch by hand a graph of the function

$$h = f(t), 0 \leq t \leq 20$$

that expresses the balloon's height (in feet) in terms of the number of seconds since the motion began. Label the axes with the appropriate variable names and units.

- (b) Activate the folder *Graph* in Line 4 and play the motion again. How did you do?
- (c) Describe how the balloon's rate of ascent varies over the time of its motion.
- (d) Are there time intervals during which the balloon's rate of ascent appears to be constant? If so, approximate these time intervals?
- (e) Use the motion to approximate the balloon's minimum and maximum rates of ascent.
- (f) Sketch by hand a graph of the function

$$r = g(t) = \frac{dh}{dt} = \frac{d}{dt}(f(t))$$

that expresses the rate of ascent in terms of the number of seconds since the motion began. Label the axes with the appropriate variable names and units.

Learning outcomes:
Author(s):

2 An Important Limit

Computing the derivatives of the trigonometric functions hinges on one very important limit,

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}.$$

Example 41. For the worksheet below, do the following.

- (a) Express the y -coordinate of P in terms of a and θ .

$$a \sin \theta$$

- (b) Express the length of arc AP in terms of a and θ .

$$a\theta$$

- (c) Drag the slider u (another name for θ in Line 1 toward zero. What do you notice about the ratio $\sin \theta / \theta$ as you do this?

Free Response:

Desmos link: <https://www.desmos.com/calculator/2ocbmfdt5r>

151: Sin theta over theta 0

Example 42. **Desmos link:**

<https://www.desmos.com/calculator/xv3rn3tyxx>

151: Sin theta over theta Limit

- (a) Find an expression for the function

$$r = m(\theta)$$

that gives the average rate of change of the function

$$f(\theta) = \sin \theta$$

over the interval $[0, \theta]$. Include the domain.

- (b) Drag the slider u in Line 6 above to sketch by hand a graph of the function m . Label the axes with the appropriate variable names and units. Then activate the folder *Average rate of change function* in Line 1 to see how you did. Zoom out to see more of the graph.

- (c) Approximate the limit

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}$$

in two ways:

- (i) Visually, by zooming in on the origin in the worksheet above.
- (ii) Numerically, with a table of average rates of change. Do this with a calculator and then open the *average rates of change* folder in Line 4 as a check.

- (d) What's your guess?

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1.$$

3 Variations on the Limit

Example 43. The purpose of this example is to evaluate the limit

$$\lim_{\theta \rightarrow 0} \frac{\sin(2\theta)}{\theta}.$$

Desmos link: <https://www.desmos.com/calculator/u1czzdozr7>

151: Sin 2theta over theta

- (a) Interpret the expressions $a \sin 2\theta$ and $a\theta$ geometrically in the worksheet above.

Free Response:

- (b) Drag the slider u (another name for θ) in Line 2 to guess the value of the limit

$$\lim_{\theta \rightarrow 0} \frac{\sin(2\theta)}{\theta}.$$

Free Response:

- (c) To compute the limit we'll make the substitution $u = 2\theta$. Then

$$\theta = u/2$$

and as $\theta \rightarrow 0$,

$$u = 2\theta \rightarrow 0.$$

So

$$\begin{aligned}\lim_{\theta \rightarrow 0} \frac{\sin(2\theta)}{\theta} &= \lim_{u \rightarrow 0} \frac{\sin u}{u/2} \\ &= 2 \lim_{u \rightarrow 0} \frac{\sin u}{u} \\ &= 2(1) \\ &= 2.\end{aligned}$$

4 Sine Proof 1

Example 44. Here's one way to show

$$\frac{d}{d\theta}(\sin \theta) = \cos \theta.$$

It uses the identity

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$$

- (a) First use the result of Example 2 to evaluate the derivative

$$\left. \frac{d}{d\theta}(\sin \theta) \right|_{\theta=0}.$$

- (b) Use the graph of the function $y = \cos \theta$ to evaluate the derivative

$$\left. \frac{d}{d\theta}(\cos \theta) \right|_{\theta=0}.$$

Explain your reasoning with a picture.

Free Response:

- (c) Explain why

$$\left. \frac{d}{d\theta}(\sin \theta) \right|_{\theta=\theta_0} = \left. \frac{d}{d\theta}(\sin(\theta + \theta_0)) \right|_{\theta=0}.$$

Do this by graphing the two functions.

Free Response:

(d) Then use parts (a)-(c).

$$\begin{aligned}
 \frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\theta_0} &= \frac{d}{d\theta} (\sin(\theta + \theta_0)) \Big|_{\theta=0} \\
 &= \frac{d}{d\theta} (\sin \theta \cos \theta_0 + \cos \theta \sin \theta_0) \Big|_{\theta=\theta_0} \\
 &= \frac{d}{d\theta} (\sin \theta \cos \theta_0) \Big|_{\theta=\theta_0} + \frac{d}{d\theta} (\cos \theta \sin \theta_0) \Big|_{\theta=\theta_0} \\
 &= (\cos \theta_0) \frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\theta_0} + (\sin \theta_0) \frac{d}{d\theta} (\cos \theta) \Big|_{\theta=\theta_0} \\
 &= (\cos \theta_0)(1) + (\sin \theta_0)(0) \\
 &= \cos \theta_0.
 \end{aligned}$$

5 Sine Proof 2

Example 45. Here's another way to show

$$\frac{d}{d\theta} (\sin \theta) = \cos \theta,$$

this time with limits. It uses the identity

$$\sin \theta_2 - \sin \theta_1 = 2 \sin \left(\frac{1}{2} (\theta_2 - \theta_1) \right) \cos \left(\frac{1}{2} (\theta_2 + \theta_1) \right).$$

- (a) Use the picture below to prove the above identity, at least when $0 < \theta_1 < \theta_2 < \pi/2$.

Desmos link: <https://www.desmos.com/calculator/yppjwpuhkz>

151: Difference of Sines

- (b) Use the identity to find an expression for

$$\frac{d}{d\theta} (\sin \theta) \Big|_{\theta=\theta_1} = \lim_{\theta_2 \rightarrow \theta_1} \frac{\sin \theta_2 - \sin \theta_1}{\theta_2 - \theta_1}.$$

Free Response:

Explanation. (a) Two key points. One is that

$$P_1 P_2 = 2 \sin \left(\frac{1}{2} (\theta_2 - \theta_1) \right).$$

The other is that

$$\angle P_1 P_2 = \angle MOQ = \frac{1}{2} (\theta_1 + \theta_2).$$

(b)

$$\begin{aligned}
 \left. \frac{d}{d\theta} (\sin \theta) \right|_{\theta=\theta_1} &= \lim_{\theta_2 \rightarrow \theta_1} \frac{\sin \theta_2 - \sin \theta_1}{\theta_2 - \theta_1} \\
 &= \lim_{\theta_2 \rightarrow \theta_1} \frac{2 \sin \left(\frac{1}{2} (\theta_2 - \theta_1) \right) \cos \left(\frac{1}{2} (\theta_2 + \theta_1) \right)}{\theta_2 - \theta_1} \\
 &= \lim_{\theta_2 \rightarrow \theta_1} \cos \left(\frac{1}{2} (\theta_2 + \theta_1) \right) \cdot \lim_{\theta_2 \rightarrow \theta_1} \frac{\sin \left(\frac{1}{2} (\theta_2 - \theta_1) \right)}{\frac{1}{2} (\theta_2 - \theta_1)} \\
 &= (\cos \theta_1) (1) \\
 &= \cos \theta_1
 \end{aligned}$$

6 The Derivative of the Cosine Function

Example 46. (a) One of the functions below is the derivative of the other. Which is which?

Free Response:

Desmos link: <https://www.desmos.com/calculator/atxecntid1>

151: Sine Cosine Derivative

(b) Drag the slider k in Line 1 to find an expression for the derivative

$$\frac{d}{d\theta} (\cos \theta).$$

Explain your reasoning. Include graphs to help with your explanation.

Free Response:

7 Exercises

Exercise 183 (a) Find a simplified equation of the tangent line to the curve

$$y = f(\theta) = 10 - 8 \sin \theta$$

at the point P with coordinates

$$(\theta_0, f(\theta_0)),$$

where

$$\theta_0 = \pi - \arcsin(3/4).$$

Do this without a calculator.

- (b) Check your work in the desmos worksheet below by entering the coordinates of P in Line 3 and an equation of the tangent line in Line 4.

Desmos link: <https://www.desmos.com/calculator/sizljxbth>

151: Tangent to Graph Sine 1

- Exercise 184** (a) Drag the slider in Line 1 in the worksheet below to approximate the y -coordinate(s) of all points on the curve

$$y = f(\theta) = 10 - 12 \cos \theta$$

where the tangent line is parallel to the line

$$3x + y = 15.$$

- (b) Find simplified expressions for the exact y -coordinates without using a calculator.

Start your solution as follows:

Let $(\theta_0, f(\theta_0))$ be the coordinates of a point on the curve $y = f(\theta)$ where the tangent line is parallel to the line

$$3x + y = 15.$$

Then ...

Desmos link: <https://www.desmos.com/calculator/8zwsudnvn0>

151: Tangent to Graph Cosine 1

- Exercise 185** (a) Use the graph below to approximate the minimum and maximum outputs of the function

$$f(\theta) = 3 \sin \theta - 5 \cos \theta.$$

Desmos link: <https://www.desmos.com/calculator/msxpibv77r>

151: Graph Sine Plus Cosine 1

- (b) Use calculus, not a trigonometric identity, to find the exact minimum and maximum outputs.

Start your solution as follows:

Let P with coordinates $(\theta_0, f(\theta_0))$ be a point on the curve

$$y = f(\theta) = 3 \sin \theta - 5 \cos \theta.$$

where the output is a minimum or a maximum. Then the tangent line to the curve at P is horizontal and

$$\left. \frac{d}{d\theta} (3 \sin \theta - 5 \cos \theta) \right|_{\theta=\theta_0} = 0 \dots$$

- (c) Enter equations of the two horizontal lines through the turning points in Lines 1 and 2 of the worksheet to check your work.

Exercise 186 A ferris wheel has a radius of 40 feet and its center is 50 feet above the ground. You ride the wheel for one revolution and then get off.

- (a) Find a function

$$h = f(\theta), \quad 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground (in feet) in terms of the rotation angle of the wheel (measured in radians with respect to the downward-pointing vertical.)

- (b) The wheel comes to a stop for the first time at some rotation angle $\theta = \theta_0$ when you are 80 above the ground for the second time.

- (i) Use numerical methods to approximate the scaling factor that converts a small change

$$\Delta\theta = \theta - \theta_0$$

in the rotation angle to an accurate approximation to the corresponding change

$$\Delta h = f(\theta_0 + \Delta\theta) - f(\theta_0)$$

in your height above the ground.

- (ii) Use the derivative, not vectors, to find the exact value of the scaling factor.
- (iii) What are the units of the scaling factor?
- (iv) Interpret the scaling factor geometrically.
- (v) Suppose the wheel turns through the angle $\Delta\theta = 0.05$ radians and stops again. Use the previous result to approximate your new height.

Derivatives of Trigonometric Functions

Working with trigonometric functions and their derivatives.

1 Introduction

Question 187 You ride a ferris wheel of radius 50 feet at a constant speed of 5 ft/sec. The wheel's center is 60 feet above the ground.

Desmos link: <https://www.desmos.com/calculator/rlxg2wbms>

151: Vector Ferris Wheel

- (a) Use precalculus and common sense to find your rate of ascent when you are
 - (i) 60 feet above the ground for the first time
 - (ii) 60 feet above the ground for the second time
 - (iii) 110 feet above the ground
 - (iv) 80 feet above the ground for the first time
 - (v) 80 feet above the ground for the second time
- (b) Express your rate of ascent in terms of the angle θ your position vector relative to the wheel's center makes with the rightward horizontal. Assume θ is measured in the same sense as the rotation of the wheel.
- (c) Express your rate of ascent in terms of the angle ϕ your position vector relative to the wheel's center makes with the upward vertical. Assume ϕ is measured in the same sense as the rotation of the wheel.

2 Velocity

Velocity is the rate of change of position with respect to time. Since position is a vector, so is velocity. The instantaneous velocity vector is the derivative

Learning outcomes:
Author(s):

(with respect to time) of the position vector. It (the velocity vector) points in the direction of motion and is therefore tangent to the path. The length of the velocity vector is the speed.

Play the slider u (time) below to watch a projectile motion in a uniform gravitational field.

Desmos link: <https://www.desmos.com/calculator/vdtsb2fc8j>

151: Projectile Motion

- (a) When is the object speeding up? Slowing down? How can you tell from looking at its velocity vector?
- (b) Suppose the position vector (measured in meters) at time t seconds past launch is

$$\langle x, y \rangle = \langle (v_0 \cos \phi)t, (v_0 \sin \phi)t - \frac{1}{2}gt^2 \rangle, \quad 0 \leq t \leq \frac{2v_0 \sin \phi}{g}.$$

Here v_0 is the initial speed (in m/sec), ϕ is the launch angle relative to the horizontal, and g is the magnitude of the gravitational acceleration (in m/sec²).

Find an expression for the velocity vector.

3 Derivatives of the Sine and Cosine Functions

We just saw we can differentiate the position vector of a motion to find the velocity vector, at least if we know how to differentiate the component functions. There is one case where we can reverse this process. The next example shows how and gives a way to find the derivatives of the sine and cosine functions.

Example 47. The idea is to look at the motion with position vector (in meters)

$$\overrightarrow{OP} = \langle 1 \cos(1t), 1 \sin(1t) \rangle,$$

expressed in terms of the number of seconds past noon. Play the slider u (another name for t) in the worksheet below to watch the motion.

Desmos link: <https://www.desmos.com/calculator/ov0azsjucq>

151: Circular Motion

- (a) What are the units of the blue 1 and the red 1 in the parameterization.

- (b) What is the speed of the motion?
- (c) Use the geometry of the circle to find the components of the velocity vector.
- (d) What does this tell us about the derivatives

$$\frac{d}{dt}(\cos t) \quad \text{and} \quad \frac{d}{dt}(\sin t).$$

- (e) Use the same idea to compute the derivatives

$$\frac{d}{dt}(\cos(3t)) \quad \text{and} \quad \frac{d}{dt}(\sin(3t)).$$

Then use the chain rule to compute the derivatives again.

4 Riding a Ferris Wheel

Question 188 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

- (a) Use the figure below to find a function

$$h = f(\theta), \quad 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground in terms of the rotation angle of the wheel (measured in radians since the time you boarded). Use the cosine function, not the sine function.

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

- (b) The function

$$\theta = a(t), \quad 0 \leq t \leq 120,$$

expresses the rotation angle (in radians, measured since you boarded the wheel) in terms of the number of seconds since you got on. Use the graph of this function below to approximate the your rate of ascent at time $t = 70$ seconds.

Desmos link: <https://www.desmos.com/calculator/kijv21gots>

151: Ferris Wheel Angle

- (c) Now suppose

$$\theta = a(t) = \pi \left(1 - \cos \left(\frac{\pi}{120} t \right) \right), \quad 0 \leq t \leq 120,$$

and compute your exact rate of ascent at time $t = 70$ seconds. Compare this with your estimate.

5 Simple Harmonic Motion

Question 189 Suppose for this problem that the earth is a ball with uniform density of radius 4000 miles. Now imagine drilling a straight tunnel through the earth from the North Pole to the South Pole.

A subway car dropped from rest at the north pole falling through the tunnel would then oscillate in simple harmonic motion between the poles and return to the north pole every 84 minutes. This means we can think of the car as being dragged along by a point moving around the earth at constant speed as illustrated below.

Desmos link: <https://www.desmos.com/calculator/ij8dqowgza>

Desmos activity available at 142: [Simple Harmonic Motion](#)

- (a) Find a function

$$s = f(t), t \geq 0,$$

that expresses the distance of the rock (measured in thousands of miles) from the South Pole in terms of the number of minutes since the rock was released. Assume the rock was dropped at noon on July 1, 2085.

- (b) Use your function and calculus to determine the car's speed as it passes the center of the earth. Then find the speed without calculus.
- (c) Express the average speed of the car during the time for one oscillation in terms of its maximum speed.
- (d) Express the car's speed when it is $2/3$ of the way from the earth's center to the South Pole in terms of its maximum speed.

Question 190 Assume for this question that each month has 30 days and that the number of hours of daylight/day in Seattle is a sinusoidal function of time. Assume also that on June 21, Seattle gets a maximum of 16 hours of daylight/day and that on December 21, Seattle gets a minimum of 8 hours of daylight/day.

- (a) Find a function

$$H = f(t), 0 \leq t \leq 12,$$

that expresses the number of daylight hours/day in Seattle in terms of the number of months since June 21. Use the cosine function. Start by sketching a graph. Explain your reasoning.

- (b) Use your function to determine the number of hours of daylight/day that Seattle gets on August 21.
- (c) Use calculus to approximate about how many more minutes of daylight/-day we get tomorrow than today.
- (d) Use your function to determine the day(s) of the year when Seattle gets 14 hours of daylight/day.

6 Two Optimization Problems

Question 191 The function

$$h = f(t) = 1200 + 500 \cos t + 300 \sin t, \quad 0 \leq t \leq 10,$$

expresses the altitude (in feet) of a balloon in terms of the number of hours past noon.

Desmos link: <https://www.desmos.com/calculator/f1ldi6yrek>

151: Trig 1

- (a) Use calculus to find the minimum and maximum heights of the balloon between noon and 6pm. Do not use a calculator.
- (b) Find the balloon's maximum rate of ascent. No calculator.

Question 192 Use calculus and some trigonometry (but not just trigonometry) to find the minimum and maximum values of the function

$$f(\theta) = 3 \cos \theta - 5 \sin \theta, \quad 0 \leq \theta \leq \pi.$$

Do not use a calculator.

7 Exercises

Question 193 The function

$$h = f(t) = 800 - 200 \tan t, \quad -1.3 \leq t \leq 1.3,$$

expresses the altitude (in feet) of a balloon in terms of the number of hours past noon.

Desmos link: <https://www.desmos.com/calculator/re6nqofgs0>

151: Trig 2

Find exact answers to the following questions without using a calculator.

- (a) When is the balloon descending at the rate of 150 ft/hour?
- (b) How high is the balloon when it is descending at the rate of 500 ft/hour?
- (c) Express the balloon's rate of ascent (in feet/hour) in terms of its altitude.

Question 194 The function

$$h = f(t) = 11 - 7 \cos t, \quad 0 \leq t \leq 6,$$

expresses the depth of the water (in feet) at the Edmonds Pier in terms of the number of two-hour periods past noon.

Desmos link: <https://www.desmos.com/calculator/4bh7kimi7f>

151: Trig 3

Find exact answers to the following questions without using a calculator.

- (a) Is the depth of the water increasing or decreasing at 2pm? At what rate?
- (b) At what rate is the depth of the water changing when the water is 6 feet deep?

Question 195 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

- (a) Use the figure below to find a function

$$h = f(\theta), 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground in terms of the rotation angle of the wheel (measured in radians since the time you boarded). Use the cosine function, not the sine function.

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

- (b) The wheel stops when you are 100 feet above the ground for the second time. It then starts again and turns through a small angle of $\Delta\theta$ radians before stopping again.

Desmos link: <https://www.desmos.com/calculator/gi8obqmnv>

151: Ferris Wheel 34

- (i) Zoom in on point P above to approximate the change in your height (in feet) if $\Delta\theta = 0.1$.
- (ii) Use a derivative to approximate the change in your height if $\Delta\theta = 0.1$.
- (iii) Use a derivative to approximate the change Δh in your height (measured in feet) as the wheel turns through the small angle $\Delta\theta$ radians.

8 Visualizing Derivatives

Question 196 You ride a ferris wheel for one revolution and get off. Let

$$h = f(s), 0 \leq s \leq ??,$$

be the function that expresses your height above the ground (measured in feet) in terms of your distance traveled, measured (in feet) along your path from your starting point.

- (a) Choose a radius for the ferris wheel that you think is reasonable and fill in the missing upper bound for the domain of f above.
- (b) Use the demonstration below to sketch by hand a graph of the function

$$r = \frac{dh}{ds} = f'(s).$$

Do not make any computations. Just use the demonstration below. Here are the key points to keep in mind:

- Approximate the derivative dh/ds by the ratio $\Delta h/\Delta s$.
- The length of the red arclength (when the ferris wheel is on the way up) is the input s .
- The length of the purple segment is the output h .
- The lengths of the orange arc and orange segments are Δs .
- The (signed) length of the green segment is Δh .

Explain your reasoning thoroughly. Be sure to include at least the following points:

- The units of the input and output to the derivative
- Scales on the vertical and horizontal axes
- A discussion of how a small change in the input to the function f changes the output at various positions along your ride.
- A discussion of where a small change in the input to f gives the greatest positive change in the output and a consideration of the ratios of these changes
- A discussion of where a small change in the input to f gives the negative change in the output with the greatest magnitude and a consideration of the ratios of these changes
- A discussion of where a small change in the input to f barely changes the output and a consideration of the ratios of these changes

(c) How would your graph of the derivative dh/ds change if you doubled the radius of the ferris wheel? Sketch the new graph.

Desmos link: <https://www.desmos.com/calculator/pxsmo04nmg>

9 The Derivative of the Sine Function

Question 197 Desmos link:

<https://www.desmos.com/calculator/jcmcyrpndw>

Desmos activity available at 151: Derivative of Sine

Question 198 (a) Describe a transformation that takes the graph of the function

$$y = f(\theta) = \sin \theta$$

to the graph of the function

$$y = g(\theta) = \cos \theta.$$

(b) Does that same transformation take the graph of the derivative of the sine function to the graph of the derivative of the cosine function? Do not use any particular facts about these functions to answer this question. Instead, give a general answer that would apply to all pairs of functions similarly related.

(c) Use your answer to part (b) and the fact that

$$\frac{d}{d\theta} (\sin \theta) = \cos \theta$$

to find an expression for the derivative

$$\frac{d}{d\theta} (\cos \theta)$$

of the cosine function.

10 Applications 1

Question 199 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You ride the wheel for one revolution and get off.

(a) Find a function

$$h = f(\theta), 0 \leq \theta \leq 2\pi,$$

that expresses your height above the ground in terms of the rotation angle of the wheel, measured in radians. Use the cosine function, not the sine function.

(b) The wheel stops when you are 100 feet above the ground and on the way up. It then starts again and turns through a small angle of $\Delta\theta$ radians before stopping again. Use the appropriate linear approximation to estimate the change Δh in your height (measured in feet) as the wheel turned through the angle θ .

Geogebra link: <https://www.geogebra.org/m/tn75cq93>

11 Transformations of the Sine Function

Question 200 (a) Describe a transformation that takes the graph of the function

$$y = f(\theta) = \sin \theta$$

to the graph of the function

$$y = g(\theta) = \sin(2\theta).$$

(b) How does that same transformation affect the slope of a line?

(c) Use your answer to part (b) to find an expression for the derivative

$$g'(\theta) = \frac{d}{d\theta}(\sin(2\theta)).$$

12 Applications 2

Question 201 The center of a ferris wheel with a radius of 50 feet is 60 feet above the ground. You travel at a constant speed of 5 ft/sec as you ride the ferris wheel.

(a) Find a function

$$h = f(t), t \geq 0$$

that expresses your height above the ground in terms of the number of seconds since you got on. Use the cosine function, not the sine function.

(b) Are you ascending or descending the second time you are 90 feet above the ground? At what rate? Use the methods of this class, not vectors, to answer this question.

(c) Find your height when you are descending at the rate of 4.8 feet/sec. Give all possibilities. Do not use a calculator except to do arithmetic.

13 MATH 142

Question 202 The graph below shows the x -coordinate function of a beetle moving around a circle at a constant speed.

Desmos link: <https://www.desmos.com/calculator/qi6c9xbhnw>

142: Edmonds Pier 2

Use the graph to answer the following questions. Be sure to include units.

- (a) Find the x -coordinate of the circle's center.
 - (b) Find the radius of the circle.
 - (c) Find the period of the motion. This the time it takes the beetle to make one revolution about the center of its circular path.
 - (d) Find a time when the beetle's x -coordinate is a maximum.
 - (e) Use (a)-(d) to find an expression $x = f(t)$ for the function that expresses the x -coordinate of the beetle (measured in feet) in terms of the number of minutes past noon. Include the domain.
 - (f) Check your expression from part (e) by substituting the two times given in the graph.
-

Trigonometry Tangent Line

Finding an equation of a tangent line.

Question 203 Find an equation of the tangent line to the curve

$$y = f(\theta) = \frac{12}{8 - 3 \cos \theta}$$

at the point where the curve intersects the line $\theta = \arccos(2/3)$.

Explanation. Let

$$\theta_0 = \arccos(2/3).$$

This is the angle between 0 and π whose cosine is $2/3$.

So $\cos \theta_0 = 2/3$ and when $\theta = \theta_0$,

$$\begin{aligned} y &= f(\theta_0) \\ &= \frac{12}{8 - 3 \cos \theta_0} \\ &= \frac{12}{8 - 3(2/3)} \\ &= 2 \end{aligned}$$

To find the slope of the tangent line at the point $(\theta_0, f(\theta_0))$ we need to first compute the derivative

$$\frac{dy}{d\theta} = \frac{d}{d\theta} \left(\frac{12}{8 - 3 \cos \theta} \right).$$

For this let

$$u = 8 - 3 \cos \theta.$$

Then

$$y = 12u^{-1}$$

and by the chain rule

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du} (12u^{-1}) \cdot \frac{d}{d\theta} (8 - 3 \cos \theta) \\ &= (-12u^{-2})(3 \sin \theta) \\ &= -\frac{36 \sin \theta}{(8 - 3 \cos \theta)^2}. \end{aligned}$$

Learning outcomes:
Author(s):

To find the slope of the tangent line to the curve $y = f(\theta)$ at the point $(\theta_0, f(\theta_0))$, we need to evaluate this derivative at $\theta = \theta_0$. This requires simplifying

$$\sin \theta_0 = \sin(\arccos(2/3)).$$

By the Pythagorean theorem

$$\cos^2 \theta_0 + \sin^2 \theta_0 = 1$$

and

$$\sin \theta_0 = \pm \sqrt{1 - (2/3)^2} = \pm \frac{\sqrt{5}}{3}.$$

Now because $0 \leq \theta_0 \leq \pi$,

$$\sin \theta_0 \geq 0$$

and

$$\sin \theta_0 = \frac{\sqrt{5}}{3}.$$

So the slope of the tangent line to the curve $y = f(\theta)$ at the point with coordinates

$$(\theta_0, f(\theta_0)) = (\arccos(2/3), 2)$$

is

$$\begin{aligned} \left. \frac{dy}{d\theta} \right|_{\theta=\theta_0} &= - \frac{36 \sin \theta}{(8 - 3 \cos \theta)^2} \Big|_{\theta=\theta_0} \\ &= - \frac{12\sqrt{5}}{36} \\ &= - \frac{\sqrt{5}}{3}. \end{aligned}$$

So an equation of the tangent line to the curve

$$y = f(\theta) = \frac{12}{8 - 3 \cos \theta}$$

at the point where the curve intersects the line $\theta = \arccos(2/3)$ is

$$y - 2 = -\frac{\sqrt{5}}{3} (\theta - \arccos(2/3)).$$

The Sliding Ladder, Part 1

Small changes in regard to a sliding ladder

Question 204 The ends of a 25-foot ladder rest against a vertical wall and a horizontal floor. The top of the ladder is 20 feet above the floor.

Desmos link: <https://www.desmos.com/calculator/syz7zherzf>

151: Ladder and ArcSine 5

- (a) Drag the slider h (the height of the ladder's top end above the floor) in Line 1 of the worksheet above to approximate the scaling factor that converts (by multiplication) a small change in the angle the ladder makes with the wall (measured in radians) to an accurate approximation of the change in the height of ladder's top end above the floor (measured in feet). The tick marks on the radian protractor are spaced at intervals of 0.1 radians.
- (b) Use calculus to find the exact value of the scaling factor. Include units. Start by defining a function with meaningful variable names. Include units in your definition.
- (c) Approximate the change in the angle the ladder makes with the wall if the ladder's top slides 3 inches down the wall. Then compare your approximation with the actual change.
- (d) How far is the bottom of the ladder from the base of the wall when the scaling factor that converts a small change in the angle the ladder makes with the wall to an accurate approximation to the change in the height of the ladder's top end above the floor is -10 ft/radian?
- (e) When the top end of the ladder is 20 feet above the floor, what is the scaling factor that converts (by multiplication) a small change in the angle the ladder makes with the floor (measured in radians) to an accurate approximation of the change in the height of ladder's top end above the floor (measured in feet).
- (f) Interpret the scaling factors in parts (b) and (e) geometrically.

Explanation. (a) Drag the slider h in Line 1 from $h = 20$ to make a small change Δh in the height. Then use the radian protractor to approximate

Learning outcomes:
Author(s):

the corresponding change $\Delta\theta$ in the angle the ladder makes with the wall. Finally, compute the average scaling factor $\Delta h/\Delta\theta$. Pay attention to the signs of these changes. I'll leave this computation for you.

(b) Let

$$h = f(\theta), \quad 0 \leq \theta \leq \pi/2,$$

be the function that expresses the height (in feet) of the top end of the ladder above the floor in terms of the angle (in radians) the ladder makes with the wall.

Then in right triangle $\triangle AOB$ above,

$$\cos \theta = \frac{h}{25}$$

and

$$h = f(\theta) = 25 \cos \theta.$$

Taking the derivative we get

$$\begin{aligned} \frac{dh}{d\theta} &= \frac{d}{d\theta} (25 \cos \theta) \\ &= -25 \sin \theta. \end{aligned}$$

This expresses the scaling factor in terms of the inclination angle θ .

Now we need to find the scaling factor when $h = 20$. For this, let θ_0 be the angle the ladder makes with the wall when its top end is 20 feet above the floor. Then using the Pythagorean theorem in right triangle $\triangle AOB$ gives the distance (in feet) between the bottom of the ladder and the wall as

$$AO = \sqrt{25^2 - 20^2} = 15.$$

So

$$\sin \theta_0 = \frac{AO}{25} = \frac{15}{25}$$

and the scaling factor is

$$\begin{aligned} \left. \frac{dh}{d\theta} \right|_{\theta=\theta_0} &= -25 \sin \theta \Big|_{\theta=\theta_0} \\ &= -25 \sin \theta_0 \\ &= -25 \left(\frac{15}{25} \right) \\ &= -15. \end{aligned}$$

The units of the scaling factor are ft/rad.

Interpretation: When the top of the ladder is 20 feet above the floor, increasing the angle between the ladder and the wall by 0.1 radians decreases the height of the ladder's top end by approximate 1.5 feet.

(c) Let

$$\Delta\theta = \theta - \theta_0 = \theta - \arccos(4/5)$$

and

$$\Delta h = h - 20$$

be the respective changes in the angle and height. Then for values of $\Delta\theta \sim 0$,

$$\begin{aligned}\Delta h &\sim \left(\frac{dh}{d\theta} \Big|_{h=20} \right) \Delta\theta \\ &= -15\Delta\theta.\end{aligned}$$

So if $\Delta h = -1/4$,

$$-\frac{1}{4} \text{ ft} \sim \left(-15 \frac{\text{ft}}{\text{rad}} \right) \Delta\theta$$

and

$$\Delta\theta \sim \left(\frac{1}{4} \text{ ft} \right) \left(\frac{1}{15} \frac{\text{rad}}{\text{ft}} \right) = \frac{1}{60} \text{ rad}.$$

So if the top of the ladder slides 0.25 feet down the wall to a height of 19.75 feet above the floor, the angle between the ladder and the wall increases by approximately 1/60 radian.

(d) In the expression for the scaling factor

$$\frac{dh}{d\theta} = -(25 \sin \theta),$$

we can interpret $25 \sin \theta$ as the length AO in right triangle AOB . This is the distance from the bottom end of the ladder to the wall.

So when the scaling factor is -10 ft/rad , the bottom end of the ladder is 10 feet from the wall and its top end is

$$OB = \sqrt{25^2 - 10^2} \text{ ft} = 5\sqrt{21} \text{ ft}$$

above the floor.

(e) When the angle between the ladder and the wall increases by $\Delta\theta$, the angle between the ladder and the floor decreases by $\Delta\theta$. So the scaling factor that converts a small change in angle the ladder makes with the floor into an accurate approximation of the change in the height is 15 ft/rad.

Question 205 This problem is similar to the previous one, except that the floor is inclined at an angle of

$$\phi = \arccos(5/6)$$

The Sliding Ladder, Part 1

below the horizontal.

The ladder is still 25 feet long and its top end is 20 feet above the base of the wall.

Desmos link: <https://www.desmos.com/calculator/q2g6qqwlzc>

151: Ladder and Sloping Ground

- (a) What effect do you think this change has on the scaling factor in part (b) of Question 1?
- (b) Answer parts (a)-(f) of Question 1 in this scenario.



The Tangent Function and its Inverse

The tangent function and its inverse.

1 The Tangent Function

The most common way to find the derivative of the function $y = \tan(\theta)$ is to write

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

and use the quotient rule. But this is not all that insightful, and we have not learned the quotient rule yet. Here are two other approaches.

Question 206 We'll use the figure below to find an expression for the derivative

$$\frac{d}{d\theta} (\tan \theta)$$

as follows.

Desmos link: <https://www.desmos.com/calculator/10538a93e8>

151: Tangent Derivative

- First convince yourself that the marked angles $\theta = \angle BOA$ and $\angle QBC$ are congruent. Well almost, assuming $\Delta\theta \sim 0$.
- Express the length OB in terms of θ .
- Express the length QB in terms of θ and $\Delta\theta$.
- Express the length BC in terms of θ and $\Delta\theta$.
- Then find an expression for

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (\tan \theta).$$

Learning outcomes:
Author(s):

Question 207 (a) Find an expression for the average rate of change of the function

$$f(\theta) = \tan \theta$$

between $\theta = \theta_1$ and $\theta = \theta_2$.

(b) Use part (a) and the algebra of limits to find an expression for the derivative

$$\left. \frac{dy}{d\theta} \right|_{\theta=\theta_1} = \left. \frac{d}{d\theta} (\tan \theta) \right|_{\theta=\theta_1}.$$

Hint: Use the identity

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

and modify the explanation to Example 6b in the chapter *Differentiating Trigonometric Functions using Limits* of our class notes.

(c) Repeat parts (a) and (b) for the function $f(\theta) = \cot \theta$.

2 Exercises

Exercise 208 (a) Find the y -coordinates of all points on the curve

$$y = f(\theta) = 10 - 4 \tan \theta$$

where the tangent line is parallel to the line

$$8x - y = 12.$$

Include a graph. Start by defining an unknown like this:

Let

$$(\theta_0, f(\theta_0))$$

be the coordinates of the point on the curve $y = f(\theta)$ where the tangent line is parallel ...

(b) Repeat part (a) to find the y -coordinates of points on the same curve where the tangent line is parallel to the line

$$x - 8y = 12$$

and then the line

$$8x + y = 12.$$

Explain what's going on and why. Include a graph.

Exercise 209 At some time in the afternoon a 40-foot tall building casts an 80-foot long shadow.

Desmos link: <https://www.desmos.com/calculator/5qqd2t4eny>

151: Building Shadow 7

- (a) Use the worksheet above to approximate the scaling factor that converts a small change in the inclination angle of the sun (in radians) to an accurate approximation to the change in the length of the shadow. Include units. The tick marks on the protractor are spaced at equal intervals of 0.1 radian.
 - (b) Use calculus to find the exact scaling factor. Include units.
 - (c) Find the length of the shadow when the scaling factor that converts a small change in the sun's inclination to an accurate approximation in the shadow's length is -500 ft/radian.
-

The Tangent Function CW1

The tangent function and its inverse, classwork.

1 The Tangent Function

Question 210 Drag the sliders v (Line 2) and θ_1 (Line 4) in the worksheet below to make some measurements. Then use only arithmetic to approximate the value of the derivative

$$\frac{d}{d\theta} (\tan \theta) \Big|_{\theta=\theta_0}.$$

Explain your reasoning.

Desmos link: <https://www.desmos.com/calculator/omoajhcokg>

151: Tangent Derivative 2

Explanation. First drag the ruler to the position shown below to approximate the x -coordinate $x_A = PA$ of point B .

Desmos link: <https://www.desmos.com/calculator/hdox48kv6y>

151: Tangent Derivative 3

Using the scale it looks like

$$x_A \sim 33 \text{ cm.}$$

Continuing like this, point B has y -coordinate

$$y_A \sim 16 \text{ cm.}$$

Point C has y -coordinate

$$y_C \sim 22 \text{ cm.}$$

The distance $r = OQ$ is

$$r \sim 37 \text{ cm.}$$

And the distance $s = BQ$ is

$$s \sim 5 \text{ cm.}$$

Learning outcomes:
Author(s):

Because $\overline{BQ}$ is *perpendicular* to $\overline{PC}$ and $\Delta\theta \sim 0$, the length $s = BQ$ is approximately equal to the length of the circular arc between B and Q centered at P . So the approximate radian measure of angle $\Delta\theta$ is

$$\Delta\theta \sim \frac{s}{r} \sim \frac{5 \text{ cm}}{37 \text{ cm}} = \frac{5}{37}.$$

Also

$$\tan \theta_0 = \frac{y_A}{x_A} \sim \frac{16 \text{ cm}}{33 \text{ cm}} = \frac{16}{33}$$

and

$$\tan(\theta_0 + \Delta\theta) = \frac{y_C}{x_C} \sim \frac{22 \text{ cm}}{33 \text{ cm}} = \frac{22}{33}.$$

So

$$\begin{aligned} \frac{d}{d\theta} (\tan \theta) \Big|_{\theta=\theta_0} &\sim \frac{\tan(\theta_0 + \Delta\theta) - \tan \theta_0}{\Delta\theta} \\ &\sim \frac{37}{5} \cdot \frac{6}{33} \end{aligned}$$

Question 211 (a) Find an expression for the average rate of change of the function

$$f(\theta) = \tan \theta$$

between $\theta = \theta_1$ and $\theta = \theta_2$. Assume $\theta_1 \neq \theta_2$.

(b) Use part (a) and the algebra of limits to find an expression for the derivative

$$\frac{dy}{d\theta} \Big|_{\theta=\theta_1} = \frac{d}{d\theta} (\tan \theta) \Big|_{\theta=\theta_1}.$$

Hints:

(i) Remember that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1.$$

(ii) Use the identity

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

and modify the explanation to Example 6b in the chapter *Differentiating Trigonometric Functions* using *Limits* of our class notes.

(c) Repeat parts (a) and (b) for the function $f(\theta) = \cot \theta$.

(d) Instead, use the result of part (b) and the chain rule to find an expression for the derivative

$$\frac{d}{d\theta} (\cot \theta).$$

2 Exercises

Exercise 212 (a) Without using a calculator, find the y -coordinates of all points on the curve

$$y = f(\theta) = 10 - 4 \tan \theta$$

where the tangent line is parallel to the line

$$8x - y = 12.$$

Start by drawing a graph of the function $y = f(\theta)$. Then define an unknown like this:

Let

$$(\theta_0, f(\theta_0))$$

be the coordinates of the point on the curve $y = f(\theta)$ where the tangent line is parallel ...

- (b) Repeat part (a) to find the y -coordinates of points on the same curve where the tangent line is parallel to the line

$$x - 8y = 12$$

and then the line

$$8x + y = 12.$$

Explain what's going on and why. Include a graph.

Exercise 213 At some time in the afternoon a 40-foot tall building casts an 80-foot long shadow.

Desmos link: <https://www.desmos.com/calculator/5qqd2t4eny>

151: Building Shadow 7

- (a) Use the worksheet above to approximate the scaling factor that converts a small change in the inclination angle of the sun (in radians) to an accurate approximation of the change in the length of the shadow. Include units. The tick marks on the protractor are spaced at equal intervals of 0.1 radian. Explain your reasoning.
- (b) Find a function that expresses the length of the shadow in terms of the inclination angle of the sun during the afternoon. Use the tangent function. Include a domain.

Choose meaningful variables, not x and y . Start with a sentence that begins like this:

Let

$??? = f(???)$, write the domain here

be the function that expresses

- (c) Use calculus to find the exact scaling factor that converts a small change in the inclination angle (measured in radians) to an accurate approximation of the change in the shadow's length (measured in feet) when the shadow is 80 feet long.

End with a concluding sentence that states the scaling factor. Include units.

- (d) Find the length of the shadow when the scaling factor is -500 ft/radian. Start by defining an unknown. Do not use θ . It is already taken.

Circular Interpolation

Circular Interpolation

1 Three Problems

The following three questions might look dramatically different, yet they are essentially identical.

Question 214 *A mass at the end of a spring oscillates in simple harmonic motion on a horizontal surface as shown below. The mass is released from rest 2 meters from its equilibrium position and it completes one oscillation every 5 seconds.*

142: Simple Harmonic motion.

Desmos link: <https://www.desmos.com/calculator/l0bp34oig>

- (a) *Which way is the mass moving 1.5 seconds after it is released? What is its speed at this time?*
- (b) *Is the mass moving left or right the second time it is 1.6 meters from its equilibrium position? At what speed?*

Question 215 *The center of a ferris wheel with radius 80 feet is 90 feet above the ground. The wheel rotates at a constant rate, completing one revolution every 100 seconds.*

- (a) *Are you ascending or descending 30 seconds after boarding. At what rate?*
- (b) *Are you ascending or descending the second time you are 42 feet above the ground? At what rate?*

Learning outcomes:
Author(s):

Question 216 Over the course of a 24-hour period, from midnight October 29 to midnight October 30, the depth of the water at the Edmonds Pier is a sinusoidal function of time. Suppose also that a high tide of 21 feet occurs at 2:00am and the following low tide of 5 feet occurs at 8:00am.

- (a) At what rate is the depth of the water changing at 5:36am?
- (b) At what rate is the depth of the water is changing the second time the water is 8.2 feet deep?

What is usually called *sinusoidal modeling* is kind of a misnomer. First, it's usually easier to model oscillatory behavior with a cosine function. Second, and more importantly, the name *sinusoidal modeling* fails to convey the main idea of replacing linear interpolation with circular interpolation.

Here's an example.

Example 48. Suppose that over the course of a 24-hour period, from midnight October 29 to midnight October 30, the depth of the water at the Edmonds Pier is a sinusoidal function of time. Suppose also that a high tide of 21 feet occurs at 2:00am and the following low tide of 5 feet occurs at 8:00am.

Our question is to determine the rate at which the depth of the water is changing the *second* time the water is 10 feet deep. One approach is to find a function that expresses the depth in terms of time and then evaluate the derivative at the appropriate time. This solution is shown below, split into two parts.

But there is an easier way. We'll look at that after the long way.

Explanation. We first find a function

$$h = f(t), 0 \leq t \leq 24,$$

that expresses the depth of the water (in feet) in terms of the number of hours past midnight, October 29.

Note that to say "sinusoidal function" means that the graph of f is generated by uniform circular motion. But the graphs of the sine and cosine functions are both generated this way, so it is ok to express $f(t)$ in terms of the cosine function, and we will do just that.

Here are the steps.

- (a) First we'll use the information above to sketch by hand a graph of the function f . Label the axes with the appropriate variable names and units. Label the coordinates of two key points on the graph.

Desmos activity available at:

142: Edmonds Pier.

Desmos link: <https://www.desmos.com/calculator/x2kocpkcfm>

- (b) Next we'll compute the mean depth of the water over the 24 hours and the maximum deviation of the depth from this mean. Including units in our computation, the mean depth is

$$h_{avg} = 0.5(21 \text{ ft} + 5 \text{ ft}) = 13 \text{ ft}.$$

And the maximum deviation from the mean is

$$21 \text{ ft} - 13 \text{ ft} = 8 \text{ ft}.$$

Activate the Amplitude folder on Line 5 of the above demonstration to draw the horizontal line showing the average depth. Note this line is labeled with its equation. There is also a vertical line that shows the the maximum deviation from the mean.

- (c) Next we'll use the graph to find the period of oscillation. Since high tide occurs at 2:00am and low tide at 8:00am, the period (the time between successive) high (or low) tides is

$$2(8 \text{ hours} - 2 \text{ hours}) = 12 \text{ hours}$$

Activate the Period folder on Line 11 to show the period on the graph.

- (d) Now we'll compute the rotation rate of a uniform circular motion that generates the sinusoidal variation in the depth of the water. This rotation rate is

$$\omega = \frac{2\pi \text{ radians}}{12 \text{ hours}} = \frac{\pi}{6} \text{ radians/hr}.$$

Activate the Uniform Circular Motion folder on Line 18 to show the period on the graph.

- (e) Next we'll use parts (a)-(d) to find an expression for the function

$$h = f(t), 0 \leq t \leq 24,$$

that gives the depth of the water (in feet) at time t hours past midnight. Use the cosine function. Keeping in mind that high tide occurs at 2:00am, our function is

$$h = f(t) = 13 + 8 \cos\left(\frac{\pi}{6}(t - 2)\right), 0 \leq t \leq 24.$$

- (f) To check that our function is correct, we'll use the given information that the depth of the water is 21 feet at 2:00am and 5 feet at 8:00am.

Substituting $t = 2$ gives the depth at 2:00am (in feet) as

$$\begin{aligned} f(2) &= 13 + 8 \cos\left(\frac{\pi}{6}(2 - 2)\right) \\ &= 13 + 8 \cos(0) \\ &= 13 + 8 \\ &= 21. \end{aligned}$$

Substituting $t = 8$ gives the depth at 8:00am (in feet) as

$$\begin{aligned} f(8) &= 13 + 8 \cos\left(\frac{\pi}{6}(8 - 2)\right) \\ &= 13 + 8 \cos(\pi) \\ &= 13 + 8(-1) \\ &= 5. \end{aligned}$$

These check out.

- (g) Finally, we'll use our function to estimate the depth (in feet) of the water at 5:30am, October 29 to be

$$\begin{aligned} f(5.5) &= 13 + 8 \cos\left(\frac{\pi}{6}(5.5 - 2)\right) \\ &= 13 + 8 \cos(7\pi/12) \\ &\sim 10.92 \end{aligned}$$

- (h) Enter the coordinates of the appropriate point in Line 27 of the Desmos Activity above to check that the depth at 5:30am is reasonable.

Example 49. This is a continuation of the previous example, and we'll start with the function

$$h = f(t) = 13 + 8 \cos\left(\frac{\pi}{6}(t - 2)\right), 0 \leq t \leq 24,$$

that expresses the depth of the water (in feet) at the Edmonds Pier in terms of the number of hours past midnight.

Our question now is to find the rate at which the depth of the water is changing the *second* time the water is 10 feet deep.

One approach would be to first determine when the water is 10 feet deep for the second time. But this is not necessary. We can determine the rate without finding the time. Here's how.

- (a) We'll first take time out of the picture by letting

$$\theta = \frac{\pi}{6}(t - 2)$$

Then we'll let t_0 be the second time when the water is 10 feet deep and define the angle θ_0 as

$$\theta_0 = \frac{\pi}{6} (t_0 - 2)$$

- (b) Our aim is to evaluate the derivative

$$\begin{aligned} \left. \frac{dh}{dt} \right|_{t=t_0} &= -\frac{4\pi}{3} \sin \left(\frac{\pi}{6} (t_0 - 2) \right) \\ &= -\frac{4\pi}{3} \sin(\theta_0), \end{aligned}$$

where I've left the computation of the derivative dh/dt to you.

Note the key point. To evaluate this derivative we need *not* know the value of θ_0 , but only the value of $\sin \theta_0$.

- (c) To determine the value of $\sin \theta_0$, we go back to the height function, which we now write in the form (since $\theta = \pi(t - 2)/6$)

$$h = g(\theta) = 13 + 8 \cos \theta.$$

When the water is 10 feet deep for the second time,

$$10 = 13 + 8 \cos \theta_0$$

and

$$\cos \theta_0 = -3/8.$$

- (d) Now since

$$\cos^2 \theta_0 + \sin^2 \theta_0 = 1,$$

we find that

$$\sin \theta_0 = \pm \frac{\sqrt{55}}{8}.$$

- (e) To choose the correct sign, we go back to the graph.

Desmos activity available at:

142: Edmonds Pier 2.

Desmos link: <https://www.desmos.com/calculator/fs9xy1lf0e>

Here we can see that at the point Q (with coordinates $(t_0, f(t_0))$,

$$8 < t_0 < 14.$$

This tells us that

$$\theta_0 = \frac{\pi}{6} (t_0 - 2)$$

is between

$$\frac{\pi}{6}(8-2) < \theta_0 < \frac{\pi}{6}(14-2)$$

or that

$$\pi < \theta_0 < 2\pi.$$

In other words, the angle θ_0 is in the third or fourth quadrant. This means that $\sin \theta_0 < 0$ so

$$\sin \theta_0 = -\frac{\sqrt{55}}{8}.$$

(f) Finally, we evaluate the derivative

$$\begin{aligned} \left. \frac{dh}{dt} \right|_{t=t_0} &= -\frac{4\pi}{3} \sin(\theta_0) \\ &= \left(-\frac{4\pi}{3} \right) \left(-\frac{\sqrt{55}}{8} \right) \\ &= \frac{\pi\sqrt{55}}{6} \end{aligned}$$

So when the water is 10 feet deep for the second time, the depth is increasing at the rate of

$$\frac{\pi\sqrt{55}}{6} \text{ ft/hr} \sim 3.88 \text{ ft/hr}.$$

2 A Quicker Approach: Circular Interpolation

Here's another approach to the same question.

Question 217 Suppose that over the course of a 24-hour period, from midnight October 29 to midnight October 30, the depth of the water at the Edmonds Pier is a sinusoidal function of time. Suppose also that a high tide of 21 feet occurs at 2:00am and the following low tide of 5 feet occurs at 8:00am. We still wish to determine exact rate at which the water's depth is changing the second time the water is 10 feet deep.

The key idea is that sinusoidal variation is generated by uniform circular motion. So we can replace our problem about tides by one about a ferris wheel.

- (a) Drag point B below to the appropriate position for the water to be 10 feet deep. Then use the circle to help approximate the rate at which the water's depth is changing at the above time.

- (b) Use a little trigonometry to find the exact rate.

Explain your reasoning and include screenshots to help with your explanations.

Free Response:

Desmos link: <https://www.desmos.com/calculator/0wxwmkzvky>

142: Circular Interpolation Tides

Review of Sine and Cosine

Some problems.

1 Derivatives

Question 218 Follow the method of Example 2 in the chapter *The Chain Rule* of these notes exactly to find expressions for each of the following derivatives.

(a) $\frac{d}{d\theta} (\sec^2 \theta)$

(b) $\sqrt{12 + 3 \sin \theta}$

(c) $\frac{d}{d\theta} (\cos^2 \theta + \sin^2 \theta)$

Question 219 Divide each side of the identity

$$\cos^2 \theta + \sin^2 \theta = 1$$

by $\cos^2 \theta$ to get

$$1 + \tan^2 \theta = \sec^2 \theta.$$

Now use the result of Question 1(a) and the chain rule to find an expression for the derivative

$$\frac{d}{d\theta} (\tan \theta).$$

2 Transformations

Question 220 (a) The point with coordinates (a, b) lies on the graph of the function $y = f(x)$. What are the coordinates of the corresponding point on the graph of the function

$$y = g(x) = f(x/2)?$$

Learning outcomes:
Author(s):

- (b) Express the derivative

$$\left. \frac{d}{dx} (g(x)) \right|_{x=2a} = \left. \frac{d}{dx} (f(x/2)) \right|_{x=2a}$$

in terms of the derivative

$$\left. \frac{d}{dx} (f(x)) \right|_{x=a}.$$

- (c) Compute the slopes of the tangent lines to the curves
- $y = \sin \theta$
- and
- $y = \sin(\theta/2)$
- at the origin. Drag the slider
- k
- in the worksheet below to visualize the transformation.

151: Tranform Sine.

Desmos link: <https://www.desmos.com/calculator/2tvvkyi1sj>

- (d) Which of the following curves share the same tangent line at the origin with the curve
- $y = \sin \theta$
- ?

Select All Correct Answers:

- (i) $y = 2 \sin \theta$
- (ii) $y = 2 \sin(2\theta)$
- (iii) $y = 2 \sin(\theta/2)$ ✓
- (iv) $y = \frac{1}{3} \sin(3\theta)$ ✓
- (v) $y = \frac{1}{3} \sin(\theta/3)$

3 Simple Harmonic Motion

Question 221 A relaxed spring has a length of 3 meters. A mass is attached to the right end of the relaxed spring and then pulled an additional 2 meters to the right. The mass is then released from rest and oscillates without friction on a horizontal surface about its equilibrium position ($x = 0$ meters) with a period of 5 seconds as shown below.

A consequence of Hooke's Law (that the force of the spring acting on the mass points toward equilibrium and has magnitude proportional to the distance of the mass from equilibrium) is that the mass oscillates in simple harmonic motion about its equilibrium position. This means that the motion of the mass is driven by uniform circular motion around the circle of radius 5 meters below.

151: Simple Harmonic motion 23.

Desmos link: <https://www.desmos.com/calculator/noxooak1au>

- (a) Find a function

$$s = f(t), t \geq 0,$$

that expresses the displacement of the mass from equilibrium in terms of the number of seconds since the mass was released.

- (b) Find a function

$$v = g(t), t \geq 0,$$

that gives the horizontal component of the velocity of the mass (in meters/sec) in terms of the number of seconds since release. Take this component to be negative when the mass is moving to the left, positive when moving to the right.

- (c) Use your function from part (b) to find the maximum speed of the mass.
- (d) Find an equation relating s and v . Use it to answer the following questions.
- Where is the mass when it is moving at $2/3$ of its maximum speed?
 - At what fraction of its maximum speed is the mass traveling when it is $3/4$ of the way from its equilibrium position to one of its extreme positions?

4 Ferris Wheels

Question 222 A ferris wheel rotates at a constant rate, making one revolution every 90 seconds. The wheel has a radius of 50 feet and its center is 60 feet above the ground.

You ride the wheel for one revolution and jump off.

- What is your speed as you ride the wheel?
- Find an equation relating your height h above the ground (in feet) and your rate of ascent r (in ft/sec) at any instant. Start by finding an expression for the height as a function of time.
- Use your equation from part (b) to determine your rate of ascent when you are 20 feet above the ground for the second time.

Question 223 The graph below shows your height (in feet) as a function of the number of seconds since you boarded a ferris wheel. The wheel does not change its sense of rotation, but its rate of rotation might vary during the time you make your first revolution.

151: Ferris Wheel 99.

Desmos link: <https://www.desmos.com/calculator/4y7ljahkwn>

- (a) Use the graph above to approximate your rate of ascent when you are 150 feet above the ground for the first time.
- (b) Use part (a) to approximate the wheel's rotation rate when you are 150 feet above the ground for the first time.

Hint: Start by expressing your height in terms of the wheel's angle of rotation since you boarded.

5 Flagpole and Track

Question 224 Let r and a be constants, with $a > r > 0$.

You run around a circular track of radius r feet, passing point A at time $t = 0$ seconds past noon (see the animation below where $r = 50$). A flagpole is located at point F , a feet from the track's center and $a - r$ feet from A as shown below (where $a = 80$).

151: Track and Flagpole.

Desmos link: <https://www.desmos.com/calculator/xymelsnabl>

Answer the following questions for general parameters a and r . Do not use the specific values in the worksheet.

- (a) Find a function

$$s = f(\theta), \theta \geq 0,$$

that expresses your distance to the flagpole (in feet) in terms of the angle $\theta = \angle AOP$.

- (b) Find an expression for derivative $ds/d\theta$.

- (c) With $r = 50$ and $a = 80$, evaluate the derivative

$$\left. \frac{ds}{d\theta} \right|_{\theta=\pi/2}.$$

Include units. Then interpret the meaning of this derivative in terms of small changes.

- (d) Now let's return to working with the general parameters a and r . Suppose you run around the track at the constant speed of v ft/sec.
- (i) Find an expression for the derivative ds/dt , where t is the number of seconds past noon.
 - (ii) Express the derivative ds/dt in terms of v and the angle $\phi = \angle OPF$ above as you run around the “top” half of the track.
 - (iii) Use the result of part (ii) to describe your position on the track when your distance from the flagpole is increasing at the greatest rate. What is this maximum rate?
 - (iv) How would your answer to part (iii) change if $0 < a < r$ (ie. if the flagpole were inside the track)?
-

Exponential Functions, Part 2

More on exponential functions and their derivatives.

QR Code

Desmos link: <https://www.desmos.com/calculator/emp4pukecm>

1 Introduction

Consider two models for the population of a bacteria colony. The first,

$$P = g(t) = 3 + \frac{1}{2}t, \quad 0 \leq t \leq 6,$$

is linear and expresses the population (in millions of bacteria) in terms of the number of hours past noon. The growth rate is easy to describe. The population increases at the constant rate of 0.5 million bacteria/hour. We know this from precalculus. We could also use calculus to compute the growth rate as

$$\frac{dP}{dt} = \frac{d}{dt} \left(3 + \frac{1}{2}t \right) = \frac{1}{2} \frac{\text{bac}}{\text{hr}}.$$

We could avoid thinking about an instantaneous growth rate by saying that the average growth rate over *any* time interval is 0.5 million bacteria/hour. This statement implies that the (instantaneous) growth rate is constant.

But saying that the population increases by 0.5 million bacteria every hour would not completely describe how the population grows. It does *not* imply a constant growth rate because it does not tell us anything about what happens to the population over, for example, a half-hour period. The function graphed below shows an example of a non-linear function that increases by 0.5 million over *every* one-hour interval. Drag the slider u in Line 1 to convince yourself of this.

Describe what you see in the animation and how this suggests the function increases by 0.5 million over *every* one-hour interval.

Free Response:

151: Not Constant Rate

Learning outcomes:
Author(s):

Desmos link: <https://www.desmos.com/calculator/ehtzdzckmy>

The second model,

$$P = f(t) = 3(2^t), 0 \leq t \leq 6,$$

is exponential. Like the first it expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon. How can we describe how this population grows?

We could say that the population doubles every hour. Equivalently, we could say that the population increases by 100% every hour. But these descriptions would be incomplete. They don't tell us, nor do they imply, anything about what happens to the population over a half-hour period for example. The function $P = h(t)$ graphed below shows an example of a non-exponential function that doubles every hour. Drag the slider u in Line 1 and keep your eye on Line 2 (where $h = 1$, and $P = f_7(t) = h(t)$) to convince yourself of this.

Question 225 Describe what you see in the animation and how this suggests the function doubles over every one-hour interval.

151: Not Constant Rate 3

Desmos link: <https://www.desmos.com/calculator/zapwlt6qju>

We might try describing what happens to the population over a shorter time interval, say Δt hours. Over the time interval from time t hours past noon to time $t + \Delta t$ hours past noon the population increases by

$$\begin{aligned} \Delta P &= f(t + \Delta t) - f(t) \\ &= 3(2^{t+\Delta t} - 2^t) \\ &= 3(2^t \cdot 2^{\Delta t} - 2^t) \\ &= 3(2^t)(2^{\Delta t} - 1) \end{aligned}$$

millions of bacteria. To get the relative change over this time interval, we divide this absolute change by the population at the start of the time interval. This gives the relative rate of change as

$$\begin{aligned} \frac{\Delta P}{P} &= \frac{f(t + \Delta t) - f(t)}{f(t)} \\ &= \frac{3(2^t)(2^{\Delta t} - 1)}{3(2^t)} \\ &= 2^{\Delta t} - 1. \end{aligned}$$

Question 226 What are the units of this rate of change?

For example, the relative rate of change in the population over a one hour period is

$$2^1 - 1 = 1 = 100\%,$$

and over a half-hour period is

$$\sqrt{2} - 1 \sim 0.414 \sim 41.4\%.$$

Question 227 Explain why this rate is less than 50%.

Exploration 228 The worksheet below shows the graph of the function $P = 3(2^t)$.

151: Exp Growth Subtangent

Desmos link: <https://www.desmos.com/calculator/zu0f4jnh4a>

- (a) Experiment with the worksheet by first dragging the slider u (another name for t) in Line 2. Watch how Lines 6, 8, 10 change, as well as how the length of segment AB changes. Describe what you see.

Notes:

- (i) Line 6

$$\frac{f(u+h)}{f(u)}$$

This is the growth factor over the interval $t = u$ to $t = u + h$ hours.

- (ii) Line 8:

$$\frac{1}{f(u)} \cdot \frac{f(u+h) - f(u)}{h}$$

This is the relative average rate of change over the interval $t = u$ to $t = u + h$ hours.

- (iii) Line 10:

$$f(u) \cdot \frac{h}{f(u+h) - f(u)}$$

This is the reciprocal of Line 8.

- (b) Now Drag $h = \Delta t$ in Line 4 to $h = 0.5$ and repeat part (a). Describe what you see.

- (c) Drag h closer to zero and describe how Lines 6, 8, and 10 change. Include units for these three expressions.
- (d) What are the units of Line 10? What is its meaning?
- (e) Summarize your observations.

2 Relative Growth Rates

We still have not answered the question about how to describe exponential growth *without* a formula. What we know so far is that over time intervals of fixed duration an exponential function increases by a fixed percentage. And Exploration 4 might have suggested that the relative instantaneous growth rate

$$\frac{1}{P} \cdot \frac{dP}{dt}$$

is constant for an exponential function. We can show this is true for our function $f(t) = 3(2^t)$ using our prior work where we found that

$$\frac{\Delta P}{P} = 2^{\Delta t} - 1.$$

So the relative instantaneous growth rate of the function $f(t) = 3(2^t)$ is

$$\begin{aligned} \frac{1}{P} \cdot \frac{dP}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{1}{P} \cdot \frac{\Delta P}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \cdot \frac{\Delta P}{P} \\ &= \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 1}{\Delta t}. \end{aligned}$$

Now we have no way to evaluate this limit algebraically (at least as of yet), but we can recognize it as the derivative

$$\left. \frac{d}{dt} (2^t) \right|_{t=0} = \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 2^0}{\Delta t}.$$

From Exploration 4 we know the approximate value of this limit, and hence the relative rate of change of our population function $P = f(t) = 3(2^t)$. It is

$$\frac{1}{P} \cdot \frac{dP}{dt} \sim 0.693 = 69.3\%/hr.$$

So our population increases at the constant relative rate of approximately 69.3%/hr. The exact relative rate turns out to be

$$\frac{1}{P} \cdot \frac{dP}{dt} = \ln 2/\text{hr}.$$

But to understand why, we first need to come to terms with the number $e \sim 2.71828$.

3 The Number e

For our purposes, the best definition of the number e is this: It is the number for which the exponential function (t in hours)

$$P = g(t) = e^t$$

increases at the constant relative rate of 100%/hr. Put another way, the number e is the one-hour growth factor for a population that increases at the relative rate of 100%/hr. But to make sense of this, we need to understand what it means to say that at any instant a population is increasing at the relative rate of 100%/hr.

It does *not* mean that the population increases 100% every hour (the function $f(t) = P_0(2^t)$ does that). Rather, like every derivative the meaning is best understood in terms of small changes.

For example, in $1/100$ of an hour the population would increase by about

$$\left(\frac{1}{100}\right)(100\%) = 1\%.$$

So the $1/100$ -hour growth factor is about 1.01. And the one-hour growth factor is about

$$\left(1 + \frac{1}{100}\right)^{100} \sim 2.705.$$

This tells us that $e \sim 2.705$ and that every hour a population increasing at the relative instantaneous rate of 100%/hr increases by about 170.5%.

We can get a better approximation to e by taking a smaller time interval. For example, a better approximation is

$$e \sim \left(1 + \frac{1}{1000}\right)^{1000} \sim 2.717.$$

And as you might have guessed,

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \sim 2.718.$$

4 An Example

When we learned about derivatives of inverse functions, we saw that it was helpful to express the derivative of a one-to-one function in terms of its *output*. For the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

for example,

$$\frac{dy}{d\theta} = \cos \theta = \sqrt{1 - y^2}.$$

When working with exponential functions, or simple transformations of these, it is almost always best to do the same. Here's an example.

Example 50. The function

$$P = 3e^{\frac{1}{5}t}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

Find the population when it is increasing at the rate of 2 million bacteria/hour.

Explanation. There's a hard way and an easy way to solve this problem. The hard way would be to first find the time when the population is increasing at the given rate. Then we could use this time to find the population.

The easy way is to ignore time altogether and express the growth rate of the population in terms of the population.

Before we do this, let's think first about the units of the factor $1/5$ in the exponent. Since any exponential function takes only dimensionless inputs and t is measured in hours, the units of $1/5$ are

Question 229 Multiple Choice:

- (a) *hours*
- (b) *dimensionless*
- (c) *millions of bacteria*
- (d) hours^{-1} ✓

The units of $1/5$ give a clue to its meaning. It is in fact the relative growth rate of the population. So at any instant this population is increasing at the relative rate of

Question 230 20%/hour .

Question 231 (a) Use the chain rule to compute the absolute growth rate

$$\frac{dP}{dt} = \frac{d}{dt} \left(3e^{\frac{1}{5}t} \right).$$

Make the explicit u -substitution. Do not skip any steps.

(b) Use part (a) to verify that the relative growth rate

$$\frac{1}{P} \cdot \frac{dP}{dt}$$

is 20%/hour.

You should have found the relative growth rate to be

$$\frac{1}{P} \cdot \frac{dP}{dt} = \frac{1}{5} \text{ hr}^{-1},$$

so that the absolute growth rate (in millions of bacteria/hour) is

$$\frac{dP}{dt} = \frac{1}{5}P.$$

Question 232 Use this to find the population when it is increasing at the rate of 2 million bacteria/hour.

5 Exercises

Question 233 The function

$$P = f(t) = 50e^{\frac{t}{10}}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past midnight. Graph shown below.

Desmos link: <https://www.desmos.com/calculator/kffjsoepmo>

151: Exp Function 5

- (a) Use the graph of the function $P = f(t)$ above to approximate the growth rate when there are 200 million bacteria.
- (b) Find a function $r = g(P)$ that expresses the growth rate (measured in millions of bacteria/hour) in terms of the population (measured in millions). Include a domain.
- (c) Find the exact growth rate when there are 200 million bacteria. No calculator.
- (d) What is the population when it is increasing at the rate of 8 million bacteria/hour?
- (e) Use the ideas of this question to describe how the population grows. Hint: Think about the relative instantaneous growth rate.

Question 234 Between noon and 6pm, a population of bacteria grows exponentially, increasing by 20% every hour. There are 5 million bacteria at noon.

- (a) Find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon.
- (b) Find a function that expresses the growth rate (in millions of bacteria/hour) in terms of the population (in millions).
- (c) Find the population when it is increasing at the rate of 2 million bacteria/hour.
- (d) What is the relative growth rate of the population?
- (e) Consider a second population that grow exponentially, increasing by 20% every four hours.
 - (i) Use common sense to find its relative growth rate.
 - (ii) Use calculus to verify your answer from part (i) with a computation.

Question 235 The function

$$N = f(t) = \frac{20}{1 + 5e^{-t/5}}, \quad 0 \leq t \leq 30,$$

models the spread of a virus throughout a population. It takes as an input the number of months since January 1st and returns as an output the number of infected individuals (measured in millions). Graph shown below.

Desmos link: <https://www.desmos.com/calculator/ikis451wxu>

151: Logistic Model

- (a) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at the rate of 800,000 people/month.
- (b) Find a function
- $$r = g(N)$$
- that expresses the infection rate (in millions of people/month) in terms of the population.
- (c) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at the rate of 750,000 people/month.
- (d) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at its fastest rate. Also approximate this rate.
- (e) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at its fastest rate. Then find this exact rate.

Question 236 Use the chain rule and your knowledge of the derivative $d(e^x)/dx$ to compute each of the following derivatives.

(a)

$$\frac{d}{dt}(100 \cdot 2^t)$$

(b)

$$\frac{d}{dt}(100 \cdot 2^{t/5})$$

Explanation. (a) The key is to realize that

$$2 = e^{\ln 2}.$$

So we can write the function to differentiate as

$$P = 100 \cdot 2^t = 100 (e^{\ln 2})^t = 100e^{t \ln 2}.$$

Now we can make a u -substitution and use the chain rule. We let

$$u = t \ln 2.$$

Then

$$P = 100e^{t \ln 2} = 100e^u$$

and

$$\begin{aligned} \frac{dP}{dt} &= \frac{dP}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du}(100e^u) \cdot \frac{d}{dt}(t \ln 2) \\ &= 100e^u(\ln 2) \\ &= P \ln 2 \\ &= (100 \cdot 2^t) \ln 2 \\ &= (100 \ln 2)(2^t) \end{aligned}$$

Question 237 A colony of bacteria has a population of 20 million at noon and a population of 25 million at 2pm. The population grows exponentially between noon and midnight.

- (a) Describe precisely how the population grows.
- (b) Use your description from part (a) to find a function

$$P = f(t), \quad 0 \leq t \leq 12,$$

that expresses the population (in millions) in terms of the number of hours past noon. Do not use e .

- (c) Find the relative instantaneous growth rate of the population.
- (d) Find a function $r = g(P)$ that expresses the (absolute) growth rate in terms of the population.
- (e) At what rate is the population changing when there are 30 million bacteria?

Explanation. (a) The population increases by

$$\frac{25 \text{ million} - 20 \text{ million}}{20 \text{ million}} = 25\%$$

every two hours.

Better yet, at least for determining the function, is to say that the two-hour growth factor is

$$\frac{25 \text{ million}}{20 \text{ million}} = 1.25.$$

This means that every two hours the population gets multiplied by 1.25.

- (b) Since the two-hour growth factor is 1.25, the one-hour growth factor is $1.25^{1/2}$. So

$$P = 20(1.25^{1/2})^t = 20(1.25)^{\frac{t}{2}}, \quad 0 \leq t \leq 12.$$

- (c) The key to computing the derivative dP/dt is to write the population function using base e . Since

$$1.25 = e^{\ln 1.25},$$

$$\begin{aligned} P &= 20(1.25)^{\frac{t}{2}} \\ &= 20(e^{\ln 1.25})^{\frac{t}{2}} \\ &= 20e^{t(\ln 1.25)/2} \end{aligned}$$

So the growth rate (in millions of bacteria/hour) is

$$\begin{aligned} \frac{dP}{dt} &= \frac{d}{dt} \left(20(1.25)^{\frac{t}{2}} \right) \\ &= 20 \frac{d}{dt} \left(e^{t(\ln 1.25)/2} \right) \\ &= \left(20e^{t(\ln 1.25)/2} \right) \frac{d}{dt} (t(\ln 1.25)/2) \\ &= P(\ln 1.25)/2. \end{aligned}$$

And the relative growth rate (units are hr^{-1}) is

$$\frac{1}{P} \cdot \frac{dP}{dt} = \frac{1}{2} \ln 1.25.$$

- (d) From part (c), the function

$$g(P) = \frac{dP}{dt} = P(\ln 1.25)/2$$

expresses the growth rate (in millions of bacteria/hour) in terms of the population.

- (e) When there are 30 million bacteria, the population is increasing at the rate of

$$g(30) = (30 \text{ million bacteria}) \left(\frac{1}{2} \ln 1.25/\text{hr} \right) = 15 \ln 1.25 \text{ million bacteria/hr.}$$

Question 238 Between ground level and an altitude of 10 km, atmospheric pressure on the planet Krypton is an exponential function of altitude. The pressure is 100 kPa at ground level and 80 kPa at an altitude of 0.75 km.

Desmos link: <https://www.desmos.com/calculator/j5h8kaj8xs>

151: Atmospheric Pressure

- (a) Describe exactly how the pressure decreases.
- (b) Use your description from part (a) to find a function

$$P = f(h), 0 \leq h \leq 10,$$

that expresses the atmospheric pressure (in kPa) in terms of the altitude (in kilometers). Do not use the number e .

- (c) Input your expression for f in Line 2 of the worksheet above.
- (d) Evaluate the derivative

$$\left. \frac{dP}{dh} \right|_{h=5}$$

and interpret its meaning.

- (e) Find a function $r = g(P)$ that expresses the derivative dP/dh in terms of P .
- (f) Use calculus to approximate the altitude at which the pressure is 0.2 kPa less at an altitude 0.1 km higher.

6 Conceptual Questions

Question 239 A population of bacteria grows exponentially. If the growth rate is 4 million bacteria/hour when the population is 20 million, determine the growth rate when the population is 30 million.

Question 240 A cup of coffee at a temperature of 90°C is brought into a room held at a constant temperature of 20°C . Suppose that the difference in temperature between the coffee and the room decreases exponentially. Suppose also that the temperature of the coffee initially decreases at the rate of $5^\circ\text{C}/\text{min}$. What is the temperature of the coffee when its temperature is decreasing at the rate of $2^\circ\text{C}/\text{min}$?

Exponential Functions, Short Version

More on exponential functions and their derivatives.

1 Introduction

To claim a population that increases by 0.5 million bacteria every hour grows at a constant rate would be a falsehood. The reason is the statement does not imply anything about what the population does over a shorter period. The graph below shows an example of a non-linear function that increases by 0.5 million over *every* one-hour interval. Drag the slider u in Line 1 to convince yourself of this.

151: Not Constant Rate

Desmos link: <https://www.desmos.com/calculator/ehtzdzckmy>

We would make a similar error in asserting a population that doubles every hour grows exponentially. The function $P = h(t)$ graphed below shows an example of a non-exponential function that doubles every hour. Drag the slider u in Line 1 and keep your eye on Line 2 (where $h = 1$, and $P = f_7(t) = h(t)$) to convince yourself of this.

151: Not Constant Rate 3

Desmos link: <https://www.desmos.com/calculator/zapwlt6qju>

We can describe linear growth by saying a population grows at a constant rate of 0.5 million bacteria/hour. Or we could avoid the idea of an instantaneous growth rate by saying that the population increases at the average rate of 0.5 million bacteria/hour over *every* time interval.

But to correct the second error and describe exponential growth, we are almost forced into thinking about the instantaneous growth rate. What characterizes exponential growth is that this rate is proportional to the population. So for example, when the population is eight times the current population, its growth rate would be eight times the current growth rate.

To understand why in a specific example, consider the graph of the function

$$P = f(t) = P_0 \cdot 2^t$$

Learning outcomes:
Author(s):

shown below.

151: Exp Growth Translate Dilate

Desmos link: <https://www.desmos.com/calculator/dmdpvei3d6>

The key point is that because

$$f(t - 3) = 2^{t-3} = (2^t)(2^{-3}) = \frac{1}{8}f(t),$$

$$8f(t - 3) = f(t).$$

Geometrically, this means that translating the graph of f three hours to the right (Drag the slider u in Line 2 above from $u = 0$ to $u = 3$) and then multiplying the outputs by 8 (drag the slider v in Line 4 from $v = 1$ to $v = 8$) sends the graph of $P = f(t)$ to itself and the point $A(0, f(0)) = A(0, P_0)$ to the point $A''(3, f(3)) = A''(3, 8P_0)$. The translation leaves the slope of the tangent line at $(0, P_0)$ unchanged and the dilation multiplies the slope by 8. So

$$\left. \frac{dP}{dt} \right|_{t=3} = 8 \left(\left. \frac{dP}{dt} \right|_{t=0} \right).$$

The same kind of reasoning shows that for any for any exponential function, the growth rate is proportional to the population. Here's a more computational proof of this fact for the function $f(t) = P_0 2^t$.

Over the time interval from time t hours past noon to time $t + \Delta t$ hours past noon the population increases by

$$\begin{aligned} \Delta P &= f(t + \Delta t) - f(t) \\ &= 3(2^{t+\Delta t} - 2^t) \\ &= 3(2^t \cdot 2^{\Delta t} - 2^t) \\ &= 3(2^t)(2^{\Delta t} - 1) \\ &= P(2^{\Delta t} - 1). \end{aligned}$$

millions of bacteria. Here $P = f(t)$ is the population at the start of the time interval.

So over the period from t hours past noon to $t + \Delta t$ hours past noon the average growth rate is

$$\frac{\Delta P}{\Delta t} = P \left(\frac{2^{\Delta t} - 1}{\Delta t} \right),$$

measured in millions of bacteria/hour.

Then at time t hours past noon, the instantaneous growth rate (in millions of bacteria/hour) is

$$\begin{aligned}\frac{dP}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{\Delta P}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} P \left(\frac{2^{\Delta t} - 1}{\Delta t} \right) \\ &= P \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 1}{\Delta t}.\end{aligned}$$

This shows what we wanted, that at any instant the growth rate is proportional to the population. Expressed differently, this says that the relative growth rate (measured in hr^{-1}) is constant and equal to

$$\frac{1}{P} \cdot \frac{dP}{dt} = \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 1}{\Delta t}.$$

The best we can do here is to approximate the above limit numerically, to get the relative growth rate as

$$\frac{1}{P} \cdot \frac{dP}{dt} \sim 0.6931 \text{ hr}^{-1} \sim 69.31\%/\text{hr}.$$

Key Point: Over a one-hour period, the population doubles. So over a one-hour period the population increases at a relative average rate of 100%/hour. This is greater than the relative instantaneous growth rate of 69.31%/hr because of compounding.

2 Relative Growth Rates

We still have not answered the question about how to describe exponential growth *without* a formula. What we know so far is that over time intervals of fixed duration an exponential function increases by a fixed percentage. And Exploration 4 might have suggested that the relative instantaneous growth rate

$$\frac{1}{P} \cdot \frac{dP}{dt}$$

is constant for an exponential function. We can show this is true for our function $f(t) = 3(2^t)$ using our prior work where we found that

$$\frac{\Delta P}{P} = 2^{\Delta t} - 1.$$

So the relative instantaneous growth rate of the function $f(t) = 3(2^t)$ is

$$\begin{aligned}\frac{1}{P} \cdot \frac{dP}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{1}{P} \cdot \frac{\Delta P}{\Delta t} \\ &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \cdot \frac{\Delta P}{P} \\ &= \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 1}{\Delta t}.\end{aligned}$$

Now we have no way to evaluate this limit algebraically (at least as of yet), but we can recognize it as the derivative

$$\left. \frac{d}{dt} (2^t) \right|_{t=0} = \lim_{\Delta t \rightarrow 0} \frac{2^{\Delta t} - 2^0}{\Delta t}.$$

From Exploration 4 we know the approximate value of this limit, and hence the relative rate of change of our population function $P = f(t) = 3(2^t)$. It is

$$\frac{1}{P} \cdot \frac{dP}{dt} \sim 0.693 = 69.3\%/\text{hr}.$$

So our population increases at the constant relative rate of approximately 69.3%/hr. The exact relative rate turns out to be

$$\frac{1}{P} \cdot \frac{dP}{dt} = \ln 2/\text{hr}.$$

But to understand why, we first need to come to terms with the number $e \sim 2.71828$.

3 The Number e

For our purposes, the best definition of the number e is this: It is the number for which the exponential function (t in hours)

$$P = g(t) = e^t$$

increases at the constant relative rate of 100%/hr. Put another way, the number e is the one-hour growth factor for a population that increases at the relative rate of 100%/hr. But to make sense of this, we need to understand what it means to say that at any instant a population is increasing at the relative rate of 100%/hr.

It does *not* mean that the population increases 100% every hour (the function $f(t) = P_0(2^t)$ does that). Rather, like every derivative the meaning is best understood in terms of small changes.

For example, in $1/100$ of an hour the population would increase by about

$$\left(\frac{1}{100}\right)(100\%) = 1\%.$$

So the $1/100$ -hour growth factor is about 1.01. And the one-hour growth factor is about

$$\left(1 + \frac{1}{100}\right)^{100} \sim 2.705.$$

This tells us that $e \sim 2.705$ and that every hour a population increasing at the relative instantaneous rate of 100%/hr increases by about 170.5%.

We can get a better approximation to e by taking a smaller time interval. For example, a better approximation is

$$e \sim \left(1 + \frac{1}{1000}\right)^{1000} \sim 2.717.$$

And as you might have guessed,

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \sim 2.718.$$

4 An Example

When we learned about derivatives of inverse functions, we saw that it was helpful to express the derivative of a one-to-one function in terms of its *output*. For the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

for example,

$$\frac{dy}{d\theta} = \cos \theta = \sqrt{1 - y^2}.$$

When working with exponential functions, or simple transformations of these, it is almost always best to do the same. Here's an example.

Example 51. The function

$$P = 3e^{\frac{1}{5}t}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

Find the population when it is increasing at the rate of 2 million bacteria/hour.

Explanation. There's a hard way and an easy way to solve this problem. The hard way would be to first find the time when the population is increasing at the given rate. Then we could use this time to find the population.

The easy way is to ignore time altogether and express the growth rate of the population in terms of the population.

Before we do this, let's think first about the units of the factor $1/5$ in the exponent. Since any exponential function takes only dimensionless inputs and t is measured in hours, the units of $1/5$ are

Question 241 Multiple Choice:

- (a) *hours*
- (b) *dimensionless*
- (c) *millions of bacteria*
- (d) hours^{-1} ✓

The units of $1/5$ give a clue to its meaning. It is in fact the relative growth rate of the population. So at any instant this population is increasing at the relative rate of

Question 242 $20\%/\text{hour}$.

Question 243 (a) *Use the chain rule to compute the absolute growth rate*

$$\frac{dP}{dt} = \frac{d}{dt} \left(3e^{\frac{1}{5}t} \right).$$

Make the explicit u -substitution. Do not skip any steps.

(b) *Use part (a) to verify that the relative growth rate*

$$\frac{1}{P} \cdot \frac{dP}{dt}$$

is $20\%/\text{hour}$.

You should have found the relative growth rate to be

$$\frac{1}{P} \cdot \frac{dP}{dt} = \frac{1}{5} \text{ hr}^{-1},$$

so that the absolute growth rate (in millions of bacteria/hour) is

$$\frac{dP}{dt} = \frac{1}{5}P.$$

Question 244 Use this to find the population when it is increasing at the rate of 2 million bacteria/hour.

5 Exercises

Question 245 The function

$$P = f(t) = 50e^{\frac{t}{10}}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past midnight. Graph shown below.

Desmos link: <https://www.desmos.com/calculator/kffjsoepmo>

151: Exp Function 5

- Use the graph of the function $P = f(t)$ above to approximate the growth rate when there are 200 million bacteria.
- Find a function $r = g(P)$ that expresses the growth rate (measured in millions of bacteria/hour) in terms of the population (measured in millions). Include a domain.
- Find the exact growth rate when there are 200 million bacteria. No calculator.
- What is the population when it is increasing at the rate of 8 million bacteria/hour?
- Use the ideas of this question to describe how the population grows. Hint: Think about the relative instantaneous growth rate.

Question 246 Between noon and 6pm, a population of bacteria grows exponentially, increasing by 20% every hour. There are 5 million bacteria at noon.

- Find a function that expresses the population (in millions of bacteria) in terms of the number of hours past noon.

- (b) Find a function that expresses the growth rate (in millions of bacteria/hour) in terms of the population (in millions).
- (c) Find the population when it is increasing at the rate of 2 million bacteria/hour.
- (d) What is the relative growth rate of the population?
- (e) Consider a second population that grows exponentially, increasing by 20% every four hours.
 - (i) Use common sense to find its relative growth rate.
 - (ii) Use calculus to verify your answer from part (i) with a computation.

Question 247 The function

$$N = f(t) = \frac{20}{1 + 5e^{-t/5}}, \quad 0 \leq t \leq 30,$$

models the spread of a virus throughout a population. It takes as an input the number of months since January 1st and returns as an output the number of infected individuals (measured in millions). Graph shown below.

Desmos link: <https://www.desmos.com/calculator/ikis451wxu>

151: Logistic Model

- (a) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at the rate of 800,000 people/month.
- (b) Find a function

$$r = g(N)$$
 that expresses the infection rate (in millions of people/month) in terms of the population.
- (c) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at the rate of 750,000 people/month.
- (d) Use the sliders in the worksheet above to approximate the number of infected individuals when the virus is spreading at its fastest rate. Also approximate this rate.
- (e) Use calculus and algebra to determine the number of infected individuals when the virus is spreading at its fastest rate. Then find this exact rate.

Question 248 Use the chain rule and your knowledge of the derivative $d(e^x)/dx$ to compute each of the following derivatives.

(a)

$$\frac{d}{dt}(100 \cdot 2^t)$$

(b)

$$\frac{d}{dt}(100 \cdot 2^{t/5})$$

Explanation. (a) The key is to realize that

$$2 = e^{\ln 2}.$$

So we can write the function to differentiate as

$$P = 100 \cdot 2^t = 100 (e^{\ln 2})^t = 100e^{t \ln 2}.$$

Now we can make a u -substitution and use the chain rule. We let

$$u = t \ln 2.$$

Then

$$P = 100e^{t \ln 2} = 100e^u$$

and

$$\begin{aligned} \frac{dP}{dt} &= \frac{dP}{du} \cdot \frac{du}{dt} \\ &= \frac{d}{du}(100e^u) \cdot \frac{d}{dt}(t \ln 2) \\ &= 100e^u(\ln 2) \\ &= P \ln 2 \\ &= (100 \cdot 2^t) \ln 2 \\ &= (100 \ln 2)(2^t) \end{aligned}$$

Question 249 A colony of bacteria has a population of 20 million at noon and a population of 25 million at 2pm. The population grows exponentially between noon and midnight.

(a) Describe precisely how the population grows.

- (b) Use your description from part (a) to find a function

$$P = f(t), 0 \leq t \leq 12,$$

that expresses the population (in millions) in terms of the number of hours past noon. Do not use e .

- (c) Find the relative instantaneous growth rate of the population.
 (d) Find a function $r = g(P)$ that expresses the (absolute) growth rate in terms of the population.
 (e) At what rate is the population changing when there are 30 million bacteria?

Explanation. (a) The population increases by

$$\frac{25 \text{ million} - 20 \text{ million}}{20 \text{ million}} = 25\%$$

every two hours.

Better yet, at least for determining the function, is to say that the two-hour growth factor is

$$\frac{25 \text{ million}}{20 \text{ million}} = 1.25.$$

This means that every two hours the population gets multiplied by 1.25.

- (b) Since the two-hour growth factor is 1.25, the one-hour growth factor is $1.25^{1/2}$. So

$$P = 20(1.25^{1/2})^t = 20(1.25)^{\frac{t}{2}}, 0 \leq t \leq 12.$$

- (c) The key to computing the derivative dP/dt is to write the population function using base e . Since

$$1.25 = e^{\ln 1.25},$$

$$\begin{aligned} P &= 20(1.25)^{\frac{t}{2}} \\ &= 20(e^{\ln 1.25})^{\frac{t}{2}} \\ &= 20e^{t(\ln 1.25)/2} \end{aligned}$$

So the growth rate (in millions of bacteria/hour) is

$$\begin{aligned} \frac{dP}{dt} &= \frac{d}{dt} \left(20(1.25)^{\frac{t}{2}} \right) \\ &= 20 \frac{d}{dt} \left(e^{t(\ln 1.25)/2} \right) \\ &= \left(20e^{t(\ln 1.25)/2} \right) \frac{d}{dt} (t(\ln 1.25)/2) \\ &= P(\ln 1.25)/2. \end{aligned}$$

And the relative growth rate (units are hr^{-1}) is

$$\frac{1}{P} \cdot \frac{dP}{dt} = \frac{1}{2} \ln 1.25.$$

- (d) From part (c), the function

$$g(P) = \frac{dP}{dt} = P(\ln 1.25)/2$$

expresses the growth rate (in millions of bacteria/hour) in terms of the population.

- (e) When there are 30 million bacteria, the population is increasing at the rate of

$$g(30) = (30 \text{ million bacteria}) \left(\frac{1}{2} \ln 1.25/\text{hr} \right) = 15 \ln 1.25 \text{ million bacteria/hr.}$$

Question 250 Between ground level and an altitude of 10 km, atmospheric pressure on the planet Krypton is an exponential function of altitude. The pressure is 100 kPa at ground level and 80 kPa at an altitude of 0.75 km.

Desmos link: <https://www.desmos.com/calculator/j5h8kaj8xs>

151: Atmospheric Pressure

- (a) Describe exactly how the pressure decreases.
 (b) Use your description from part (a) to find a function

$$P = f(h), \quad 0 \leq h \leq 10,$$

that expresses the atmospheric pressure (in kPa) in terms of the altitude (in kilometers). Do not use the number e .

- (c) Input your expression for f in Line 2 of the worksheet above.
 (d) Evaluate the derivative

$$\left. \frac{dP}{dh} \right|_{h=5}$$

and interpret its meaning.

- (e) Find a function $r = g(P)$ that expresses the derivative dP/dh in terms of P .
 (f) Use calculus to approximate the altitude at which the pressure is 0.2 kPa less at an altitude 0.1 km higher.

6 Conceptual Questions

Question 251 A population of bacteria grows exponentially. If the growth rate is 4 million bacteria/hour when the population is 20 million, determine the growth rate when the population is 30 million.

Question 252 A cup of coffee at a temperature of 90°C is brought into a room held at a constant temperature of 20°C . Suppose that the difference in temperature between the coffee and the room decreases exponentially. Suppose also that the temperature of the coffee initially decreases at the rate of $5^{\circ}\text{C}/\text{min}$. What is the temperature of the coffee when its temperature is decreasing at the rate of $2^{\circ}\text{C}/\text{min}$?

Practice Quiz 2

Practice quiz, Weeks 4-6

Directions:

- (a) Show all work.
- (b) Give brief explanations for each problem. Include these explanations in the flow of the solution.
- (c) Show all units in all computations.
- (d) No calculators.
- (e) Show each step when using the chain, product, and quotient rules and use the Leibniz notation when doing so. Here is an example.

Example 52. Find an expression for the derivative

$$\frac{d}{d\theta} (4 + 5 \cos^3(2\theta)) .$$

Explanation. With $y = 4 + 5 \cos^3(2\theta)$, the derivative $dy/d\theta$ is

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{d}{d\theta} (4 + 5 \cos^3(2\theta)) \\ &= \frac{d}{d\theta} (4) + 5 \frac{d}{d\theta} ((\cos(2\theta))^3) \\ &= 5(3)(\cos(2\theta))^2 \frac{d}{d\theta} (\cos(2\theta)) \\ &= 15(\cos(2\theta))^2 (-\sin(2\theta)) \frac{d}{d\theta} (2\theta) \\ &= -30(\cos(2\theta))^2 \sin(2\theta). \end{aligned}$$

Question 253 Find expressions for each of the following derivatives.

(a)

$$\frac{d}{dt} (4 - t^4 e^{-5t})$$

(b)

$$\frac{d}{d\theta} \left(\frac{12}{4 + 5 \tan(4\theta)} \right)$$

Learning outcomes:
Author(s):

(c)

$$\frac{d}{dw} \left(\sqrt{40 - 5w^2} \right)$$

(d)

$$\frac{d}{dt} \left(2^{t/8} \right)$$

Question 254 (a) Find an expression for the derivative

$$\frac{d}{dx} (\sin(\arcsin x)).$$

Do this directly using the chain rule. Do not simplify the function first.

- (b) Find an expression for the same derivative by first simplifying the function $y = \sin(\arcsin x)$ and then using the chain rule.
- (c) Find an expression for the same derivative by first simplifying the function $y = \sin(\arcsin x)$ and then using geometry.

Question 255 (a) Find an equation of the tangent line to the curve

$$y = 4 - t^4 e^{-5t}$$

at the point on the curve with t -coordinate $t = 0$.

- (b) Find an equation of the tangent line to the curve

$$y = \frac{12}{4 + 5 \tan(4\theta)}$$

at the point on the curve with θ -coordinate $\theta = 0$.

Question 256 Assume for this question that each month has 30 days and that the number of hours of daylight/day in Seattle is a sinusoidal function of time. Assume also that on June 21, Seattle gets a maximum of 16 hours of daylight/day and that on December 21, Seattle gets a minimum of 8 hours of daylight/day.

- (a) Find a function

$$H = f(t), \quad 0 \leq t \leq 12,$$

that expresses the number of daylight hours/day in Seattle in terms of the number of months since June 21. Use the cosine function. Start by sketching a graph. Explain your reasoning.

- (b) Evaluate the derivative

$$\left. \frac{dH}{dt} \right|_{t=2}.$$

- (c) What are the units of the derivative above?
- (d) Explain the meaning of the above derivative.

Question 257 The function

$$T = f(m) = 20 + 70e^{-m/20}, \quad 0 \leq m \leq 60,$$

expresses the temperature (in Celsius degrees) of a cup of coffee in terms of the number of minutes past noon.

- (a) At what rate is the temperature changing at 12:40pm?
- (b) Does the temperature change at a constant relative rate? Justify your assertion.
- (c) Find a function $r = g(T)$ that expresses the rate of change in the temperature (measured in $^{\circ}\text{C}/\text{min}$) in terms of the temperature (measured in Celsius degrees).
- (d) Find the temperature when it is decreasing at the rate of $3^{\circ}\text{C}/\text{min}$.
- (e) Find a function $s = h(T)$ that expresses the relative rate of change in the temperature in terms of the temperature (measured in Celsius degrees).
- (f) Find the temperature when it is decreasing at the rate of $3\%/\text{min}$.

Question 258 Assume for this problem that over the course of a 24-hour period beginning at midnight of July 21, the temperature in Shoreline is a sinusoidal function of time. Assume also that the temperature reaches its minimum of 50°F at 5am and its maximum of 80° at 5pm.

At what rate (with respect to time) is the temperature changing when it is 68°F for the second time? Start by finding a function that expresses the temperature in terms of time. Be sure to define your variables.

Question 259 The function

$$h = f(t) = 3 + 5 \tan\left(\frac{t-9}{12}\right), \quad 0 \leq t \leq 12,$$

expresses the height (in thousands of feet) of a balloon in terms of the number of hours past noon.

Determine the possible height(s) of the balloon when it is ascending at the rate 500 ft/hour.

Question 260 The function

$$G = f(s) = 5e^{-s/20}, \quad 0 \leq s \leq 40,$$

expresses the number of gallons of gas in a truck in terms of the trip odometer reading (measured in miles).

Find a function

$$m = h(s), \quad 0 \leq s \leq 40,$$

that takes expresses the dashboard reading for the number of miles left to drive before running out of gas in terms of the trip odometer reading (in miles). The dashboard reading assumes the truck burns gas at the current rate (ie. the rate at odometer reading s miles) for the remainder of the trip.

Derivatives of Inverse Functions, Shortest Version

Derivatives of inverse functions, the main ideas.

1 The Main Idea

The main idea here is simple. That the derivative of the *inverse* of a function is the *reciprocal* of the function's derivative. Or expressed more succinctly, *the derivative of the inverse is the reciprocal of the derivative*.

We don't really need to know much about calculus to understand why.

Question 261 Take for example, the one-to-one function

$$G = f(s), 0 \leq s \leq 50,$$

that expresses the number of gallons of gas in your car in terms your distance from home as you drive home from school. The distance is measured along your route in miles.

Then G is an increasing function of s . Explain why.

Free Response:

The derivative dG/ds gives the rate (in gal/mile) at which your car is burning gas when you are s miles from home.

The derivative ds/dG of the inverse function

$$s = f^{-1}(G)$$

gives your gas mileage (in miles/gal) when you are s miles from home.

It should be clear, from units, notation, and the meanings of the derivatives in this particular scenario, that

$$\frac{ds}{dG} = \frac{1}{dG/ds}.$$

For example, suppose

$$f(30) = 8$$

Learning outcomes:
Author(s):

and

$$\left. \frac{dG}{ds} \right|_{s=30} = 0.025.$$

This means when you are 30 miles from home, your car is burning gas at the rate of 0.025 gal/mile. So at this moment your gas mileage is

$$\left. \frac{ds}{dG} \right|_{G=8} = \frac{1}{0.025 \text{ gal/mile}} = 40 \text{ miles/gal}.$$

2 Theorem

So if $y = f(x)$ is a one-to-one differentiable function of x , then the inverse

$$x = f^{-1}(y)$$

is a differentiable function of y and

$$\left. \frac{dx}{dy} \right|_{y=f(a)} = \frac{1}{\left. (dy/dx) \right|_{x=a}}$$

3 Examples

The difficulty in finding expressions for the inverse of a function is algebraic. The key is to express the derivative of the original function in terms of its *output* instead of its input.

Here are some examples.

Question 262 (a) Explain in words, without any mathematical notation, the meaning of $\arctan(4)$.

Free Response:

(b) Let

$$x = f(\theta) = \tan \theta.$$

Find an expression for the derivative

$$\frac{d}{dx} (\arctan x)$$

of the inverse function

$$\theta = f^{-1}(x) = \arctan x$$

in terms of x .

Explanation. The derivative of the function

$$x = f(\theta) = \tan \theta$$

is

$$\frac{dx}{d\theta} = \frac{d}{dx} (\tan \theta) = \sec^2 \theta.$$

The key step is here, to express the derivative $\sec^2 \theta$ in terms of the output $x = \tan \theta$ of the function f . Since

$$\sec^2 \theta = 1 + \tan^2 \theta = 1 + x^2,$$

$$\frac{dx}{d\theta} = 1 + x^2$$

and

$$\begin{aligned} \frac{d}{dx} (\arctan x) &= \frac{d\theta}{dx} \\ &= \frac{1}{dx/d\theta} \\ &= \frac{1}{\sec^2 \theta} \\ &= \frac{1}{1 + x^2}. \end{aligned}$$

Question 263 (a) Explain in words, without any mathematical notation, the meaning of $\arcsin(-1/3)$.

(b) The function

$$\theta = \arcsin y, \quad -1 \leq y \leq 1,$$

is not the inverse of the function

$$y = g(\theta) = \sin \theta.$$

Explain why.

Free Response:

(c) But the function

$$\theta = f^{-1}(y) = \arcsin y, \quad -1 \leq y \leq 1,$$

is the inverse of the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2.$$

Use the method of Question 2 to find an expression for the derivative

$$\frac{d}{dy} (\arcsin y).$$

Note: When solving the equation

$$\cos^2 \theta + \sin^2 \theta = 1$$

for $\cos \theta$, there are two solutions. You need to choose one and explain the logic behind your choice.

4 Exercises

Exercise 264 Use the method of Question 2 to find expressions for the derivatives of the following functions.

- (a) The function

$$\theta = \arccos x.$$

- (b) The inverse of the function

$$y = \sin \theta, \pi/2 \leq \theta \leq 3\pi/2.$$

- (c) The inverse of the function

$$y = \tan \theta, \pi/2 < \theta < 3\pi/2.$$

Exercise 265 Use the chain rule and the results above to find simplified expressions for each of the following derivatives. Do not simplify any of the functions before differentiating.

- (a)

$$\frac{d}{dx} (\arctan x + \arctan(1/x))$$

- (b)

$$\frac{d}{dx} \left(\arcsin \left(\sqrt{1-x^2} \right) \right)$$

- (c)

$$\frac{d}{dx} (\sin(\arccos x))$$

Derivatives of Inverse Functions, Shortest Version

(d)

$$\frac{d}{dx} (\arcsin(\sin x))$$

(e)

$$\frac{d}{dx} \left(\sin \left(\arccos \left(\sqrt{1 - x^2} \right) \right) \right)$$

Exercise 266 First simplify the function in Exercises 5a, b, c, e and then compute their derivatives.

Derivatives of Inverse Functions, Short Version

Derivatives of inverse functions.

1 The Main Idea

The main idea here is simple. That the derivative of the *inverse* of a function is the *reciprocal* of the function's derivative. Or expressed more succinctly, *the derivative of the inverse is the reciprocal of the derivative.*

We don't really need to know much about calculus to understand why.

Question 267 Take for example, the function

$$G = f(s), 0 \leq s \leq 20,$$

that expresses the number of gallons of gas in your car in terms of your distance from home. The distance is measured in miles along your route and the graph of the function is shown below.

Desmos link: <https://www.desmos.com/calculator/oltnpzth9>

151: Inverse Function 1

- (a) Use the slider s above to approximate (with units) the derivative

$$\left. \frac{dG}{ds} \right|_{s=2}.$$

Explain the derivative's meaning.

- (b) Use the slider s above to approximate (with units) the derivative

$$\left. \frac{ds}{dG} \right|_{G=f(2)}.$$

Explain the derivative's meaning.

- (c) Write an equation that expresses the relationship between the derivatives in parts (a) and (b). Explain why this relationship holds.

Learning outcomes:
Author(s):

Question 268 This question is a continuation of Question 1. Now we are given an expression

$$G = f(s) = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}, \quad 0 \leq s \leq 20,$$

for the function f .

- (a) Use calculus to find an expression for the derivative dG/ds and to compute the exact value of the derivative

$$\left. \frac{dG}{ds} \right|_{s=2}.$$

- (b) Use the idea of Question 1 to find an expression for the derivative ds/dG and the exact value of the derivative

$$\left. \frac{ds}{dG} \right|_{G=f(2)}.$$

- (c) For a more algebraic way to find an expression for the derivative ds/dG , we'll differentiate both sides of the equation

$$G = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}$$

with respect to G . Keeping in mind that this equation, for values of s between 0 and 20, defines s implicitly as a function of G (why?). Then using the chain rule tells us that

$$\frac{dG}{dG} = \left(\frac{s + 23}{1000} \right) \left(\frac{ds}{dG} \right).$$

And solving for the derivative ds/dG gives

$$\frac{ds}{dG} = \frac{1000}{s + 23}.$$

2 Derivatives of Inverse Power Functions

Let's use what we just learned to verify something we already know, but in a more meaningful context.

Example 53. The function

$$A = f(s) = s^2, \quad s \geq 0,$$

expresses the area of a square (measured in square feet) in terms of its side length (in feet).

Since

$$\frac{dA}{ds} = 2s,$$

we know that the derivative of the inverse function

$$s = f^{-1}(A) = \sqrt{A}, \quad A \geq 0,$$

expressing the side length of a square in terms of its area is

$$\begin{aligned} \frac{ds}{dA} &= \frac{1}{dA/ds} \\ &= \frac{1}{2s} \\ &= \frac{1}{2\sqrt{A}}. \end{aligned}$$

We could verify this using the power rule:

$$\begin{aligned} \frac{ds}{dA} &= \frac{d}{dA} (\sqrt{A}) \\ &= \frac{1}{2\sqrt{A}}. \end{aligned}$$

To make this more meaningful, let's look at the derivative

$$\begin{aligned} \left. \frac{dA}{ds} \right|_{s=5} &= (2s) \Big|_{s=5} \\ &= 10 \frac{\text{ft}^2}{\text{ft}} \\ &= 10 \text{ ft}. \end{aligned}$$

Although it's almost never a good idea to simplify the units of the derivative, it is here. The derivative is 10 feet and is equal to *half* the perimeter of our square with side length $s = 5$ feet. To get an idea why this is so, we'll interpret the derivative in terms of small changes.

Suppose we make a small change

$$\Delta s = s - 5 \sim 0$$

in the side length of the square. What can we say about the change

$$\Delta A = f(s) - f(5) = A - 25$$

in its area?

Well since $\Delta s \sim 0$,

$$\begin{aligned}\frac{\Delta A}{\Delta s} &\sim \left. \frac{dA}{ds} \right|_{s=5} = 10, \\ \Delta A &\sim \left(\left. \frac{dA}{ds} \right|_{s=5} \right) \Delta s = 10 \Delta s.\end{aligned}$$

This tells us that *to approximate the change in the area we multiply the change in the side length by 10 feet.*

- (a) The picture below suggests why this is a good approximation. Explain how.
- (b) Do some algebra and compare the actual change

$$\Delta A = f(5 + \Delta s) - f(5) = (5 + \Delta s)^2 - 25$$

in the area with the approximate change. Reconcile the difference with the picture.

Free Response:

Desmos link: <https://www.desmos.com/calculator/h2fm6mm8ua>

151: Square Error

We can interpret the derivative

$$\left. \frac{ds}{dA} \right|_{A=25} = \frac{1}{10} \text{ ft}^{-1}$$

in much the same way by rewriting the above approximation as

$$\Delta s \sim \left(\left. \frac{ds}{dA} \right|_{A=25} \right) \Delta A = \left(\frac{1}{10} \right) \Delta A.$$

This tells us *to approximate the change in the side length we divide the change in the area by 10 feet.*

This interpretation of the derivative is useful in approximating square roots. For example, the side length of a square with an area of 25.2 ft^2 is

$$\begin{aligned}\sqrt{25.2 \text{ ft}^2} &= 5 \text{ ft} + \Delta s \text{ ft} \\ &\sim 5 \text{ ft} + \left(\frac{1}{10 \text{ ft}} \right) (0.2 \text{ ft}^2) \\ &= 5.02 \text{ ft},\end{aligned}$$

giving us a pretty good approximation to

$$\sqrt{25.2} \sim 5.01996.$$

Exercise 269 Carry out a similar analysis with the function

$$V = s^3, s \geq 0,$$

expressing the volume of a cube (in m^3) in terms of its edge length (in meters). Try to include a picture as well.

3 Derivative of the Arc Sine Function

Our goal here is to find an expression for the derivative

$$\frac{d\theta}{dy} = \frac{d}{dy}(\arcsin y)$$

of the inverse sine function

$$\theta = \arcsin y = \sin^{-1}(y), -1 \leq y \leq 1.$$

The first step is to recognize that the inverse sine function is *not* the inverse of the sine function. Because the sine function is not one-to-one, its inverse is not a function.

Rather, the inverse sine function $\theta = \arcsin y$ is the inverse of the function

$$y = f(\theta) = \sin \theta, -\pi/2 \leq \theta \leq \pi/2.$$

Example 54. Before computing the derivative of the inverse sine function in general, we'll look at a specific example and evaluate the derivative

$$\left. \frac{d}{dy}(\arcsin y) \right|_{y=4/5}.$$

Desmos link: <https://www.desmos.com/calculator/alp2mnxqhc>

151: Inverse Sine Function

The idea to computing the above derivative is to first find the slope of the tangent line to the curve $y = \sin \theta$, $-\pi/2 \leq \theta \leq \pi/2$, at the point P on the graph above. Since $d(\sin \theta)/d\theta = \cos \theta$, the slope is the derivative

$$\begin{aligned} \left. \frac{d}{d\theta}(\sin \theta) \right|_{\sin \theta=4/5} &= \left. \frac{d}{d\theta}(\sin \theta) \right|_{\theta=\arcsin(4/5)} \\ &= \cos \theta \Big|_{\theta=\arcsin(4/5)} \\ &= \cos(\arcsin(4/5)). \end{aligned}$$

To evaluate this composition, we need to find the cosine of the angle

$$\theta_0 = \arcsin(4/5)$$

between $-\pi/2$ and $\pi/2$ whose sine is $4/5$. But since

$$\cos^2 \theta_0 + \sin^2 \theta_0 = 1,$$

$$\begin{aligned} \cos \theta_0 &= \pm \sqrt{1 - \sin^2 \theta_0} \\ &= \pm \sqrt{1 - \left(\frac{4}{5}\right)^2} \\ &= \pm 3/5. \end{aligned}$$

To choose the correct sign, we use the fact that $-\pi/2 \leq \theta_0 \leq \pi/2$. This tells us $\cos \theta_0 \geq 0$, so $\cos \theta_0 = 3/5$.

We now know that for the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

$$\left. \frac{dy}{d\theta} \right|_{y=4/5} = 3/5.$$

So for the inverse function

$$\theta = \arcsin y,$$

$$\begin{aligned} \left. \frac{d\theta}{dy} \right|_{y=4/5} &= \left(\left. \frac{dy}{d\theta} \right|_{y=4/5} \right)^{-1} \\ &= \frac{1}{\sqrt{1 - (4/5)^2}} \\ &= 5/3. \end{aligned}$$

Next we'll compute the derivative of the arcsine function for a general input.

Example 55. The inverse sine function

$$\theta = \arcsin(y) = \sin^{-1}(y) \quad -1 \leq y \leq 1,$$

gives the angle between $-\pi/2$ and $\pi/2$ whose sine is equal to y . The key to expressing its derivative

$$\frac{d\theta}{dy} = \frac{d}{dy} (\arcsin y)$$

in terms of y , is to express the derivative of its inverse

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

in terms of $y = \sin \theta$.

Now

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (\sin \theta) = \cos \theta.$$

To express this derivative in terms of $y = \sin \theta$, we use the fact that

$$\cos^2 \theta + \sin^2 \theta = 1.$$

So

$$\frac{dy}{d\theta} = \cos \theta = \pm \sqrt{1 - \sin^2 \theta} = \pm \sqrt{1 - y^2}.$$

To choose between $\pm$, we must remember that by the definition of the arcsine function (see equation (1)),

$$\pi/2 \leq \theta = \arcsin y \leq \pi/2.$$

This tells us that $\cos \theta \geq 0$ and so $\cos \theta = \sqrt{1 - y^2}$. So the derivative of the function

$$y = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

is

$$\frac{dy}{d\theta} = \cos \theta = \sqrt{1 - y^2}.$$

And the derivative of its inverse

$$\theta = \arcsin y$$

is

$$\begin{aligned} \frac{d}{dy} (\arcsin y) &= \frac{d\theta}{dy} \\ &= \left(\frac{dy}{d\theta} \right)^{-1} \\ &= \frac{1}{\sqrt{1 - y^2}}. \end{aligned}$$

4 Derivative of the Arc Cosine Function

The inverse of the function

$$x = \cos \theta, \quad 0 \leq \theta \leq \pi,$$

is the arccosine function

$$\theta = \arccos x, \quad -1 \leq x \leq 1.$$

It returns the angle between 0 and π whose cosine is x .

Exercise 270 (a) Follow the method of Example 5 above to evaluate the derivative

$$\left. \frac{d}{dx} (\arccos x) \right|_{x=-3/4}$$

Desmos link: <https://www.desmos.com/calculator/hgxqjixv4n>

151: Inverse Cosine Function

(b) Follow the method of Example 6 above to find an expression for the derivative

$$\frac{d}{dx} (\arccos x).$$

5 Exercises

Exercise 271 Express the derivative of each of the following functions in terms of its output.

(a)

$$x = \cos \theta, 0 \leq \theta \leq \pi$$

(b)

$$x = \cos \theta, \pi \leq \theta \leq 2\pi$$

(c)

$$y = \tan \theta, -\pi/2 < \theta < \pi/2$$

(d)

$$P = e^{t/5}, t \in \mathbb{R}$$

(e)

$$y = \frac{e^x - e^{-x}}{2}, x \in \mathbb{R}$$

(f)

$$y = \frac{e^x - e^{-x}}{e^x + e^{-x}}, x \in \mathbb{R}$$

(g)

$$y = \frac{e^x + e^{-x}}{2}, x \leq 0$$

(h)

$$G = \frac{s^2}{2000} + \frac{23s}{1000} + \frac{1}{10}, 0 \leq s \leq 20,$$

Exercise 272 Use the results of Exercise 5 to express the derivative of the inverse of each of the functions in that exercise in terms of its independent variable.

Exercise 273 • Find simplified expressions for each of the following derivatives.

- Do not simplify a function before taking its derivative.
- Show all steps in using the chain rule. Use the Leibniz notation.
- Use Desmos to graph each function (before taking the derivative).
- Explain how each graph is related to the derivative.

(a)

$$\frac{d}{dx} (\sin(\arcsin x))$$

(b)

$$\frac{d}{d\theta} (\arcsin(\sin \theta))$$

(c)

$$\frac{d}{dx} (\arctan x + \arctan(1/x))$$

(d)

$$\frac{d}{dx} \left(\arccos x + \arccos \left(\sqrt{1 - x^2} \right) \right)$$

(e)

$$\frac{d}{dx} \left(\arcsin x + \arcsin \left(\sqrt{1 - x^2} \right) \right)$$

Derivatives of Inverse Functions

Using the chain rule to compute the derivative of the inverse of a function.

Introduction

We saw at the beginning of the class the relationship between the derivative of a function and the derivative its inverse. Recall the two problems:

- You measure the edge length of a cube to be 2 cm and use this measurement to compute its volume. How is the error in computing the volume related to the error (assumed to be small) in your measurement of the edge length?
- You submerge a cube in water and measure its volume to be 8 cm³. You then use this measurement to compute the edge length of the cube. How is the error in computing the edge length related to the error (assumed to be small) in your measurement of the volume?

The key to the first problem was to differentiate the function

$$V = f(s) = s^3, s > 0$$

that expresses the volume (in cubic cm) of the cube in terms of its edge length (in cm). We used limits and found that

$$\left. \frac{dV}{ds} \right|_{s=2} = 12.$$

Question 274 (a) What are the units of the above derivative? How do you know?

(b) Interpret the meaning of the above derivative in terms of the geometry of the cube.

Question 275 From here, we let

$$\Delta s = s - 2$$

Learning outcomes:
Author(s):

be the error in our measurement (measured in cm) and

$$\Delta V = s^3 - 8$$

the error in the computed volume (measured in cubic cm). Then for small errors $\Delta s \sim 0$,

$$\frac{\Delta V}{\Delta s} \sim \left. \frac{dV}{ds} \right|_{s=2} = 12,$$

and so

$$\Delta V \sim 12\Delta s.$$

For the second problem we used limits to differentiate the function

$$s = f^{-1}(V), \quad V > 0,$$

and found that

$$\left. \frac{ds}{dV} \right|_{V=8} = 1/12.$$

Question 275.1 What are the units of the above derivative? How do you know?

Then for small errors ΔV in our measurement of the volume,

$$\Delta s \sim \frac{1}{12}\Delta V$$

But all this work took us back to the same approximate relation between the errors ΔV and Δs that we had already found in solving the first problem.

Expressed another way, having found that

$$\left. \frac{dV}{ds} \right|_{s=2} = 12,$$

we should have known immediately (without any computations) that

$$\left. \frac{ds}{dV} \right|_{V=8} = 1/12.$$

For the same reason, this same relationship between the derivative of any function and the derivative of its inverse holds. That is, for any differentiable function $y = f(x)$, the derivative dx/dy of the inverse function $x = f^{-1}(y)$ is

$$\begin{aligned} \frac{d}{dy} (f^{-1}(y)) &= \frac{dx}{dy} \\ &= \frac{1}{dy/dx}. \end{aligned}$$

Well almost. We need to take some care in evaluating the above derivatives at the correct inputs.

Supposing that $b = f(a)$, a more precise statement of the relation between the derivative of a function $y = f(x)$ and the derivative dx/dy of its inverse is

$$\begin{aligned}\frac{d}{dy} (f^{-1}(y)) \Big|_{y=b} &= \frac{dx}{dy} \Big|_{y=b} \\ &= \frac{1}{\frac{dy}{dx} \Big|_{x=a}}.\end{aligned}$$

That is, the derivative of the inverse (of a function) is the reciprocal of the derivative (of that function).

Well, not quite. The above relationship is true if

$$\frac{dy}{dx} \Big|_{x=a} \neq 0.$$

This condition also guarantees that the function $y = f(x)$ is one-to-one in a sufficiently small neighborhood near $x = a$ and is therefore invertible in that neighborhood.

Here are some examples.

The Derivative of the Natural Log Function

Example 56. (a) Use the graph of the function $y = f(x)$ below to estimate the value of the derivative

$$\frac{dy}{dx} \Big|_{x=1.1} = \frac{dy}{dx} \Big|_{y=3} \sim 3$$

(b) Use the result of part (a) and the graph below to estimate the value of the derivative

$$\frac{dx}{dy} \Big|_{x=1.1} = \frac{dx}{dy} \Big|_{y=3} \sim 1/3.$$

Desmos link: <https://www.desmos.com/calculator/nshzdh6jpp>

(c) Set $u = 0.6931$ and $n = 90$ in the demonstration above and estimate the value of the derivative

$$\frac{dy}{dx} \Big|_{x=0.6931} = \frac{dy}{dx} \Big|_{y=2} \sim 2$$

(d) Use the result of part (c) and the graph above to estimate the value of the derivative

$$\left. \frac{dx}{dy} \right|_{x=0.6931} = \left. \frac{dx}{dy} \right|_{y=2} \sim 0.5$$

Question 276 Parts (a) and (c) and perhaps a few more derivatives suggest that

$$f(x) = e^x.$$

(e) Use the results of parts (b) and (d) to evaluate the derivatives

$$\left. \frac{dx}{dy} \right|_{y=3} = \left. \frac{d}{dy} (\ln y) \right|_{y=3} = 1/3,$$

$$\left. \frac{dx}{dy} \right|_{y=2} = \left. \frac{d}{dy} (\ln y) \right|_{y=2} = 1/2,$$

and more generally

$$\left. \frac{dx}{dy} \right|_{y=a} = \left. \frac{d}{dy} (\ln y) \right|_{y=a} = 1/a.$$

Question 277 Our conclusion is that

$$\frac{d}{dx} (\ln x) = \frac{1}{x}.$$

Geogebra activity available at [151: Magnification Factor 3](#)

Question 278 Here's an equivalent, but more computational way to show that

$$\frac{d}{dx} (\ln x) = 1/x.$$

The key is to recognize that the chain rule tells us that if $u = g(x)$ is a differentiable function of x , then

$$\frac{d}{dx} (e^u) = e^u \cdot \frac{du}{dx}.$$

Now to compute the derivative above, we know that since the functions $f(x) = \ln x$ and $g(x) = e^x$ are inverses of one another,

$$e^{\ln x} = x.$$

Then differentiate both sides of this equation with respect to x to get

$$\frac{d}{dx} (e^{\ln x}) = \frac{dx}{dx}.$$

And by the chain rule we can rewrite this equation as

$$(e^{\ln x}) \frac{d}{dx} (\ln x) = 1.$$

And since $e^{\ln x} = x$,

$$\frac{d}{dx} (\ln x) = 1/x.$$

Question 279 Find an equation of the tangent line to the curve $y = \ln x$ at the point $(4, \ln 4)$.

$$y - \ln 4 = \frac{1}{4} (x - 4).$$

Question 280 (a) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln(4x)).$$

(b) Compute the same derivative without using the chain rule.

Explanation. Let

$$y = \ln(4x)$$

and

$$u = 4x.$$

Then

$$y = \ln u$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du} (\ln u) \cdot \frac{d}{dx} (4x) \\ &= \frac{1}{u} \\ &= \frac{1}{4x}. \end{aligned}$$

(b) We can compute the derivative

$$\frac{d}{dx} (\ln(4x))$$

without the chain rule as follows:

$$\begin{aligned}\frac{d}{dx} (\ln(4x)) &= \frac{d}{dx} (\ln 4 + \ln x) \\ &= \frac{d}{dx} (\ln 4) + \frac{d}{dx} (\ln x) \\ &= 0 + \frac{1}{x}.\end{aligned}$$

Question 281 (a) Compute the derivative

$$\frac{d}{dx} (\ln(x^2))$$

both with and without the chain rule. Follow the steps exactly as in the previous example when using the chain rule.

(b) What are the domains of the functions $f(x) = \ln x$ and $g(x) = \ln(x^2)$ and their derivatives.

(c) Graph the function $g(x) = \ln(x^2)$ and its derivative on the same coordinate system by hand.

Question 282 (a) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln |x|).$$

Do this by noting that

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

and using the chain rule to compute

$$\frac{d}{dx} (-x).$$

(b) A reflection about the y -axis takes the graph of a function $y = f(x)$ to the graph of a function $y = g(x)$. Describe the transformation that takes the graph of $y = f'(x)$ to the graph of $y = g'(x)$. Explain your reasoning.

Question 283 (a) Find a function

$$T = f(k), \quad k \geq 1,$$

that expresses the time (in years) it takes an investment to grow by a factor of k . Assume the investment grows at a constant relative instantaneous rate of $i\%/yr$. So, for example if $i = 5$, $f(3)$ would be the time it takes an investment to triple at an relative instantaneous growth rate of $5\%/yr$.

Hint: Let B_0 be the initial investment. Then the value of the investment T years later is

$$kB_0 = B_0 e^{\frac{i}{100}T}.$$

Solve this equation for T to find an expression for the function f . You should get that

$$T = f(k) = (100 \ln k)/i$$

(b) Suppose $i = 5$ and evaluate the derivative

$$\left. \frac{dT}{dk} \right|_{k=3}.$$

(c) What are the units of the above derivative? Explain its meaning in terms of small changes.

I'll leave the units up to you. But the meaning is that it takes an investment (increasing at the relative instantaneous rate of $5\%/yr$) about **0.067** years longer to increase by 201% than it does to triple.

(d) Use the result of parts (b) and (c) to estimate how much longer it would take an investment growing at an relative instantaneous rate of $5\%/yr$ to increase by 210% than it would take to triple. Just use simple arithmetic (ie. multiplication), no calculator.

Question 284 The function

$$G = f(v) = 40 - 0.08 \left(\frac{v}{2} - 25 \right)^2, \quad 25 \leq v \leq 65,$$

expresses the gas mileage of a car (in miles/gallon) in terms of its speed (in miles/hr).

(a) Explain the meaning of the derivative

$$\frac{d}{dv} (\ln(f(v))).$$

Include units in your explanation. Also, what are the units of 25 in the above expression? How do you know?

(b) Evaluate the above derivative at $v = 30$ and explain its meaning in terms of small changes.

Question 285 The function

$$v = g(h), \quad 50 \leq h \leq 200,$$

expresses the speed (in ft/sec) of a hawk in terms of its altitude (in feet) during a portion of its flight.

Suppose that $f(150) = 80$ and

$$\left. \frac{dv}{dh} \right|_{h=150} = -2.$$

(a) What are the units of the above derivative? Explain the meaning of the derivative in the context of small changes in this particular scenario. Be specific.

I'll leave the units up to you, but if we assume the hawk is descending, then its speed would *increase* by about 2 ft/sec as it falls from 150 to 149 feet.

(b) Evaluate the derivative

$$\left. \frac{d}{dh} (\ln(g(h))) \right|_{h=150}.$$

Show all work and explain your reasoning.

The key here is to use the chain rule to evaluate the derivative. So with

$$v = g(h)$$

and

$$w = \ln(g(h)),$$

we have

$$w = \ln v.$$

Then by the chain rule,

$$\begin{aligned} \frac{dw}{dh} &= \frac{dw}{dv} \cdot \frac{dv}{dh} \\ &= \left(\frac{1}{v} \right) \left(\frac{dv}{dh} \right), \end{aligned}$$

and continue from here by evaluating the above expression at $h = 150$.

(c) What are the units of the derivative in part (b)? Explain the meaning of the derivative in the context of small changes in this particular scenario. Be specific by answering this question much like part (a) above (appropriately modified).

The Inverse Sine Function

Let

$$\theta = g(y) = \arcsin y = \sin^{-1}(y).$$

- (a) What is the domain of the function g ?
- (b) What is its range?
- (c) Explain the meaning of $\arcsin y$.
- (d) True or false: The inverse sine function is the inverse of the sine function.
- (e) What function is the inverse of the inverse sine function?

Question 286 The graph of the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

is shown below.

Desmos link: <https://www.desmos.com/calculator/lxwoeir1pt>

Worksheet available at 151: Arc Sine

- (a) Use the graph to estimate the derivative

$$\left. \frac{d}{dy} (\arcsin y) \right|_{y=0.8}.$$

Explain your reasoning.

- (b) Use the graph and the slider u to estimate the derivative

$$\left. \frac{d}{dy} (\arcsin y) \right|_{y=0.8}.$$

- (c) Express the derivative of the function

$$y = f(\theta) = \sin \theta, \quad -\pi/2 \leq \theta \leq \pi/2,$$

in terms of y .

- (d) Use part (c) and the ideas in parts (a),(b) to find an expression for the derivative

$$\frac{d}{dy} (\arcsin y).$$

- (e) Check your answer to part (d) by evaluating the derivative at $y = 0.6, 0.8$.
 (f) What is the domain of the derivative in part (d)?

Question 287 The top of a 25-foot long ladder slides down a vertical wall at the constant rate of 4 ft/sec.

(a) Find a function

$$\theta = f(t), \quad 0 \leq t \leq 6.25,$$

that expresses the angle the ladder makes with the ground (measured in radians) in terms of the number of seconds since the ladder was in the vertical position.

(b) Find the rotation rate of the ladder when the top of the ladder is

(i) 15 feet above the ground.

(ii) 24 feet above the ground.

(iii) 24.9 feet above the ground.

(c) Solve this problem again by working directly with the sine function, not the arcsine function.

Desmos link: <https://www.desmos.com/calculator/5c4lssovbi>

Worksheet available at 151: Ladder and ArcSine

Question 288 The bottom end of a 25-foot long ladder slides across a horizontal floor at the constant rate of 4 ft/sec as the top end slides down a vertical wall.

(a) Find a function

$$\theta = g(u), \quad 0 \leq u \leq 25,$$

that expresses the angle the ladder makes with the wall (measured in radians) in terms of the distance (in feet) between the wall and the bottom end of the ladder.

Hint: Let A be the bottom end of the ladder, B the top end, and point O the point on the wall closest to A (as in the demonstration below). Use right triangle trigonometry in $\triangle AOB$.

(b) Use the slider u in the animation below to approximate each of the following derivatives. Include units. Note that the tick marks on the radian protractor are spaced at intervals of 0.1 radians.

(i) $\left. \frac{d\theta}{du} \right|_{u=1}$

(ii) $\left. \frac{d\theta}{du} \right|_{u=15}$

(iii) $\left. \frac{d\theta}{du} \right|_{u=24.9}$

- (c) What do the above derivatives tell you?
- (d) Find an expression for the derivative $d\theta/du$. Use your expression to evaluate the three derivatives in part (b) and compare these with your estimates.
- (e) Find a function

$$\theta = f(t), \quad 0 \leq t \leq 6.25,$$

that expresses the angle the ladder makes with the ground (measured in radians) in terms of the number of seconds since the ladder was in the vertical position.

Hint: Express the distance u (in feet) between the bottom end of the ladder and the wall in terms of t . Then substitute this expression for u in your function from part (a).

- (f) Find an expression for the derivative

$$r = \omega(t) = d\theta/dt.$$

Interpret its meaning. Include units.

- (g) Find the rotation rate of the ladder when the bottom end of the ladder is
- (i) 1 foot from the wall.
- (ii) 15 feet from the wall.
- (iii) 24.9 feet above the wall.

- (h) Evaluate the limit

$$\lim_{t \rightarrow 6.25} \omega(t)$$

and interpret its meaning.

Desmos link: <https://www.desmos.com/calculator/egolipj5qg>

Worksheet available at 151: Ladder and ArcSine 2

Exercises

Question 289 (a) Simplify the derivative

$$\frac{d}{d\theta} (\arcsin(\sin \theta))$$

- (b) Use the result of part (a) to graph the function

$$y = \arcsin(\sin \theta).$$

Explain your reasoning.

Question 290 Find the measure of the acute angle that the tangent line to the curve

$$y = f(\theta) = \ln |\sec \theta|$$

at the point $(\pi/7, f(\pi/7))$ makes with the x -axis Do not use a calculator.

Question 291 The function

$$q = f(p) = 0.2(2p - 40)^2, \quad 5 \leq p \leq 12,$$

expresses the average number of burgers sold per day at Five Guys in Edmonds in terms of the price (in dollars/burger).

(a) Evaluate the derivative

$$\left. \frac{d}{dp} (\ln(f(p))) \right|_{p=7.5}$$

(b) What are the units of the derivative above? Explain its meaning.

Question 292 The bottom end of a 25-foot ladder lies 24 feet from the base of a vertical wall. Use the appropriate linear approximation to estimate the angle through which the ladder rotates when the bottom end is pulled an additional 0.1 feet away from the wall along a horizontal floor.

Solve this problem twice, first using an inverse trig function and again without an inverse trig function.

Start by defining your variables, with units.

Use a calculator if need be, but only for arithmetic and not to evaluate any trigonometric functions.

Compare your estimate with the actual angle of rotation.

Question 293 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a 14-foot ladder leans against the trunk, the other rests on the horizontal ground. The bottom end of the ladder is pulled away from the trunk at the constant speed of 4 ft/sec. At what rate is the ladder rotating when the bottom and tops ends are respectively 16 and 10 feet from the base of the trunk?

Hint: Use the law of sines.

Desmos link: <https://www.desmos.com/calculator/rpms2jqfpm>

Desmos activity available at 151: *Tree and Ladder*

Question 294 (a) The animation below shows water draining from a tank. Play the animation and sketch by hand a graph of the function $V = f(t)$ that expresses the depth of the water as a function of time. Explain your reasoning. Label the axes with units and the appropriate variable names.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/pdghky6tie>

(b) Torricelli's law says that the rate, say in cm^3/sec , at which water drains out of a small hole in the bottom of a tank is proportional to the square root of the depth of the water. So if $V = f(h)$ is a function that expresses the volume (in cm^3) of water in the tank in terms of the depth (in feet) and $h = g(t)$ is a function that expresses the depth of the water (in feet) in terms of number of seconds past noon, then

$$\frac{dV}{dt} = -k\sqrt{h}$$

for some positive constant k .

For a cylindrical tank of radius r cm,

$$V = f(h) = \pi r^2 h$$

and

$$\frac{dV}{dh} = \pi r^2.$$

So by the chain rule

$$\begin{aligned}\frac{dV}{dt} &= \frac{dV}{dh} \cdot \frac{dh}{dt} \\ &= \pi r^2 \frac{dh}{dt}.\end{aligned}$$

So for the cylindrical tank we can write Torricelli's law as

$$\frac{dV}{dt} = \pi r^2 \frac{dh}{dt} = -k\sqrt{h}.$$

Or equivalently as

$$\frac{dh}{dt} = -\frac{k}{\pi r^2} \sqrt{h} = -k_2 \sqrt{h},$$

where

$$k_2 = \frac{k}{\pi r^2}$$

is a positive constant.

(c) Which of the following functions might express the depth of water in a cylindrical tank (in terms of time) as the water drains out of a small hole in the bottom of the tank? Justify your reasoning.

(i) $h = g(t) = 2(5 - t)^3, 0 \leq t \leq 5$

(ii) $h = g(t) = 2(5 - t)^2, 0 \leq t \leq 5$

Solution:

The key idea is to express the derivative dh/dt as a function of the depth h of the water.

(i) If $h = 2(5 - t)^3$, then

$$\begin{aligned} \frac{dh}{dt} &= \frac{d}{dt} (2(5 - t)^3) \\ &= 2(3)(5 - t)^2 \cdot \frac{d}{dt} (5 - t) \\ &= -6(5 - t)^2. \end{aligned}$$

Now to express dh/dt in terms of h , solve the equation

$$h = 2(5 - t)^3$$

for t to get

$$t = 5 - \left(\frac{h}{2}\right)^{1/3}.$$

Then, substitute this expression for t into the derivative dh/dt :

$$\begin{aligned} \frac{dh}{dt} &= -6(5 - t)^2 \\ &= -6 \left(\frac{h}{2}\right)^{2/3} \\ &= -\left(\frac{6}{2^{2/3}}\right) h^{2/3}. \end{aligned}$$

This tells us that the rate of change in the depth of the water is not proportional to the square root of the water's depth as Torricelli's law requires. So the function

$$h = 2(5 - t)^3, 0 \leq t \leq 5$$

is not a possible depth function for water draining from a cylindrical tank.

Free Response: Give a similar analysis for the depth function of part (ii).

Question 295 This is a continuation of the previous problem.

Now we pour water into a cylindrical tank at a constant rate, while at the same time water leaks out through a small hole in the bottom of the tank. We'll suppose that the tank starts with some initial volume of water.

Free Response: (a) What do you think happens to the water level in the tank initially?

(b) What do you think happens to the water level in the long run?

To model this situation, we need to modify Torricelli's law. For this, let's suppose that we pour water into the cylindrical tank (of radius r) at the constant rate of k_3 cm³/sec. Then with the same notation as before,

$$\frac{dV}{dt} = k_3 - k\sqrt{h}.$$

But since

$$\begin{aligned}\frac{dV}{dt} &= \frac{dV}{dh} \cdot \frac{dh}{dt} \\ &= \pi r^2 \cdot \frac{dh}{dt},\end{aligned}$$

the above modification of Torricelli's law becomes

$$\begin{aligned}\frac{dh}{dt} &= \frac{k_3}{\pi r^2} - \frac{k}{\pi r^2} \sqrt{h} \\ &= k_1 - k_2 \sqrt{h},\end{aligned}$$

where

$$k_1 = \frac{k_3}{\pi r^2}$$

and

$$k_2 = \frac{k}{\pi r^2}$$

are positive constants.

Free Response: (a) What are the units of k_1 and k_2 ? How do you know?

(b) Explain the meaning of k_1 .

The equation

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}$$

expresses the rate of change in the water's depth as a function of the depth. It is called a differential equation and you will learn a little about how to solve equations like this next quarter.

For this particular differential equation, it is not possible to express the depth of the water explicitly as a function of time. But assuming that the depth of the water is h_0 cm at time $t = 0$, it turns out that the function

$$t = g^{-1}(h) = -\frac{2}{k_2} \left(\sqrt{h} - \sqrt{h_0} + \frac{k_1}{k_2} \ln \left| \frac{k_1 - k_2\sqrt{h}}{k_1 - k_2\sqrt{h_0}} \right| \right), \quad t \geq 0,$$

expresses the time (in seconds) in terms of the depth of the water (in cm). We can check that this is indeed correct as follows:

- (a) Show algebraically that the depth of the water at time $t = 0$ is $h = h_0$.
- (b) Use the above expression for $t = g^{-1}(h)$ to compute and then simplify the derivative dt/dh .
- (c) Use the result of part (b) to show that

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}.$$

Desmos link: <https://www.desmos.com/calculator/c78kv7wifv>

Worksheet available at 151: *Draining Cylinder 2*

- (d) Experiment with the sliders above and summarize your observations about how the graph of the function $h = g(t)$ changes depending on the initial depth of the water and the constants k_1 and k_2 .
- (e) Express the equilibrium depth in terms of k_1 and k_2 . Check that your expression has the correct units. The equilibrium depth is the depth at which the water level remains constant. It is also the depth which the water level approaches (independent of the initial depth).
- (f) What happens to the equilibrium depth when k_1 increases (and k_2) is held constant? When k_2 changes and k_1 is held constant?

Looking Down on the Moon, Part 2

Looking at a spherical cap on the lunar surface.

Question 296 This question is about looking on the moon from a spaceship Artemis II.

The idea is that the astronauts can see only a portion of the moon from any altitude above its surface. We wish to analyze how a small change in altitude changes the radius of the spherical cap in view. The radius is measured on the moon's surface. It is the orange arc shown below. Drag the slider h (altitude, in thousands of miles) below to see how small changes in the altitude affect the radius of the spherical cap.

Desmos3D link: <https://www.desmos.com/3d/paqpozwxm>

151: Spaceship above earth

Here is a two-dimensional cross-section of that picture.

Desmos link: <https://www.desmos.com/calculator/b5c4c7ba17>

151: Distance to Horizon 1

Drag point P_2 toward P_1 in the worksheet below to change the change Δh in the altitude. Keep your eye on the change in the radius Δs of the visible spherical cap and the ratio $\Delta s/\Delta h$ of small changes.

Desmos link: <https://www.desmos.com/calculator/url0dnbaai>

151: Distance to Horizon 3

(a) Find a function

$$s = f(h), \quad h \geq 0$$

that expresses the radius of the visible spherical cap (in miles) in terms of the altitude above the surface (in miles). Take the moon to be a perfect sphere of radius 2000 km.

$$s = f(h) = 2000 \arccos(2000/(2000 + h)), \quad h \geq 0.$$

Click the Hint button above for help.

Learning outcomes:
Author(s):

Looking Down on the Moon, Part 2

Hint: Use right triangle $\triangle OTP$ to find an expression for the radian measure of angle $\angle POT$. Then use this angle to find an expression for the arclength AT . Here are more details.

- (i) Enter the two side lengths, measured in feet, in right triangle $\triangle OPT$ below.

$$OT = 2000$$

and

$$OP = h + 2000.$$

- (ii) Let θ be the radian measure of $\angle TOP$. Write an equation with a trigonometric function of θ that relates the two lengths in part (i). Use the Math Editor tab to enter the trig function and the angle θ .

$$\cos \theta = \frac{2000}{h + 2000}.$$

- (iii) Now solve the equation from part (ii) for θ in terms of h . Then use what you know about measuring arclength along a circle to find an expression for the function f . Use the Math Editor tab to help.

$$s = f(h) = 2000 \arccos \left(\frac{2000}{h + 2000} \right).$$

- (b) Now suppose we are 500 km above the lunar surface. Fill in the missing entries in the table below to approximate the scaling factor that converts a small change

$$\Delta h = h - 500$$

in altitude to a good approximation of the change

$$\Delta s = f(h) - f(500)$$

in the radius of the visible spherical cap.

h (km)	$s = f(h)$ (km)	$\Delta h = h - 500$ (km)	$\Delta s = f(h) - f(500)$ (km)	$\Delta s / \Delta h$ (units?)
497				
498				
499				
500				
501				
502				
503				

- (c) *Do the data above suggest that the quotients $\Delta s/\Delta h$ approach some number as h approaches 500? If so, use the data to approximate that number. If not, explain why not.*
 - (d) *Use calculus to determine the exact scaling factor. Include units.*
 - (e) *Explain the meaning of the derivative as a scaling factor. Give an example with specific small changes to explain the meaning of the derivative.*
-

Inverse Trig CW

Applications of inverse trig derivatives.

Question 297 The ends of a 25-foot ladder rest against a vertical wall and a floor inclined at an angle of

$$\phi = \arccos(5/6)$$

below the horizontal.

Desmos link: <https://www.desmos.com/calculator/q2g6qqwlzc>

151: Ladder and Sloping Ground

- (a) Find a function that expresses the acute angle the ladder makes with the wall in terms of the distance from the top of the ladder to the base of the wall. Start with a sentence that defines the output, the input, and the function name. Include units in your definition and choose meaningful variable names. Include a picture to help with your explanation.
- (b) Now suppose the ladder's top end is 20 feet above the base of the wall. Find the scaling factor that converts (by multiplication) a small change in the height of the ladder's top end above the base of the wall (in feet) to an accurate approximation of the corresponding change in the acute angle the ladder makes with the wall. Include units.
- (c) Interpret the meaning of the scaling factor with an example using specific small changes.

Question 298 A rocket orbits a planet of radius 2000km at an altitude of 400 km above the surface. We wish to find a scaling factor that converts (by multiplication) a small change in orbit's radius to an accurate approximation of the corresponding change in radius of the portion of the planet's surface visible to an astronaut in the rocket. The radius is measured along the planet's surface.

For help see

[Looking Down on the Moon Part 2](#)

Learning outcomes:
Author(s):

- (a) Start by writing a complete sentence that defines a function expressing the radius of the visible portion in terms of the altitude above the surface. Use meaningful variable names. Include units in your definition.
- (b) Use your function to compute the exact value of the scalar factor described above.
- (c) Explain the meaning of the scaling factor by giving an example with specific small changes.

Question 299 An elastic cord runs from the top of an 80-foot tall tower to a point on the ground.

- (a) Find a function that expresses the acute angle (in radians) the cord makes with the ground in terms of the cord's length (in feet). Start with a complete sentence that defines a function with meaningful variable names and units.
- (b) Suppose the cord is attached to the ground 40 feet from the base of the tower. Find the scaling factor that converts a small change in the cord's length (in feet) to an accurate approximation of the change in the angle the cord makes with the ground.
- (c) Explain the meaning of the scaling factor with an example using specific changes.
- (d) How far from the base of the tower is the end of the cord when the scaling factor in part (b) is $-1/30$ rad/ft?

Question 300 The function

$$h = f(\phi) = \frac{24}{\pi} \arccos(-\sqrt{3} \tan \phi)$$

expresses the number of hours of daylight per day at a latitude of 60°N in terms of the declination of the sun. The declination is the angle the sun's rays make with the plane of the equator. It is positive between the spring and fall equinoxes in the northern hemisphere, negative in the fall and winter months.

- (a) Find an expression for the derivative $dh/d\phi$.
- (b) Evaluate the derivative

$$\left. \frac{dh}{d\phi} \right|_{\phi=\arctan(1/3)}.$$

- (c) Interpret the derivative in part (b) as a scaling factor. Give an example with specific small changes.

Question 301 The function

$$h = f(s), \quad 0 \leq s \leq 10,$$

expresses the altitude (in km) of a trail in terms of the distance from the trailhead (in km).

- (a) Find a function

$$\phi = g(\theta), \quad -\pi/2 < \theta < \pi/2$$

that expresses the acute angle the trail makes with the horizontal at the point s miles from the trailhead in terms of the acute angle the curve $h = f(s)$ makes with the s -axis at the point $(s, f(s))$.

- (b) Evaluate the derivative

$$\left. \frac{d\phi}{d\theta} \right|_{\theta=\arctan(2)}.$$

- (c) Interpret the derivative in part (b) as a scaling factor. Give an example with specific small changes.

Question 302 The function

$$T = f(r) = \frac{R^{3/2}}{\sqrt{2GM}} \left(\frac{\pi}{2} - \arcsin \left(\sqrt{\frac{r}{R}} \right) + \sqrt{\frac{r}{R}} \left(1 - \frac{r}{R} \right) \right)$$

The Quotient Rule

An introduction to the quotient rule.

Discussion Questions

Question 303 Consider the following two questions.

- (a) During the course of one year, the national debt of a country increases by 20% and its population increases by 5%. Find the relative change in the per-capita (ie. per-person) share of the national debt over the one-year period.
- (b) At some instant the national debt of a country is increasing at the rate of 20%/yr and its population is increasing at the rate of 5%/yr. At what relative rate is the per-capita share of the national debt changing at this instant?
- (a) Are these questions identical?
- (b) Answer the first question.
- (c) Use numerical methods to approximate the answer to the second question. Hint: Approximate the relative changes in the national debt and population over a small time interval. Then use these to approximate the relative change and the relative average rate of change in the per-capita share of the debt over this time interval. Use the worksheet below to help.

Desmos link: <https://www.desmos.com/calculator/rk4aocmccz>

Desmos activity available at 151: Relative Quotient Rule 1

- (d) For a more algebraic approach to the second question, let's suppose that at the moment in question, say time $t = 0$ measured in years, the national debt is D_0 dollars, the population is P_0 , and the per-capita share is $C_0 = D_0/P_0$. We'll also suppose at this instant that the debt is increasing at the relative rate of $100p\%/yr$ and the population at the relative rate of $100q\%/yr$. We wish to determine the relative rate at which the per-capita share of the debt is changing at this time. We'll do this using limits.

Learning outcomes:
Author(s):

To start let $D = f(t)$ and $P = g(t)$ be the debt (in dollars) and population functions expressed in terms of the number of years since the moment in question. Also, let

$$C = \frac{D}{P} = \frac{f(t)}{g(t)}$$

be the per-capita share of the debt.

Now the relative average rate of change in the per-capita share over the time interval $t \in [0, \Delta t]$ is

$$\begin{aligned} \frac{1}{C_0} \cdot \frac{\Delta C}{\Delta t} &= \frac{1}{\Delta t} \cdot \frac{P_0}{D_0} \cdot \Delta C \\ &= \frac{1}{\Delta t} \cdot \frac{P_0}{D_0} \left(\frac{f(\Delta t)}{g(\Delta t)} - \frac{f(0)}{g(0)} \right). \end{aligned}$$

The key now is to approximate the debt

$$D = f(\Delta t)$$

and the population

$$P = g(\Delta t)$$

at time Δt . For this we need to use the respective relative rates of change $100p\%/yr$ and $100q\%/yr$ in the national debt and population at time $t = 0$.

At time $t = \Delta t$ hours, the national debt is approximately

$$D = f(\Delta t) \sim D_0(1 + p\Delta t)$$

and the population approximately

$$P = g(\Delta t) \sim P_0(1 + q\Delta t).$$

So the change in the per-capital share of the debt over this time interval is approximately

$$\begin{aligned} \Delta C &= \frac{D}{P} - \frac{D_0}{P_0} \\ &\sim \frac{D_0(1 + p\Delta t)}{P_0(1 + q\Delta t)} - \frac{D_0}{P_0} \\ &= \frac{D_0}{P_0} \left(\frac{(p - q)\Delta t}{1 + q\Delta t} \right) \end{aligned}$$

and the relative change approximately

$$\begin{aligned} \frac{\Delta C}{C_0} &= \frac{P_0}{D_0} \cdot \Delta C \\ &= \frac{(p - q)\Delta t}{1 + q\Delta t}. \end{aligned}$$

So over the time interval $t \in [0, \Delta t]$, the relative average rate of change in the per-capita share of the debt is

$$\begin{aligned}\frac{1}{C_0} \cdot \frac{\Delta C}{\Delta t} &= \frac{1}{\Delta t} \cdot \frac{\Delta C}{C_0} \\ &= \frac{p - q}{1 + q\Delta t}.\end{aligned}$$

Finally, the relative instantaneous rate of change in the per-capita share of the debt at time $t = 0$ is

$$\begin{aligned}\frac{1}{C_0} \cdot \frac{dC}{dt} \Big|_{t=0} &= \lim_{\Delta t \rightarrow 0} \frac{p - q}{1 + q\Delta t} \\ &= p - q.\end{aligned}$$

- (e) For yet another way to approach the second question, assume the debt and population functions increase at constant relative rates. Then find expressions for these functions and use these to answer the question.
- (f) What is your guess about the relative (instantaneous) rate of change in the quotient of two functions? How is it related to the relative rates of change of the functions?

Question 304 At some instant the per-capita share of a country's national debt is increasing at the rate of 4%/yr and its population is increasing the rate of 6%/yr.

- (a) Find exponential functions $P = f(t)$ and $C = g(t)$ that model the population and per-capita share of the debt, each expressed in terms of the number of years since the moment described above.
- (b) Use your expressions from part (a) to find a function $D = h(t)$ that expresses the national debt in terms of the same variable t . Simplify the expression for $D = h(t)$.
- (c) Use your simplified expression for the function $D = h(t)$ to find the relative rate at which the national debt is changing at the instant in question.
- (d) Repeat parts (a)-(c) using linear functions for the population and per-capita share of the debt instead.

Question 305 Use your guess for the relative rate of change in the quotient of two functions to find an expression for the derivative of the quotient of two functions.

- Question 306** (a) At 9am on May 29, the balance in an account is increasing at the relative rate of 8%/yr. At the same time, the price of a stock is increasing at the rate of 5%/yr. Is the number of shares you can buy with the balance in the account increasing or decreasing at this time? At what relative rate? No computations. Just explain what you think.
- (b) How would your answer to part (a) change if instead the stock price were decreasing at the rate of 5%/yr?

Question 307 Let $B = f(t)$ and $P = g(t)$ be functions that respectively express the balance (in dollars) in an account and the price (in dollars/share) of a stock in terms of the number of years past 9am on May 29. Let $S = h(t)$ be the number of shares of the stock you can buy with the balance in the account at time t years past 9am on May 29.

- (a) Interpret the meanings of the following derivatives. Include units.

(i) $\frac{d}{dt}(\ln(f(t))) = \frac{d}{dt}(\ln B)$

(ii) $\frac{d}{dt}(\ln(g(t))) = \frac{d}{dt}(\ln P)$

(iii) $\frac{d}{dt}(\ln(h(t))) = \frac{d}{dt}(\ln S)$

item Express the derivative

$$\frac{d}{dt}(\ln S) = \frac{d}{dt} \left(\ln \left(\frac{B}{P} \right) \right)$$

in terms of the derivatives

$$\frac{d}{dt}(\ln B)$$

and

$$\frac{d}{dt}(\ln P).$$

The Relative Quotient Rule

Had we never learned about relative changes and relative rates of change, the following questions would have almost forced these ideas upon us.

Question 308 (a) Suppose over a six-month period, the national debt of a small country increases from \$4 billion to \$4.16 billion while the population increases from 20 million to 21 million. Does the per-capita (ie. per-person) share of the national debt increase or decrease during this period?

(b) At noon on July 1, the national debt of a small country with a population of 20 million is \$4 billion. At that same instant the population is increasing at the rate of 1 million people/yr while the national debt is increasing at the rate of \$0.16 billion/yr. Is the per-capita share of the national debt increasing or decreasing at this instant? At what rate?

Explanation. (a) The question is really to determine which is greater,

$$c_1 = \frac{4 \times 10^9 \text{ dollars}}{2 \times 10^7 \text{ person}}$$

or

$$c_2 = \frac{4.16 \times 10^9 \text{ dollars}}{2.1 \times 10^7 \text{ person}} = \frac{4 \times 10^9 \times 1.04 \text{ dollars}}{2 \times 10^7 \times 1.05 \text{ person}} = \left(\frac{1.04}{1.05} \right) c_1.$$

And because the total debt increased by 4% while the population increased by 5%, the per-capita share of the national debt decreased during the six-month period.

But the per-capita share of the debt did *not* decrease by 1%, but rather by about (to the nearest thousandth of a percent) **0.952%**.

(b) For the instantaneous rate, let

$$Q = f(t)$$

be the total national debt in billions of dollars at time t years past noon on July 1 and

$$P = g(t)$$

the population in millions at time t . Also, let

$$C = h(t) = \frac{f(t)}{g(t)}$$

be the per-capita share of the debt in thousands of dollars/person at time t years past noon on July 1. We wish to find the value of the derivative

$$\left. \frac{dC}{dt} \right|_{t=0}.$$

Part (a) suggests we express the relative instantaneous rate of change

$$\frac{1}{C} \cdot \left. \frac{dC}{dt} \right|_{t=0}$$

in terms of the relative rates

$$\frac{1}{Q} \cdot \left. \frac{dQ}{dt} \right|_{t=0} = \mathbf{0.04}$$

and

$$\frac{1}{P} \cdot \frac{dP}{dt} \Big|_{t=0} = 0.05$$

Perhaps the easiest way to do this, although not entirely complete, is to assume that the population and national debt each grow exponentially. In that case, the relative growth rates of each are constant and

$$Q = f(t) = 4e^{0.04t}$$

and

$$P = g(t) = 20e^{0.05t}.$$

This tells us that

$$C = \frac{f(t)}{g(t)} = 0.2e^{-0.01t}$$

and

$$\frac{1}{C} \cdot \frac{dC}{dt} \Big|_{t=0} = -0.01$$

So at noon on July 1, we suspect that the per-capita share of the debt is decreasing at the relative rate of 1%/yr and at the absolute rate of \$2/person/yr.

To be sure this is true even if the relative rates of change in the population and total debt are *not* constant, we can be more general and compute the relative instantaneous rate of change in the per-capita share of the national debt as

$$\begin{aligned} \frac{1}{C} \cdot \frac{dC}{dt} &= \frac{d}{dt} \left(\ln \left(\frac{Q}{P} \right) \right) \\ &= \frac{d}{dt} (\ln Q - \ln P) \\ &= \frac{d}{dt} (\ln Q) - \frac{d}{dt} (\ln P) \\ &= \frac{1}{Q} \cdot \frac{dQ}{dt} - \frac{1}{P} \cdot \frac{dP}{dt}. \end{aligned}$$

Theorem 2. (a) (The Relative Quotient Rule) If $Q = f(t)$ and $P = g(t)$ are differentiable functions of t , then if $g(t) \neq 0$,

$$C = \frac{f(t)}{g(t)} = \frac{Q}{P}$$

is a differentiable function of t and

$$\frac{1}{C} \cdot \frac{dC}{dt} = \frac{1}{Q} \cdot \frac{dQ}{dt} - \frac{1}{P} \cdot \frac{dP}{dt}.$$

The relative rate of change in a quotient of two functions is equal to the difference in the relative rates of change in the functions.

(b) (The Quotient Rule) With the same hypotheses (and obtained by multiplying both side of the previous equation by $C = Q/P$),

$$\frac{d}{dt} \left(\frac{Q}{P} \right) = \frac{1}{P} \cdot \frac{dQ}{dt} - \frac{Q}{P^2} \cdot \frac{dP}{dt}.$$

The Tangent Fuction

Question 309 Compute the derivative

$$\frac{d}{d\theta} (\tan \theta)$$

Explanation. We compute this derivative from scratch by letting

$$y = \tan \theta = \frac{\sin \theta}{\cos \theta}.$$

Then

$$\ln |y| = \ln |\sin \theta| - \ln |\cos \theta|$$

and

$$\frac{d}{d\theta} (\ln |y|) = \frac{d}{d\theta} (\ln |\sin \theta| - \ln |\cos \theta|).$$

So

$$\begin{aligned} \frac{1}{y} \cdot \frac{dy}{d\theta} &= \frac{1}{\sin \theta} \cdot \frac{d}{d\theta} (\sin \theta) - \frac{1}{\cos \theta} \cdot \frac{d}{d\theta} (\cos \theta) \\ &= \cot \theta + \tan \theta. \end{aligned}$$

Then multiplying both sides by $y = \tan \theta$ gives

$$\frac{d}{d\theta} (\tan \theta) = 1 + \tan^2 \theta.$$

Question 310 (a) Use the graph of the function

$$y = f(\theta) = \tan(2\theta)$$

below to estimate the y -coordinates of all points on the curve where the tangent lines are parallel to the lines

(i) $6x - y = 12$.

(ii) $6x + y = 12$.

(b) Find the exact y -coordinates without using a calculator.

Desmos link: <https://www.desmos.com/calculator/obz6ghw3ej>

Desmos activity available at 151: Tangent Graph

Question 311 You stand 50 feet from the base of a tree and measure the angle of elevation to the top of the tree with an error of at most $\pm 2^\circ$. You then compute the height of the tree above eye level to be 100 feet.

Desmos link: <https://www.desmos.com/calculator/yjyghsoeog>

Desmos activity available at 151: Angle of Elevation 1

- (a) Use the demonstration above to approximate your error in computing the tree's height.
- (b) Use the appropriate linear approximation to estimate your error in computing the tree's height. Compare this with your estimate from part (a). Do not use a calculator. Do this as follows:

- Find a function

$$h = f(\theta), 0 < \theta < \pi/2,$$

that expresses the computed height of the tree above eye level (measured in feet) in terms of the measured angle of elevation (in radians). Draw a picture to help with your explanation.

This function is

$$h = f(\theta) = 50 \tan \theta, 0 < \theta < \pi/2.$$

- Next find an expression for the derivative $dh/d\theta$ and evaluate the derivative

$$\left. \frac{dh}{d\theta} \right|_{h=100}.$$

- Now we'll take the exact height of the tree above eye level to be 100 feet and let

$$\Delta h = h - 100$$

be the error in the computed height and

$$\Delta \theta = \theta - \arctan(2)$$

be the error in the measured angle.

Then if $\Delta \theta \sim 0$,

$$\left. \frac{dh}{d\theta} \right|_{h=100} \sim \frac{\Delta h}{\Delta \theta}$$

and so

$$\Delta h \sim 250 (\Delta \theta).$$

- I'll let you continue from here.

The Inverse Tangent Function

Question 312 (a) Explain the meaning of the function

$$\theta = f(y) = \arctan y.$$

In particular, what does it take as an input and what does it return as an output? Include the domain and range of the function as part of your explanation.

(b) Express the derivative

$$\frac{d}{d\theta} (\tan \theta) = 1 + \tan^2 \theta$$

of the function

$$y = \tan \theta$$

in terms of y .

(c) Use part (b) to find an expression for the derivative

$$\frac{d\theta}{dy} = \frac{d}{dy} (\arctan y) = \frac{d}{dy} (\tan^{-1} y).$$

(d) Evaluate the derivatives

$$\left. \frac{d}{d\theta} (\tan \theta) \right|_{\theta=\pi/4}$$

and

$$\left. \frac{d}{dx} (\arctan x) \right|_{x=1}.$$

Comments?

Question 313 You stand 60 feet from the base of a tree and measure the angle of elevation to the top of an 85-foot tall tree. You then move 4 feet closer to the tree and measure the same angle. Assume your eyes are five feet above the ground.

Desmos link: <https://www.desmos.com/calculator/qgqvq3noah>

Desmos activity available at 151: Tree 3

(a) Drag the slider u in the demonstration above to estimate the change in the angle of elevation after you move 4 feet closer to the tree. Note that the angles of elevation are $\angle TEB$ and $\angle TFB$ and that $\angle ETF$ measures their difference. Consecutive tick marks on the protractor are spaced at intervals of 0.01 radians.

(b) Use derivatives to approximate the change in the angle of elevation and compare your approximation with your estimate in part (a) and with the actual change. Use a calculator with only addition, subtraction, multiplication, and division.

Go about this by first finding a function

$$\theta = f(s), s > 0,$$

that expresses the angle of elevation to the top of the tree (measured in radians) in terms of your distance from the tree (measured in feet). Use the arctangent function in your expression. Do not use the inverse cotangent function.

The function is

$$\theta = f(s) = \arctan(80/s), s > 0.$$

Then continue in a manner similar to Question 4.

Two Motions

Question 314 (a) Play the slider s in the demonstration below to show the motion of a beetle crawling along the y -axis as it leaves behind tracks spaced at equal time intervals.

(b) Use the animation to sketch (by hand) a graph of the function

$$s = f(t)$$

that expresses the position (in this case the y -coordinate) of the beetle as a function of time. Label the axes with the appropriate variable names and units. Then activate the folder in Line 2 to see how you did.

(c) Use the animation to sketch (by hand) a graph of the function

$$v = g(t)$$

that expresses the beetle's velocity (in this case the rate of change, with respect to time, in the beetle's y -coordinate). Label the axes with the appropriate variable names and units. Then activate the folder in Line 7 to see how you did.

Desmos link: <https://www.desmos.com/calculator/srotstrdzm>

Desmos activity available at 151: *Tangent Motion*

Question 315 Repeat parts (a)-(c) of the previous question for the motion below.

Desmos link: <https://www.desmos.com/calculator/h6vq21lfql>

Desmos activity available at 151: *ArcTangent Motion*

Exercises

Question 316 (a) Use the website below to compute an accurate estimate of the current rate (in dollars/year) at which the U.S. national debt is changing. Explain your method.

National Debt Clock

(b) Use the website below to compute an accurate estimate (in people/yr) at which the U.S. population is currently increasing.

Population Clock

(c) Use your estimates above and the current national debt and U.S. population to compute

(i) the current rate of change in the per-capita share of the national debt.

(ii) the current relative rate of change in the per-capita share of the national debt.

Include units in every number in each step of your computations.

Question 317 The function

$$C = f(t) = At^4 e^{-kt}, \quad t \geq 0,$$

expresses the concentration of a drug (measured in mg/L) in the bloodstream in terms of the number of hours since the drug was injected. Here A and k are positive constants.

Desmos link: <https://www.desmos.com/calculator/lvsxu3wa9a>

Desmos activity available at [151: Drug Concentration](#)

- (a) What are the units of the constant A ? How do you know?
- (b) What are the units of k ? How do you know?
- (c) Use the graph of $C = f(t)$ above to sketch by hand a graph of the derivative $r = dC/dt$. Be sure to label the axes with the appropriate variable names and units. Do not use technology. Explain your reasoning.
- (d) Use calculus and algebra to find an expression in terms of k for the time when the concentration is a maximum. Work in general. Do not use a specific value of k . Check your work by following the directions, Lines 4 and 5, in the desmos demonstration. Label also the coordinates of the corresponding point on your hand-drawn graph of the derivative.
- (e) Find expressions (in terms of k) for the times when the concentration is increasing and decreasing at the maximum rates. Check that your expressions have the correct units. Work in general. Do not use a specific value of k . Check your work by following the directions, Lines 6-9, in the desmos demonstration. Label also the coordinates of the corresponding points on your hand-drawn graph of the derivative.
- (f) Find an expression (in terms of k) for the relative rate at which the concentration is changing t hours after the injection. Check that your expression has the correct units. Work in general. Do not use a specific value of k .
- (g) Suppose the concentration is at its maximum five hours after injection and determine when the concentration is increasing at the rate of 50%/hr. Determine also when the concentration is decreasing at the rate 50%/hr.
- (h) Observations?

Question 318 The function

$$P = f(t) = 5 - 3t + t^2, \quad 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) Use the graphs of the function $P = f(t)$ and the function $r = f'(t)/f(t)$ to estimate when the stock price is increasing at the greatest relative rate.
- (b) Use algebra to find the exact time when the stock price is increasing at the greatest relative rate.

Hint: What is the value of the derivative dr/dt at this time? But start by finding an expression for the instantaneous relative rate of change in the stock price.

Desmos link: <https://www.desmos.com/calculator/y78fnny7s3>

Desmos activity available at [151: Stock Price 4](#)

Question 319 You jog once around a circular track of radius r meters at the constant speed of v m/sec. A flagpole lies b meters due east of the track's center.

(a) Find a function

$$s = f(t), 0 \leq t \leq 2\pi r/v,$$

that expresses your distance (in meters) to the flagpole in terms of the time (measured in seconds) since you started running. Assume you start at the point A on the track nearest the flagpole. Explain your reasoning. Work with the general parameters r , v , and b , not with any specific values for these parameters.

(b) Find an expression for the time when your distance to the flagpole is increasing at the greatest rate. Try to give a geometric interpretation of your position at this time.

Desmos link: <https://www.desmos.com/calculator/bxofhvfbs>

Demonstration available at [Math 151: Jogger 3](#)

Moving Averages and their Rates of Change

Moving averages and the quotient rule.

- Question 320** (a) Your average for the first four exams in a class is 90%. How would a score of 100% on the fifth exam change your average?
- (b) Your average for the first nine exams in a class is 90%. How would a score of 100% on the tenth exam change your average?
- (c) Over the first four hours of a ten-hour car trip, your average speed was 90 km/hour. During the fifth hour you drove at a constant speed of 100 km/hour. Compare your average speeds over the first four hours and over the first five hours of your trip.
- (d) Over the first nine hour of a ten-hour car trip, your average speed was 90 km/hour. During the tenth hour you drove at a constant speed of 100 km/hour. Compare your average speeds over the first nine hours and over the entire ten hours of your trip.
- (e) A computer display keeps track of your average speed since the start of your trip. Suppose four hours into a ten-hour car trip, your trip odometer reading (set to zero at the start of your trip) reads 360km and your speedometer reads 100 km/hour. Find the rate of change, with respect to time (measured in hours), in your average speed as shown on the display at this instant.
- (f) A computer display keeps track of your average speed since the start of your trip. Suppose nine hours into a ten-hour car trip, your trip odometer reading (set to zero at the start of your trip) reads 810km and your speedometer reads 100 km/hour. Find the rate of change, with respect to time (measured in hours), in your average speed as shown on the display at this instant.

Implicit Differentiation

An introduction to implicit differentiation.

Discussion Questions

Question 321 Which of the following equations define y implicitly as a function of x in a sufficiently small neighborhood of the given point? Supplement your reasoning with a graph of each relation.

- (a) $x^2 + y^2 = 25$ near the point $(4, -3)$
- (b) $x^2 + y^2 = 25$ near the point $(0, -5)$
- (c) $x^2 + y^2 = 25$ near the point $(-5, 0)$
- (d) $x^2 - xy + y^2 = 1$ near the point $(1, 1)$.
- (e) $x^2 - xy + y^2 = 3$ near the point $(1, 2)$.
- (f) $x^2 - xy + y^2 = 3$ near the point $(2, 1)$.

Introduction to Implicit Differentiation

Question 322 Let $p = f(t)$ be a differentiable function of t . Find expressions for each of the following derivatives, first supposing that

$$p = f(t) = t^3 + 1,$$

and then more generally, not assuming any particular expression for the function f .

- (a) $\frac{d}{dp} (p^5)$
- (b) $\frac{d}{dt} (p^5)$
- (c) $\frac{d}{dt} (e^{2p})$
- (d) $\frac{d}{dt} (t^4 \sin(p))$

Learning outcomes:
Author(s):

Question 323 (a) Find equations of the tangent lines to the ellipse

$$x^2 - xy + 2y^2 = 9$$

at the points where the ellipse intersects the x -axis.

(b) Find equations of all horizontal tangent lines to the ellipse.

(c) Find equations of all vertical tangent lines.

Desmos link: <https://www.desmos.com/calculator/ox3wvfitqk>

Desmos activity available at 151: *Implicit Ellipse*

Question 324 (a) Find an equation of the tangent line to the curve

$$4xe^y - 3y \cos(2xy) = 0$$

at the origin.

(b) Enter your equation in Line 5 of the worksheet below to check your answer.

Desmos link: <https://www.desmos.com/calculator/1qprxg226m>

Desmos activity available at 151: *Implicit 1*

Astroids

Question 325 (a) The coordinate axes cut out a segment from the tangent line to the curve

$$x^{2/3} + y^{2/3} = 5$$

at the point $P(8, 1)$. Find the length of that segment. Do this by using implicit differentiation to help find an equation of the tangent line at P and go from there. But first show that P lies on the curve.

(b) As a bug crawls around the above curve and passes the point $P(8, 1)$, it is moving away from the x -axis at the rate of 4 cm/sec. Is the bug moving toward or away from the y -axis at this time? At what rate?

(c) Find the slope of the tangent line in part (a) without using implicit differentiation.

Question 326 (a) Use implicit differentiation to show that segments cut by the coordinate axes from the tangent lines to the astroid

$$x^{2/3} + y^{2/3} = a^{2/3}$$

all have the same length. Here $a > 0$ is a constant.

(b) Prove the same result by using trigonometric functions to parameterize the astroid.

Desmos link: <https://www.desmos.com/calculator/vrythrvjuc>

Desmos activity available at 151: Astroid

Ellipses and Related Curves

Question 327 (a) Use implicit differentiation to find an equation of the tangent line to the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

at the point P with coordinates (x_0, y_0) . Here a, b are positive constants.

(b) Input your equation from part (a) on Line 3 of the desmos worksheet below. Drag point P to check your equation is correct.

(c) Solve part (c) again without calculus by considering a composition of transformations that takes the circle

$$x^2 + y^2 = 1$$

to the ellipse in part (a).

Desmos link: <https://www.desmos.com/calculator/t1e9v7ncpv>

Desmos activity available at 151: Tangents to Ellipse

Question 328 An ellipse through the point $P(0, b)$ has focal points F_1 at the origin and F_2 at the point $A(a, 0)$.

(a) Use the definition of an ellipse as the set of points whose distances to the foci have a constant sum to find an equation of the ellipse.

(b) Use implicit differentiation to find the slope of the tangent line to the ellipse at P .

- (c) Find an equation of the line normal to the ellipse at Q .
- (d) Find the coordinates of the point Q where the normal line intersects the x -axis.
- (e) Express the ratio $F_1Q : QF_2$ in terms of a and b . Interpret the ratio geometrically.

The ratio is

$$\frac{F_1Q}{QF_2} = \frac{b}{\sqrt{a^2 + b^2}}.$$

- (f) Check your work in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/xqywotxxf9>

Desmos activity available at 151: Normal to an Ellipse

Question 329 Let F_1 and F_2 be respectively the origin and the point with coordinates $(a, 0)$. The curve $\mathcal{C}$ passes through the point $P(0, b)$. The curve is defined by the property that the sum of the distance from a point Q of $\mathcal{C}$ to F_1 and k times its distance to F_2 is a constant.

- (a) Find an equation of the curve $\mathcal{C}$.
- (b) Find an equation of the tangent line to $\mathcal{C}$ at P .
- (c) Find an equation of the line normal to $\mathcal{C}$ at P .
- (d) Find the coordinates of the point Q where the normal line intersects the x -axis.
- (e) Express the ratio $F_1Q : QF_2$ in terms of a and b . Interpret the ratio geometrically.

The ratio is

$$\frac{F_1Q}{QF_2} = \frac{bk}{\sqrt{a^2 + b^2}}.$$

- (f) Check your work in the demonstration below.

Desmos link: <https://www.desmos.com/calculator/uz7w5uh0v1>

Desmos activity available at 151: Generalized Ellipse

The Sliding Ladder

Question 330 You walk along a straight path, moving either directly toward or directly away from a tree at the end of the path.

Let $s = f(t)$ be a function that expresses your distance to the tree in terms of the number of minutes past noon and suppose that

$$\left. \frac{ds}{dt} \right|_{t=5} = -240.$$

- (a) Interpret the meaning of the above derivative.
- (b) What is your speed at 12:05pm?
- (c) Find an expression for your speed at time t minutes past noon.

Question 331 A tree leans precariously with its trunk making an angle of ϕ radians with the ground. One end of a ladder with length L feet leans against the trunk, the other rests on the horizontal ground. You slide the bottom end of the ladder along the ground at random.

- (a) Let s and t be the respective distances (measured in feet) from the base of the trunk to the bottom and top ends of the ladder. Find an equation that relates these distances. Use the parameters L and ϕ , not their particular values in the demonstration below.
- (b) Let u be time measured in seconds and suppose that s and t are (unknown) functions of u . Express the derivative dt/du in terms of the derivative ds/du . Interpret the meanings of these derivatives.
- (c) Switch s and t in your equation from part (b). What do you notice? Explain why.
- (d) Use part (b) to express the speed of the top end of the ladder in terms of the speed of its bottom end.
- (e) Suppose $\phi = \pi/3$ and that at some instant the bottom end of the ladder is three times as far from the trunk's base as the top end. What can you say about the speeds of the ends of the ladder at this instant?
- (f) Still supposing $\phi = \pi/3$, find all possible angles the ladder makes with the ground when the top end of the ladder is moving four times as fast as the bottom end. Characterize each angle in one of two ways:
 - (i) One end of the ladder is moving toward the trunk's base and the other end away from the base.
 - (ii) Both ends of the ladder are either moving toward or away from the trunk's base.

Give exact angles. Then use a calculator to approximate their radian measures to the nearest hundredth.

(g) Use the animation below to check that your answers to parts (e) and (f) are reasonable. The graph is of the relation

Desmos link: <https://www.desmos.com/calculator/58jqjo0inl>

Desmos activity available at 151: *Tree and Ladder*

Question 332 A tree leans precariously with its trunk making an angle of ϕ radians with the ground. One end of a ladder with length L feet leans against the trunk, the other rests on the horizontal ground. You slide the bottom end of the ladder along the ground at random.

Suppose at some instant the ladder makes an angle of β_0 radians with the trunk and that the bottom end is s_0 feet from the trunk's base. Suppose also that the ladder is rotating at the rate of ω_0 rad/sec at this same moment.

- (a) Find an expression for rate of change in the distance from the bottom end of the ladder to the base of the trunk at this time.
- (b) Find an expression for rate of change in the distance from the top end of the ladder to the base of the trunk at this time.
- (c) Find expressions for the speeds of the ends of the ladder at this time.

Hint: Use the law of sines in $\triangle BOT$ in the demonstration from the previous question.

Question 333 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere.

Geogebra link: <https://www.geogebra.org/m/vnhrutwu>

Geogebra activity available at 151: *Declination of Sun 2*

(a) Use the graph of the function f below (at latitude $\phi \sim 1.1$) to sketch a graph of the derivative

$$\frac{dh}{d\delta} = f'(\delta).$$

(b) Suppose the latitude ϕ is held constant and find an expression for the derivative

$$\frac{dh}{d\delta} = \frac{d}{d\delta} (-\tan \phi \tan \delta).$$

(c) What are the units of the derivative in part (b)?

(d) Input your expression for the derivative in Line 4 of the worksheet below (follow the directions there). Then vary the slider ϕ to see how the function f and its derivative vary with latitude. Summarize your observations.

(e) Find an expression for the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0} = \left. \frac{d}{d\delta} (-\tan \phi \tan \delta) \right|_{\delta=0}$$

in terms of the latitude ϕ .

(f) Evaluate the derivative in part (e) at a latitude of $\phi = \pi/4$. Interpret its meaning in terms of small changes.

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at [151: Length of Day 1](#)

(g) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

Find an expression for a function

$$\delta = k(t), \quad t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

(h) Find a function

$$r = g(\phi), \quad -\pi/2 < \phi < \pi,$$

that expresses the rate of change in the number of minutes of daylight per day (measured in (minutes of daylight/day)/day) on the spring equinox in terms of the latitude ϕ . Input this function in Line 1 of the demonstration below.

(i) Use the result of part (f) to approximate the rate in (hours of daylight/-day)/day at which the number of daylight hours per day is changing at a latitude of $\phi = \pi/4$ radians on the spring equinox and on the fall equinox.

(j) Evaluate the rates from part (i) in Fairbanks, Alaska, latitude 64.8°N .

Desmos link: <https://www.desmos.com/calculator/nf8n5uphhl>

Desmos activity available at 151: Length of Day 2

Waves

Question 334 The function

$$y = f(x, t) = a \sin(kx - \omega t), t \geq 0, \quad (3)$$

describes a wave on a string. The functions expresses the displacement (in meters) of a point on the string in terms of the position x (in meters) of the point and time t , measured in seconds since the motion began.

(a) Experiment with the sliders k , ω , in the demonstration below, playing the slider u (u is just another name for t). Summarize your observations. In particular, be sure to turn off Line 1 to be better able to see the motion of the individual points of the string.

(b) What are the units of k and ω ? How do you know?

(c) Find an expression for the wavelength λ in terms of k , ω .

(d) Find an expression for the period of oscillation T in terms of k , ω .

(e) Hold y constant and differentiate each side of equation (3) with respect to t to find an expression for the speed of the wave. Turn on the graph in Line 9. Explain the logic behind the computation.

Desmos link: <https://www.desmos.com/calculator/9xmkg9hwi>

Desmos activity available at 151: Traveling Wave 1

The Ladder and the Tree

Implicit differentiation.

Question 335 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground. We analyze how a small change in the distance between the bottom of the ladder and the base of the trunk changes the distance between the top of the ladder and the base of the trunk.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/u8bxskxyhl>

151: Ladder and Tree 20

We'll let u be the distance OT between the top of the ladder and the base of the trunk (measured in feet) and s the distance OB between the bottom of the ladder and the base of the trunk (also measured in feet).

We'll focus on the particular position of the ladder when the bottom end is 5 feet from the trunk's base and the top end is 8 feet from the base as illustrated above.

- (a) Use the slider s approximate the derivative

$$\left. \frac{ds}{du} \right|_{(s,u)=(5,8)}.$$

Be sure to include units.

- (b) Find the length of the ladder
- (c) Find an equation that relates s and u . Then activate the Folder in Line 20. Explain the significance of the curve and the slope of the tangent line to the curve at P .
- (d) Find an expression for the derivative ds/du .
- (e) Use your expression for ds/du to evaluate the derivative

$$\left. \frac{ds}{du} \right|_{(s,u)=(5,8)}.$$

Learning outcomes:
Author(s):

- (f) Interpret the meaning of the derivative using the language of small changes.
- (g) Approximate the distance of the bottom of the ladder from the base of the tree when the top of the ladder is 7.95 feet from the tree's base.

Question 336 This is a continuation of the previous question.

- (a) Use the slider s to approximate the distance $s = OB$ when moving the top of the ladder a small distance away from the base of the tree makes the bottom end move twice as far toward the base. Then use calculus and algebra to approximate that distance.
- (b) Activate the folder in Line 31 of the worksheet above. Let θ be the angle (measured in radians) the ladder makes with the ground as marked. Take $\theta = 2\pi/3$ when the ladder lies along the tree.

- (i) Use the slider s to approximate the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=5}.$$

Include units.

- (ii) Write an equation relating θ and s .
- (iii) Find an expression for the derivative $ds/d\theta$ and use it to compute the exact value of the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=5}.$$

- (iv) Interpret the meaning of the above derivative using the language of small changes.
- (v) Find the exact value of the derivative

$$\left. \frac{d\theta}{du} \right|_{s=5}.$$

Statuary Hall and Implicit Differentiation

Statuary Hall and John Quincy Adams

Question 337 YouTube link:

https://www.youtube.com/watch?v=watch?v=FX6rUU_74kk

Statuary Hall

Up until the middle of the 19th century, Statuary Hall in the U.S. Capitol was the meeting place of the House of Representatives. The room is in the shape of an ellipse and John Quincy Adams had his desk at one of the foci. Legend has it that he was able to eavesdrop on the whispered conversations of his political opponents when they were standing at the other focus.

The aim of this problem is to explain this and prove the reflective property of ellipses.

A light wave (or a sound wave) emitted from one focus of an ellipse passes through the other focus after reflecting off the ellipse.

We need to know the definition of an ellipse:

An ellipse is the set of points, the sum of whose distances from two fixed points (the foci) is constant.

Desmos link: <https://www.desmos.com/calculator/6kxtojk72r>

Access Desmos interactives through the online version of this text at

151: Statuary Hall.

In Calculus 3 you'll learn about the gradient vector and a short way to prove the reflective property of ellipses in general. But we'll work with a specific example, and take our ellipse to have focal points $F_1(0, 0)$ and $F_2(4, 0)$. We'll also suppose the ellipse passes through $P(0, 3)$. Our problem is to first find an equation of the normal line to the ellipse at P . Then we'll show that this line bisects angle F_1PF_2 .

- (a) Start by using the definition of the ellipse to write an equation of the ellipse.

Learning outcomes:
Author(s):

- (b) Use implicit differentiation to find the slope of the tangent line to the ellipse at $P(0, 3)$.
- (c) Find an equation of the normal line at P . Enter this equation on Line 16 in the worksheet above.
- (d) For a shorter way to describe the normal line, do the following.
 - (i) Use vector arithmetic to find a vector parallel to the normal line at P .
 - (ii) Use your vector from part (i) to parameterize the normal line at P .

Question 338 Another way to prove the reflective property of ellipse is related to an optimization problem.

We are given an ellipse (dotted) with focal points A and B , a point P on the ellipse, and the tangent line L to the ellipse at P , as shown below.

Desmos link: <https://www.desmos.com/calculator/ncyrefzwwg>

Access Desmos interactives through the online version of this text at

151: Minimization Property of Ellipses.

To prove the reflective property of the ellipse, it's enough to show that the marked angles at P are congruent.

To do this we temporarily forget about the ellipse and the point P . All that's left are the points A , B , and the line L . Now our optimization problem is this. Given points A and B , to find the point on the given line L that minimizes the sum

$$\text{dist}(P, A) + \text{dist}(P, B)$$

of the distances from P to the points A and B .

There are two key ideas.

- (a) The first is to show that the point P that minimizes the above sum lies on the given ellipse with focal points A and B that is tangent to L . To see why, drag the slider s in the worksheet above. Then explain the logic behind this claim.
- (b) The second is to forget about the ellipse and solve the optimization problem to show that the marked angles at P are congruent for the point P on the line that minimizes the above sum. Try this.

The Ladder and the Tree, Part 2

Implicit differentiation.

Question 339 The bottom end of a seven-foot ladder slides across a horizontal floor as its top end slides down a vertical wall.

Desmos link: <https://www.desmos.com/calculator/4nmxshey0e>

151: Ladder and Tree 23

- (a) Let θ (in radians) be the angle the ladder makes with the ground and s the distance (in feet) between the bottom of the ladder and the wall.

Drag the slider s in the worksheet above to approximate the derivative

$$\left. \frac{d\theta}{ds} \right|_{s=3}.$$

Include units. The markings on the protractor are spaced 0.01 radians apart.

- (b) Suppose the bottom of the ladder is sliding toward the wall at a speed of 5 ft/sec when the bottom is 3 feet from the wall. At what rate is the ladder rotating at this instant?
- (c) Drag the slider s in Line 1 of the worksheet below to approximate the angle the ladder makes with the ground when its bottom end is moving twice as fast as its top end.
- (d) Find the exact angle in part (c).

Access Desmos interactives through the online version of this text at

.

Question 340 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground.

At the moment the bottom end of the ladder is 8 feet from the tree's base (in the position shown below), its bottom end is moving toward the tree's base at a

Learning outcomes:
Author(s):

speed of 5 ft/sec. At this same moment the top end of the ladder is 3 feet from the tree's base.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/xrgftlr9dx>

151: Ladder and Tree 29

- (a) Is the ladder rotating clockwise or counterclockwise at this moment? Guess first, then drag the slider s above to see if you were correct.
- (b) At what rate? Do not assume that the bottom end moves at a constant speed. Solve this problem twice. First, with implicit differentiation and then again without. Do not use a calculator.

Question 341 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a seven-foot ladder leans against the trunk, the other rests on the horizontal ground.

- (a) Find all possible angles the ladder makes with the ground when its bottom end is moving twice as fast as its top end.
- (b) Drag the slider ϕ_2 in Line 2 of the worksheet below to check your answer.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/oftz4vb9qj>

151: Ladder and Tree 22

Question 342 A tree leans precariously with its trunk making an angle of $\phi = \pi/3$ radians with the ground. One end of a ladder leans against the trunk, the other rests on the horizontal ground.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/bscbpblji1>

151: Ladder and Tree 21

- (a) Find a function

$$v_B = f(\theta), 0 \leq \theta \leq 2\pi/3,$$

that expresses the speed of the ladder's bottom end in terms of the angle $\theta = \angle OTP$ the ladder makes with the tree and the ladder's rotation rate ω (measured in rad/sec). Do not assume the rotation rate is constant.

- (b) Check that your expression for the speed has the correct units.

- (c) Similarly find a function

$$v_T = g(\theta), 0 \leq \theta \leq 2\pi/3,$$

that expresses the speed of the ladder's top end in terms of θ and ω .

- (d) Play the slider in Line 2 of the worksheet above. Then use the animation to sketch by hand graphs of the speed functions in parts (a) and (c) assuming the ladder rotates at a constant rate.
- (e) Activate the speed functions folder in Line 41 to check your graphs.
- (f) Find the ratio v_B/v_T of the speeds when the bottom end of the ladder is five feet from the tree's base.
- (g) What angle does the ladder make with the tree when the ladder's bottom end is moving twice as fast as its top end? Find all possibilities.
- (h) Answer part (g) when the tree is perpendicular to the ground instead.

More Related Rates Problems

Related rates.

Directions:

- Start by defining all variables, each in a complete sentence with units. Be sure to precisely define the variable for time as well.
- Answer each problem with a concluding sentence.

1 Shadow of a Building

Question 343 *On the vernal equinox at a location on the equator, the sun rises due east at 6am, passes directly overhead at noon, and sets due west at 6pm.*

151: Building Shadow

Desmos link: <https://www.desmos.com/calculator/55pixnvvqz>

- (a) *Play the slider u in Line 2 of the worksheet to see how the length of building's shadow changes during the afternoon. Summarize your observations.*

Free Response:

- (b) *Use the slider in the worksheet above to approximate the relative rate of change (with respect to time) in the length of the shadow when the shadow's length is twice the building's height*
- (c) *Find a function*

$$s = f(\theta), 0 < \theta \leq \pi/2,$$

that expresses the length of the shadow cast by an h -foot tall building in terms of the angle of elevation of the sun (this is the angle the sun's rays may with the ground).

- (d) *Find the relative rate of change (with respect to time) in the length of the shadow when the shadow's length is twice the building's height and the sun is setting. How does this compare with your estimate from part (b)?*

Learning outcomes:
Author(s):

- (e) During the afternoon, when is the length of the shadow increasing at its minimum relative rate? Justify your assertion and find this rate.

2 Kite Flying

Question 344 A kite drifts due east as it maintains a constant altitude. Play the slider u in Line 1 of the worksheet below to watch the motion.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/qdrzdt4erc>

151: Kite

- Which do you think is greater, the rate at which the string is being let out or the speed of the kite? No computations, just go with your intuition.
- Express the speed of the kite (not assumed constant) in terms of the rate at which the string is being let out (not assumed constant) and the angle θ the string makes with the horizontal.
- What can you say if $\theta = \pi/3$?
- Was your intuition correct?

3 A Crawling Beetle

Question 345 As a beetle crawling in the xy -plane passes the point $(-4, -3)$ (coordinates in centimeters) its distance from the x -axis is increasing at the rate of 2 cm/sec and its distance from the origin is increasing at the rate of 3 cm/sec.

Is the beetle rotating clockwise or counterclockwise at this instant? At what rate?

4 A Mechanical Motion

Question 346 As a rod OP of length R meters rotates about its endpoint O , the other endpoint P drags along a second rod PA of length L meters as illustrated below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/r20xwnok5r>

151: Engine

- (a) Express the speed of A in terms of R , L , the angle θ shown above, and the rotation rate ω (measured in radians/sec) of rod OP . Do not assume ω is constant. Work in general, not with the specific values of L and R in the worksheet.
- (b) Check that your expression in part (a) has the correct units.
- (c) Express the signed rotation rate of PA in terms of the same parameters. Measure the rate to be positive when PA rotates counterclockwise. Do not assume ω is constant.
- (d) Check that your expression in part (c) has the correct units.

5 Tracking a Helicopter

Sensors on the ground four hundred feet apart track a helicopter. At the instant shown below, the sensors at A and B are rotating counterclockwise at the respective rates of 2 rad/min and 3 rad/min. And the marked angles at A and B have respective measures $\angle A = \pi/6$ and $\angle B = \pi/3$.

Is the helicopter ascending or descending at this instant? At what rate?

Desmos link: <https://www.desmos.com/calculator/xl8t3toppg>

151: Tracking a Helicopter

6 Distance to the Horizon

Question 347 The distance to the horizon is limited by the curvature of the earth as illustrated in the demonstration below. This distance is the length of the (red) arc AT on the earth's surface.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/8qwt6mfirt>

Desmos activity available at

151:Distance to Horizon 11

Let

$$s = f(h)$$

be the function that expresses the distance to the horizon (in thousands of miles) in terms of your altitude above the surface (in thousands of miles).

- Sketch by hand a graph of the function f . Drag the slider h in the worksheet above to help with your sketch. Then activate the Graph folder in Line 17 to see how you did.
- Find an expression for the function f .
- Use the graph of the function f to approximate the derivative

$$\left. \frac{ds}{dh} \right|_{h=1}.$$

Include units.

- Use your expression for the function $s = f(h)$ to find the exact value of the derivative above.
- Interpret the meaning of the derivative in terms of small changes.
- Suppose at some instant you are in a rocket 1000 miles above the earth and descending at the rate of v miles/hour. Find an expression (in terms of v) for the rate (in miles/hour) at which your distance to the horizon (the length of the red arc AT above) is changing at this instant. Do not assume v is constant.

7 Jar Lid and Rubber Band

Question 348 You wrap a rubber band around a circular jar lid and pull the band tight as illustrated below.

Desmos link: <https://www.desmos.com/calculator/jbku3wrtdq>

151: String and Jar Lid

Suppose at some instant you are pulling the point P of the rubber band directly away from the center of the lid at a speed of v inches/sec. Express the rate (in inches/sec) at which the length of the rubber band is changing at this instant in terms of v and the marked angle θ between the straight segments of the band at P .

Hint: Express the length of the rubber band in terms of θ and the distance h (measured in inches) between P and the center of the lid. Keep in mind the band has a curved section in addition to the straight parts.

8 Disco Dancing

Question 349 A spotlight in a circular dance hall of radius r meters is located b meters from the center of the hall.

At some instant the light is rotating at the rate of ω rad/sec.

Desmos link: <https://www.desmos.com/calculator/m2o267u9ur>

151: Disco Dancing

- (a) Find an expression for the speed of the light beam as it moves along the wall at this instant. Your expression should be in terms of ω , r , b , and the angle θ marked below.
- (b) Check that your expression in part (a) has the correct units.

The Natural Log Function

Log functions and their derivatives.

1 The Derivative of the Natural Log Function

Question 350 (a) *Explain the meaning of*

$$\ln 5 = \log_e 5.$$

(b) *Simplify the function*

$$f(x) = e^{\ln x}.$$

Include the appropriate domain.

(c) *Use the result of part (b) to find an expression for the derivative*

$$\frac{d}{dx}(\ln x).$$

Explanation. To compute the derivative in part (c), we know that since the functions $f(x) = \ln x$ and $g(x) = e^x$ are inverses of one another,

$$e^{\ln x} = x.$$

Then differentiate both sides of this equation with respect to x to get

$$\frac{d}{dx}(e^{\ln x}) = \frac{dx}{dx}.$$

And by the chain rule we can rewrite this equation as

$$(e^{\ln x}) \frac{d}{dx}(\ln x) = 1.$$

And since $e^{\ln x} = x$,

$$\frac{d}{dx}(\ln x) = 1/x.$$

Learning outcomes:
Author(s):

Question 351 Find simplified expressions for each of the following derivatives. Include an appropriate domain for each.

(a)

$$\frac{d}{dx} (\ln(x/2))$$

(b)

$$\frac{d}{dx} (\ln(x^2))$$

Question 352 Find an equation of the tangent line to the curve

$$y = 3 \ln \left(\frac{x^2 + 1}{x^3 + 1} \right)$$

at the point on the curve with x -coordinate $x = 1$.

Question 353 (a) Graph the function

$$y = f(x) = \ln|x|$$

by hand.

(b) Use the graph above to sketch (by hand) a graph of the derivative

$$y = \frac{d}{dx} (\ln|x|).$$

(c) Use the chain rule to compute the derivative

$$\frac{d}{dx} (\ln|x|).$$

Do this by noting that

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

and using the chain rule to compute

$$\frac{d}{dx} (-x).$$

- (d) A reflection about the y -axis takes the graph of a function $y = f(x)$ to the graph of a function $y = g(x)$. Describe the transformation that takes the graph of $y = f'(x)$ to the graph of $y = g'(x)$. Explain your reasoning.

Question 354 Find an equation of the tangent line to the curve

$$y = 3 \ln \left| \frac{x^2 + 1}{x^3 - 1} \right|$$

at the point on the curve with x -coordinate $x = -1$.

Question 355 Let $\mathcal{L}$ be the tangent line to the curve

$$y = \ln |\sec \theta|, \quad -\pi/2 < \theta < \pi/2,$$

at the point on the curve with $\theta = \pi/7$.

Find the acute angle $\mathcal{L}$ makes with the x -axis.

2 Relative Rates of Change

Question 356 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d}{dt} (\ln P) \right|_{t=4} = \left. \frac{d}{dt} (\ln(f(t))) \right|_{t=4} = \frac{3}{10}.$$

- What are the units of this derivative?
- Explain the meaning of the derivative with and without the language of small changes.

Question 357 At a certain instant the population of a country is decreasing at the rate of 3%/yr. At this same instant the country's national debt is increasing at the rate of 5%/yr.

- At what relative rate is the per capita (ie. per person) share of the national debt changing at this time? Hint: Use the idea of the previous question.

- (b) If the per capita share of the national debt is \$200,000/person at this time, at what (absolute) rate is the per-capita share of the debt changing at this instant?

Question 358 The function

$$G = f(v) = 40 - 0.08 \left(\frac{v}{2} - 25 \right)^2, \quad 25 \leq v \leq 65,$$

expresses the gas mileage of a car (in miles/gallon) in terms of its speed (in miles/hr).

- (a) Explain the meaning of the derivative

$$\frac{d}{dv} (\ln(f(v))).$$

Include units in your explanation. Also, what are the units of 25 in the above expression? How do you know?

- (b) Evaluate the above derivative at $v = 30$ and explain its meaning in terms of small changes.

Derivatives and the Shapes of Curves

First and second derivatives and their implications.

1 A Surge Function

Question 359 *The function*

$$C = f(t) = Ae^{-kt}, t \geq 0,$$

expresses the concentration of a drug (in mg/L) in terms of the number of hours since the drug was injected. Here A and k are positive constants.

- (a) *What are the units of A ? Of k ? Explain how you know.*
- (b) *Find an expression for the time when the concentration is a maximum.*
 - (i) *Justify your assertion without relying on a graph.*
 - (ii) *Check that your expression has the correct units.*
- (c) *Find expressions for the times when*
 - (i) *the concentration is increasing at its maximum rate.*
 - (ii) *the concentration is decreasing at its maximum rate.*

Justify your assertions without relying on a graph.

- (d) *Determine all time intervals during which*
 - (i) *the concentration is increasing at an increasing rate.*
 - (ii) *the concentration is increasing at a decreasing rate.*
 - (iii) *the concentration is decreasing at an increasing rate.*
 - (iv) *the concentration is decreasing at a decreasing rate.*

Justify your assertions without relying on a graph.

Learning outcomes:

Author(s):

2 Changes and Rates of Change, Relative Changes and Relative Rates of Change

Question 360 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{dP}{dt} \right|_{t=4} = 400,000.$$

- What are the units of this derivative?
- Explain the meaning of the derivative without using the language of small changes.
- Explain the meaning of the derivative using the language of small changes by completing the following sentence.
Between 4:00pm and 4:01pm the population ...

Question 361 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d}{dt} (\ln P) \right|_{t=4} = \left. \frac{d}{dt} (\ln(f(t))) \right|_{t=4} = \frac{3}{10}.$$

- What are the units of this derivative?
- Explain the meaning of the derivative without using the language of small changes.
- Explain the meaning of the derivative using the language of small changes by completing the following sentence.
Between 4:00pm and 4:01pm the population ...

Question 362 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d^2 P}{dt^2} \right|_{t=4} = -6,000. \quad (4)$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.
- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.
Between 4:00pm and 4:01pm ...

- (d) Suppose also that

$$\left. \frac{dP}{dt} \right|_{t=4} = 400,000.$$

Explain the meaning of the second derivative above without using the language of small changes by completing the following sentence.

At 4:01pm ...

Question 363 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d^2(\ln P)}{dt^2} \right|_{t=4} = -\frac{3}{25}. \quad (5)$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.
- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.
Between 4:00pm and 4:01pm ...

- (d) Suppose also that

$$\left. \frac{d}{dt}(\ln P) \right|_{t=4} = \left. \frac{d}{dt}(\ln(f(t))) \right|_{t=4} = \frac{3}{10}.$$

Explain the meaning of the second derivative above without using the language of small changes by completing the following sentence.

At 4:01pm ...

Question 364 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\frac{d}{dt} \left(\ln \left| \frac{d}{dt} (\ln P) \right| \right) \Big|_{t=4} = -\frac{3}{5}. \quad (6)$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.
- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.

Between 4:00pm and 4:01pm ...

- (d) Suppose also that

$$\frac{d}{dt} (\ln P) \Big|_{t=4} = \frac{d}{dt} (\ln(f(t))) \Big|_{t=4} = \frac{3}{10}.$$

Explain the meaning of the second derivative above without using the language of small changes by completing the following sentence.

At 4:01pm ...

Optimization

Optimization.

Question 365 Use calculus to find an expression for the x -coordinate of the turning point of the parabola

$$y = ax^2 + bx + c, \quad c \neq 0.$$

Question 366 Determine the minimum and maximum values of the function

$$y = f(x) = x^3 - 9x, \quad -3 \leq x \leq 0.$$

Question 367 The function $G = f(v)$ graphed below expresses the gas mileage (in miles/gal) of a car in terms of its speed (in miles/hour).

Desmos link: <https://www.desmos.com/calculator/bcj0k1fymu>

151: Burning Gas 2

- Use the graph to determine the rate (in gal/hr) at which the car burns gas at a speed of 50 miles/hour.
- Use the slider v to sketch by hand a graph of the function $r = g(v)$ that expresses the rate (in gal/hour) at which your car burns gas in terms of its speed (in miles/hour). Then activate the folder in Line 8 to see how you did.
- Drag the slider v in Line 1 to approximate the speeds between 35 miles/hour and 70 miles/hour at which the car burns gas at the maximum and minimum rates (measured in gal/hr). Explain your reasoning.
- Now suppose

$$G = f(v) = -\frac{v^2}{20} + 5v - 90, \quad 35 \leq v \leq 70.$$

Use calculus to determine the exact speeds in part (c).

Learning outcomes:
Author(s):

Question 368 The function

$$P = f(t) = 5 - 3t + t^2, 0 \leq t \leq 4,$$

expresses the price in \$/share of a stock in terms of the number of hours past 9am.

- (a) Use the graphs of the function $P = f(t)$ and the function $r = f'(t)/f(t)$ to estimate when the stock price is increasing at the greatest relative rate.
- (b) Use calculus and algebra to find the exact time when the stock price is increasing at the greatest relative rate. Justify your assertion.

Desmos link: <https://www.desmos.com/calculator/y78fnny7s3>

Desmos activity available at 151: Stock Price 4

Question 369 The function

$$P = f(t) = t^2 - 4t + 8, 3 \leq t \leq 6,$$

expresses the price (in dollars/share) of a stock in terms of the number of hours past noon.

- (a) Use calculus, not precalculus, to show that the stock price increases between 3pm and 6pm.
- (b) When is the stock price increasing at the maximum relative rate? When is it increasing at the minimum relative rate?
- (c) Find the minimum and maximum relative rates of change in the stock price.
- (d) Justify your answers to parts (b) and (c). Explain your reasoning in complete sentences.
- (e) Preface each computation block with a brief description of what you are about to compute.

Question 370 A vertical wall b feet high runs parallel to a tall building. The wall is a feet from the building. A ladder reaching from the ground to the building rests on the top of the wall as shown below.

Desmos link: <https://www.desmos.com/calculator/4ak46ub8ay>

151: Shortest Ladder

- (a) Find a function that express the length of the ladder (measured in feet) in terms of the angle the ladder makes with the ground. Include an appropriate domain.
- (b) Use part (a) to express the length of the shortest such ladder in terms of a and b . Justify your assertion.
- (c) Check that your expression has the correct units.

Work in general and not with the particular values of a and b in the worksheet above.

Question 371 The bottom and top edges of a painting are respectively a and b feet above eye level.

Desmos link: <https://www.desmos.com/calculator/dkqndsegod>

151: Viewing Angle

- (a) Find a function that expresses the viewing angle in terms of your distance from the painting. Include an appropriate domain.
 - (b) How far from the painting should you stand to maximize the viewing angle marked above? Justify your assertion.
 - (c) Check that your expression has the correct units.
-

Measuring Distances on the Earth

Shortest paths on the earth.

1 Trigonometry: Along a Circle of Latitude

Question 372 Pick points A and B at the same latitude ϕ on the earth's surface and let θ (measured in radians) be the difference in their longitudes, with $0 < \theta \leq \pi$.

Desmos3D link: <https://www.desmos.com/3d/hhpog6ijnr>

151: Distances on Earth

The purpose of this problem is to compare two distances in traveling from A to B , one along their common circle of latitude, the other along the great circle through A and B . We'll assume the earth is a sphere of radius R miles.

To compute each distance, we need to see inside the earth. You can do this by deactivating the Sphere folder in Line 3.

- (a) First the distance along the circle of latitude.
 - (i) Start by expressing the radius of the circle of latitude in terms of the latitude ϕ and the radius of earth R . Work in general, not with the specific values in the worksheet above.
 - (ii) Then express the distance between A and B along the circle of latitude in terms of ϕ , R , and θ .
- (b) Next we'll compute the distance along the great circle through A and B . The center of this circle coincides with the center of the sphere.
 - (i) Try to do this on your own. Here are two triangles (from inside the sphere as illustrated above) shown in two dimensions to help.

Desmos link:

<https://www.desmos.com/calculator/wkdhhbojii>

151: Distances Earth 2D

Learning outcomes:

Author(s):

- (c) Use the results of parts (a) and (b) to compare these two distances between San Francisco (latitude 38°N , longitude 122°W) and Washington DC (latitude 38°N , longitude 77°W). Take the radius of the earth to be 3960 miles.
- (d) Find two other locations at approximately the same latitude and compare the two distances.

2 Calculus: Along a Circle of Latitude

This is a continuation of the previous problem.

The question is this: Fix the difference in longitude θ between two points at the same latitude and determine an expression for the latitude at which the positive difference in the two distances (one along the circle of latitude, the other along the great circle through the points) is a maximum.

Desmos link: <https://www.desmos.com/calculator/muhahdawza>

151: Distances Earth 2D Part 2

The difference in the two distances is a maximum at latitude

$$\phi = \arccos \left(\sqrt{\csc^2 \left(\frac{\theta}{2} \right) - \frac{4}{\theta^2}} \right).$$

Disco Dancing, an Optimization Problem

Revolving light in a dance hall.

Question 373 A spotlight in a circular dance hall of radius r meters is located b meters from the center of the hall.

The light rotates at the constant rate of ω rad/sec.

Desmos link: <https://www.desmos.com/calculator/vtzsgtxswm>

151: Disco Dancing 25

- (a) Find an expression for the speed of the light beam as it moves along the wall at this instant. Your expression should be in terms of ω , r , b . Hint: Use the law of sines in $\triangle LOB$ to first find an expression for the measure of angle $\angle OLB$.
- (b) Check that your expression in part (a) has the correct units.
- (c) Activate the Distance Function folder in Line 27 of the worksheet above. Then try to guess the position of the spotlight on the wall when its speed is increasing at the greatest rate.
- (d) Determine the position of the spotlight on the wall when its speed is increasing at the greatest rate.

Clock Hands

A twist on a familiar problem.

Question 374 The question is to determine the angle between the minute and hour hands of a clock when the distance between the tips of the hands is increasing at the fastest rate.

We'll assume the hour and minute hands have respective lengths a cm and b cm.

Desmos link: <https://www.desmos.com/calculator/efhex45zkr>

151: Hands of a Clock 3

- (a) Play the slider u in Line 2 above and reconcile the graph of the function $c = f(t)$ that expresses the distance between the hands (in cm) in terms of the number of minutes past noon.
- (b) Change the lengths of the hands in Lines 4 and 6 to see how the graph of the function f changes.
- (c) The distance function looks sinusoidal, but open the folder in Line 26 to see the graph of the derivative dispels that notion.
- (d) Find an expression for the function

$$c = h(\theta), 0 \leq \theta \leq 2\pi,$$

that expresses the distance between the tips of the hands in terms of the angle between the hands. Work in general with the parameters a and b , not with their specific values in the worksheet.

- (e) Find an expression for the derivative

$$\frac{dc}{d\theta} = \frac{d}{d\theta}(h(\theta)).$$

Include units.

- (f) Use your expression for the derivative $dc/d\theta$ to find an expression for the derivative

$$\frac{dc}{dt} = \frac{d}{dt}(f(t)).$$

Learning outcomes:
Author(s):

- (g) Use your expression for the derivative $dc/d\theta$ to determine the angle between the hands when the distance between their tips is increasing at the fastest rate. There are three cases to consider.
- (i) the minute hand is longer than the hour hand ($b > a$)
 - (ii) the hour hand is longer ($a > b$)
 - (iii) the hands have the same length ($a = b$)
- (h) What can you say about the angle between the shorter hand and the segment AB when the distance is increasing at the fastest rate?

A Computational Solution

Question 375 Determine the angle between the hour and minute hands of a clock when the distance between the tips of these hands is increasing at the fastest rate. Suppose the hour and minute hands have respective lengths a and b inches, with $a < b$.

Explanation. We'll let a and b denote the respective lengths of the hour and minute hands, θ the angle (in radians) between the hands, and c the distance. Since the hands each turn at a constant rate, our aim is to maximize the derivative

$$\frac{dc}{d\theta} = \frac{ab \sin \theta}{c} = \frac{ab \sin \theta}{\sqrt{a^2 + b^2 - 2ab \cos \theta}}.$$

Differentiating again and doing some algebra gives

$$\frac{d^2c}{d\theta^2} = \frac{ab}{c^3} (c^2 \cos \theta - ab \sin^2 \theta).$$

Then setting the second derivative equal to zero and substituting

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

leads to the equation (symmetric in a and b)

$$ab \cos^2 \theta - (a^2 + b^2) \cos \theta + ab = 0$$

or

$$(a \cos \theta - b)(b \cos \theta - a) = 0.$$

Assuming the minute hand is longer than the hour hand ($b > a$) and $0 \leq \theta \leq 2\pi$, we conclude that the distance is increasing at the fastest rate when $\theta = \arccos(a/b)$ and decreasing at the fastest rate when $\theta = 2\pi - \arccos(a/b)$.

Question 376 (a) What can you say if $a < b$? If $a = b$?

(b) Supposing $a \neq b$, find an expression for the maximum rate at which the distance between the tips of the hands is increasing.

A More Geometrical Solution

We could have saved ourselves the trouble of taking the second derivative by using the law of sines in $\triangle ABC$. Then

$$\left| \frac{dc}{d\theta} \right| = \frac{ab \sin C}{c} = \frac{ab \sin B}{B} = a \sin B$$

so that the rate of change is maximized when $B = \pi/2$ or when $\theta = \arccos(a/b)$ as above.

A Solution with Vectors

Mars and the Outer Planets

The retrograde motion of Mars and the outer planets.

Exploration 377 YouTube link:

<https://www.youtube.com/watch?v=https://www.youtube.com/watch?v=1nVSzzYCAYk>

Retrograde Motion of Mars

Exploration 378 Assume for this problem that the planets rotate about the sun at constant rates in circular orbits that lie in the same plane. Suppose also that earth has an orbital period of 365 days and an orbital radius of 1 astronomical unit.

Mars has an orbital period of Mars to be 687 earth days and an orbital radius of 1.524 astronomical units. Earth has an orbital period of 365 days and an orbital radius of 1 astronomical unit.

- (a) Parameterize the motions of earth and mars about the sun in terms of the number of earth days. Assume
 - The sun is at the origin.
 - The planets rotate counterclockwise at constant rates about the sun.
 - The earth has coordinates $(1, 0)$ a time $t = 0$ days.
 - The vector from the sun to Mars has polar angle θ_0 at time $t = 0$.
- (b) Express the position of Mars relative to Earth in terms of the vectors $\overrightarrow{SE}$ and $\overrightarrow{SM}$ that respectively give the positions of Earth and Mars relative to the sun.
- (c) Use part (b) to parameterize the motion of mars relative to the earth (with the earth fixed at the point $(0, -4)$).
- (d) Check your work using the demonstration below by inputting the correct functions in Lines 3, 4, 6, 7, 9, and 10.

Access Desmos interactive at

[Mars Retrograde Motion](#)

Learning outcomes:
Author(s):

Desmos link: <https://www.desmos.com/calculator/htc4xgrjxs>

- (e) Compute the synodic period of Mars as seen from Earth. This is the time, measured in earth days, it takes Earth to “lap” Mars. It is the time it takes Mars to return to the same position relative to the sun as seen from Earth.
- (f) Sketch by hand graphs of the x -coordinate functions for the motions of Earth and Mars on the same coordinate system. Label the exact coordinates of at least two turning points on each graph. Label the axes with the appropriate variable names and units.
- (g) Sketch by hand graphs of the y -coordinate functions for the motions of Earth and Mars on the same coordinate system. Label the exact coordinates of at least two turning points on each graph. Label the axes with the appropriate variable names and units.

Access Geogebra interactive at
[Mars Retrograde Motion](#)

Geogebra link: <https://www.geogebra.org/m/addm38j6>

Hours of Daylight

Modeling the number of hours of daylight.

Question 379 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere. The graph of the function $h = f(\delta)$ for latitude $\phi \sim 1.1$ is shown below.

Don't be confused by the angle δ (delta). It's just the lower case Greek letter for Δ .

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at 151: Length of Day 1

- (a) Use the graph to approximate the number of hours of daylight/day at a latitude of $\phi = 1.1$ radians on March 1, when the sun is about 8° below the plane of the equator. What about at our latitude? At the equator?

- (b) Evaluate the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0}.$$

- (c) What are the units of the derivative above? Interpret its meaning.

- (d) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

- (i) Find an expression for a function

$$\delta = k(t), \quad t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

Learning outcomes:

Author(s):

- (ii) Evaluate the derivative

$$\left. \frac{d}{dt} (f(k(t))) \right|_{t=0} = \left. \frac{dh}{dt} \right|_{t=0}.$$

- (iii) What are the units of the derivative above? Interpret its meaning.

Question 380 The function

$$h = f(\delta) = \frac{24}{\pi} \arccos(-\tan \phi \tan \delta), \quad -\pi/2 + \phi < \delta < \pi/2 - \phi,$$

expresses the number of hours of daylight per day at latitude ϕ in terms of the declination of the sun. The declination of the sun (δ) is the angle the sun's rays make with the plane of the equator, taken to be positive between the spring and fall equinoxes in the northern hemisphere. The latitude ϕ , $-\pi/2 \leq \phi \leq \pi/2$, is positive at points in the northern hemisphere.

Geogebra link: <https://www.geogebra.org/m/vnhrutwu>

Geogebra activity available at 151: Declination of Sun 2

- (a) Use the graph of the function f below (at latitude $\phi \sim 1.1$) to sketch a graph of the derivative

$$\frac{dh}{d\delta} = f'(\delta).$$

- (b) Suppose the latitude ϕ is held constant and find an expression for the derivative

$$\frac{dh}{d\delta}$$

- (c) What are the units of the derivative in part (b)?
- (d) Input your expression for the derivative in Line 4 of the worksheet below (follow the directions there). Then vary the slider ϕ to see how the function f and its derivative vary with latitude. Summarize your observations.
- (e) Find an expression for the derivative

$$\left. \frac{dh}{d\delta} \right|_{\delta=0}$$

in terms of the latitude ϕ .

- (f) Evaluate the derivative in part (e) at a latitude of $\phi = \pi/4$. Interpret its meaning in terms of small changes.

Desmos link: <https://www.desmos.com/calculator/ifomatkcta>

Desmos activity available at 151: *Length of Day 1*

- (g) Suppose now that each month has 30 days so that there are 360 days in one year. Suppose also that the declination of the sun varies sinusoidally as a function of time, that the maximum declination of $\delta = 23.5^\circ$ occurs on the summer solstice (say June 21st) and the minimum declination $\delta = -23.5^\circ$ occurs on the winter solstice (December 21st).

Find an expression for a function

$$\delta = k(t), t \geq 0,$$

that gives the declination of the sun (measured in radians) in terms of the number of days since the spring equinox.

- (h) Find a function

$$r = g(\phi), -\pi/2 < \phi < \pi,$$

that expresses the rate of change in the number of hours of daylight per day (measured in (hours of daylight/day)/day)) on the spring equinox in terms of the latitude ϕ . Input this function in Line 1 of the demonstration below.

- (i) On the spring equinox, all latitudes receive 12 hours of daylight/day. Use the result of part (h) to approximate how many extra minutes of daylight we would get on the following day living in

(i) Shoreline, latitude $47.75^\circ N$

(ii) Fairbanks, Alaska, latitude $64.8^\circ N$

Desmos link: <https://www.desmos.com/calculator/nf8n5uphhl>

Desmos activity available at 151: *Length of Day 2*

Motion

An introduction to motion.

Question 381 Play the slider u in the activity below to see the motion of a balloon. Use the animation to sketch graphs of

- (a) the altitude of the balloon as a function of time.
- (b) the balloon's rate of ascent as a function of time.
- (c) Activate the folders in Lines 13 and 18 to see how you did.

Desmos link: <https://www.desmos.com/calculator/amv52b9ljt>

Desmos activity available at 151: Balloon

Question 382 The function

$$h = f(t) = 10 - \frac{1}{2}t^2e^{-t/5}, \quad 0 \leq t \leq 40,$$

expresses the altitude of a balloon (in thousands of feet) in terms of the number of hours since noon on August 31, 2023.

Use the graph of this function in Question 1 to first approximate answers to the following questions. Then use calculus and algebra to determine the exact times.

- (a) When is the balloon at its minimum height? At its maximum height?
- (b) When is the balloon ascending at the fastest rate? Descending at the fastest rate?

To compute these times you will end up solving a quadratic equation that may be written in the form

$$t^2 - 20t + 50 = 0.$$

So the balloon is ascending at its fastest rate at time (give the exact times and then approximations to the nearest hundredth of an hour)

$$t = 10 + \sqrt{50} \sim 17.07$$

Learning outcomes:
Author(s):

hours past noon and descending at its fastest rate at time

$$t = 10 - \sqrt{50} \sim 2.93$$

hours past noon.



Linear Approximation

The derivative as a scaling factor.

1 The Derivative as a Scaling Factor

We started this class by interpreting the derivative as a dimensionless, local stretching factor of a rubber band. Perhaps stretching factor was not the best description because a rubber band might get stretched in some regions but compressed in others.

As we saw many times, a better, more universal description would be to think of the derivative as a *local scaling factor*. Multiply a small change in the input to a function by the derivative (at that input) and the result is a good approximation to the change in the function's output. Good in the sense that the error in the approximation approaches zero faster than the change in the input.

This interpretation of the derivative is often more useful than interpreting the derivative as the slope of a tangent line or an instantaneous rate of change. For one, the idea of a slope does not generalize to higher dimensions. And if you pause to really think about things, you might find it hard to make precise sense of what exactly anyone means by an instantaneous rate of change.

The idea of linear approximation is not new. We've been talking about it all quarter. And you knew about linear approximation well before you took this class. There's no need to learn anything more, or to use a special formula. It's really just common sense.

Here's a simple example.

Example 57. The function

$$h = f(t), \quad 0 \leq t \leq 60,$$

expresses the altitude of the balloon (measured in feet) in terms of the number of minutes past noon.

Suppose

(a) $f(10) = 1500$ and

(b) $\left. \frac{dh}{dt} \right|_{t=10} = -50.$

Learning outcomes:
Author(s):

Approximate the height of the balloon at 12:08pm.

Explanation. We know at 12:10pm the balloon is descending at the rate of 50 ft/min. So during the time interval (backward in time) from 12:10pm to 12:08pm of

$$\Delta t = (8 - 10) \text{ min} = -2 \text{ min},$$

the change in the balloon's altitude is approximately

$$\begin{aligned} \Delta h &= f(8) - f(10) \\ &\sim \left(\left. \frac{dh}{dt} \right|_{t=10} \right) (\Delta t) \\ &= \left(-50 \frac{\text{ft}}{\text{min}} \right) (-2 \text{ min}) \\ &= 100 \text{ ft.} \end{aligned}$$

Since the balloon's altitude is 1500 feet at 12:10pm, its altitude at 12:08pm was approximately

$$\begin{aligned} f(8) &= f(10) + \Delta h \\ &\sim (1500 + 100) \text{ ft} \\ &= 1600 \text{ ft.} \end{aligned}$$

2 Interpreting the Derivative

2.1 Speed

A key idea in this class has been to interpret *speed* as a derivative. This has been a bit difficult, especially with the mindset that the derivative is always a rate of change. We typically don't think about speed this way.

But speed is a rate of change.

Speed is the rate of change, with respect to time, of the the function that records the cumulative distance traveled.

The image of a cumulative distance function easiest for me to understand is the odometer reading on a car. It records the distance the car has traveled since it came off the lot. The speedometer records the derivative of this function with respect to time (measured in hours).

Here's an example. Don't be confused by the variable s . It does not stand for speed. It's commonly used, as it is here, to measure distance. Or as we'll see later, and as you'll see in integral calculus and in your physics classes, for *signed distance*.

Don't be confused either by the variable $v = ds/dt$. It is *not* velocity, but rather speed. Speed is a scalar, velocity is a vector. Speed is the length of the velocity vector.

Example 58. The function

$$s = f(t) = 2t + 10t^2 - t^3, \quad 0 \leq t \leq 6,$$

expresses the mileage on the trip odometer reading in terms of the number of hours since the start of a car trip.

Desmos link: <https://www.desmos.com/calculator/t1ruocrgm4>

151: Odometer Reading 34

- (a) Evaluate the derivative

$$\left. \frac{ds}{dt} \right|_{t=2}$$

and interpret its meaning. Include units.

- (b) Interpret the meaning of the above derivative in terms a scaling factor and small changes.
- (c) Use the result of part (b) to approximate the reading on the odometer at 2:01pm.
- (d) Use the result of part (b) to approximate when the odometer reads 35 miles.
- (e) When is the speed of the car a maximum? Use the graphs above to approximate the time. Then find the exact time and maximum speed.

3 Weight and Distance

Example 59. The function

$$W = f(h) = \frac{2000}{(h + 4)^2}, \quad h \geq 4,$$

expresses the weight of an astronaut (in pounds) in terms of her distance above the earth's surface (measured in thousands of miles).

- (a) Evaluate the derivative

$$\left. \frac{dW}{dh} \right|_{h=1}.$$

Include units.

- (b) Interpret the meaning of the above derivative in terms a scaling factor and small changes.
- (c) Approximate the astronaut's weight at an altitude of 1010 miles.
- (d) At what approximate altitude does the astronaut weigh 126 pounds?

4 Speed and Height

Example 60. The function

$$v = f(h), 0 \leq h \leq 100,$$

expresses the speed of a rock (measured in meters/sec) dropped from rest on the planet Krypton in terms of its height above the surface (measured in meters).

- (a) Which of the following is more likely?

(i)

$$\left. \frac{dv}{dh} \right|_{h=60} = 0.5$$

or

(ii)

$$\left. \frac{dv}{dh} \right|_{h=60} = -0.5?$$

Explain your reasoning. Include units for the derivative.

- (b) Interpret the meaning of the more likely derivative in terms of a scaling factor and small changes.
- (c) Suppose that $f(60) = 40$.
 - (i) Interpret the meaning of this statement.
 - (ii) Assume the more likely derivative from part (a) and approximate the rock's speed at a height of 59.9 meters.
 - (iii) Suppose the speed of the rock increases at the constant rate of 20 (m/sec)/sec as it falls and find the rock's exact speed when it is 59.9 meters above the surface. Compare this with your estimate.

5 Distance to the Horizon

Example 61. An astronaut above the surface of a planet sees only a fraction of the surface as suggested by the figure below.

Desmos link: <https://www.desmos.com/calculator/8shf1msp4m>

151: Distance to Horizon 44

The visible part of the surface is a spherical disk with spherical radius BC above. We can think of this distance (an arc of a circle) as the distance to the horizon.

- (a) Find a function

$$s = f(h), \quad h \geq 0,$$

that expresses the distance to the horizon on a planet of radius R kilometers in terms of the altitude of the astronaut (in km) above the surface.

- (b) Evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=2R/3}.$$

Include units.

- (c) Explain the meaning of the derivative in terms of a scaling factor and small changes.
- (d) Approximate the change in the distance to the horizon when the altitude above the surface increases from $h = (2/3)R$ to $h = (0.01 + 2/3)R$.

6 Gas Mileage

Example 62. The one-to-one function

$$G = f(v), \quad 50 \leq v \leq 80,$$

expresses the gas mileage of a car (in miles/gal) in terms of its speed (in miles/hr).

- (a) Which of the following is more likely?

(i)

$$\left. \frac{dv}{dG} \right|_{v=70} = 2$$

or

(ii)

$$\left. \frac{dv}{dG} \right|_{v=70} = -2?$$

Explain your reasoning. Include units for the derivatives.

(b) Suppose $f(70) = 24$

- (i) Interpret the meaning of this statement.
- (ii) Assume the more likely derivative from part (a) and approximate the gas mileage at a speed of 73 miles/hour.
- (iii) Simplify the units of the derivative in part (a). What do these units suggest about its meaning? Is this correct? Explain.

Linear Approximation, Part 2

Linear approximation and relative changes.

1 Review of the Natural Log Function

Question 383 Find expressions for each of the following derivatives.

(a) $\frac{d}{dx}(\ln x)$

(b) $\frac{d}{dx}(\ln(4x))$

(c) $\frac{d}{dx}(\ln(-x))$

(d) $\frac{d}{dx}(\ln |x|)$

(e) $\frac{d}{dx}(\ln(x^5))$

(f) $\frac{d}{dx}(\ln(1 + x^2))$

2 First Derivatives

Question 384 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{dP}{dt} \right|_{t=4} = 400,000.$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.

Learning outcomes:
Author(s):

- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.

Between 4:00pm and 4:01pm the population ...

Question 385 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d}{dt} (\ln P) \right|_{t=4} = \left. \frac{d}{dt} (\ln(f(t))) \right|_{t=4} = \frac{3}{10}.$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.
- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.

Between 4:00pm and 4:01pm the population ...

Question 386 The function

$$G = f(s), \quad 0 \leq s \leq 130,$$

graphed below expresses the number of gallons of gas in a car in terms of the trip odometer reading (measured in miles).

Desmos link: <https://www.desmos.com/calculator/gdeybih5t2>

151: Relative Rates Gas

- (a) A reading on the car's dashboard shows the number of miles left to drive before running out of gas assuming the car continues to burn gas at the current rate. Drag the slide u (another name for s) in Line 1 to approximate this reading when the trip odometer reads 80 miles.
- (b) Drag the slider u in Line 1 to approximate the derivative

$$\left. \frac{d}{ds} \left(\ln \left(\frac{G}{1} \right) \right) \right|_{s=80}.$$

- (c) What are the units of the derivative above?

- (d) Explain the meaning of the derivative.
- (e) Approximate the derivative $d/ds(\ln G)$ for some other values of s . What do you notice? What does this suggest about the function f ?
- (f) In light of part (e), how would you describe what it means for a population of bacteria to decrease at the constant relative rate of 5%/hour? The rate is instantaneous.

3 Second Derivatives

Question 387 The function $P = f(t)$, $0 \leq t \leq 12$, expresses the population of a colony of bacteria (measured in the number of bacteria) in terms of the number of hours past noon.

Suppose that

$$\left. \frac{d^2 P}{dt^2} \right|_{t=4} = -6,000. \quad (7)$$

- (a) What are the units of this derivative?
- (b) Explain the meaning of the derivative without using the language of small changes.
- (c) Explain the meaning of the derivative using the language of small changes by completing the following sentence.
Between 4:00pm and 4:01pm ...
- (d) Suppose also that

$$\left. \frac{dP}{dt} \right|_{t=4} = 400,000.$$

Explain the meaning of the second derivative above without using the language of small changes by completing the following sentence.

At 4:01pm ...

Review of Differentiation

Derivative Review.

Examples

Example 63. Find an equation of the tangent line to the curve

$$y = f(x) = (2x^3 + 1)^2$$

at the point $(1, 9)$.

Explanation. Let

$$y = (2x^3 + 1)^2$$

and

$$u = 2x^3 + 1.$$

Then

$$y = u^2$$

and

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(u^2) \cdot \frac{d}{dx}(2x^3 + 1) \\ &= 2u(6x^2) \\ &= 2(2x^3 + 1)(6x^2). \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 2(3)(6) = 36,$$

and an equation of the tangent line is

$$y - 9 = 36(x - 1).$$

Example 64. Find an expression for the derivative

$$\frac{d}{d\theta}(\theta \cos(5\theta)).$$

Learning outcomes:
Author(s):

Explanation. We use the product rule first to get

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)).\end{aligned}$$

Now we use the chain rule to compute

$$\frac{d}{d\theta}(\cos(5\theta)).$$

For this we hide the composition by letting

$$y = \cos(5\theta)$$

and

$$u = 5\theta.$$

Then

$$y = \cos u$$

and by the chain rule

$$\begin{aligned}\frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du}(\cos u) \cdot \frac{d}{d\theta}(5\theta) \\ &= (-\sin u)(5) \\ &= -5 \sin(5\theta).\end{aligned}$$

The final result is that

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) - 5\theta \sin(5\theta).\end{aligned}$$

Here's a shorter version of the same solution.

We use the product rule first to get

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)).\end{aligned}$$

Then we use the chain rule to differentiate $\cos(5\theta)$, giving

$$\begin{aligned}\frac{d}{d\theta}(\theta \cos(5\theta)) &= \frac{d}{d\theta}(\theta) \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta \frac{d}{d\theta}(\cos(5\theta)) \\ &= \cos(5\theta) + \theta(-\sin(5\theta)) \frac{d}{d\theta}(5\theta) \\ &= \cos(5\theta) - 5\theta \sin(5\theta).\end{aligned}$$

Exercises

Directions: Follow the method of Example 1 *exactly* for each of the following problems.

Question 388 Find expressions for each of the following derivatives. Simplify your expressions for the derivatives. Do not simplify the function being differentiated.

Part 1:

- (a) $\frac{d}{dw}(\arctan w + \arctan(1/w))$
 (b) $\frac{d}{dy}(\arcsin(\sqrt{1-y^2}))$

Part 2:

Any comments or observations on the derivatives above?

Question 389 *Desmos link:*

<https://www.desmos.com/calculator/zjqsdhvvz4p>

Desmos activity available at 151: Building Temperature

Question 390 Water is poured into a cylindrical tank at a constant rate. At the same time, water flows out of a small hole in the bottom of the tank. The tank is empty at noon.

The function

$$t = f(h) = -\frac{2}{k_2} \left(\sqrt{h} + \frac{k_1}{k_2} \ln \left| \frac{k_1 - k_2 \sqrt{h}}{k_1} \right| \right), \quad t \geq 0,$$

expresses the time (measured in minutes past noon) in terms of the depth (measured in cm) of water in the tank. Here k_1, k_2 are positive constants.

- (a) Use the above function to verify that the tank is empty at noon.
- (b) Find a simplified expression for the derivative dt/dh .
- (c) Use the result of part (b) to show that

$$\frac{dh}{dt} = k_1 - k_2\sqrt{h}.$$

- (d) What are the units of k_1, k_2 ? How do you know?
- (e) For much more on this problem, see Questions 23 and 24 of the chapter Derivatives of Inverse Functions.

Question 391 The function

$$s = f(t) = Ae^{-k_1 t} \cos(k_2 t), \quad t \geq 0,$$

expresses the displacement (in meters) from equilibrium of an oscillating mass on a spring in terms of the number of seconds since the mass was released from rest.

- (a) What are the units of the constants A, k_1 , and k_2 ? Explain how you know.
- (b) Find an expression for the velocity ds/dt of the mass.
- (c) Find an expression for the acceleration

$$\frac{d}{dt} \left(\frac{ds}{dt} \right) = \frac{d^2 s}{dt^2}$$

of the mass.

- (d) Use algebra to show that

$$\frac{d^2 s}{dt^2} = - \left(2k_1 \frac{ds}{dt} + (k_1^2 + k_2^2)s \right).$$

Desmos link: <https://www.desmos.com/calculator/ygikqgj7af>

Desmos activity available at 151: Damped Harmonic Oscillator

Question 392 (a) Find a function

$$s = f(h), \quad h \geq 0,$$

that expresses the distance (in miles) to the horizon in terms of your altitude (in miles). We'll suppose the earth to be a perfect sphere of radius $R = 4000$ miles. The distance to the horizon is the arclength AT below, measured along the surface of the earth (you can think of this distance as the radius of the spherical disk visible to us). Our height is the distance AP .

The function is

$$s = f(h) = 4000 \arccos \left(\frac{4000}{4000 + h} \right), \quad h \geq 0.$$

Desmos link: <https://www.desmos.com/calculator/ewowig5sgk>

(b) Use the graph of the function $s = f(h)$ show below to sketch a graph of the derivative

$$y = \frac{ds}{dh} = f'(h).$$

by hand. What is the domain of the derivative? Note that the curve $s = f(h)$ has a horizontal asymptote. What is an equation of this asymptote?

Desmos link: <https://www.desmos.com/calculator/nof5l3mtfy>

Then check your sketch by activating the folder in Line 1 of the worksheet below.

Desmos link: <https://www.desmos.com/calculator/pdbjfao316>

(c) Find an expression for the derivative

$$\frac{ds}{dh} = \frac{d}{dh} \left(\arccos \left(\frac{R}{R + h} \right) \right).$$

Hint: Start by using the graphs of the functions $y = \arcsin x$ and $y = \arccos(x)$ below and your knowledge of the derivative $d/dx(\arcsin x)$ to find an expression for $d/dx(\arccos x)$. Or equivalently, recognize that

$$\arcsin x + \arccos x = \pi/2.$$

Desmos link: <https://www.desmos.com/calculator/zuq9rf1j4d>

(d) With $R = 4000$, evaluate the derivative

$$\left. \frac{ds}{dh} \right|_{h=16}.$$

(e) Interpret the meaning of the derivative in part (d) in terms of specific small changes.

(f) You take a ride on Blue Origin and in two minutes are boosted straight up to an altitude of 32 miles. Suppose that the function

$$h = g(t) = \begin{cases} 10t^2 + 6t^3, & 0 \leq t \leq 1 \\ 32 - 10(2 - t)^2 - 16(2 - t)^3, & 1 < t \leq 2, \end{cases}$$

expresses your altitude (in miles) in terms of the number of minutes since launch for the first two minutes of your flight.

(i) At what rate (with respect to time) is your distance to the horizon changing when you are one minute into the flight? Use the graph of the function $s = f(g(t))$ shown below to first approximate this rate.

(ii) At what rate (with respect to time) is your distance to the horizon changing at the start of the flight? Use the graph of the function $s = f(g(t))$ shown below to first approximate this rate. Are you surprised given your graph in part (b)?

Desmos link: <https://www.desmos.com/calculator/jchwjpxugf>

Final, Math 151

Directions for our final.

1 Directions

Here are the directions for our final.

- Use only the material of our class. In particular, *do not use vectors or other ideas that were not part of our class.*
- Use the Leibniz differential notation for derivatives and their evaluations. Do *not* use the prime notation.
- Use the Leibniz notation to show *all* steps when using the the chain, product, and quotient rules. See the next section *Examples of How to Show Work* for more details.
- Define, in complete sentences and with units any variables or unknowns you introduce.
- Include a domain for each function you introduce.
- Include units for *each* number in *each* numerical computation.
- Give explanations of your reasoning along with your solutions to each problem.
- Show all your work.
- Simplify each of your answers as much as possible.
- Answer each word problem with a concluding sentence.
- Write LARGE and neatly.
- Leave plenty of space.
- Work vertically. Do *not* split a page into multiple columns.
- No technology permitted. Put away all cell phones and calculators. An open cell phone will result in an automatic score of 0 for the exam.

Learning outcomes:
Author(s):

- A few derivatives:

$$\frac{d}{dx} (\arctan x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} (\arccos x) = -\frac{1}{\sqrt{1-x^2}}$$

2 Examples of How to Show Work

Here are some examples of how to show your work when taking derivatives. These examples will *not* be included with the exam.

Example 65. The chain rule (do *not* use the quotient rule for a derivative like this one):

$$\begin{aligned} \frac{d}{d\theta} \left(\frac{4}{1+\tan \theta} \right) &= 4 \cdot \frac{d}{d\theta} ((1+\tan \theta)^{-1}) \\ &= -4(1+\tan \theta)^{-2} \cdot \frac{d}{d\theta} (\tan \theta) \\ &= -4(1+\tan \theta)^{-2} (\sec^2 \theta). \end{aligned}$$

Or, if you prefer, you could make the u -substitution explicit by letting

$$y = 4(1+\tan \theta)^{-1}$$

and

$$u = 1 + \tan \theta.$$

Then

$$y = u^{-1}$$

and by the chain rule

$$\begin{aligned} \frac{dy}{d\theta} &= \frac{dy}{du} \cdot \frac{du}{d\theta} \\ &= \frac{d}{du} (u^{-1}) \cdot \frac{d}{d\theta} (\tan \theta) \\ &= (-u^{-2}) (\sec^2 \theta) \\ &= -4(1+\tan \theta)^{-2} (\sec^2 \theta). \end{aligned}$$

Example 66. The quotient rule with the chain rule:

$$\begin{aligned}\frac{d}{dw} \left(\frac{w^2}{1 + e^{3w}} \right) &= \frac{(1 + e^{3w}) \frac{d}{dw}(w^2) - w^2 \frac{d}{dw}(1 + e^{3w})}{(1 + e^{3w})^2} \\ &= \frac{(1 + e^{3w})(2w) - w^2 e^{3w} \frac{d}{dw}(3w)}{(1 + e^{3w})^2} \\ &= \frac{(1 + e^{3w})(2w) - 3w^2 e^{3w}}{(1 + e^{3w})^2}.\end{aligned}$$

Review Problems

Some problems for review.

Question 393 This question is about how the temperature inside a building changes in response to changes in the outdoor temperature. We assume the building has no internal heating or cooling system.

We'll suppose that the function

$$f(t) = M - B \cos\left(\frac{\pi}{12}t\right), \quad t \geq 0,$$

expresses the outdoor temperature (in Fahrenheit degrees) in terms of the number of hours past 4am.

Newton's law of cooling models the rate at which the indoor temperature is changing at any time. It says that this rate of change is proportional to the difference in the indoor and outdoor temperatures. So if the function

$$T = g(t) \quad t \geq 0,$$

expresses the outdoor temperature (in Fahrenheit degrees) in terms of the number of hours past 4am, Newton's law says that

$$\frac{dT}{dt} = k(f(t) - g(t)) \tag{8}$$

for some constant k .

- (a) What are the units of k ? How do you know?
- (b) Is k positive or negative? How do you know?
- (c) Experiment with the sliders in the demonstration below. Summarize your observations.

Desmos link: <https://www.desmos.com/calculator/oag9lhvgo5>

Desmos activity available at 151: *Building Temperature*

Next quarter you will learn how to use the above equation to determine the indoor temperature at any time given the temperature at some specific time.

Learning outcomes:

Author(s):

For now, we'll just claim that the indoor temperature is given by the function

$$\begin{aligned}T &= g(h) \\&= M + Ce^{-kt} - \frac{B}{1 + (\frac{\pi}{12k})^2} \left(\cos\left(\frac{\pi}{12}t\right) + \frac{\pi}{12k} \sin\left(\frac{\pi}{12}t\right) \right) \\&= M + Ce^{-kt} - \frac{B}{\sqrt{1 + (\frac{\pi}{12k})^2}} \cos\left(\frac{\pi}{12}t - \phi\right),\end{aligned}$$

where

$$\phi = \arctan\left(\frac{\pi}{12k}\right).$$

Hanging Chains

Comparing a catenary with a weighted chain.

Question 394 *Desmos link:*

<https://www.desmos.com/calculator/cifsqaas5j>

Desmos activity available at 151: Weighted Chain

Question 395 *Desmos link:*

<https://www.desmos.com/calculator/sv70z21j2j>

Desmos activity available at 151: Catenary

Learning outcomes:
Author(s):

Electron in a Crossed Field

Electron.

Crossed Fields and the Hodograph

The force on an a charge q with velocity $\mathbf{v}$ in a crossed magnetic/electric field is given by

$$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}).$$

Our problem is to parameterize the motion of a charged particle when the fields are uniform and mutually orthogonal. We suppose the magnetic and electric fields point in the respective directions $\mathbf{k}$ and $\mathbf{j}$, and that the charge has velocity $\mathbf{v}_0$ perpendicular to $\mathbf{k}$ at time $t = 0$.

Then the charge, assumed to be positive, has acceleration

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = k_1\mathbf{j} + k_2\mathbf{v} \times \mathbf{k},$$

for some constants $k_1, k_2 \geq 0$, with respective units m/sec^2 and sec^{-1} .

This is a differential equation that is easy enough to solve algebraically. But our aim here is to take a geometric approach that uses some key ideas from Calculus 3.

In sketching motion we typically draw the path and perhaps draw a few position vectors at equally-spaced time intervals. These vectors allow us to approximate a few velocity vectors, or at least their relative lengths. But instead of drawing the velocity vectors tangent to the path, it is usually more useful to draw them with their tails pinned at a common point. The curve traced by these tips of these pinned vectors is called the *hodograph* of a motion.

For starters, the hodograph gives us a way to visualize the motion's acceleration as it is tangent to the hodograph. But to get some idea of the acceleration's magnitude, we need to consider not only the direction and magnitude of the velocity vector but also its rotation rate.

For a uniform circular motion that rotates around a circle of radius r meters at a constant rate of ω rad/sec, for example, we *see* from the hodograph (a circle of radius $v = \omega r$ that the acceleration vector points directly toward the center of the path and rotates at the same rate as both the position and velocity vectors.

Learning outcomes:
Author(s):

So without any derivatives, we get the acceleration's magnitude to be

$$|\mathbf{a}| = \omega|\mathbf{v}| = \omega^2 r.$$

Turning off the Electric Field

Returning to our charged particle, with the electric field turned off ($k_1 = 0$), the acceleration and velocity vectors are perpendicular, so the charge moves with constant speed $|\mathbf{v}_0| = v_0$. But then because $\mathbf{v}$ is perpendicular to $\mathbf{k}$,

$$|\mathbf{a}| = k_2|v| = k_2 v_0$$

is also constant.

So we are looking for a plane motion having constant speed and an acceleration with constant magnitude. One choice would be uniform circular motion, where

$$|\mathbf{a}| = v_0 \omega,$$

where $\omega = d\theta/dt$ is the (constant) rotation rate of the pinned velocity vector. Then $k_2 = \omega$ and the trajectory has radius

$$r = \frac{v_0}{\omega} = \frac{v_0}{k_2}.$$

To see that there are no other possible motions we could probably appeal to some uniqueness theorem of differential equations. But for a more geometric approach, consider what we know.

- (a) that the charge moves with constant speed and
- (b) that the acceleration vector rotates at a constant rate.

Question 396 What do these conditions imply about the trajectory?

Hint: Think about curvature.

Explanation. Click the arrow to the lower right for the solution.

Let θ be the angle from $\mathbf{i} = \langle 1, 0, 0 \rangle$ to $\mathbf{v}$. Then because $\mathbf{v}$ and $\mathbf{a}$ are orthogonal, these vectors rotate at the same rate $\omega = d\theta/dt$. And because the speed is constant, the path's radius of curvature

$$r = \frac{|v|}{\left| \frac{d\theta}{dt} \right|} = \frac{v_0}{k_2}$$

is also constant. And because the path is a plane curve, it is a circle with this radius.

1 Newton's Law of Cooling

Exploration 397 *Desmos link:*

<https://www.desmos.com/calculator/yebumxuwms>

Electron 11

Question 398 *The function*

$$P = 40e^{-t/20}, \quad 0 \leq t \leq 15,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

(a) *Find a function*

$$r = g(P)$$

that expresses the growth rate of the population (in millions of bacteria/hour) in terms of the population. Include a domain.

(b) *Use the result of part (a) to determine the population when it is decreasing at the rate of 1,820,000 bacteria/hour. End with a concluding sentence. No credit for other methods.*

Short Quizzes Math 151

Short Quizzes Math 151.

Question 399 The function

$$h = f(v), \quad 80 \leq v \leq 120,$$

expresses the height of a helicopter (measured in feet) in terms of its speed (measured in ft/sec).

Suppose

$$\left. \frac{dh}{dv} \right|_{v=100} = -20.$$

- (a) What are the units of the above derivative?
- (b) Explain the meaning of the derivative using the language of small changes.

Question 400 The function

$$P = f(t) = 40e^{-t/20}, \quad 0 \leq t \leq 20,$$

expresses the population (in millions) of a colony of bacteria in terms of the number of hours past noon.

- (a) Find a function

$$r = g(P),$$

that expresses the growth rate (in millions of bacteria/hour) of the population in terms of the population (in millions of bacteria).

- (b) Use the result of part (a) to determine the population when it is decreasing at the rate of 1,820,000 bacteria/hour.
 - (i) No credit for other methods.
 - (ii) End with a concluding sentence.

Learning outcomes:
Author(s):

Question 401 Find an equation of the tangent line to the curve

$$x^3 \sin(5y) + 4e^{xy} = 4x$$

at the point $(1, 0)$.

- (a) Show all work.
 - (b) Use the Leibniz notation for derivatives and their evaluations.
 - (c) Include a brief explanation with your work.
 - (d) End with a concluding sentence.
-

Short Quizzes Math 142

Short Quizzes Math 142

Question 402 Let Q be the point on the circle of radius 40 meters that is 90 meters from the point $A(40, 0)$ (coordinates measured in meters). The distance is measured clockwise around the circle from A to Q .

Between 12:06pm and 1:00pm a beetle crawls counterclockwise around this circle at a constant speed of 5 meters/min, passing the point Q at 12:43pm.

- (a) Find an expression for the function

$$\theta = f(t), \quad 6 \leq t \leq 60,$$

that expresses the polar angle (measured in radians) of the beetle (more precisely of the vector giving the beetle's position relative to the origin) in terms of the number of minutes past noon being sure to do the following:

- (i) Explain your reasoning. This means to include a brief description of what you are computing for each of your computations.
- (ii) Include units for each number in each computation.
- (iii) Include a graph (drawn by hand) of the function $\theta = f(t)$ to help with your explanation.
- (iv) Include a sketch (drawn by hand) of the circle on a coordinate system with appropriately labeled points and arclengths. Label the axes with the appropriate variable names and units.

- (b) Find functions

$$x = g_1(t), \quad 6 \leq t \leq 60,$$

and

$$y = g_2(t), \quad 6 \leq t \leq 60,$$

that express the coordinates (in meters) of the beetle in terms of the number of minutes past noon. Include a brief explanation.

- (c) Input the correct coordinate functions on Lines 2 and 3 of the worksheet below. Then play the slider u (another name for t , the number of minutes past noon) in Line 1 to see if your functions are correct.

Desmos link: <https://www.desmos.com/calculator/lkrunhfgxi>

Learning outcomes:
Author(s):

142: Short Quiz 1

- (i) *Explain how you know that your functions are correct.*
 - (ii) *Include at least two screenshots of the animation at different times to help with your explanation.*
-

Question 403 *Solve the equation*

$$9 - 3 \sin \theta = 11.$$

- (a) *Explain your reasoning thoroughly, in complete sentences.*
 - (b) *Include a picture as in class to help with your explanation.*
 - (c) *End with a concluding sentence that expresses the solution as a set.*
-

Sinusoidal Modeling, Math 142

Implicit differentiation.

Question 404 Suppose that during a five-day period, beginning at midnight Monday morning, the depth of the water at the Edmond's Pier is a sinusoidal function of time.

Suppose also that a low tide of 7 feet occurs at 1:00am Monday morning and that the following high tide of 23 feet occurs at 6:45am that same morning.

- (a) Sketch by hand a graph of the function

$$h = f(t)$$

that expresses the depth of the water (in feet) as a function of the number of hours since midnight, Monday morning.

- (b) Find an expression for the above function. Include a domain.
 (c) Find all times during the week when the water is 10 feet deep. Give exact times. Do not use a calculator.
 (d) Approximate the clock times (to the nearest minute) on Friday when the water is 10 feet deep.

Question 405 Assume for this question that each month has 30 days and that the number of hours of daylight/day in Seattle is a sinusoidal function of time. Assume also that on June 21, Seattle gets a maximum of 16 hours of daylight/day and that on December 21, Seattle gets a minimum of 8 hours of daylight/day.

- (a) Find a function

$$H = f(t), 0 \leq t \leq 12,$$

that expresses the number of daylight hours/day in Seattle in terms of the number of months since June 21. Use the cosine function. Start by sketching a graph. Explain your reasoning.

- (b) Use your function to determine the number of hours of daylight/day that Seattle gets on March 1.

Learning outcomes:
 Author(s):

- (c) *About how many more minutes of daylight/day do we get tomorrow than today?*
 - (d) *Use your function to determine the day(s) of the year when Seattle gets 14 hours of daylight/day.*
-

Simple Harmonic Motion, Law of Cosines, Math 142

Simple harmonic motion.

Question 406 Suppose for this problem that the earth is a ball with uniform density of radius 4000 miles. Now imagine drilling a straight tunnel through the earth from the North Pole to the South Pole.

A rock dropped from rest at the north pole falling through the tunnel would then oscillate in simple harmonic motion between the poles and return to the north pole every 84 minutes. This means we can think of the rock as being dragged along by a point moving around the earth at constant speed as illustrated below.

Desmos link: <https://www.desmos.com/calculator/ij8dqowgza>

Desmos activity available at 142: *Simple Harmonic Motion*

- (a) Activate the Protractor folder in Line 15 to see the protractor, where consecutive tick marks subtend equal angles of $\pi/100$ radians about the earth's center. Use the protractor but not any trigonometry to answer the following questions. Hint: Think proportionately.
 - (i) Estimate the distance of the rock from the South Pole at time $t = 16.8$ minutes after the rock is released.
 - (ii) Estimate the first two times when the rock is 1000 miles from the South Pole.
- (b) Sketch by hand one period of the graph of the function

$$s = f(t) \quad t \geq 0,$$

that expresses the distance of the rock (measured in thousands of miles) from the South Pole in terms of the number of minutes since the rock was released. Assume the rock is dropped at midnight on July 1, 2085.

- (c) Find an expression for the function $s = f(t)$. Start by expressing the distance s in terms of the polar angle θ marked above.

Learning outcomes:
Author(s):

- (d) Use your function to determine the exact distance between the rock and the South Pole at time $t = 16.8$ minutes past noon. Then use a calculator to approximate this distance to the nearest mile and compare it with your estimate in part (i).
- (e) Use your function to find all (exact) times when the rock is 1000 miles from the South Pole. Do this by first finding all (exact) polar angles when the rock is 1000 miles from the South Pole.
- (f) Find the (exact) first two times when the rock is 1000 miles from the South Pole. Then use a calculator to estimate these clock times to the nearest minute.
- (g) Find all (exact) times between 8pm and 12pm, July 1, 2085 when the rock is 1000 miles from the South Pole. Then use a calculator to estimate these clock times to the nearest minute.

Question 407 In $\triangle MAT$, angle $\angle MAT$ has measure $\pi/3$ radians. Sides AM and AT have respective lengths 2 and 5 inches. Determine the exact measure of angle $\angle AMT$. Do not use a calculator.

Question 408 You measure the angle of elevation to the top of a tree to be θ_1 radians. You then walk an additional c feet directly away from the tree and measure the angle of elevation to be θ_2 radians.

- (a) Express the height of the tree above eye level in terms of θ_1 , θ_2 , and c .
- (b) Check that your expression has the correct units.

Question 409 A tree leans precariously with its trunk inclined at an angle of $\pi/3$ radians to the ground. The top of a seven-foot ladder rests against the tree and the bottom of the ladder lies eight feet from the tree's base. How far is the top of the ladder from the base of the tree? Do not use a calculator except for arithmetic. Do not rely on the worksheet below.

Desmos link: <https://www.desmos.com/calculator/sjmjseyqyp>

Desmos activity available at 142: Ladder and Tree 45

Quiz Solutions

Solutions to daily quizzes.

1 Quiz 5A

Question 410 The function $G = f(s)$, $0 \leq s \leq 80$, expresses the number of gallons of gas in a car in terms of the trip odometer reading (measured in miles).

- (a) Use the Leibniz evaluation notation to write an expression for the derivative of G with respect to s evaluated at an odometer reading of 10 miles.
- (b) Now suppose

$$G = f(s) = \frac{1}{2000} (s^2 - 200s + 10000), \quad 0 \leq s \leq 80.$$

- (i) Evaluate the derivative in part (a). Include units. Interpret the meaning of the derivative in a way that someone without any knowledge of calculus would understand. In particular, do not use the phrasing rate of change of ... with respect to
- (ii) At what odometer reading(s), if any, is the (instantaneous) gas mileage 17 miles/gal? Explain your reasoning in a few complete sentences. End with a concluding sentence.

Explanation. (a) The derivative is

$$\left. \frac{dG}{ds} \right|_{s=10}.$$

- (b) Differentiating, we get

$$\begin{aligned} \frac{dG}{ds} &= \frac{d}{ds} \left(\frac{1}{2000} (s^2 - 200s + 10000) \right) \\ &= \frac{1}{2000} \left(\frac{d}{ds} (s^2 - 200s + 10000) \right) \\ &= \frac{2s - 200}{2000} \\ &= \frac{s - 100}{1000}. \end{aligned}$$

Learning outcomes:
Author(s):

So

$$\begin{aligned}\left.\frac{dG}{ds}\right|_{s=10} &= \left.\frac{s-100}{1000}\right|_{s=10} \\ &= \frac{10-100}{1000} \\ &= -0.09 \frac{\text{gal}}{\text{mile}}\end{aligned}$$

This means that at an odometer reading of 10 miles, the car is burning gas at the rate of 0.09 gal/mile. The derivative is negative because as the odometer reading increases the volume of gas decreases.

Alternatively, we could say that as you drive one mile, between odometer readings $s = 10$ and $s = 11$, the car burns *approximately* 0.09 gallons of gas.

- (c) When the car is getting 17 miles/gal, it is burning gas at the rate of $(1/17)$ gal/mile and

$$\frac{dG}{ds} = \frac{s-100}{1000} = -\frac{1}{17}.$$

Solving this equation for s gives

$$17s - 1700 = -1000$$

and

$$s = \frac{700}{17}.$$

So when the car is getting 17 miles/gal, the odometer reads $s = 700/17$ miles.

Introduction to Integration, Part 1

Introduction to integral calculus.

The *differential* in Differential Calculus has the same root as *difference*, and *calculus* the same root as *compute*. Differential calculus is all about *computing differences*. Expressed more simply, it is about *subtraction*.

Integral Calculus, on the other hand, is about addition. *Integral* shares the same root as *integer* and comes from the Latin *integrare*, to make whole.

Addition and subtraction are inverse operations. They undo each other. Similarly, differentiation undoes integration. But we need to be more precise to claim that integration undoes differentiation. More about this later.

It would be a mistake to think integral calculus is primarily about finding anti-derivatives. While anti-differentiation plays a role in integration, this class is more about recognizing *when* to integrate rather than *how* to integrate. Many of your STEM classes use definite integrals, with an emphasis on setting up and interpreting integrals, not computing their values.

It would also be a mistake to think this class is about using definite integrals to compute *numbers*, like the distance travelled over a certain time interval or the period of a pendulum. Rather, the emphasis here will be on using definite integrals to create *functions*. Expressing the position of a pendulum as a function of time, for example, gives a lot more information than just computing its period.

Using definite integrals to write functions generalizes the point-slope equation of a line. You used this equation in differential calculus to approximate a function in the neighborhood of a point. To capture the local nature of the approximation (ie. the approximation is usually accurate in just a small neighborhood of the point of tangency), it's the point-slope equation, *not* slope-intercept, that works best. Here's an example that illustrates this idea.

Example 67. Between 12:03pm and 1:00pm a balloon ascends at a constant rate of 73 ft/min. The balloon is 1900 feet high at 12:10pm.

- (a) Find an expression for the function

$$h = g(t), \quad 0 \leq t \leq 60,$$

that expresses the balloon's height (measured in feet) in terms of the number of minutes past noon.

Learning outcomes:
Author(s):

- (b) Explain the logic behind your expression.

Explanation. The function is

$$h = g(t) = 1900 + 73(t - 10), \quad 0 \leq t \leq 60.$$

The logic behind the expression is this. To find the height at time t minutes past noon, we start with the height (1900 feet) at 12:10pm and add the change in height from 12:10pm to time t minutes past noon. The change in height,

$$\Delta h = 73(t - 10)$$

is the product of the rate of ascent (73 ft/min) and the number of minutes

$$\Delta t = t - 10$$

from 12:10pm to time t minutes past noon.

In the context of the last example, the main topic for this class is how to compute the height function *if the rate of ascent is not constant*.

The logic behind the solution is almost identical to point-slope. Suppose we're given a function

$$r = f(t), \quad 0 \leq t \leq 60,$$

that gives the balloon's rate of ascent (in ft/min) at a general time t minutes past noon. To find the height at time t , we start with the height of the balloon at 12:10 pm and add to that the change in height from 12:10pm to time t . But because the rate of ascent is not necessarily constant, we'll almost certainly have (infinitely) many rates. Our job is to use these rates to compute, or at least approximate the change in height. The next example illustrates this idea.

Example 68. Suppose we know the balloon is 1900 feet high at 12:10pm and that we're given a graph of the function

$$r = f(t), \quad 0 \leq t \leq 60,$$

(assumed continuous) that expresses its rate of ascent (measured in ft/min) in terms of the number of minutes past noon. This graph is shown below.

Desmos link: <https://www.desmos.com/calculator/h6cworakdw>

152:Balloon Introduction 2

Our problem is to approximate the balloon's height at 12:50pm.

We'll start by computing a rough approximation of the change in height from 12:10pm and 12:50pm. For this, we'll partition the 40-minute interval into five equal subintervals of length

$$\Delta t = \frac{40 \text{ min}}{5} = 8 \text{ min.}$$

Now we'll suppose that during each subinterval the balloon ascends at a constant rate. While this is not true, the rate certainly varies less over an 8-minute interval than it does over the entire 40-minute interval. So we should get a better approximation this way than if we assumed just one constant rate of ascent between 12:10pm and 12:50pm.

What rate we choose for each 8-minute interval doesn't really matter as long as we choose the actual rate of ascent at some time during that interval. To make things simple, we'll choose the rate at the start of each subinterval.

Then between 12:10pm and 12:18pm, the approximate change in height (measured in feet) is

$$f(10)(8) \sim (31 \text{ ft/min})(8 \text{ min}) = 248 \text{ ft.}$$

Between 12:18pm and 12:26pm, the approximate change in height (measured in feet) is

$$f(18)(8) \sim (12 \text{ ft/min})(8 \text{ min}) = 96 \text{ ft.}$$

And we continue similarly to approximate the changes in height (this time negative) over the remaining three 8-minute intervals. So from 12:10pm to 12:50pm the approximate change in height (in feet) is

$$\begin{aligned} \Delta h_5 &= h - 1900 \\ &\sim (f(10) + f(18) + f(26) + f(34) + f(42))(8) \\ &\sim -448. \end{aligned}$$

Verify this computation by typing this expression in Line 38 of the worksheet above.

This sum is on Line 7 of the worksheet.

To see the previous computation geometrically, move the slider a (the start of our time interval) in Line 2 above to $a = 10$ and the slider b (the end of our time interval) in Line 3 to $b = 50$. Then the signed area of a shaded rectangle measures the approximate change in height during that subinterval. And the approximate change in height from 12:10pm to 12:50pm is just the sum of these signed areas. The sign of the area and the rate of ascent are the same. So rectangles below the horizontal axis contribute a negative signed area (ie. a decrease in height), while those above make a positive contribution (ie. an increase in height).

Our first approximation to the balloon's height at 12:50pm is then

$$(1900 - 448) \text{ ft} \sim 1452 \text{ ft.}$$

To get a better approximation, we partition the 40-minute interval from 12:10pm to 12:50pm into more subintervals of equal length. To do this, drag the slider n on Line 5 (the number of equal subintervals) to the right.

- (a) Describe what happens to the approximate change in height (s_1 in Line 5) and how the picture changes as n grows without bound.

- (b) Then approximate the exact change in height and interpret that change geometrically.

Free Response:

Takeaways:

Here's some notation along with a summary of the main idea for the last example.

The exact height of the balloon at 12:50pm is

$$1900 + \int_{10}^{50} f(t) dt.$$

This expression starts with the height at 12:10pm (1900 feet) and adds to it the change in height

$$\int_{10}^{50} f(t) dt$$

from 12:10pm to 12:50pm. The integral sign $\int$ stands for the letter *S* and means *sum*. You can think of the product

$$f(t) dt$$

as a small change in height. It multiplies the rate of ascent $f(t)$ and a small (differential) time interval dt to give a small (differential) change in height. The integral then sums these changes to get the exact change in height.

More generally, the height of the balloon as a function of time (measured in minutes past noon) is

$$h = f(t) = 1900 + \int_{10}^t f(u) du$$

and the logic is the same.

Compare this with the expression (point-slope) from Example 1 for the height

$$h = 1900 + 73(t - 10).$$

Free Response:

Introduction to Integration, Part 2

Introduction to numerical integration.

This chapter is a continuation of the introduction where we saw how to approximate a balloon's change in height over an interval of time with a sum of small approximate changes. We develop that theme here with summation notation and take advantage of technology to get accurate approximations.

But first a question for review.

1 Review

Example 69. Between 12:14pm and 1:00pm the temperature of a beaker of water increases at a constant rate of $\frac{33^\circ}{13}$ C/sec. The temperature is 23°C at 12:23pm.

- (a) Find an expression for the function

$$C = f(t), \quad 14 \leq t \leq 60,$$

that expresses the temperature of the water (in Celsius degrees) in terms of the number of minutes past noon.

- (b) Explain the logic behind your expression.

Free Response:

2 Summation Notation and Technology

Back to the problem from the introduction. But first a few conceptual questions.

Example 70. The function

$$r = f(t), \quad 0 \leq t \leq 60,$$

expresses a balloon's rate of ascent (measured in ft/min) in terms of the number of minutes past noon. Its graph is shown below.

Learning outcomes:
Author(s):

Desmos link: <https://www.desmos.com/calculator/tgi5yiuzab>

152:Balloon Introduction

- (a) When is the balloon ascending? Descending? Give all times for each and explain your reasoning.
- (b) Do you think the balloon is higher at noon or at 1:00pm? Explain your reasoning. Try to prove your assertion using only arithmetic and simple reasoning.

Free Response:

Now suppose the balloon is 1900 feet above the ground at 12:10pm.

- (a) Use the graph to approximate the balloon's height 30 seconds later. Explain your reasoning.
- (b) Is your estimate greater or less than the actual height? Explain your reasoning.

Free Response:

We can turn the previous question of differential calculus into an addition problem of integral calculus by estimating the balloon's height at a time farther removed from 12:10.

Example 71. We'll assume as before that the balloon is 1900 feet above the ground at 12:10pm, and we'll approximate its height at 12:50pm.

Desmos link: <https://www.desmos.com/calculator/h6cworakdw>

152:Balloon Introduction 2

We'll start by computing a rough approximation of the change in height between 12:10pm and 12:50pm. For this, we'll partition the 40-minute time interval into five equal subintervals and suppose that during each subinterval the balloon ascends at a constant rate.

What rate we choose for each interval doesn't really matter as long as we choose the actual rate of ascent at some time during that interval. To make things simple, we'll choose the rate at the start of each subinterval as illustrated in the worksheet above.

- (a) Use summation notation to express the approximate change in height from 12:10pm to 12:50pm. Type this expression in Line 5 of the worksheet above. Compare this value with your estimate from part (a).

Using the rate of ascent at the start of each 8-minute time interval, the approximate change in height from 12:10pm to 12:50pm is

$$\sum_{i=0}^4 8f(10 + 8i)$$

and the approximate height at 12:50pm is

$$1900 + \sum_{i=0}^4 8f(10 + 8i).$$

- (b) Repeat part (a) using the rate of ascent at the *end* of each subinterval instead. Turn off the folder in Line 11 and activate the folder in Line 18 above. As a check, type your expression for the approximate height in Line 5 above.

The approximate change in height from 12:10pm to 12:50pm is

$$\sum_{i=1}^5 8f(10 + 8i)$$

and the approximate height at 12:50pm is

$$1900 + \sum_{i=1}^5 8f(10 + 8i).$$

- (c) Repeat part (a) using the rate of ascent at the *midpoint* of each subinterval instead. Turn off the folder in Line 18 and activate the folder in Line 24 above. As a check, type your expression for the approximate height in Line 5 above.

The approximate change in height from 12:10pm to 12:50pm is

$$\sum_{i=0}^4 8f(14 + 8i)$$

and the approximate height at 12:50pm is

$$1900 + \sum_{i=0}^4 8f(14 + 8i).$$

- (d) Compare your three estimates for the change in height. Which do you think is most accurate? Explain why.

Example 72. (a) Drag the slider in Line 4 of Example 3 to $n = 20$ and repeat parts (a) - (c) of Example 4 with $n = 20$ equal subintervals.

(b) Try parts (c) - (e) of Example 3 in general, with n -equal subintervals.

Using left endpoints, the approximate height at 12:50pm is

$$f(50) \sim 1900 + \frac{40}{n} \sum_{i=0}^{n-1} f\left(10 + \frac{40}{n}i\right).$$

With right endpoints,

$$f(50) \sim 1900 + \frac{40}{n} \sum_{i=1}^n f\left(10 + \frac{40}{n}i\right).$$

And with midpoints,

$$f(50) \sim 1900 + \frac{40}{n} \sum_{i=0}^{n-1} f\left(10 + \frac{20}{n} + \frac{40}{n}i\right).$$

Addition

When do we need to add?

1 Two Balloon Problems

Exercise 411 (a) *The table below shows the height of a balloon at several times.*

Time	Height (feet)	Average Rate of Ascent (ft/min)
12:30	420	—
12:40	500	
12:44	550	
12:50	400	

Find the balloon's average rate of ascent over each of the three time intervals between consecutive times.

(b) *The table below shows the rate of ascent (in ft/min) of another balloon at several times.*

Time	Rate of ascent (ft/min)	Approximate height (ft)
12:30	40	
12:40	25	1800
12:44	-30	
12:50	-50	

The balloon is 1800 feet high at 12:40pm.

Estimate the heights of the balloon at 12:30pm, 12:44pm, and 12:50pm. Do this three times. First assume the balloon ascends at a constant rate over each time interval equal to the rate of ascent at the start of each interval. Then at a constant rate equal to the rate at the end of each interval. And finally at a constant rate equal to the average of the rates at the endpoints of each interval.

(c) *Compare your computations in parts (a) and (b). How are they related?*

Learning outcomes:
Author(s):

2 Time, Distance, and Mileage Problems

Exercise 412 The table below shows your speed at several times as you drive north on I-5 on part of a two-hour trip.

Time	Speed (miles/hour)
12:30	42
12:40	50
12:44	55
12:50	40

The trip odometer reading is 110 miles at 12:30pm.

- Use the data to give two approximations to the trip odometer reading at 12:50pm. Once assuming a constant speed over each subinterval equal to the speed at the start of the interval. And again assuming a constant speed over each subinterval equal to the speed at the end of the interval. Explain your reasoning.
- Activate the folder in Line 5 of the worksheet below and explain how what you see is related to part (a). Then turn the folder off and activate the folder in Line 11. Explain.
- To practice summation notation, let t_i , $i = 1, 2, 3, 4$ be the times, in order, in the above table measured in hours past 12:30pm and let v_i be the corresponding speeds. Then using the speed at start of each subinterval, the trip odometer reading at 12:50pm is approximately

$$110 + \sum_{i=1}^4 (t_{i+1} - t_i) v_i.$$

Using the speed at the end of each subinterval, the odometer reading is approximately

$$110 + \sum_{i=1}^4 (t_{i+1} - t_i) v_{i+1}.$$

- Input the sums from part (c) in Lines 16 and 18 of the worksheet below to check your arithmetic. You'll need to write $t[i]$ in place of t_i and so on.

Desmos link: <https://www.desmos.com/calculator/jjobuctzsa>

152: Distance Travelled

Exercise 413 The table below shows your gas mileage on a car trip as you pass different exits while driving north on I-5. The exit numbers record the distance to the Columbia river along I-5, measured in miles.

Exit Number	Mileage (miles/gallon)
176	40
200	32
210	30
248	25

- Is this enough information to compute upper and lower bounds for the number of gallons of gas your car burned between Exits 176 and 248? If so, explain why. If not, what other information do you need?
- Using the given information and your additional hypothesis from part (a), if needed, find the best possible upper and lower and upper bounds for the number of gallons of your car burned between Exits 176 and 248.

Desmos link: <https://www.desmos.com/calculator/csjpg0u3sw>

152: Gas Consumed 34

- Let e_i and g_i , $i = 1, 2, 3, 4$ denote the respective exit numbers and gas mileages in the table above. Use summation notation to write your estimates in part (b) above.

Assuming the gas mileage is a decreasing function, the number of gallons of gas burned between Exits 176 and 248 is at least

$$\sum_{i=1}^3 (e_{i+1} - e_i) \left(\frac{1}{g_i} \right).$$

This is the best possible (ie. greatest possible) lower bound.

With the same assumption, the number of gallons of gas burned is at most

$$\sum_{i=1}^3 (e_{i+1} - e_i) \left(\frac{1}{g_{i+1}} \right).$$

This is the best possible (ie. least possible) upper bound.

- Use the desmos worksheet above to represent the sums in parts (a) and (b) geometrically by activating the appropriate folder in Line 5 (Left endpoints) or Line 11 (Right endpoints).

Explanation

- (a) The table does not give enough information to determine such bounds because we don't know how the gas mileage varies during each interval. But with the additional assumption that the gas mileage is a decreasing function of the exit number we can compute these bounds.
- (b) The key, as always in this class, is to consider the constant case. Here that means to assume the gas mileage is constant, say 40 miles/gallon. Then over a 200-mile trip, for example, your car would burn

$$\frac{200 \text{ miles}}{40 \text{ miles/gal}} = 5 \text{ gallons}$$

of gas.

The car burns less gas when its gas mileage is greater. So for the greatest possible lower bound (ie. to find the minimum number of gallons of gas the car burned), we assume the gas mileage is the greatest possible during each interval. Here that means choosing left endpoints. That is, during each interval we assume the gas mileage is constant and equal to the gas mileage at the start of the interval (ie. equal to its maximum over the interval).

So between Exits 176 and 248 the car burns at least

$$\frac{(200 - 176) \text{ miles}}{40 \text{ miles/gal}} + \frac{(210 - 200) \text{ miles}}{32 \text{ miles/gal}} + \frac{(248 - 210) \text{ miles}}{30 \text{ miles/gal}} > 2.179 \text{ gallons.}$$

Similar reasoning shows that between Exits 176 and 248 the car burns at most

$$\frac{(200 - 176) \text{ miles}}{32 \text{ miles/gal}} + \frac{(210 - 200) \text{ miles}}{30 \text{ miles/gal}} + \frac{(248 - 210) \text{ miles}}{25 \text{ miles/gal}} < 2.604 \text{ gallons.}$$

Exercise 414 The table below shows your speed and gas mileage at several times during the course of a two-hour trip.

Time	Speed (miles/hr)	Mileage (miles/gal)
1:00pm	40	30
1:45pm	48	36
2:15pm	60	40
3:00pm	70	28

- (a) Give two estimates for the distance travelled during the trip. Explain your reasoning.
- (b) Give two estimates for the number of gallons of gas your car burned during the trip. Explain your reasoning.

3 On Trails

Exercise 415 The table below shows the inclination angle of the Mt. Washington Cog Railway at several distances from Waumbek Station (the departure point, elevation 4000 feet) on its way to the summit (elevation 6288 feet).

Distance (miles)	Inclination Angle (radians)
0.5	0.25
1.0	0.30
1.5	0.32
2.0	0.38

- (a) Use the data to find two estimates for the elevation change from mile marker 0.5 to mile marker 2.0. Explain your reasoning.
- (b) Let s_i and ϕ_i denote the respective distances and angles in the table above. Use summation notation to write the estimate in part (a)

Assuming the steepness of the track increases, a lower bound for the change in height (in miles) is

$$h_1 = \sum_{i=1}^3 (s_{i+1} - h_i) (\sin \phi_i).$$

An upper bound is

$$s_1 = \sum_{i=1}^3 (h_{i+1} - h_i) (\sin \phi_{i+1}).$$

- (c) Input your expressions for the lower and upper bounds in the worksheet below.

Desmos link: <https://www.desmos.com/calculator/ehnbprdu6q>

152: Cog Railroad

Exercise 416 The table below shows the slope along a section of a trail at several altitudes.

Altitude (feet)	Slope (ft/ft)
1000	0.05
1200	0.10
1350	0.12
1500	0.15

- (a) Use the data to find two estimates for the length of the trail between altitudes 1000 ft and 1500 feet. Explain your reasoning.
- (b) Let h_i and m_i , $i = 1, 2, 3, 4$ denote the respective altitudes and slopes in the table above. Use summation notation to write your estimates in part (a).

Assuming the steepness of the trail increases between altitudes 1000 feet and 1500 feet, a lower bound for the trail's length (in feet) is

$$s_1 = \sum_{i=1}^3 (h_{i+1} - h_i) \left(\sqrt{1 + (1/m_{i+1})^2} \right).$$

An upper bound is

$$s_1 = \sum_{i=1}^3 (h_{i+1} - h_i) \left(\sqrt{1 + (1/m_i)^2} \right).$$

- (c) Input your expressions for the lower and upper bounds in the worksheet below.

Desmos link: <https://www.desmos.com/calculator/mmi4rx2rqp>

152: Length of Trail

4 A Rollercoaster Problem

Exercise 417 The table below shows several speeds of a roller coaster as it slides down a circular loop-the-loop with a radius of 80 feet. The bottom of the loop is 20 feet above the ground.

Height above ground	Speed (ft/sec)
100	40
80	48
50	64
20	80

- (a) Is this enough information to compute upper and lower bounds for the time it takes for the roller coaster to fall from a height of 100 feet to a height of 20 feet? If so, explain why. If not, what other information do you need? Do not assume any particular values for the speed at other heights.

- (b) Using the given information and your additional hypothesis from part (a) if needed, find the best possible upper and lower and upper bounds for the time it takes for the roller coaster to fall from a height of 100 feet to a height of 20 feet. Do not use a calculator for this part. But your bounds should be in a form that could be input directly into a calculator.
- (c) Use a calculator to find approximations to your bounds from part (b). Then give your best estimate of the time together with bounds for the error in your estimate.

Explain your reasoning thoroughly in the form of a letter to a friend with no knowledge of calculus or the ideas of Section 5.1. Include units in all steps of your computations. Be careful. In particular, take note that the roller coaster moves along a circular arc. It does not fall straight down.

Exercise 418 Assume for this problem that each month has 30 days and that the number of hours of daylight/day in Seattle throughout the course of a year is a sinusoidal function of time. Suppose also that Seattle receives a maximum of 16 hours of daylight/day on June 21 and a minimum of 8 hours of daylight/day on December 21.

- (a) Find an expression for a function

$$h = f(t), 0 \leq t \leq 360,$$

that expresses the number of hours of daylight/day in terms of the number of days since June 21. Use the cosine function.

The function is

$$h = f(t) = 12 + 4 \cos \left(\frac{\pi}{180} t \right).$$

- (b) Use summation notation to write an expression for the total number of hours of daylight Seattle gets during the summer.

The total number of hours of daylight during the summer is

$$\sum_{i=0}^{89} f(i).$$

- (c) Enter the sum in Line 1 of the worksheet below.

Desmos link: <https://www.desmos.com/calculator/jvwyqn0vbj>

152: Hour of Daylight 1

- (d) Compute the average number of hours of daylight/day in Seattle during the summer?
 - (e) Is the number of hours of daylight/day in Seattle increasing or decreasing on March 21st? At what rate?
 - (f) Does the previous question make sense? Explain.
-

5 The Mercator Map

Look at a map of Seattle and you'll see a scale, probably near the bottom. It might be 0.5 miles/inch, or something like that. But you will not see a scale like this on a map of the earth for the simple reason that one does not exist. The scale varies from place to place and is usually a function of latitude. Even at a specific location, there is often not one but infinitely many scaling factors that depend on direction.

But for some maps, the *conformal* ones, the scaling factor is independent of direction and depends only on latitude. The Mercator map is one of these maps. The next few exercises are about this map.

Exercise 419 The worksheet below shows a Mercator map of the earth. The evenly-spaced vertical lines are the meridians (half-circles running between the north and south poles) and divide the earth into its 24 time zones. The horizontal lines are the parallels (circles of latitude). These are unevenly spaced.

We'll suppose the map is $2\pi \sim 6.3$ feet wide and that the earth is a perfect sphere of radius 3960 miles. The scale along the equator is then

$$\frac{2\pi(3960 \text{ miles})}{2\pi \text{ ft}} = 3960 \frac{\text{miles}}{\text{ft}}$$

So the 24 meridians are spaced at intervals of

$$\frac{2\pi \text{ ft}}{24} = \frac{\pi}{12} \text{ ft}$$

on the map. And along the equator of the earth, they are spaced at intervals of

$$\frac{2\pi(3960) \text{ miles}}{24} = 330\pi \text{ miles} \sim 1037 \text{ miles}.$$

Desmos link: <https://www.desmos.com/calculator/cr8vg7urnc>

152: Mercator Map Riemann Sum 2

Segment AB indicates the map scale. It represents a length of $330\pi \sim 1037$ miles along any parallel. Along the equator AB has length $\pi/12 \sim 0.26$ feet. But drag point A vertically and you'll see how the scale varies with latitude as the length of AB changes.



Integrating Power Functions

Power functions.

1 Two Balloon Problems

Riemann Sums

Riemann sums and summation notation.

Question 420 The function

$$v = g(s) = 40 + \frac{s}{5}, \quad 20 \leq s \leq 100,$$

expresses the speed (in miles/hr) of a car in terms of the trip odometer reading (measured in miles) during part of a 200-mile trip.

- (a) Use summation notation for a Riemann sum with 10 equal intervals of distance to find an upper bound for the time it takes to travel the 80 miles in this portion of the trip. Enter this in Line 1 of the worksheet below to get an approximation for the time. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (b) Use summation notation for a Riemann sum with 10 equal intervals of distance to find a lower bound for the time it takes to travel the 80 miles in this portion of the trip. Enter this in Line 1 of the worksheet below to get an approximation for the time. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (c) Use summation notation for a Riemann sum with n equal intervals of distance to find an upper bound for the time it takes to travel the 80 miles in this portion of the trip. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (d) Use summation notation for a Riemann sum with n equal intervals of distance to find a lower bound for the time it takes to travel the 80 miles in this portion of the trip. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (e) Use either part (c) or part (d) to write an expression (as a limit) for the exact time it takes to travel the 80 miles.
- (f) Write a definite integral for the time it takes to travel the 80 miles and enter this on Line 39 of the worksheet below.

Learning outcomes:
Author(s):

Desmos link: <https://www.desmos.com/calculator/k10phbukon>

152: Distance Speed 2

Question 421 The function

$$r = f(t) = 70 \cos^3(\sqrt{t+40}), \quad 0 \leq t \leq 48,$$

expresses a balloon's rate of ascent (in ft/min) in terms of the number of minutes past noon. The balloon was 300 feet high at 12:04pm.

- (a) Use summation notation for an expression with 10 equal time intervals to find a lower bound for the balloon's height at 12:44pm. Enter this expression in Line 1 of the worksheet below to get an approximation for the height. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (b) Use summation notation for an expression with 10 equal time intervals to find an upper bound for the balloon's height at 12:44pm. Enter this expression in Line 1 of the worksheet below to get an approximation for the height. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (c) Use summation notation for an expression with n equal time intervals to find a lower bound for the balloon's height at 12:44pm. Enter this expression in Line 1 of the worksheet below to get an approximation for the height. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (d) Use summation notation for an expression with n equal time intervals to find an upper bound for the balloon's height at 12:44pm. Enter this expression in Line 1 of the worksheet below to get an approximation for the height. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (e) Use either part (c) or part (d) to write an expression (as a limit) for the exact height at 12:44pm.
- (f) Write an expression with a definite integral for the height at 12:44pm and enter this on Line 39 of the worksheet below.

Desmos link: <https://www.desmos.com/calculator/qwuqprd9vf>

152: Balloon 56

Question 422 The function

$$\theta = g(s) = \frac{s}{2} - \frac{s^3}{12}, \quad 0 \leq s \leq 3,$$

expresses the inclination angle (in radians) of the trail to Nada Lake in terms of the distance from the trailhead (measured along the trail in miles). The inclination angle is the angle the trail makes with the horizontal. It is positive (negative) when the trail slopes upward (downward) in the direction away from the trailhead.

The trail is at an elevation of 3200 feet 0.5 miles from the trailhead.

- (a) Use summation notation for an expression with 10 equal intervals of distance to find a lower bound for the trail's elevation (in feet) at Nada Lake (3 miles from the trailhead). Enter this in Line 1 of the worksheet below to get an approximation for the elevation. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (b) Use summation notation for an expression with 10 equal intervals of distance to find an upper bound for the trail's elevation (in feet) at Nada Lake. Enter this in Line 1 of the worksheet below to get an approximation for the elevation. Activate the appropriate folder in Line 12 (Left endpoints) or Line 19 (Right Endpoints) to interpret the sum geometrically.
- (c) Use summation notation for an expression with n equal intervals of distance to find an upper bound for the trail's elevation (in feet) at Nada Lake. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (d) Use summation notation for an expression with n equal intervals of distance to find a lower bound for the trail's elevation (in feet) at Nada Lake. Enter this in Line 1 below. Then drag the slider n in Line 2 to get a more accurate upper bound.
- (e) Use either part (c) or part (d) to write an expression (as a limit) for the exact elevation of Nada Lake.
- (f) Write an expression with a definite integral for the elevation of Nada Lake.

Desmos link: <https://www.desmos.com/calculator/qhyco8cuou>

152: Inclination Angle 34

Question 423 Repeat Question 3 with the function

$$m = g(s) = \frac{s}{2} - \frac{s^3}{12}, \quad 0 \leq s \leq 3,$$

that expresses the slope of a trail in terms of the distance from the trailhead (measured along the trail in miles). The slope is measured with respect to the horizontal. It is positive (negative) when the trail slopes upward (downward) in the direction away from the trailhead.

Question 424 The function

$$v = g(t) = 20 + 40t - 3t^2, \quad 0 \leq t \leq 1,$$

expresses the speed of a car (in miles/hr) in terms of the number of hours past noon.

The function

$$G = h(t) = 30 + 10t, \quad 0 \leq t \leq 1$$

expresses the gas mileage (in miles/gal) of that same car in terms of the number of hours past noon.

The car has 10 gallons of gas in its tank at 12:15pm.

Answer the six parts of the previous questions in regard to approximating the number of gallons of gas in the tank at 1:00pm.

Geometric Integration

Evaluating or approximating definite integrals geometrically.

Directions: Do *not* use the Fundamental Theorem of Calculus for these problems. Instead evaluate an integral geometrically or approximate an integral with a Riemann sum with 100 equally-spaced intervals.

Question 425 For each definite integral, draw a graph of the appropriate function and shade the region of integration. Then use geometry to evaluate the integral.

(a) $\int_3^8 12 \, du$

(b) $\int_8^3 12 \, dt$

(c) $\int_a^b k \, dx$, where a, b, k are constants

(d) $\int_4^9 \frac{w}{2} \, dw$

(e) $\int_a^b \frac{z}{k} \, dz$, where a, b, k are constants

(f) $\int_{-10}^2 (4 - |x + 3|) \, dx$

(g) $\int_0^4 \sqrt{25 - x^2} \, dx$

(h) $\int_{-5}^5 \left(2 - \sqrt{25 - x^2}\right) \, dx$

Question 426 Make up a question in an applied scenario whose answer is the integral in Question 1(b) above and explain why the answer makes sense.

Learning outcomes:
Author(s):

Question 427 The function

$$r = f(t), 0 \leq t \leq 60,$$

expresses a balloon's rate of ascent (measured in ft/min) in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/tgi5yiuzab>

152:Balloon Introduction

The balloon is 1900 feet high at 12:10pm.

- (a) Write an expression for the balloon's height at 12:40pm. Then use the graph to approximate this height.
- (b) Write an expression for the balloon's height at 12:04pm. Then use the graph to approximate this height.
- (c) Find an expression for the function

$$h = g(t), 0 \leq t \leq 60,$$

that gives the balloon's height at time t minutes past noon.

Question 428 The function

$$r = f(t), 0 \leq t \leq 60,$$

expresses a balloon's rate of ascent (measured in ft/min) in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/vfk9toeayo>

152:Balloon 53

The balloon is 3200 feet high at 12:40pm.

- (a) Ignore the graph and write an expression with a for the balloon's height at 12:30pm. Then use the graph to find this height.
- (b) Ignore the graph and write an expression for the balloon's height at 12:10pm. Then use the graph to find this height.
- (c) When is the balloon the same height as it is at 12:09pm?

- (d) When is the balloon 20 feet higher than it is at 12:18pm?
- (e) Ignore the graph and find an expression for the function

$$h = g(t), 0 \leq t \leq 60,$$

that gives the balloon's height at time t minutes past noon. Then use the graph and find an explicit expression for this piece-wise defined function.

The Fundamental Theorem of Calculus, Part 2

FTC

Question 429 The function

$$v = f(t), 0 \leq t \leq 8,$$

expresses the speed (in feet/min) of a beetle in terms of the number of minutes past noon.

- (a) Write an expression for the distance the beetle crawls from 12:03pm to 12:07pm. Use the graph of the function $v = f(t)$ below to approximate this distance.
- (b) Write an equation you would solve to determine when the beetle is halfway to its destination. Start by defining an unknown (in a complete sentence, with units). Do not use t , it is already taken.
- (c) Use the graph of the function $v = f(t)$ below to approximate the time in part (b). But first determine whether this time is before or after 12:04pm.

Desmos link: <https://www.desmos.com/calculator/gd54bqgk6l>

152:Beetle 1

- (d) Suppose

$$v(t) = \frac{20}{(t+2)^2}.$$

and determine the exact time when the beetle is halfway to its destination.

Question 430 The function

$$r = f(t), 2 \leq t \leq 36,$$

gives the net rate (in gal/min) at which water flows into a tank at time t minutes past noon.

Learning outcomes:
Author(s):

The Fundamental Theorem of Calculus, Part 2

- (a) Write an expression that gives the change in volume from 12:15pm to 12:32pm.
- (b) Write an equation you would solve to find when the tank holds 22 fewer gallons than it does at 12:09pm. Start by defining an unknown (in a complete sentence, with units). Do not use t , it is already taken.
- (c) Use the graph of the function

$$r = f(t), 2 \leq t \leq 36,$$

shown below to approximate the time(s) when the tank holds 22 fewer gallons than it does at 12:09.

Desmos link: <https://www.desmos.com/calculator/rcsfzdhsax>

152:Tank 1

- (d) Suppose

$$f(t) = 12 - 3\sqrt{t}, 2 \leq t \leq 36.$$

Use calculus and algebra to find the exact time(s) in part (c). Then use a calculator to approximate these times to the nearest second.

Question 431 The function

$$m = f(s), 30 \leq s \leq 300$$

expresses your gas mileage (in miles/gallon) in terms of your distance from home. The distance is measured along your route in miles. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/v34aycmz93>

152:Mileage 89

Your tank has 12 gallons of gas when you are 50 miles from home.

- (a) Suppose you are driving away from home.
 - (i) Use the graph above to approximate the number of gallons of gas in your tank when you are 150 miles from home.
 - (ii) Find a function $G_1 = g_1(s)$ that expresses the number of gallons in your tank in terms of your distance (measured along your route in miles) from home.

- (b) Suppose you are driving home.
- (i) Use the graph above to approximate the number of gallons of gas in your tank when you are 150 miles from home.
 - (ii) Find a function $G_2 = g_2(s)$ that expresses the number of gallons in your tank in terms of your distance (measured along your route in miles) from home.

Question 432 The function

$$r = f(t) = 100/t, \quad a \leq t \leq b$$

expresses the rate (in gal/min) at which water flows into a tank in terms of the number of minutes past noon. The tank is empty at time $t = a$ minutes past noon and full at time $t = b$ minutes past noon.

- (a) Sketch a graph of the function $r = f(t)$. Label the axes with the appropriate units and variable names.
- (b) Would you expect the tank to be half-full before or after time $t = (a+b)/2$? Explain.
- (c) When is the tank half-full?
- (d) When does water flow into the tank at a rate equal the average rate at which it flows into the tank between $t = a$ and $t = b$?

Question 433 The function

$$r = f(t), \quad 0 \leq t \leq 12,$$

expresses a balloon's rate of ascent (in ft/min) in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/yp3awkyck8>

152:Balloon 11

- (a) When is the balloon the same height as it is at 12:02pm? Explain.
- (b) Use the graph to approximate the time(s) when the balloon is 20 feet higher than it was at 12:02pm.

The Fundamental Theorem of Calculus, Part 2

- (c) Write an equation whose solution(s) give the time(s) when the balloon is 20 feet higher than it was at 12:02pm.

- (d) Suppose

$$f(t) = 80 \sin\left(\frac{\pi}{6}t\right).$$

Use algebra to find exact solution(s) to your equation from part (c). Then use a calculator to approximate the clock time(s) to the nearest second.

The Fundamental Theorem of Calculus, Part 1

FTC

Question 434 The function $f(t)$, $0 \leq t \leq 12$, gives the rate (in ft/hour) at which the depth of the water in a large tank is changing at time t hours past noon. The water is 40 feet deep at 5:00pm.

Desmos link: <https://www.desmos.com/calculator/gnwlraiqhd>

152:Beetle 1

- (a) The graph of the function $r = f(t)$ is shown above. Label the axes with the appropriate variable names and units.
- (b) Find a function $h = g(t)$ that expresses the depth of the water (in feet) in terms of the number of hours past noon.
- (c) Sketch the graph of the function

$$\frac{dh}{dt} = \frac{d}{dt}(g(t)).$$

Label the axes with the appropriate variable names and units.

- (d) Find expressions that gives the minimum and maximum depths of water in the tank. Use the graph to approximate these depths.
- (e) Sketch a graph of the function $h = g(t)$. Label the axes with the appropriate variable names and units.
- (f) Write an equation you would solve to determine the time(s) when the water level is 10 feet lower than it is at 10:00pm. Use the graph to estimate this time(s). Explain your reasoning.

Question 435 The function

$$v = f(t), 0 \leq t \leq 6,$$

expresses the speed (in inches/sec) of a beetle in terms of the number of seconds past noon.

Learning outcomes:
Author(s):

- (a) Find a function $s = g(t)$ that expresses the distance (in inches) of the beetle to its destination in terms of the number of seconds past noon. The distance is measured along the route.
- (b) Find an expression for the derivative ds/dt . Give two answers to this question. First, an intuitive explanation that someone without any knowledge of calculus might understand. Then again, using the Fundamental Theorem.

- (c) Now Suppose

$$v = f(t) = \sqrt{6t - t^2}, \quad 0 \leq t \leq 6.$$

Desmos link: <https://www.desmos.com/calculator/cr9oays9zo>

152:Beetle Destination

- (i) Use the graph above to make a reasonably accurate graph of the function $s = g(t)$. Explain your reasoning.
- (ii) Use the geometry of the curve $v = f(x)$ to find an explicit expression (without the integral) for the function $s = g(t)$.
- (iii) Differentiate the function $s = g(t)$ to confirm your expression is correct.

Question 436 The function

$$V = \int_0^h A(w) dw, \quad 0 \leq h \leq 6,$$

expresses the volume (in ft^3) of water in a tank in terms of the water's depth (measured in feet). The graph of the function A is shown below.

Desmos link: <https://www.desmos.com/calculator/ae516hl7j6>

152:Beetle 1

- (a) Is the volume V above a function of w or h ? Explain.
- (b) What are the units of $A(w)$? How do you know?
- (c) Label the axes on the graph with their appropriate variable names and units.

Suppose the function $h = p(t)$, $0 \leq t \leq 12$, expresses the depth of the water (in ft) in terms of the number of hours past noon.

- (d) Find a function $V = k(t)$ that expresses the volume of water in terms of the number of hours past noon.
- (e) Find an expression for the rate of change in the volume of the water with respect to time. Explain this expression intuitively.
- (f) Now suppose

$$p(t) = 3 + 2 \cos\left(\frac{\pi}{12}t\right), \quad 0 \leq t \leq 12.$$

- (i) Is the volume increasing or decreasing at 6 pm? At what rate?
- (ii) Is the volume increasing or decreasing when the tank holds 44 ft³ of water? At what approximate rate?

1 Gas Consumption

Question 437 The function

$$r = g(s), \quad 0 \leq s \leq 200,$$

expresses the rate, in gallons/mile, at which your car burns gas in terms of the distance (in miles) from your home, as measured along your route on a 200-mile road trip to Spokane. When you are 50 miles from home, your tank has 10 gallons of gas. The trip takes four hours, and the function

$$v = w(t) = 64 - 6(t - 3)^2, \quad 0 \leq t \leq 4.$$

expresses your speed (in miles/hour) in terms of the number of hours past noon.

- (a) Find a function $G = f(s)$ that expresses the volume of gas (in gallons) your tank holds in terms of your distance from home. Include its domain.
- (b) Find a function $G = Q(t)$ that expresses the volume of gas in terms of the number of hours past noon. Include its domain.
- (c) Use the Fundamental Theorem to find an expression for the derivative dG/dt . Then explain this expression intuitively, in a way that a student without a knowledge of calculus could understand.
- (d) Use the graph of the function $r = g(s)$ shown below to find approximate answers to the following questions.

Desmos link: <https://www.desmos.com/calculator/tzmqpsznou>

152: FTC1 Gas

- (i) Label the axes above with the appropriate variable names and units.
- (ii) How many gallons of gas are in your tank at 2:00pm?
- (iii) At what rate, with respect to time, is your car burning gas when you are 62 miles from Spokane?
- (iv) What is your gas mileage at 2:00pm?
- (v) At what rate (with respect to time) is your gas mileage changing at 2:00pm?

2 On Trails

Question 438 The function

$$h = f(s) = 5 \sin \left(\frac{s}{5} \right), \quad 0 \leq s \leq 15,$$

expresses the altitude (in meters) of a beetle's trail in terms of the distance from the start (measured along the trail in meters).

The function

$$v = g(t) = \frac{t}{90} + \frac{t^2}{900}, \quad 0 \leq t \leq 30,$$

expresses the speed (in meters/min) of the beetle in terms of the number of minutes past noon (when it began its journey).

- (a) How far from the start is the beetle at 12:15pm?
- (b) Find the inclination angle of the trail beneath the beetle at 12:15pm.
- (c) How fast is the beetle crawling at 12:15pm?
- (d) Is the beetle gaining or losing altitude at 12:15pm? At what rate?
- (e) Express the rate at which the beetle's altitude is changing at 12:15pm in terms of some of the following derivatives. Include units for each derivative and explain their meanings.

- (i) $\left. \frac{dh}{ds} \right|_{s=15}$
- (ii) $\left. \frac{dh}{ds} \right|_{s=g(15)}$
- (iii) $\left. \frac{dh}{ds} \right|_{s=2.5}$

$$\begin{aligned} \text{(iv)} \quad & \left. \frac{dv}{dt} \right|_{t=15} \\ \text{(v)} \quad & \frac{d}{dt} \int_0^t g(u) du \end{aligned}$$

Question 439 The function

$$\theta = f(s), 0 \leq s \leq 15,$$

expresses the inclination angle (in radians) of a beetle's trail in terms of the distance from the start (measured along the trail in meters). The trail is 200 meters above sea level at the start.

The function

$$v = g(t) = 5 \sin(t/15), 0 \leq t \leq 30,$$

expresses the speed (in meters/min) of the beetle in terms of the number of minutes past noon (when it began its journey).

- (a) Find an expression for the function

$$h = w(t), 0 \leq t \leq 30,$$

that expresses the beetle's elevation in terms of the number of minutes past noon.

- (b) Find an expression for the derivative dh/dt and interpret its meaning.
 (c) Compare this question with the Question 5.

Question 440 The function

$$h = f(s), 0 \leq s \leq 60,$$

expresses the elevation (in meters) of a mountain road in terms of the distance (in kilometers) from the start.

The function

$$T = g(h), 0 \leq h \leq 5000,$$

expresses the temperature (in Celsius degrees) in terms of altitude (in meters) along the road.

- (a) Use the functions f and g to write an expression for the function

$$T = w(s), 0 \leq s \leq 60,$$

that expresses the temperature (in Celsius degrees) along the road in terms of the distance (in kilometers) from the start.

The Fundamental Theorem of Calculus, Part 1

- (b) Use the Leibniz differential notation to write an expression for the derivative

$$\left. \frac{dT}{ds} \right|_{s=54}$$

in terms of the appropriate derivatives of the functions f and g .

- (c) Use the graphs of the functions f and g below to approximate the temperature at the point on the road 54 kilometers from the start.
- (d) Use the sliders in Lines 1 and the graphs of the functions f and g below to approximate the derivative

$$\left. \frac{dT}{ds} \right|_{s=54}.$$

Include units.

- (e) Use the language of small changes to interpret the meaning of the above derivative.

Desmos link: <https://www.desmos.com/calculator/gcmfa2a6jz>

152: Elevation and Temperature

Desmos link: <https://www.desmos.com/calculator/ohqk3ak539>

152: Elevation and Temperature 2

Question 441 The function

$$h = f(s), 0 \leq s \leq 60,$$

expresses the elevation (in meters) of a mountain road in terms of the distance (in kilometers) from the start.

The function

$$r = k(h), 0 \leq h \leq 5000,$$

expresses the temperature gradient (in Celsius degrees/meter) in terms of altitude (in meters) along the road.

The temperature is 16.5°C at an elevation of 3400 meters.

- (a) Use the functions f and k to write an expression for the function

$$T = w(s), 0 \leq s \leq 60,$$

that expresses the temperature (in Celsius degrees) along the road in terms of the distance (in kilometers) from the start.

The Fundamental Theorem of Calculus, Part 1

- (b) Use the graphs of the functions f and k below to approximate the temperature at the point on the road 54 kilometers from the start.
- (c) Find an expression for the derivative

$$\left(\frac{d}{ds} \int_{3400}^{f(s)} k(h) dh \right) \Big|_{s=54}$$

and interpret its meaning. Do this twice. First using common sense. Then again using the Fundamental Theorem of Calculus, Part 1. Include units.

- (d) Use the sliders in Lines 1 and the graphs of the functions f and g below to approximate the derivative

$$\frac{dT}{ds} \Big|_{s=54}.$$

Include units.

Desmos link: <https://www.desmos.com/calculator/gcmfa2a6jz>

152: Elevation and Temperature

Desmos link: <https://www.desmos.com/calculator/vvr0qrgoh1>

152: Elevation and Temperature 3

Substitution and Parametrically-defined Curves

Substitution and parameterized curves

1 The Basics

Question 442 Which of the following definite integrals are equal to $\int_2^6 f(x) dx$?
 Avoid using a u -substitution. Instead think geometrically by describing the transformations that take the graph of $y = f(x)$ to the graph of each integrand. Assume f is continuous.

(a) $\int_4^{12} 2f(2x) dx$

(b) $\int_4^{12} (1/2)f(x/2) dx$

(c) $\int_1^3 (1/2)f(2x) dx$

(d) $\int_1^3 2f(2x) dx$

Question 443 The function

$$r = f(t), 0 \leq t \leq 12,$$

expresses a balloon's rate of ascent (in ft/min) in terms of the number of minutes past noon. Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/yp3awkyck8>

152:Balloon 11

(a) When is the balloon the same height as it is at 12:02pm? Explain.

Learning outcomes:
 Author(s):

Substitution and Parametrically-defined Curves

- (b) Use the graph to approximate the time(s) when the balloon is 20 feet higher than it was at 12:02pm.
- (c) Write an equation whose solution(s) give the time(s) when the balloon is 20 feet higher than it was at 12:02pm.
- (d) Suppose

$$f(t) = 80 \sin\left(\frac{\pi}{6}t\right).$$

Use a u -substitution and algebra to find exact solution(s) to your equation from part (c). Then use a calculator to approximate the clock time(s) to the nearest second.

Question 444 Between noon and 8pm the growth rate of a population of bacteria decreases exponentially. At noon the population increases at the rate of 700,000 bac/hr and at 3pm the population increases at the rate of 400,000 bac/hr.

The population is 2 million at noon.

- (a) Sketch by hand a rough graph of the function

$$r = f(t), \quad 0 \leq t \leq 8,$$

that expresses the growth rate (in hundreds of thousand of bacteria/hr) in terms of the number of hours past noon.

- (b) Find an expression for the function in part (a). Do not use e .
- (c) Use part (a) to sketch by hand a graph of the function

$$P = g(t), \quad 0 \leq t \leq 8,$$

that expresses the population in terms of the number of hours past noon.

- (d) Find an expression for the function in part (c).

Question 445 Identify for which of the following definite integrals a u -substitution would work. Make the substitution for these and evaluate the integral. Do this by changing the lower and upper bounds. Do not substitute back to the original variable of integration.

(a) $\int_0^{\sqrt{\pi}} \theta \sin(\theta^2) d\theta$

$$(b) \int_0^{\pi} \theta^2 \sin(\theta) d\theta$$

$$(c) \int_0^{\pi/2} \sin \theta \cos^3 \theta d\theta$$

$$(d) \int_0^{\pi/4} \tan \theta \sec^2 \theta d\theta$$

$$(e) \int_0^{\pi/4} \tan \theta \sec^3 \theta d\theta$$

$$(f) \int_0^2 x^2(1+x^2)^2 dx$$

$$(g) \int_0^2 x^2(1+x^3)^2 dx$$

$$(h) \int_1^2 \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$$

$$(i) \int_1^2 \sqrt{x} \cos(\sqrt{x}) dx$$

$$(j) \int_0^1 \frac{e^x}{(1+e^x)^2} dx$$

$$(k) \int_0^1 \frac{e^x}{1+e^x} dx$$

$$(l) \int_0^1 \frac{e^x}{1+e^{2x}} dx$$

$$(m) \int_2^3 \frac{x}{\ln x} dx$$

$$(n) \int_2^3 \frac{\ln x}{x} dx$$

2 A Geometric Interpretation

To understand what u -substitution does geometrically, it helps to think about making a *reverse* u -substitution. This means starting with a simple integral and making it more complicated with a substitution. Here's an example.

Example 73. We'll start with the integral

$$I = \int_0^{\pi} \sin u \, du$$

and make the substitution

$$u = t^2.$$

This substitution really reparameterizes the curve

$$y = f(u) = \sin u, \, 0 \leq u \leq \pi$$

from the standard parameterization

$$(x, y) = (u, \cos u), \, 0 \leq u \leq \pi$$

to the new parameterization

$$(x, y) = (t^2, \cos(t^2)), \, 0 \leq t \leq \sqrt{\pi}.$$

The substitution maps the rectangle with height $y = \sin u$ and width du to the rectangle with the same height $y = \sin(t^2)$ and width

$$du = 2t \, dt.$$

The effect is to transform a partition of the interval of integration, originally $u \in [0, \pi]$ with equal subintervals into a partition of the new interval of integration $t \in [0, \sqrt{2\pi}]$ with *unequal* subintervals $du = 2t \, dt$. You can see this transformation by dragging the slider w in Line 2 of the worksheet below from $w = 0$ to $w = 1$.

Desmos link: <https://www.desmos.com/calculator/toiwzzqkva>

152: Sub 2

The next example puts this idea in more a more meaningful context.

Example 74. The function

$$r = f(s), \, 0 \leq s \leq 240,$$

expresses the rate at which a car burns gas (in gal/mile) in terms of the trip odometer reading (in miles) during a 240-mile trip.

The car has 14 gallons of gas at odometer reading 100 miles.

The function

$$s = g(t), \, 0 \leq t \leq 4,$$

expresses the car's trip odometer reading (in miles) in terms of the number of hours since the start of the trip.

Substitution and Parametrically-defined Curves

- (a) Find a function

$$G = w(s), 0 \leq s \leq 240,$$

that expresses the number of gallons of gas in the car in terms of the trip odometer reading.

- (b) Make a substitution in your expression from part (a) to find a function

$$G = h(t), 0 \leq s \leq 240,$$

that expresses the number of gallons of gas in the car in terms of the number of hours since the start of the trip.

- (c) Your function for part (b) should have been

$$G = h(t) = 14 - \int_{100}^t f(g(u))g'(u) du.$$

Interpret the integral in this expression in two different ways.

- (i) By writing it as

$$\int_{100}^t f(g(u))(g'(u) du)$$

and describing the meaning of the product $g'(u) du$.

- (ii) By keeping its original form and describing the meaning of the product $f(g(u))g'(u)$.

- (d) Drag the slider w in Line 2 of the worksheet below to change the odometer reading function g . Describe what happens.

Desmos link: <https://www.desmos.com/calculator/gsdzd01bva>

Substitution and Gas Consumption 2

Example 75. The previous example assumed the car's gas mileage depended only on the trip odometer reading and not on its speed.

A more realistic example would look at the elevation change over a section of trail, the change being independent of how you traverse the trail.

Make up your own example like this. Start with the function $\theta = f(s)$ that expresses the trail's inclination angle in terms of the distance from the trailhead (measured along the trail in km).

The Limits to Growth

Depletion of the world's resources.

The Project on the Predicament of Mankind was initiated in 1968 by a group of 30 individuals having a wide range of expertise with the intent of examining the complex problems troubling all nations at that time and that still persist today. Within a year they had created a model of the five basic factors they identified as limiting growth - population, agricultural production, natural resources, industrial production, and pollution. The model provided valuable insights into how these areas interact with each other to limit growth. The findings are summarized in the 1972 classic *The Limits to Growth*, available at

Limits to Growth .

Here we focus on their analysis of the depletion of the world's chromium reserves as illustrated in the figure below.

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/9kqmbczco7>

Chromium is just one of 19 non-renewable natural resources analyzed. For each, the authors provide the following information.

- The known global reserves in 1970; for chromium 7.75×10^8 tons
- The static index. This is the number of years it the global reserves would last at the current global consumption rate. The static index for chromium is 420 years.
- The projected relative growth rate in the global consumption rate. For chromium, this relative instantaneous growth rate is 2.6%/yr, compounded continuously.

For each resource, the authors compute the *exponential index*. This is the number of years the known current reserves (in 1970) would last with consumption growing exponentially at assumed relative rate. They also compute the exponential index using five times the known reserves.

Learning outcomes:
Author(s):

Question 446 Let's work in general (like the authors) and let r be the relative instantaneous growth rate (measured in yr^{-1}) in the global consumption rate and let s be the static index. Our goal here is to express the exponential index in terms of r and s .

Access Desmos interactives through the online version of this text at

.

Desmos link: <https://www.desmos.com/calculator/er2jq9vzeq>

Access this worksheet at

Limits to Growth 2

We'll let R_0 be the 1970 known global reserves (in units of hundreds of millions of tons). Use this parameter along with the parameters r and s as need be, to find expressions for the following:

- (a) The usage rate function

$$u = f(t), 0 \leq t \leq 240,$$

that expresses the usage rate (measured in 10^8 tons/year) in terms of the number of years since 1970. Recall the assumption that the usage rate increases at the relative instantaneous rate of $r \text{ yr}^{-1}$. This means that

$$u = f(t) = u_0 e^{rt}, 0 \leq t \leq 240$$

where u_0 is the usage rate in 1970. All that you need to do is to express u_0 in terms of R_0 and the static index s .

This function is graphed above, but it does not really belong on the same coordinate system as the others. Why not?

- (b) The remaining reserves function

$$R = g(t), 0 \leq t \leq 240,$$

that expresses the remaining global reserves (in hundreds of millions of tons) in terms of the number of years since 1970, assuming the usage rate function in part (a).

- (c) The exponential index in terms of r and s .

- (i) Check that your expression for the exponential index is dimensionally correct.
- (ii) Use the values of r and s for Chromium given above to determine when the model predicts the world's Chromium reserves will run out. Check your result with the appropriate graph above.

Area

Computing the area of regions in the plane bounded by graphs of functions.

Directions: When computing the area of a region, be sure to do the following.

- State the direction in which you decide to slice the region.
- Sketch a differential rectangle and label the coordinates of its endpoints.
- State the area of the differential rectangle.
- Explain the role of the definite integral in computing the area of the region.

1 Parabolas and Power Functions, Part 1

Question 447 Let $a, b > 0$ be constants, measured in meters.

- (a) Find the area of the region bounded by the x -axis and the parabola

$$y = b - \frac{x^2}{a},$$

where x and y are measured in meters.

Include units and check that your answer is dimensionally correct. Do this twice:

- first, by slicing the region perpendicular to the y -axis,
- and again, by slicing the region perpendicular to the x -axis

Question 448 Express the area of the region shown below bounded by a parabola and a line perpendicular to its axis of symmetry in terms of the height h and the length of the base b .

Desmos link: <https://www.desmos.com/calculator/kkdfpr66rt>

152: Parabola 1

Learning outcomes:
Author(s):

Question 449 Use ideas of differential calculus, not the Fundamental theorem to answer Question 2 by comparing the areas of the differential rectangles shown below.

Desmos link: <https://www.desmos.com/calculator/pvskgbpnew>

152: Parabola 2

Desmos link: <https://www.desmos.com/calculator/0w37bixe79>

152: Parabola 3

Question 450 Let $b, h > 0$ be constants with units of meters and let $a = b/2$. Suppose the curve $y = cx^8$ passes through the point A with coordinates $(b/2, h) = (a, h)$. We wish to express the area of the shaded region below bounded by the curve and the line $y = h$ in terms of its base b and height h . Do this as follows.

- Express c in terms of a and h .
- Express the area of the region first in terms of a and h , and then in terms of b and h .
- Check that your expression for the area has the correct units.
- Interpret your expression for the area geometrically. Does it seem reasonable? Compare it with the area of a triangle and with the area of a rectangle.
- Find an equation of the horizontal line that bisects the area of the region. Type your equation in Line 2 above as a check.

Desmos link: <https://www.desmos.com/calculator/egdpe7cccl>

152: Power Function 3

Question 451 Let $a, b > 0$ be constants with units of meters.

- Use the Fundamental Theorem to evaluate the definite integral

$$\int_0^b \frac{x^2}{a} dx.$$

Include units.

- (b) Interpret the result of part (a) geometrically.

Question 452 Let $a, b, c \in \mathbb{R}$ be constants with units of meters.

Find the area of the region bounded by the x -axis and the parabola

$$y = c + \frac{(x - b)^2}{a},$$

where the coordinates (x, y) are also measured in meters.

2 Parabolas, Part 2

Question 453 (a) Find the area of the region bounded by the parabola $y = x^2$ and the line $y = 2x + 3$. Start by deciding how to slice the region and computing the area of a differential slice.

- (b) Drag the slider w in Line 1 of the worksheet below from $w = 0$ to $w = 1$ and describe what you see.

Desmos link: <https://www.desmos.com/calculator/vci24vc4f1>

152: Area Parabolic Segment

- (c) Find an expression for the area of the region bounded by the parabola $y = x^2$ and the line through the points (a, a^2) and (b, b^2) .

Question 454 Let $\mathcal{P}$ be the parabola symmetric about the y -axis with its vertex at the origin that passes through the point (a, h) . Assume $a, h > 0$.

- Express the area of the region bounded by the parabola and the line $y = h$ in terms of its height h and its base $b = 2a$.
- Check that your expression for the area has the correct units.
- Interpret your expression for the area geometrically. Does it seem reasonable? Compare it with the area of the surrounding rectangle.
- Find an equation of the horizontal line that bisects the area of the region bounded by $\mathcal{P}$ and the line $y = h$.
- Check that your equation is dimensionally correct.

- (f) Enter your equation for the bisecting line in Line 18 of the worksheet below as a check.

Desmos link: <https://www.desmos.com/calculator/mbxrsmkho8>

152: Parabola 23

Question 455 Let $\mathcal{P}$ be the parabola symmetric about the y -axis with its vertex at the origin that passes through the point (a, h) . Assume $a, h > 0$.

- Find an expression (in terms of a and h) for the area of the region bounded by the x -axis, the parabola $\mathcal{P}$, and the tangent line to the parabola at P . Check that your expression is dimensionally correct. Start by slicing the region perpendicular to the y -axis.
- Find an expression for the area of the region in part (a) in terms of its base $b = OQ$ and height h .
- What fraction of triangle $\triangle AQP$ does the region in part (a) occupy?
- Compare the area of the region in part (a) with the area of the region bounded by the parabola and the segment OP .

Desmos link: <https://www.desmos.com/calculator/avl2r9dubt>

152: Parabola and Tangent Line 3

Explanation. (a) Let $y = cx^2$ be an equation of the parabola. Then since the point (a, h) lies on the parabola

$$h = ca^2$$

and

$$c = \frac{h}{a^2}.$$

So the parabola has equation

$$y = \frac{h}{a^2}x^2.$$

The left and right endpoints of a horizontal slice AB of the region share the same y -coordinate. Call it y . Since the left endpoint $A(x, y)$ lies on the parabola,

$$x = \pm \sqrt{\frac{ya^2}{h}}.$$

But because A lies in the first quadrant, $x \geq 0$ and A has coordinates

$$\left(\sqrt{\frac{ya^2}{h}}, y \right).$$

To express the x -coordinate of the right endpoint B in terms of y , we need an equation of the tangent line to the parabola at the point $P(a, h)$.

The slope of this tangent line is

$$\begin{aligned} \frac{d}{dx} \left(\frac{h}{a^2} x^2 \right) \Big|_{x=a} &= \frac{2h}{a^2} x \Big|_{x=a} \\ &= \frac{2h}{a}. \end{aligned}$$

So an equation of the tangent line is

$$y = h + \frac{2h}{a} (x - a) = \frac{2h}{a} x - h$$

and B has coordinates

$$\left(\frac{a}{2h} y + \frac{a}{2}, y \right).$$

Now the differential rectangle AB has length

$$\frac{a}{2h} y + \frac{a}{2} - \sqrt{\frac{ya^2}{h}},$$

width dy and area

$$dA = \left(\frac{a}{2h} y + \frac{a}{2} - \sqrt{\frac{ya^2}{h}} \right) dy.$$

Finally, the area of the region bounded by the x -axis, the parabola, and the tangent line at (a, h) is

$$\begin{aligned} \int_0^h \left(\frac{a}{2h} y + \frac{a}{2} - \sqrt{\frac{ya^2}{h}} \right) dy &= \left(\frac{a}{4h} y^2 + \frac{a}{2} y - \frac{2}{3} \sqrt{\frac{a^2}{h}} y^{3/2} \right) \Big|_{y=0}^{y=h} \\ &= \frac{ah}{4} + \frac{ah}{2} - \frac{2ah}{3} \\ &= \frac{1}{12} ah. \end{aligned}$$

3 Which Area is Greatest? Least?

Question 456 The vertical segments below are equally-spaced. The horizontal segments cut the shaded region into 20 smaller regions. Which of these has the greatest area? The least?

Desmos link: <https://www.desmos.com/calculator/wsgteg55ta>

152: Self-calibrating area

Question 457 The vertical segments below are equally-spaced. The horizontal segments cut the shaded region into 20 smaller regions. Which of these has the greatest area? The least?

Desmos link: <https://www.desmos.com/calculator/dir9hpszvse>

152: Self-calibrating area 2

4 Trigonometric Functions

Question 458 Let $a, b, k > 0$ be constants with units of cm.

Find a simplified expression for the area of the shaded region below bounded by the y -axis and the curves $y = a \cos(x/k)$ and $y = b \sin(x/k)$. Check your expression is dimensionally correct.

The area is

$$k(\sqrt{a^2 + b^2} - b).$$

Desmos link: <https://www.desmos.com/calculator/ps1m0hksl0>

152: Sine Cosine Area

Explanation:

Suppose the point of intersection has coordinates

$$(x_0, y_0) = (x_0, a \cos \theta_0) = (x_0, b \sin \theta_0),$$

where

$$\theta_0 = x_0/k.$$

Then

$$a \cos \theta_0 = b \sin \theta_0$$

and

$$\tan \theta_0 = \frac{a}{b}.$$

Since $0 < \theta_0 < \pi/2$, we can draw a right triangle with θ_0 as one of the acute angles, a the length of the side opposite this angle and b the length of the adjacent side. From this we know that

$$\cos \theta_0 = \frac{b}{\sqrt{a^2 + b^2}}$$

and

$$\sin \theta_0 = \frac{a}{\sqrt{a^2 + b^2}}.$$

Now slice the region into differential rectangles perpendicular to the x -axis. Then the general differential rectangle PQ with endpoints $P(x, a \cos(x/k))$ and $Q(x, b \sin(x/k))$ has differential area

$$dA = (a \cos(x/k) - b \sin(x/k)) dx.$$

So the area of the region is

$$\begin{aligned} A &= \int_0^{x_0} \left(a \cos\left(\frac{x}{k}\right) - b \sin\left(\frac{x}{k}\right) \right) dx \\ &= \left(ak \sin\left(\frac{x}{k}\right) + bk \cos\left(\frac{x}{k}\right) \right) \Big|_{x=0}^{x=x_0} \\ &= k \left(\frac{a^2}{\sqrt{a^2 + b^2}} + \frac{b^2}{\sqrt{a^2 + b^2}} - b \right) \\ &= k(\sqrt{a^2 + b^2} - b). \end{aligned}$$

5 The Rectangular Elastica

Let $a > 0$ be a constant with units of meters and let

$$f(x) = \int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du.$$

Question 459 *Desmos link:*

<https://www.desmos.com/calculator/c3vderglut>

152: Elastica

- (a) Express the coordinates of the x intercept of the curve $y = f(x)$ as some multiple of a . Use a Riemann sum with 1000 equal subintervals to approximate the multiple.
 - (b) Express the exact and approximate coordinates of the y -intercept in terms of a .
 - (c) Find an equation of the tangent line to the curve $u = f(x)$ at the point where the line $x = a/2$ intersects the curve.
 - (d) Find an expression for the area bounded by the curve $y = f(x)$ and the y -axis. Activate the Hint folder in Line 1 of the worksheet above for a hint.
-

Volumes of Solids in Three-Dimensions

Volume.

1 Square Cross Sections

Question 460 *The base of a solid is a disk of radius a meters. Parallel cross-sections perpendicular to the base are squares.*

- (a) *Find the volume of the solid.*
- (b) *Compare the solid's volume with the volume of the surrounding cylinder.*

Desmos3D link: <https://www.desmos.com/3d/q5jo9tip5q>

152: Circular Base Square Cross Sections

Explanation.

It is not necessary to visualize the solid to compute its volume. Instead, the best approach is to sketch its base.

Desmos link: <https://www.desmos.com/calculator/mnjevsoffp>

152: Solid Base 1

We can image the square cross-sections coming out of the page, but the key is to start by computing the area of the cross-section. The square cross-section has side length $s = 2y$ and area

$$A = s^2 = 4y^2.$$

The differential slice of the solid has width dx and differential volume

$$dV = A dx = 4y^2 dx.$$

Before integrating, we'll need to eliminate the mixed variables and express the differential volume in terms of x and dx . To do this we need to use the fact that the base is bounded by the circle with equation

$$x^2 + y^2 = a^2$$

Learning outcomes:
Author(s):

in the above coordinate system. So

$$y^2 = a^2 - x^2$$

and the differential slice has volume

$$\begin{aligned} dV &= 4y^2 dx \\ &= 4(a^2 - x^2) dx. \end{aligned}$$

We add these differential volumes to get the volume of the solid,

$$\begin{aligned} V &= \int_{-a}^a 4(a^2 - x^2) dx \\ &= 8 \int_0^a (a^2 - x^2) dx \\ &= 8 \left(a^2 x - \frac{1}{3} x^3 \right) \Big|_{x=0}^{x=a} \\ &= \frac{16}{3} a^3. \end{aligned}$$

Activate the folder in Line 1 above to see the cylinder that surrounds the solid.

This cylinder has volume

$$W = (2a)(\pi a^2) = 2\pi a^3 \sim 6.28a^3.$$

So the solid occupies

$$\frac{V}{W} = \frac{16}{6\pi} \sim 84.9\%$$

the volume of the cylinder.

Question 461 (a) Solve Question 1 if the cross-sections are semi-circles (with their diameters in the base) instead of squares.

(b) Describe the solid in this case.

(c) What fraction of the cylinder's volume below is occupied by the tennis balls?

Desmos link: <https://www.desmos.com/calculator/5ip866uqox>

152: Tennis Balls

Question 462 Let $a, b > 0$ be constants.

The base of a solid is the region bounded by the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

Cross-sections perpendicular to the x -axis are squares.

Desmos3D link: <https://www.desmos.com/3d/zbcupsnbjv>

152: Elliptical Base Square Cross Sections

- (a) Find the volume of the solid.
- (b) Check that your expression for the volume is dimensionally correct.
- (c) Compare the solid's volume with the volume of the surrounding cylinder.

Question 463 The base of a tent is a regular hexagon with side length a meters. Three poles bent into semicircles run between opposite vertices. Cross sections parallel to the base are regular hexagons having their vertices on the poles.

Desmos3D link: <https://www.desmos.com/3d/wczn2fm141>

152: Tent

- (a) Find an expression for the volume of the tent.
- (b) Compare the volume with the volume of the surrounding cylinder.
- (c) The tent is partially filled with water that takes up half the tent's volume. What is the approximate depth of the water?

Question 464 The base of a solid is the parabolic region below with base $b = 2a$ and height h . Cross sections perpendicular to the y -axis are rectangles half as tall as they are wide

Desmos link: <https://www.desmos.com/calculator/mbxrsmkho8>

152: Parabola 23

Desmos3D link: <https://www.desmos.com/3d/gdebhncchrz>

152: Parabolic Base

- (a) Find an expression for the volume of the solid.
- (b) Check that your expression is dimensionally correct.
- (c) Compare the solid's volume with the volume of the surrounding cylinder.

2 The Hoof of Archimedes

Question 465 A solid circular cylinder of radius a and height h is sliced by a plane passing through a diameter of the base and intersecting the top of the cylinder in exactly one point. The plane cuts the cylinder into two pieces. Our goal here is to find an expression in h and a for the volume of the lower (smaller) piece.

Desmos3D link: <https://www.desmos.com/3d/cnspcoiggo>

152: Archimedes Hoof 2

We'll determine the volume of the lower piece (the hoof of Archimedes) by taking slices perpendicular to the coordinate axes. Turn off the folder in Line 2 and activate the appropriate folder in Lines 5, 13, or 22 if you need help visualizing the cross sections.

Slice the solid

- (a) perpendicular to the x -axis (activate the folder in Line 5),
 - (b) perpendicular to the y -axis (activate the folder in Line 13), and
 - (c) perpendicular to the z -axis (activate the folder in Line 22).
- (a) Check that your expression for the volume is dimensionally correct.
 - (b) What fraction of the original cylinder does the hoof occupy? Activate the folder in Line 33 to see the cylinder.

3 Archimedes' Bicylinder

Question 466 Two solid right circular cylinders of radius a cm intersect orthogonally as shown below (the cylinders' axes of symmetry also intersect). We'll determine the volume of the solid common to both (Archimedes' bicylinder).

Desmos3D link: <https://www.desmos.com/3d/wzzhod4ya6>

152: Bicylinder

- (a) Find an expression for the volume of the bicylinder and check that it is dimensionally correct.
- (b) Compare the volume with its surrounding cylinders to make sure your answer is reasonable.
- (c) Compare the volume of the bicylinder with the volume of the solid in Question 1 above. Then drag the slider w_2 in the worksheet below and describe what you see.

Desmos3D link: <https://www.desmos.com/3d/cwcjqkjndw>

152: Square Cross-sections to Bicylinder

Question 467 Turn off the folders in Line 1 (Bicylinder) and Line 5 (Hoofs) below. Then use your expression for the volume of the hoof of Archimedes to compute the volume of the bicylinder.

Desmos3D link: <https://www.desmos.com/3d/phy3gfnhcj>

152: Bicylinder and Hoof

4 Triangles and Cones

Question 468 Use integration to find an expression for the area of $\triangle ABC$ below in terms of its base $b = AC$ and height $h = BZ$.

Do this as follows:

Volumes of Solids in Three-Dimensions

- (a) Establish a one-dimensional coordinate system as shown below.
- (b) Use similar triangles to express the length QR in terms of $w = BT$, h , and b .
- (c) Find an expression for the differential area of the slice QR .
- (d) Integrate to find the area of the triangle.

Desmos link: <https://www.desmos.com/calculator/jly5a1jr9d>

152: Triangle Area

Question 469 Use integration to express the volume of the red cone in terms of the area A of its base and its height h .

Do this as follows:

- (a) Establish a one-dimensional coordinate system as shown below.
- (b) Use similar cones to express the area of the cross-section at coordinate w in terms of w , h , and A .
- (c) Find an expression for the differential volume of the slice.
- (d) Integrate to find the volume of the cone.

Desmos3D link: <https://www.desmos.com/3d/6pzlieihca>

152: Cone

Desmos3D link: <https://www.desmos.com/3d/mwkyamaixz>

Proof Without Words: Pyramid with Square Base

The Hoof of Archimedes

Finding the volume of the hoof, several ways.

Question 470 A solid circular cylinder of radius a and height h is sliced by a plane passing through a diameter of the base and intersecting the top of the cylinder in exactly one point. The plane cuts the cylinder into two pieces. Our goal here is to find an expression in h and a for the volume of the lower (smaller) piece.

Desmos3D link: <https://www.desmos.com/3d/cnspcoiggo>

152: Archimedes Hoof 2

We'll determine the volume of the lower piece (the hoof of Archimedes) by taking slices perpendicular to the coordinate axes. Turn off the folder in Line 2 and activate the appropriate folder in Lines 5, 13, or 22 if you need help visualizing the cross sections.

- (a) Compute the volume by slicing the solid perpendicular to the x -axis. To see the cross-sections, turn off the folder Slanted Face in Line 2 and activate the folder Cross-sections perpendicular to the x -axis in Line 5).

Explanation. Let the top vertex of the red triangle have coordinates (x, y, z) . Then the triangle has area

$$A = yz/2$$

and the thin slice has differential volume

$$dV = yz/2 \, dx.$$

To express the differential volume in terms of x , first use the Pythagorean theorem to express y in terms of x to get

$$y = \sqrt{a^2 - x^2}.$$

Then use similar triangles (red and green) to get

$$\frac{z}{y} = \frac{h}{a}.$$

Learning outcomes:
Author(s):

and express z in terms of y as

$$z = \frac{yh}{a}.$$

Now substitute the previous expression for y to express the area of the cross-section in terms of x ,

$$A = \frac{h}{2a} (a^2 - x^2).$$

So the differential volume of the thin slice is

$$dA = \frac{h}{2a} (a^2 - x^2) dx.$$

The volume of the hoof is then

$$V = 2 \int_0^a \frac{h}{2a} (a^2 - x^2) dx = \frac{2}{3} a^2 h.$$

- (b) *Now compute the volume by slicing perpendicular to the y -axis. To see the slices turn off the folder in Line 5 and activate the folder in Line 13.*

Explanation. In terms of the coordinates (x, y, z) of the top right vertex of the rectangular cross-section, its area is

$$A = 2xz.$$

So the volume of the differential slice is

$$dV = 2xz dy.$$

In terms of y , the differential volume is

$$dV = \frac{2h}{a} y \sqrt{a^2 - y^2} dy.$$

So the volume of the hoof is

$$V = \frac{2h}{a} \int_0^a y \sqrt{a^2 - y^2} dy = \frac{2}{3} a^2 h.$$

- (c) *Lastly, compute the volume by slicing perpendicular to the z -axis. To see the slices turn off the folder in Line 13 and activate the folder in Line 22.*

Explanation. A common error is to think the cross-sections are semicircles. They are not. Drag the slider z in Line 2 of the worksheet below to see how the cross-sections vary with the z -coordinate of the slice.

Desmos link: <https://www.desmos.com/calculator/gxhq1yelvv>

152: Hoof Cross Sections

The first step in finding an expression for the area of the cross-section in terms of the coordinates (x, y, z) of the point on the surface directly above A is to find an expression for the angle $\theta = \angle AOW$ above.

This angle has measure

$$\theta = \arccos\left(\frac{y}{a}\right).$$

So the sector of the circle subtended by $\angle AOB$ has area

$$A_1 = a^2 \arccos\left(\frac{y}{a}\right)$$

and $\triangle AOB$ area

$$A_2 = y\sqrt{a^2 - y^2}.$$

Then the slice with differential thickness dz has differential volume

$$dV = a^2 \left(\arccos\left(\frac{y}{a}\right) - y\sqrt{a^2 - y^2} \right) dz.$$

Now to get rid of the mixed variables, we'll do something different and express dz in terms of dy . We know

$$z = \frac{yh}{a},$$

so

$$dz = \frac{h}{a} dy.$$

Then the hoof has volume

$$V = \frac{h}{a} \int_0^a \left(\arccos\left(\frac{y}{a}\right) - y\sqrt{a^2 - y^2} \right) dy.$$

- (d) Check that your expression for the volume is dimensionally correct.
- (e) What fraction of the original cylinder does the hoof occupy? Activate the folder in Line 33 to see the cylinder.

Volumes, Part 2

Volume.

1 Paraboloid of Revolution

Question 471 (a) Find an expression for the area of the region below bounded by the parabola and the line $y = h$ in terms of the height h and base b of the region.

Desmos link: <https://www.desmos.com/calculator/ftsereyryj4>

152: Parabola 23B

- (b) What fraction of the area of the surrounding rectangle does the parabolic region occupy?
- (c) Rotate the region above about the y axis and you get a paraboloid of revolution as illustrated below.

Desmos3D link: <https://www.desmos.com/3d/1bculfxxtt>

152: Paraboloid

- (d) Find an expression for the volume of the paraboloid in terms of its height h and the radius a of its base.
- (e) Click the wrench in the upper right corner of the worksheet above and check the Translucent Surfaces box. Then activate the folder Cylinder in Line 3.
- (f) What fraction of the volume of the surrounding cylinder does the paraboloid occupy? Does this look correct?
- (g) How can you explain the difference between the fractions in parts (b) and (f)?
- (h) One last question. Suppose the paraboloid is a tank that is partly filled with water that takes up half the volume of the tank. Express the depth of the water in terms of the height h of the tank.

Drag the slider b in Line 3 of the worksheet below to check your work.

Learning outcomes:
Author(s):

Desmos3D link: <https://www.desmos.com/3d/ejuarpozjx>

152: Paraboloid 2

Explanation. To answer part (h), let b be the depth of the the water (in meters) that takes up half the volume of the tank and let $y = cx^2$ be an equation of the parabola in the first worksheet above.

The segment AB with endpoints $A(0, y)$ on the y -axis and $B(x, y)$ on the parabola sweeps out a disk with radius x and area

$$A = \pi x^2.$$

So the corresponding differential slice with thickness dy has volume

$$dV = \pi x^2 dy.$$

But since $B(x, y)$ lies on the parabola $y = cx^2$,

$$x^2 = y/c.$$

and

$$dV = \pi x^2 dy = \frac{\pi}{c} y dy.$$

Summing these differential volumes gives the volume

$$V_t = \int_0^h \frac{\pi}{c} y dy$$

of the tank.

Similarly the volume of the water with depth b is

$$V_w = \int_0^b \frac{\pi}{c} y dy.$$

Since the water takes up half the volume of the tank,

$$\int_0^b \frac{\pi}{c} y dy = \frac{1}{2} \int_0^h \frac{\pi}{c} y dy.$$

So

$$\int_0^b y dy = \frac{1}{2} \int_0^h y dy$$

and

$$\frac{b^2}{2} = \frac{h^2}{4}.$$

Then since $b, h > 0$,

$$b = \frac{h}{\sqrt{2}}.$$

So the tank is half full when the depth of the water is $h/\sqrt{2} \sim 0.707h$ meters.

2 Napkin Ring

Question 472 (a) Start with a solid sphere of radius a as shown below.

Desmos3D link: <https://www.desmos.com/3d/dw5gjr0es4>

152: Napkin Ring

- (b) Click the wrench in the upper right corner of the worksheet above and check the Translucent Surfaces box.
- (c) Now drill a circular hole of height $2h$ through the center of the sphere. To see this:
 - (i) Activate the folder Cylinder in Line 3 above.
 - (ii) Then activate the folder Napkin Ring in Line 5.
 - (iii) Then turn off the folders Sphere and Cylinder in Lines 1 and 3.
- (d) We now go about expressing the volume of the remaining solid (a napkin ring) in terms of a and h .
 - (i) We start by slicing the solid perpendicular to the axis of rotation as illustrated below. Drag the slider w in Line 2 to translate the cross-section. Turn off the folder Napkin Ring in Line 7 to better see the cross-sections.

Desmos3D link: <https://www.desmos.com/3d/ubyhmxtccz>

152: Napkin Ring 2

- (ii) To find the area of a cross-section, it helps to drop down a dimension and work in the plane.

Desmos link:

<https://www.desmos.com/calculator/6gylqexlh5>

152: Napkin Ring Cross Section

- (iii) Express the area of the annulus swept out by segment AB as it rotates about the z -axis in terms of the common z -coordinate of points A and B .
- (iv) Then find an expression for the differential volume swept out by the differential rectangle AB with thickness dz .
- (v) Finally, find an expression for the volume of the napkin ring.
- (e) Drag the slider a in Line 12 of the first worksheet to change the radius of the original sphere. How does the volume of the napkin ring change as you vary a ? Try to explain this geometrically by opening the Tangent Cluster folder and dragging the same slider a in Line 6 of the second worksheet.

The Mylar Balloon

The mylar balloon.

1 The Rectangular Elastica

Question 473 Let $a > 0$ be a constant with units of meters and for $x > 0$ define

$$f(x) = \int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du.$$

Desmos link: <https://www.desmos.com/calculator/c3vderglut>

152: Elastica

- Find the coordinates of the x -intercept of the curve $y = f(x)$.
- Find the exact coordinates of the y -intercept. Then make a substitution to express the x -intercept as a multiple of a . Then use Line 3 in the worksheet above to approximate the multiple.
- Find an equation of the tangent line to the curve $u = f(x)$ at the point where the line $x = a/2$ intersects the curve. Enter this equation in Line 4 above. Use the point-slope equation of a line, not slope-intercept.
- Find an expression for the area bounded by the curve $y = f(x)$ and the coordinate axes. Activate the Hint folder in Line 1 of the worksheet above for a hint.
- Compare the area in part (d) with the area of the surrounding rectangle. Does the ratio seem reasonable?
- Activate the folder Rectangle in Line 5. Compare the area of this rectangle with the area in part (d).

Explanation. (a) The coordinates of the x -intercept are the roots of the equation

$$f(x) = \int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du = 0, \quad x \geq 0.$$

Since the integrand is positive for $u > 0$, the integral is zero if and only if $x = a$. So the x -intercept has coordinates $(a, 0)$.

Learning outcomes:
Author(s):

- (b) The
- y
- intercept has
- y
- coordinate

$$f(0) = \int_0^a \frac{u^2}{\sqrt{a^4 - u^4}} du = \frac{1}{a^2} \int_0^a \frac{u^2}{\sqrt{1 - \left(\frac{u}{a}\right)^4}} du.$$

Making the substitution $v = u/a$ and some algebra leads to

$$f(0) = a \int_0^1 \frac{v^2}{\sqrt{1 - v^4}} dv \sim 0.599a.$$

So the y -intercept has coordinates

$$\left(0, a \int_0^1 \frac{v^2}{\sqrt{1 - v^4}} dv\right) \sim (0, 0.599a).$$

- (c) Since

$$y = f(x) = \int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du = - \int_a^x \frac{u^2}{\sqrt{a^4 - u^4}} du,$$

by the fundamental theorem

$$\begin{aligned} \frac{dy}{dx} &= - \frac{d}{dx} \int_a^x \frac{u^2}{\sqrt{a^4 - u^4}} du \\ &= - \frac{x^2}{\sqrt{a^4 - x^4}}. \end{aligned}$$

So at the point $(a/2, f(a/2))$, the tangent line has slope

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{x=a/2} &= - \frac{\left(\frac{a}{2}\right)^2}{\sqrt{a^4 - \left(\frac{a}{2}\right)^4}} \\ &= -1/\sqrt{15}. \end{aligned}$$

So at the point $(a/2, f(a/2))$ the tangent line to the curve has equation

$$y = \int_{a/2}^2 \frac{u^2}{\sqrt{a^4 - u^4}} du - \frac{1}{\sqrt{15}} \left(x - \frac{a}{2}\right).$$

- (d) Slicing the region perpendicular to the
- x
- axis, the area of a differential slice is

$$dA = y dx = \left(\int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du \right) dx.$$

So the area of the region bounded by the curve $y = f(x)$ and the coordinate axes is

$$A = \int_0^a \left(\int_x^a \frac{u^2}{\sqrt{a^4 - u^4}} du \right) dx.$$

But now we run into two problems. The first is that double integrals are not part of this class. The second is that even if they were, it's still not clear how we would evaluate this double integral.

So instead we slice the region perpendicular to the y -axis. Then the area of a differential slice is

$$dA = x \, dy.$$

But since

$$\frac{dy}{dx} = \frac{-x^2}{\sqrt{a^4 - x^4}},$$

$$dy = \frac{-x^2}{\sqrt{a^4 - x^4}} dx$$

and

$$dA = x \, dy = \frac{-x^3}{\sqrt{a^4 - x^4}} dx.$$

Letting the y -intercept have coordinates $(0, b) \sim (0, 0.599a)$, the area of the region is

$$\begin{aligned} A &= \int_0^b x \, dy \\ &= \int_a^0 \frac{-x^3}{\sqrt{a^4 - x^4}} dx \\ &= \frac{1}{2} a^2, \end{aligned}$$

where the last equality follows by making the substitution $u = x^4$ (the details are left for you).

2 Happy Birthday!

Question 474 Rotate two copies of the rectangular elastica of Question 1 about the y -axis and you get a surface that models a mylar balloon.

Desmos link: <https://www.desmos.com/calculator/wfmpdd2tk>

152: Balloon

Desmos link: <https://www.desmos.com/calculator/u1vk5xr0ep>

152: Elastica 2

Desmos3D link: <https://www.desmos.com/3d/ztdgovbaqd>

152: Mylar Balloon 2

- (a) Find an expression for the volume of the balloon in the form $V = ka^3$, for some constant k .
- (b) Use Line 9 of the worksheet above to approximate the constant k .

Explanation. Slicing the solid perpendicular to the y -axis through the point $(x, f(x), 0)$, the cross-section is a circle with radius x and area

$$A = \pi x^2.$$

So the differential slice with thickness dy has differential volume

$$\begin{aligned} dV &= \pi x^2 dy \\ &= \frac{-x^4}{\sqrt{a^4 - x^4}} dx. \end{aligned}$$

And with $(0, b)$ the coordinates of the y -intercept of the curve $y = f(x)$, the solid has volume

$$\begin{aligned} V &= \pi \int_0^b x^2 dy \\ &= \pi \int_a^0 \frac{-x^4}{\sqrt{a^4 - x^4}} dx \\ &= \pi a^3 \int_0^1 \frac{v^4}{\sqrt{1 - v^4}} dv \\ &\sim 0.437\pi a^3. \end{aligned}$$

As a check, we compare this with the volume V_c of the surrounding right circular cylinder with radius a and height $b \sim 0.6a$. The ratio of volumes is

$$\frac{V}{V_c} \sim \frac{0.437\pi a^3}{0.6\pi a^3} \sim 0.73.$$

To better understand this ratio, we compare it with the ratio of the area $A = 0.5a^2$ of the region of rotation and the area $A_R \sim 0.6a^2$ of the surrounding rectangle. This ratio of areas is

$$\frac{A}{A_R} \sim \frac{0.5a^2}{0.6a^2} \sim 0.83.$$

I'm not quite sure about this, but it should be that

$$\left(\frac{A}{A_R}\right)^{3/2} < \frac{V}{V_c} < \frac{A}{A_R},$$

which is the case here since

$$\left(\frac{A}{A_R}\right)^{3/2} \sim 0.54.$$

Which Tanks Empty Faster?

Draining tanks.

1 Preliminaries

2 Thoughts on Differential Equation in Math 152

It is possible to integrate (ha!) differential equations in a meaningful way rather than saving it for the end.

For me, this approach starts with changing the focus from using definite integrals and Part 2 (?) of the Fundamental Theorem to compute *numbers* to using definite integrals and Part 1 of the FT to find expressions for functions. It makes generalizing the point-slope equation of a line a fundamental part of the course.

Here are two examples.

Example 76. (a) *Math 99* During a 50-mile trip a car gets a constant gas mileage of 40 miles/gal. When the trip odometer reads 32 miles, the car has 9 gallons of gas in its tank.

Find a function

$$G = h(s), 0 \leq s \leq 50,$$

that expresses the number of gallons of gas in the tank in terms of the trip odometer reading (in miles).

(b) *Math 152* The function

$$m = f(s) = ???, 0 \leq s \leq 50,$$

expresses the gas mileage of a car (measured in miles/gal) in terms of the trip odometer reading (measured in miles) during a 50-mile trip. When the trip odometer reads 32 miles, the car has 9 gallons of gas in its tank.

Find a function

$$G = h(s), 0 \leq s \leq 50,$$

that expresses the number of gallons of gas in the tank in terms of the trip odometer reading (in miles).

Learning outcomes:
Author(s):

Explanation. (a)

$$G = h(s) = 9 - \frac{1}{40}(s - 32), \quad 0 \leq s \leq 50.$$

- (b) Two approaches. One starts by adding the differential number of gallons of gas

$$dG = -\frac{ds}{f(s)}$$

the car burns over the intervals $s \in [u, u + du]$ miles as the trip odometer ranges from $s = 32$ miles to $s = u$ miles.

The other starts with an initial value problem

$$\frac{dG}{ds} = -\frac{1}{f(s)}, \quad G = 9 \text{ when } s = 32.$$

Either way we get

$$G = h(s) = 9 - \int_{32}^s \frac{du}{f(u)}, \quad 0 \leq s \leq 50.$$

The two approaches are logically equivalent if we use definite integrals to solve the differential equation. Using indefinite integrals misses the point. The integral stops being a sum and the computation loses meaning.

Here's a more interesting example.

3 Which Tanks Drain Faster?

Fill a tank with water, punch a small hole in the bottom of the tank, and let the water drain out. The problem is to find a function that expresses the depth of water in terms of time.

Desmos link: <https://www.desmos.com/calculator/zltxdekv7z>

152: Draining Cylinders

The given information:

- (a) The cross-sectional area $A = f(h)$ of the tank as a function of depth.
- (b) The area a of the small hole.
- (c) The assumption (or theoretical fact) that the water exits the hole with speed

$$v = g(h) = \sqrt{2gh},$$

where h is the depth of the water and g the magnitude of the gravitational acceleration near the earth's surface.

Which Tanks Empty Faster?

4 A Hemispherical Tank

Desmos link: <https://www.desmos.com/calculator/ggqodhz3jv>

152: Draining Cylinders 2

Parametrically-defined curves and Area

Curves defined parametrically and signed area.

1 The Basics

In a sense there is nothing new here. Making a u -substitution is equivalent to reparameterizing the graph of a function.

To see why and to understand what u -substitution does geometrically, it helps to think about making a *reverse* u -substitution. This means starting with a simple integral and making it more complicated with a substitution. Here's an example.

Example 77. We'll start with the integral

$$I = \int_0^{\pi} \sin u \, du$$

and make the substitution

$$u = t^2.$$

This substitution really reparameterizes the curve

$$y = f(u) = \sin u, \quad 0 \leq u \leq \pi$$

from the standard parameterization

$$(x, y) = (u, \cos u), \quad 0 \leq u \leq \pi$$

to the new parameterization

$$(x, y) = (t^2, \cos(t^2)), \quad 0 \leq t \leq \sqrt{\pi}.$$

The substitution maps the rectangle with height $y = \sin u$ and width du to the rectangle with the same height $y = \sin(t^2)$ and width

$$du = 2t \, dt.$$

The effect is to transform a partition of the interval of integration, originally $u \in [0, \pi]$ with equal subintervals into a partition of the new interval of integration $t \in [0, \sqrt{\pi}]$ with *unequal* subintervals $du = 2t \, dt$. You can see this transformation by dragging the slider w in Line 2 of the worksheet below from $w = 0$ to $w = 1$.

Learning outcomes:
Author(s):

Desmos link: <https://www.desmos.com/calculator/toiwzzqkva>

152: Sub 2

2 The Area of a Circle

Computing the area of a circle of radius a with integration can be a challenging problem when approached in the standard way.

Starting with an equation

$$\sqrt{x^2 + y^2} = a$$

of a circle centered at the origin, we'll find the area of the region bounded by the semicircle

$$y = f(x) = \sqrt{a^2 - x^2}$$

and the x -axis. On the one hand, the area of a differential rectangle having width dx and endpoints $(x, 0)$ and $(x, y) = (x, f(x))$ is

$$dA = y dx = \sqrt{a^2 - x^2} dx.$$

So the area of the semicircle is

$$A = 2 \int_0^a \sqrt{a^2 - x^2} dx.$$

Now we're faced with the problem of finding an anti-derivative of $\sqrt{a^2 - x^2}$. The standard method is to use a trigonometric substitution. But this really amounts parameterizing the circle.

So let's start over and parameterize the circle $x^2 + y^2 = a^2$ in terms of the polar angle θ of the vector $\vec{OP}$ from the origin to a point P on the circle with coordinates (x, y) .

From the *definitions* of the cosine and sine functions we have

$$x = a \cos \theta \text{ and } y = a \sin \theta, 0 \leq \theta \leq 2\pi.$$

We'll slice the upper-semicircle perpendicular to the x -axis as before. Now since $0 \leq \theta \leq \pi$ for the upper semicircle,

$$dx = d(a \cos \theta) = -a \sin \theta d\theta,$$

the area of the differential rectangle is

$$\begin{aligned} dA &= -y dx \\ &= -(a \sin \theta) d(a \cos \theta) \\ &= -(a \sin \theta)(-a \sin \theta d\theta) \\ &= a^2 \sin^2 \theta d\theta. \end{aligned}$$

Parametrically-defined curves and Area

Geometrically, the effect of this parameterization is to transform the original parametrization

$$(x, y) = (x, \sqrt{a^2 - x^2}), \quad -a \leq x \leq a$$

of the upper-semicircle

Desmos link: <https://www.desmos.com/calculator/eu8hz9nmgz>

152: Circle Area

Parametrically-defined Curves, CW

Curves defined parametrically and signed area.

1 The Basics

Not all curves are the graphs of functions.

The green curve below in the first quadrant, for example, is neither the graph of y as a function of x (ie. $y = f(x)$) nor of x as a function of y (ie. $x = f(y)$).

Why not?

Free Response:

Desmos link: <https://www.desmos.com/calculator/vio0b8ztnl>

152: Parametric 2

But we can describe the curve *parametrically*. This means to give functions

$$x = f(t) \text{ and } y = g(t)$$

that express the coordinates (x, y) of a point on the curve in terms of some parameter (t here) which is often helpful to think of as being time. We can then think dynamically of the *motion* of a point along its path instead of just a static path.

These coordinate functions for the green curve above are graphed in the fourth ($x = f(t)$) and second ($y = g(t)$) quadrants and share the domain $0 \leq t \leq 4$ (you should ignore the negative signs on the vertical and horizontal axes). To see the motion of the point along its path, play the slider $u = t$ in Line 2.

The question we would like to address here is to compute the unsigned (ie. positive) area of the region bounded by the curve

$$(x, y) = (f(t), g(t)), 0 \leq t \leq 4.$$

We'll use with the demonstration below, where the graphs of the coordinate functions are hidden.

Desmos link: <https://www.desmos.com/calculator/z9ndh3wvvv>

Learning outcomes:
Author(s):

152: Parametric 2B

As usual, our first choice is to decide which way to slice the region and we'll start by slicing perpendicular to the x -axis. But this is not really an accurate description because the differential slices run from the x -axis to the curve and do not always lie within the region.

To see this, drag the slider the slider u in Line 2 to move the differential rectangle. Sometimes the rectangle lies below the region, other times it lies partly within and partly below the region, and at other times completely within the region. Nevertheless, we can use these differential rectangles to compute the area of the region.

The key idea is to work with their *signed* areas. Cutting perpendicular to the x -axis, the differential rectangles have signed area

$$\begin{aligned} dA &= y \, dx \\ &= g(t) d(f(t)) \\ &= g(t)(f'(t) \, dt) \\ &= g(t)f'(t) \, dt. \end{aligned}$$

And so the region bounded by the green curve and the x -axis has signed area

$$A = \int_0^4 g(t)f'(t) \, dt.$$

Let's stop and think a minute about what we mean by signed area. The sign of the differential area dA depends upon *both* the signs of $y = g(t)$ and $dx = f'(t) \, dt$. For our example $y = g(t) \geq 0$, $dt > 0$, and so dA has the same sign as dx .

- (a) Drag the slider u in Line 2 from $u = 0$ to $u = b = 4$. For what values of u is dA positive? Negative?
- (b) Sketch by hand a rough graph of the cumulative signed area function

$$A_1(t) = \int_0^t g(u)f'(u) \, du.$$

Then drag the slider u in Line 2 to $u = 0$ and activate the folder *Graph of signed area function, slices perp to x-axis* in Line 7. Drag u to $u = b = 4$ and see the graph of the signed area function. Compare it with your graph.

- (c) How is the integral

$$A_1(4) = \int_0^4 g(t)f'(t) \, dt$$

related to the area of the region bounded by the x -axis and the curve $(x, y) = (f(t), g(t))$? Explain your reasoning.

Free Response:

We can also take our differential rectangles perpendicular to the y -axis.
Then

$$dA = x \, dy = f(t)g'(t) \, dt.$$

- (d) Repeat parts (a)-(c) above for these differential rectangles and the signed area function

$$A_2(t) = \int_0^t f(u)g'(u) \, du.$$

- (e) Now use the actual coordinate functions

$$x = f(t) = t^2 - 3t + 3, \, 0 \leq t \leq 4$$

and

$$y = g(t) = 4t - t^2, \, 0 \leq t \leq 4$$

to compute the area of the region bounded by the x -axis and the curve

$$(x, y) = (f(t), g(t)), \, 0 \leq t \leq 4.$$

Do this twice. Once with slices perpendicular to the x -axis and again with slices perpendicular to the y -axis.

2 An Application: Gas Consumption

Example 78. The function

$$r = f(t), \, 0 \leq t \leq 4,$$

expresses the rate (in gal/mile) at which a car burns gas in terms of the number of hours past noon during a 240-mile trip.

The function

$$s = g(t), \, 0 \leq t \leq 4,$$

expresses the car's trip odometer reading (in miles) in terms of the number of hours past noon.

The car has 14 gallons of gas at 1:00pm.

The functions f (second quadrant) and g (fourth quadrant) along with the curve $(x, y) = (g(t), f(t))$ (first quadrant) are graphed below.

Desmos link: <https://www.desmos.com/calculator/akedzpal8z>

Substitution and Gas Consumption 22

- (a) Approximate the rate, measured in gal/hour, at which the car burns gas at 1:54pm. Explain your reasoning and include units for all numbers in your computation.
- (b) Approximate the number of gallons of gas in the tank at 1:54pm.
- (c) Find a function

$$G = w(s), 0 \leq t \leq 4,$$

that expresses the number of gallons of gas in the car in terms of the number of hours past noon.

3 Exercises

Exercise 475 This question is about the circle

$$(x, y) = (f(\theta), g(\theta)) = (a \cos \theta, b + a \sin \theta), 0 \leq \theta \leq 2\pi.$$

Here a, b are positive constants measured in meters.

Desmos link: <https://www.desmos.com/calculator/weiqr1fxu>

152: Parametric 3

- (a) Use the worksheet below to sketch by hand the signed area functions

$$A_1(\theta) = \int_0^t g(t)f'(t) dt$$

and

$$A_2(\theta) = \int_0^t f(t)g'(t) dt.$$

Then check your graphs by activating the folders in Lines 7 and 23.

Desmos link: <https://www.desmos.com/calculator/bg3tmflc1f>

152: Parametric 3B

- (b) Evaluate the integrals

$$A_1(2\pi) = \int_0^{2\pi} g(\theta)f'(\theta) d\theta$$

and

$$A_2(2\pi) = \int_0^{2\pi} f(\theta)g'(\theta) d\theta$$

geometrically by using the graph of the function

$$A = a^2 \sin^2 \theta, 0 \leq \theta \leq \pi$$

shown below. Work in general, not with the specific value of a in the worksheet.

Desmos link: <https://www.desmos.com/calculator/dsoqsyfba>

152: Sine Squared

Exercise 476 Use the parameterization

$$(x, y) = (a \cos \theta, a \sin \theta), 0 \leq \theta \leq 2\pi,$$

of the circle with radius a meters centered at the origin to find an expression for the volume of a sphere with the same radius.

Exercise 477 The circle with center $(x, z) = (b, 0)$ and radius a is rotated about the z -axis to sweep out the torus shown below.

Desmos3D link: <https://www.desmos.com/3d/baapxxf50y>

152: Torus

The aim of this problem is to express the volume of the torus in terms of the parameters a and b , measured in meters. We'll do this by parameterizing the circle in terms of the polar angle of the vector $\overrightarrow{CP}$ from the center of the circle to a point P on the circle with coordinates (x, y) . See the figure below.

Desmos link: <https://www.desmos.com/calculator/qjpoe06x3y>

152: Torus 2D

- Express the coordinates of point P in terms of the polar angle θ .
- Activate the folder in Line 1 of the worksheet above to see the cross-section of the torus swept out by PQ . The cross-section should really be drawn perpendicular to the page. You can see the cross-section in the top worksheet by deactivating the folder Torus in Line 4.
- Express the area of the cross-section in terms of a , b , and θ .

- (d) Express the differential volume of the corresponding slice in terms of a , b , and θ .
- (e) Add these differential volumes to find the volume of the torus.

Exercise 478 Do Question 2 in the chapter

Volumes, Part 2

of our class notes by using trig functions to parameterize the circle of radius a centered at the origin.

Exercise 479 Find the area bounded by the astroid

$$(x, y) = (a \cos^3 \theta, a \sin^3 \theta), \quad 0 \leq \theta \leq 2\pi,$$

where $a > 0$ is a constant measured in meters.

Exercise 480 Find the area bounded by the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

where $a, b > 0$ are constants measured in meters.

Start by parameterizing the ellipse.

Integration by Parts

Integration by parts, geometrically.

1 The Basics

Learning outcomes:
Author(s):

Arclength of Curves in the Plane

Arclength.

1 The Ideas

We'll speak of our curves as trails and assume they lie in a vertical plane (they don't twist around a mountain like a real trail usually does).

The constant case for computing the length of a trail is when the trail makes a constant angle ϕ with the horizontal. This means that the trail is a line segment and looks just like a ramp.

In this case, if the endpoints of the ramp have coordinates (x_1, y_1) and (x_2, y_2) , the ramp has length

$$\Delta s = \sqrt{(\Delta x)^2 + (\Delta y)^2},$$

where

$$\Delta x = x_2 - x_1$$

and

$$\Delta y = y_2 - y_1.$$

It can be helpful to express the length in terms of the inclination angle ϕ of the ramp. Here we have two possibilities assuming $\Delta s \geq 0$. They are

$$\Delta s = |\Delta x| \sec \phi$$

or

$$\Delta s = |\Delta y| \csc \phi.$$

For a general trail, where the inclination angle is not necessarily constant, the Pythagorean theorem tells us the differential arclength is

$$ds = \sqrt{(dx)^2 + (dy)^2}.$$

We can write this in terms of the inclination angle ϕ (that now can vary along the curve) as

$$ds = \sec \phi \, dx$$

or

$$ds = \csc \phi \, dy.$$

Learning outcomes:
Author(s):

When the equation of a trail is expressed as a function $y = f(x)$, we usually write

$$\begin{aligned} ds &= \sqrt{(dx)^2 + (dy)^2} \\ &= \sqrt{1 + \left(\frac{dy}{dx}\right)^2} |dx| \\ &= \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \end{aligned}$$

if $dx > 0$.

I like to think of

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} \geq 1,$$

as a dimensionless scaling factor that converts a differential change dx along the horizontal to a differential arclength along the trail.

Question 481 (a) Express the derivative dy/dx in terms of the inclination angle ϕ .

(b) Express the scaling factor $\sqrt{1 + \left(\frac{dy}{dx}\right)^2}$ in terms of ϕ .

Finally, the length of the trail between points (a, b) and (c, d) (with $c > a$) is the sum

$$\int_a^c \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

of these differential arclengths.

2 Circles

Example 79. This problem is about the semicircle

$$y = f(x) = \sqrt{a^2 - x^2},$$

where the constant $a > 0$ is measured in meters.

(a) Use integration to find an expression for the function

$$s = g(w), -a \leq w \leq a,$$

that gives the distance from the point $A(a, 0)$ to the point P with coordinates $(w, f(w))$ on the semicircle

$$y = f(x) = \sqrt{a^2 - x^2}.$$

Measure the distance from A to P along the semicircle.

Start by expressing the differential arclength ds along the semicircle in terms of the differential change dx .

Desmos link: <https://www.desmos.com/calculator/rmeuvbnjds>

152: ArcLength Semicircle

- (b) Interpret your result in part (a) geometrically.
- (c) Sketch by hand a graph of the function $s = g(x)$. Then activate the folder *Arclength Function* in Line 11 above to see how you did.
- (d) Express $g'(x)$ in terms of $f'(x)$.
- (e) Express the coordinates of P in terms of its distance s from $(a, 0)$ as measured along the semi-circle.

Example 80. Let $a > 0$ be a constant measured in meters.

Use integration to find an expression for the function

$$s = f(\phi), \quad 0 \leq \phi < 2\pi,$$

that gives the distance from the point $A(a, 0)$ to the point P with coordinates $(a \cos \phi, a \sin \phi)$ on the circle

$$(x, y) = (a \cos \theta, a \sin \theta), \quad 0 \leq \theta < 2\pi,$$

in terms of ϕ . Measure the distance from A to P counterclockwise along the circle.

Start by expressing the differential arclength ds along the semicircle in terms of the differential change $d\theta$.

3 Parameterizing an Astroid by Arclength

Example 81. This example is about the curve (an astroid)

$$(x, y) = (a \cos^3 \theta, a \sin^3 \theta), \quad 0 \leq \theta \leq 2\pi,$$

where $a > 0$ is a constant measured in meters.

Desmos link: <https://www.desmos.com/calculator/up6jvtmspl>

152: Astroid

- (a) Find an expression for the function

$$s = f(\phi), 0 \leq \phi < 2\pi,$$

that gives the distance from the point $A(a, 0)$ to the point P with coordinates $(a \cos^3 \phi, a \sin^3 \phi)$ on the astroid

$$(x, y) = (a \cos^3 \theta, a \sin^3 \theta), 0 \leq \theta < 2\pi,$$

in terms of ϕ . Measure the distance from A to P counterclockwise along the curve.

Start by expressing the differential arclength ds along the astroid in terms of the differential change $d\theta$.

- (b) Evaluate $f(\pi/2)$ and interpret its meaning. Compare this distance with the straight-line distance from $(a, 0)$ to $(0, a)$.
- (c) Express the coordinates of a point P on the curve in terms of its distance s from A measured counterclockwise along the astroid.

4 Trail Profiles

The Cycloidal Pendulum

Pendulums.

1 The Cycloid

Example 82. A cycloid is the curve traced by a point attached to a circle as the circle rolls on a line.

Drag the slider θ_2 in Line 2 of the worksheet below to roll the circle.

On which line does the circle roll? Explain how you know.

Free Response:

Desmos link: <https://www.desmos.com/calculator/rs6esnvlfi>

152: Cycloidal Pendulum 2

We'll parameterize the cycloid above traced by the point P in terms of the marked rotation angle θ (called θ_2 in the worksheet above) of the rolling circle. with radius a meters. Do this as follows.

- (a) Express the components of the vector $\overrightarrow{OC}$ from the origin to the center of the circle in terms of θ and the radius a of the circle.

$$\overrightarrow{OC} = \langle a\theta, a \rangle.$$

- (b) Express the components of the vector $\overrightarrow{CP}$ in terms of θ and a .

$$\overrightarrow{CP} = \langle a \sin \theta, -a \cos \theta \rangle.$$

- (c) Use parts (a) and (b) to express the components of the vector $\overrightarrow{OP}$ in terms of θ and a .

$$\begin{aligned} \overrightarrow{OP} &= \overrightarrow{OC} + \overrightarrow{CP} \\ &= \langle a\theta + a \sin \theta, a - a \cos \theta \rangle. \end{aligned}$$

Learning outcomes:
Author(s):

2 The Cycloidal Track

Watch the video below.

What is the main point of the video?

Free Response:

YouTube link: https://www.youtube.com/watch?v=jTN8UY_9xV0

Cycloid Track

Example 83. There are several ways to show that the cycloid is a *tautochrone* (ie. that the period of oscillation is independent of the release point). The most insightful uses vectors and we'll do this next quarter. But for now, here's what might be the second best approach.

We'll start with a parameterization

$$(x, y) = (a\theta + a \sin \theta, a - a \cos \theta), \quad -\pi \leq \theta \leq \pi,$$

of the cycloid shown in the rectangular coordinate system below, where the parameter a has units of meters.

Desmos link: <https://www.desmos.com/calculator/x6qb4zs2rq>

152: Cycloidal Pendulum 1

We'll start by computing the time for a mass released from the top of the cycloid to reach the low point (at the origin).

For this, we need some physics. Supposing the ball to slide along the cycloid without friction implies the ball's mechanical energy

$$E = mgh + \frac{1}{2}mv^2$$

is constant. Here m (in kg) is the ball's mass, h its height (in meters), and v (in m/sec) its speed. The constant $g \sim 9.8$ (m/sec)/sec is the magnitude of acceleration due the force of gravity near the surface of the earth.

We'll measure the ball's gravitational potential energy mgh upward from the y -axis, so that $h = y$.

We'll assume the the ball is released from the top of the cycloid at parameter $\theta = \pi$ and first express the speed of the ball in terms of its height y . In this case the mechanical energy of the ball (in terms of m , g , and a) is always

$$E = 2mga.$$

So when the mass has height y meters,

$$mgy + \frac{1}{2}mv^2 = 2mga$$

and since $v \geq 0$,

$$v = \sqrt{2g(2a - y)}.$$

Now to compute the time of descent, we first analyze the constant case, the one where speed is constant. Then to travel Δs meters at a constant speed of v meters/sec takes

$$\Delta t = \frac{\Delta s}{v} \text{ sec.}$$

So to compute the descent time from the release point $(a\pi, 2a)$ to the low point $(0, 0)$, we'll sum the differential times

$$\begin{aligned} dt &= \frac{ds}{v} \text{ sec} \\ &= \frac{ds}{\sqrt{2g(2a - y)}} \end{aligned}$$

for the mass to travel the differential distances ds meters along the cycloid at speeds of $v = \sqrt{g(4a - y)}$ m/sec.

Here we have a choice. We could either

- (a) express both ds and y in terms of θ , or
- (b) express y in terms of the arclength parameter s , measured from the top of the cycloid at height $2a$ to height y .

We'll do both.

For the first, the parameterization tells us

$$y = a - a \cos \theta, \quad -\pi \leq \theta \leq \pi.$$

So it remains only to express the differential arclength s in terms of the parameter θ . Here's the computation.

$$\begin{aligned} ds &= \sqrt{(dx)^2 + (dy)^2} \\ &= \sqrt{(d(a\theta + a \sin \theta))^2 + (d(a - a \cos \theta))^2} \\ &= \sqrt{(a + a \cos \theta)^2 + a^2 \sin^2 \theta} d\theta, \end{aligned}$$

where the last equality assumes $d\theta \geq 0$.

A little algebra shows

$$ds = a\sqrt{2 + 2\cos\theta} d\theta.$$

Now using the double angle formula

$$\cos\theta = 2\cos^2(\theta/2) - 1$$

gives

$$ds = 2a\cos(\theta/2) d\theta, \quad -\pi \leq \theta \leq \pi.$$

So the time (in seconds) for the mass to traverse the differential distance ds (meters) at height $y = a(1 - \cos\theta)$ meters is

$$\begin{aligned} dt &= \frac{ds}{v} \\ &= \frac{2a\cos(\theta/2) d\theta}{\sqrt{2ga(1 + \cos\theta)}}. \end{aligned}$$

And since $-\pi \leq \theta \leq \pi$, we know $\sqrt{\cos^2(\theta/2)} = \cos(\theta/2)$ and using the double-angle formula for $\cos\theta$ gives the surprising result that

$$dt = \sqrt{\frac{a}{g}} d\theta.$$

So the quarter-period of oscillation (in seconds) for a mass released from the top of the cycloid is

$$T = \int_0^\pi \sqrt{\frac{a}{g}} d\theta = \pi\sqrt{a/g}$$

and full period

$$4T = 4\pi\sqrt{\frac{a}{g}} \text{ sec.}$$

Or we can get the same result by expressing the differential time

$$\begin{aligned} dt &= \frac{ds}{v} \\ &= \frac{ds}{\sqrt{2g(2a - y)}} \end{aligned}$$

in terms of s . This amounts to expressing y in terms of s .

For this, we'll first express s in terms of θ by summing the differential distances $ds = 2a\cos(\theta/2) d\theta$ along the cycloid measured from A (where $\theta = \pi$) to the point P with parameter θ . This shows the distance AP along the cycloid is

$$s = 2a \int_0^\theta \cos(\theta/2) d\theta$$
$$4a \sin(\theta/2), -\pi \leq \theta \leq \pi.$$

And using the double-angle formula

$$\cos \theta = 1 - 2 \sin^2(\theta/2),$$

tells us that

$$\begin{aligned} y &= a(1 - \cos \theta) \\ &= 2a \sin^2(\theta/2) \\ &= \frac{s^2}{8a}. \end{aligned}$$

152 Review

Some problems for review.

Question 482 The function

$$r = f(t), t \geq 0,$$

expresses the rate (in gal/hr) at which water flows into a tank in terms of the number of hours past noon, July 23, 2025.

The tank holds 500 gallons at 3pm on July 23, 2025.

(a) Find a function

$$V = g(t), t \geq 0,$$

that expresses the volume of water (measured in gallons) in the tank in terms of terms of the number of hours past noon, July 23, 2025.

(b) Suppose now that

$$r = f(t) = \frac{200}{(2 + \frac{t}{10})^2}, t \geq 0.$$

(i) Find an explicit expression for the function g .

(ii) Assuming the water flows forever, determine if there is a tank that can hold all the water without overflowing? If so, compute the minimum volume of such a tank. If not, explain why.

Use limits to justify your response.

Question 483 The continuous function

$$v = f(s), 20 \leq s \leq 70,$$

expresses the speed of a car (in miles/hour) in terms of the trip odometer reading (in miles) during a portion of a 200-mile trip.

The trip odometer reads 50 miles at 4:00pm.

Learning outcomes:

Author(s):

- (a) Find a function

$$t = g(s), \quad 20 \leq s \leq 70$$

that expresses the time (measured in hours past noon) in terms of the trip odometer reading. Begin your solution by analyzing the constant case.

- (b) Suppose now that the speed is a linear function of the odometer reading during this 50-mile section of the trip. Suppose also the speed is 60 miles/hour when the odometer reads 20 miles and 40 miles/hour when the odometer reads 70 miles.

- (i) Find an explicit expression for the function $t = g(s)$.
 (ii) Find the exact clock time when the odometer reads 70 miles. Then use a calculator to approximate this time to the nearest minute.

Explanation. (a) If the speed is constant at v miles/hour, then the time it takes to travel Δs miles is

$$\Delta t = \frac{\Delta s \text{ miles}}{v \text{ miles/hr}} \text{ hrs} = \frac{\Delta s}{v} \text{ hrs.}$$

If the speed is not constant, we divide the stretch of the trip from odometer reading 50 miles to odometer reading s miles into small intervals of length ds miles over which the speeds are constant. The time to travel the distance ds miles at a speed of $v = f(s)$ miles/hr is

$$dt = \frac{ds}{v} \text{ hrs.}$$

Adding these differential times gives the time (in hours) to travel from odometer reading 50 miles to odometer reading s miles as

$$\Delta t = \int_{50}^s \frac{ds}{f(s)}.$$

Since the odometer reads 50 miles at time $t = 4$ hours past noon, when the odometer reading is s miles, the time is

$$\begin{aligned} t &= g(s) \\ &= 4 + \Delta t \\ &= 4 + \int_{50}^s \frac{1}{f(x)} dx. \end{aligned}$$

- (b) The function

$$\begin{aligned} v &= f(s) \\ &= 60 - \frac{2}{5}(s - 20) \\ &= 68 - \frac{2}{5}s, \quad 20 \leq s \leq 70, \end{aligned}$$

expresses the speed (in miles/hr) in terms of the odometer reading (in miles). So by part (a)

$$t = g(s) = 4 + \int_{50}^s \frac{1}{68 - \frac{2}{5}x} dx.$$

Making the substitution

$$u = 68 - \frac{2}{5}x$$

gives (details left for you)

$$\begin{aligned} t &= g(s) \\ &= 4 - \int_{50}^s \frac{1}{68 - \frac{2}{5}x} dx \\ &= 4 - \frac{5}{2} \int_{48}^{68 - \frac{2}{5}s} \frac{1}{u} du \\ &= 4 - \left(\frac{5}{2} \ln |u| \right) \Big|_{u=48}^{u=68 - \frac{2}{5}s} \\ &= 4 - \frac{5}{2} \ln \left(\frac{68 - \frac{2}{5}s}{48} \right). \end{aligned}$$

So at the odometer reading $s = 70$ miles, the time is

$$\begin{aligned} t &= g(70) \\ &= 4 - \frac{5}{2} \ln \left(\frac{68 - \frac{2}{5}(70)}{48} \right) \\ &= 4 - \frac{5}{2} \ln \left(\frac{40}{48} \right) \\ &\sim 4.45 \end{aligned}$$

hours past noon.

Question 484 Use trigonometric functions to parameterize the circle of radius a cm centered at the origin. Then use the parameterization to compute

- (a) the sphere's volume and
- (b) the sphere's surface area.

Explanation. The parameterization in terms of the polar angle θ of the vector $\overrightarrow{OP}$ from the origin to a point $P(x, y)$ on the circle is

$$(x, y) = (a \cos \theta, a \sin \theta), \quad 0 \leq \theta \leq 2\pi.$$

- (a) Slice the sphere perpendicular to the y -axis, the cross section is a circle with radius $x = a \cos \theta$ and area

$$A = \pi a^2 \cos^2 \theta.$$

The differential slice has width

$$dy = d(a \sin \theta) = a \cos \theta d\theta$$

and volume

$$dV = A dy = \pi a^3 \cos^3 \theta d\theta.$$

So the volume of the sphere is

$$\begin{aligned} V &= 2\pi a^3 \int_0^{\pi/2} \cos^3 \theta d\theta \\ &= 2\pi a^3 \int_0^{\pi/2} \cos^2 \theta \cos \theta d\theta \\ &= 2\pi a^3 \int_0^{\pi/2} (1 - \sin^2 \theta) \cos \theta d\theta \end{aligned}$$

Making the substitution $u = \sin \theta$ gives

$$\begin{aligned} V &= 2\pi a^3 \int_0^{\pi/2} (1 - \sin^2 \theta) \cos \theta d\theta \\ &= 2\pi a^3 \int_0^1 (1 - u^2) du \\ &= \frac{4\pi}{3} a^3. \end{aligned}$$

- (b) Slicing the sphere the same way, the differential surface swept out by the arclength $ds = a d\theta$ at distance $x = a \cos \theta$ from the y -axis has area

$$\begin{aligned} dA &= (2\pi x) ds \\ &= (2\pi a \cos \theta)(a d\theta) \\ &= 2\pi a^2 \cos \theta d\theta. \end{aligned}$$

So the surface area of the sphere is

$$\begin{aligned} S &= 4\pi a^2 \int_0^{\pi/2} \cos \theta d\theta \\ &= 4\pi a^2. \end{aligned}$$

Question 485 Use trigonometric functions to parameterize the circle of radius a cm centered at the origin. Then use the parameterization to do the following.

- (a) Express the surface area of the spherical cap with height h cm in terms of a and h .

Desmos3D link: <https://www.desmos.com/3d/mlx6kjtjlm>

152: Golden Sphere 2

- (b) A sphere is cut into 10-zones by nine equally-spaced planes as shown below. Which zone has the greatest surface area?

Desmos3D link: <https://www.desmos.com/3d/0jqijhbw6z>

152: Golden Sphere

Explanation. (a) From the last question, since $y = a \sin \theta$,

$$dy = a \cos \theta d\theta.$$

So

$$dA = 2\pi a(a \cos \theta d\theta) = 2\pi a dy$$

and the area of the spherical cap of height h is

$$A = 2\pi a \int_{a-h}^h dy = 2\pi ah.$$

- (b) Part (a) tells us that the zones all have the same area. Why?

Question 486 This example is about the curve (an astroid)

$$(x, y) = (a \cos^3 \theta, a \sin^3 \theta), \quad 0 \leq \theta \leq 2\pi,$$

where $a > 0$ is a constant measured in meters.

Desmos link: <https://www.desmos.com/calculator/up6jvtmspl>

152: Astroid

- (a) Find an expression for the function

$$s = f(\phi), \quad 0 \leq \phi < \pi/2,$$

that gives the distance from the point $A(a, 0)$ to the point P with coordinates $(a \cos^3 \phi, a \sin^3 \phi)$ on the astroid

$$(x, y) = (a \cos^3 \theta, a \sin^3 \theta), \quad 0 \leq \theta < 2\pi,$$

in terms of ϕ . Measure the distance from A to P counterclockwise along the curve.

Start by expressing the differential arclength ds along the astroid in terms of the differential change $d\theta$.

- (b) Evaluate $f(\pi/2)$ and interpret its meaning. Compare this distance with the straight-line distance from $(a, 0)$ to $(0, a)$.
- (c) Express the coordinates of a point P on the curve in terms of its distance s from A measured counterclockwise along the astroid.

Explanation. (a) Since

$$dx = -3a \cos^2 \theta \sin \theta d\theta$$

and

$$dy = 3a \sin^2 \theta \cos \theta d\theta,$$

the differential arclength is

$$\begin{aligned} ds &= \sqrt{(dx)^2 + (dy)^2} \\ &= 3a \sqrt{\cos^2 \theta \sin^2 \theta (\cos^2 \theta + \sin^2 \theta)} d\theta \\ &= 3a \sqrt{\cos^2 \theta \sin^2 \theta} d\theta \\ &= 3a \cos \theta \sin \theta \text{ if } 0 \leq \theta \leq \pi/2. \end{aligned}$$

Then the arclength from $\theta = 0$ to $\theta = \phi$ is

$$s = 3a \int_0^\phi \cos \theta \sin \theta d\theta.$$

Make the substitution $u = \sin \theta$. Then

$$\begin{aligned} s &= f(\phi) \\ &= 3a \int_0^\phi \cos \theta \sin \theta d\theta \\ &= 3a \int_0^{\sin \phi} u du \\ &= \frac{3a}{2} \sin^2 \phi. \end{aligned}$$

- (b) The arclength of the astroid from $(a, 0)$ to $(0, a)$ in the first quadrant is

$$s = f(\pi/2) = 3a/2 = 1.5a.$$

This is a bit more than the straight-line distance $\sqrt{2}a \sim 1.4a$ between these points.

- (c) We can skip this one.

Question 487 The function

$$r = f(t), 0 \leq t \leq 4,$$

expresses the rate (in gal/mile) at which a car burns gas in terms of the number of hours past noon during a 240-mile trip.

The function

$$s = g(t), 0 \leq t \leq 4,$$

expresses the car's trip odometer reading (in miles) in terms of the number of hours past noon.

The car has 14 gallons of gas at 1:00pm.

Which of the following functions express the number of gallons in the tank in terms of the number of hours past noon? Select all that apply and explain your reasoning.

Select All Correct Answers:

(a) $w(t) = 14 + \int_1^t f'(u)g(u) du$

(b) $w(t) = 14 + \int_1^t f(t)g'(t) dt$

(c) $w(t) = 14 + \int_1^t f(u)g'(u) du$ ✓

(d) $w(t) = 14 + \int_{g(1)}^{g(t)} f(u)g'(u) du$

(e) $w(t) = 14 + \int_{g(1)}^{g(t)} f(g^{-1}(u)) du$ ✓

(f) $w(t) = 14 + f(t)g(t) - f(1)g(1) - \int_1^t f'(u)g(u) du$ ✓

Explanation. For the constant case, suppose the car burns gas at a constant rate of $r = f(t) = 0.03$ gal/mile and the car travels at a constant speed of 60 miles/hr. Then

$$s = g(t) = 60t, \quad 0 \leq t \leq 4$$

and the car burns gas at the constant rate of

$$\left(0.03 \frac{\text{gal}}{\text{mile}}\right) \left(60 \frac{\text{miles}}{\text{hour}}\right) = f(t)g'(t) \frac{\text{gal}}{\text{hour}}.$$

So in the general case, during the differential time dt hours the car burns

$$-dG = f(t)g'(t) dt$$

gallons of gas.

And from 1:00pm to time t hours past noon, the car burns

$$-\Delta G = \int_1^t f(v)g'(v) dv$$

gallons. Since the car has 14 gallons of gas at 1:00pm,

$$G = w(t) = 14 - \int_1^t f(v)g'(v) dv$$

and (c) is correct.

Integration by parts then tells us that (f) is also correct.

Make the substitution $u = g(v)$ in the expression for $-\Delta G$ above. Then

$$du = g'(v) dv$$

and

$$v = g^{-1}(u).$$

So

$$\begin{aligned} -\Delta G &= \int_1^t f(v)g'(v) dv \\ &= \int_{g(1)}^{g(t)} f(g^{-1}(u)) du \end{aligned}$$

and (e) is correct.

To understand this, note that $f(g^{-1}(u))$ gives the rate (in gal/hr) at which the car burns gas when the odometer reads u miles. And the product

$$f(g^{-1}(u)) du$$

gives the number of gallons of gas the car burns as it travels du miles. Summing these differential volumes from odometer readings $g(1)$ to $g(t)$ miles gives the number of gallons of gas the car burns from odometer reading $g(1)$ to odometer reading $g(t)$.

Question 488 The function

$$u = f(t) = u_0 e^{rt}, \quad 0 \leq t \leq 240,$$

expresses the consumption rate of Chromium (measured in 10^8 tons/yr) in terms of the number of years since 1970. Here u_0 is the consumption rate at the start of 1970 and $r > 0$ is a constant.

At the start of 1970 the global reserves of Chromium totaled R_0 hundreds of millions of tons and would have been depleted in s years had the consumption rate remained constant.

(a) Express u_0 in terms of R_0 and s .

(b) Find a function

$$R = g(t), \quad 0 \leq t \leq 240,$$

that expresses the remaining Chromium reserves (measured in 10^8 tons/yr) in terms of the number of years since 1970. Assume the consumption rate described by the function $u = f(t)$.

(c) Find an expression in terms of R_0 , s , and r for the number of years since 1970 when the Chromium reserves were projected to be depleted had the consumption rate grew exponentially as described by the function $u = f(t)$.

(d) Check that your expression in part (c) has the correct units.

Explanation. (a) $u_0 = R_0/s$.

(b)

$$\begin{aligned} R &= g(t) \\ &= R_0 + \int_0^t f(v) dv \\ &= R_0 + \frac{R_0}{s} \int_0^t e^{rv} dv \\ &= R_0 + \frac{R_0}{rs} (1 - e^{rt}). \end{aligned}$$

(c) If $R = 0$, then

$$R_0 + \frac{R_0}{rs} (1 - e^{rt}) = 0$$

and

$$t = \frac{\ln(1 + rs)}{r}.$$

Question 489 *Practice integration.*

Loxodromes and Equiangular Spirals

Rhumb lines.

1 Loxodromes

Desmos3D link: <https://www.desmos.com/3d/jamvjfsxrv>

152: Loxodrome 53

Circles, Spheres, Area, Volume

Thoughts on the hoof of Archimedes, spheres, circles.

1 The Area of a Circle

Desmos link: <https://www.desmos.com/calculator/2pe3pmtjyv>

152: Unwrapping Semicircle

Desmos link: <https://www.desmos.com/calculator/9a3oluopqk>

152: Unwrapping Circle

2 The Hoof of Archimedes

A solid circular cylinder of radius a and height h is sliced by a plane passing through a diameter of the base and intersecting the top of the cylinder in exactly one point. The plane cuts the cylinder into two pieces. Our goal here is to find an expression in h and a for the volume of the lower (smaller) piece.

Here are three ways to find the volume.

Desmos3D link: <https://www.desmos.com/3d/j4cadgw4ct>

152: Hoof of Archimedes 3

The volume of the hoof turns out to be

$$V = \frac{2}{3}a^2h.$$

This looks like the volume of a cone and suggests a quicker way to find the volume.

Desmos3D link: <https://www.desmos.com/3d/ddcsbxgbrl>

152: Hoof of Archimedes Unwrapped

Learning outcomes:
Author(s):

3 Archimedes' Bicylinder

The hoof of Archimedes comes up in unexpected places. Here's one.

Desmos3D link: <https://www.desmos.com/3d/wzzhod4ya6>

152:Bicylinder

Desmos3D link: <https://www.desmos.com/3d/phy3gfnhcj>

152: Bicylinder and Hoof

The bicylinder has close ties to one of our solids, the one with the circular base and square cross-sections.

Desmos3D link: <https://www.desmos.com/3d/cwcjqkjndw>

152: Square Cross-Sections to Bicylinder

4 Tori

The hoof is also connected to tori.

The unwrapping preserves both surface area and volume.

Desmos3D link: <https://www.desmos.com/3d/9gcd0sjhxr>

152: Torus Unwrapped

5 The Sphere

We can also unwrap a sphere, preserving both its volume and surface area.

Desmos3D link: <https://www.desmos.com/3d/wlqzsd2xp>

152: Sphere Unwrapped

A second unwrapping should take this truncated cylinder to a cone. To be continued...

Hyperbolic Trigonometry

An introduction to the hyperbolic trigonometric functions.

1 The Hyperbolic Cosine and Sine Functions

The function *hyperbolic cosine* is defined as

$$\cosh(x) = 0.5(e^x + e^{-x}).$$

Its graph is shown below.

Desmos link: <https://www.desmos.com/calculator/nubdwzgsa9>

152: Hyp Cosine

- (a) Drag the slider u in Line 2 of the worksheet above. Explain how to get the graph of the function $y = \cosh x$ from the graphs of the functions $y = e^x$ and $y = e^{-x}$.

Free Response:

- (b) What is the range of the function $f(x) = \cosh x$?
- (c) Activate the folder *Hyperbolic Sine Function* in Line 15 above to see the graph of the hyperbolic sine function

$$\sinh(x) = 0.5(e^x - e^{-x}).$$

- (d) Drag the slider u in Line 2 of the worksheet above. Explain how to get the graph of the function $y = \sinh x$ from the graphs of the functions $y = e^x$ and $y = e^{-x}$.

Free Response:

- (e) What is the range of the function $g(x) = \sinh x$?
- (f) Prove that

$$\cosh^2 x - \sinh^2 x = 1.$$

Hint: For a quick way, first factor the above expression.

Learning outcomes:
Author(s):

(g) Show that

$$\frac{d}{dx}(\cosh x) = \sinh x$$

and

$$\frac{d}{dx}(\sinh x) = \cosh x$$

2 Hanging Chains

Hold the ends of a chain and let the chain sag under its own weight. The resulting curve looks like a parabola, and so it was thought. But it turns out that this is not correct.

Desmos link: <https://www.desmos.com/calculator/tj3dz2cnf0>

152: Hanging Chain 1

- (a) To see that the dashed (orange) chain in the worksheet above is *not* a parabola drag the slider c in Line 2.
- (b) It turns out that a chain with uniform density hanging in a uniform gravitational field assumes the shape of the hyperbolic cosine function $y = \cosh x$ (we'll see why later in the course). All that's needed to describe a particular chain is to scale the size of the graph by some factor. Test this by dragging the slider a in Line 6 to make the graph of the function

$$f(x) = -a + a \cosh(x/a)$$

match the highlighted (orange) chain below.

3 The Hyperbolic Tangent Function

The hyperbolic tangent function is defined as

$$\begin{aligned} \tanh x &= \frac{\sinh x}{\cosh x} \\ &= \frac{e^x - e^{-x}}{e^x + e^{-x}}. \end{aligned}$$

- (a) Find the domain and range of the function $y = \tanh x$.
- (b) Sketch by hand a graph of the function $y = \tanh x$.
- (c) Use your graph from part (b) to sketch a graph of the derivative

$$y = \frac{d}{dx}(\tanh x).$$

- (d) Find an expression for the derivative in part (c) in terms of $\cosh x$ and/or $\sinh x$.

4 Inverse Functions

The key idea here is that *the derivative of the inverse of a function is the reciprocal of the function's derivative*. This should be intuitive.

So to find the derivative of the inverse of a one-to-one function $y = f(x)$, we should first express dy/dx in terms of y . Then take the reciprocal.

- (a) Express the derivative of each function below in terms of its output.

(i)

$$y = \sinh x$$

(ii)

$$y = \cosh x, x \geq 0$$

(iii)

$$y = \tanh x$$

- (b) Use the results of part (a) to find expressions for the derivatives

(i)

$$\frac{d}{dy}(\sinh^{-1} y)$$

(ii)

$$\frac{d}{dy}(\cosh^{-1} y)$$

(iii)

$$\frac{d}{dy}(\tanh^{-1} y)$$

Review of the Chain Rule

Derivative review.

1 Example

Here's an example of how to use u -substitution with the chain rule. Just as importantly, it's an example of how to show your work clearly and precisely.

Example 84. (a) Use the chain rule to find an expression for the derivative

$$\frac{d}{dx} ((2x^3 + 1)^2).$$

(b) Use part (a) to find an equation of the tangent line to the curve

$$y = (2x^3 + 1)^2$$

at the point $(1, 9)$.

(c) Answer part (a) without the chain rule.

Explanation. (a) Let

$$y = (2x^3 + 1)^2$$

and

$$u = 2x^3 + 1.$$

Then

$$y = u^2$$

and from the chain rule,

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du} (u^2) \cdot \frac{d}{dx} (2x^3 + 1) \\ &= 2u(6x^2) \\ &= 2(2x^3 + 1)(6x^2). \end{aligned}$$

Learning outcomes:
Author(s):

- (b) The slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 2(3)(6) = 36,$$

and an equation of the tangent line is

$$y - 9 = 36(x - 1).$$

- (c) We can get the same result without the chain rule, by rewriting the original function as

$$y = f(x) = (2x^3 + 1)^2 = 4x^6 + 4x^3 + 1.$$

Then

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} (4x^6 + 4x^3 + 1) \\ &= 4 \frac{d}{dx} (x^6) + 4 \frac{d}{dx} (x^3) + \frac{d}{dx} (1) \\ &= 24x^5 + 12x^2. \end{aligned}$$

Then the slope of the tangent line to the curve $y = (2x^3 + 1)^2$ at the point $(1, 9)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 14 + 12 = 36,$$

as before.

2 Exercises

Directions: Follow all the steps of *Example 1* (a) exactly for each of the following. Show all work as above. Do not skip steps.

Exercise 490 Find simplified expressions for each of the following derivatives.

(a)

$$\frac{d}{d\theta} (\ln |\cot(\theta/2)|)$$

(b)

$$\frac{d}{d\theta} (\ln |\sec \theta + \tan \theta|)$$

(c)

$$\frac{d}{d\theta} (\operatorname{arctanh}(\sin(\theta)))$$

(d)

$$\frac{d}{d\theta}(\operatorname{arcsinh}(\tan(\theta)))$$

(e)

$$\frac{d}{dt}(\tan(k \arctan(t/2))),$$

where k is a constant.

Exercise 491 Find the acute angle the tangent line to the curve

$$y = \ln |\sec \theta|$$

at the point $P(\pi/7, \ln(\sec(\pi/7)))$ makes with the x -axis.

Exercise 492 Find simplified expressions for each of the following derivatives. Follow all steps of Example 1(a).

(a)

$$\frac{d}{dx} \left(x \arctan x - \ln \sqrt{1+x^2} \right)$$

(b)

$$\frac{d}{dt} (t \ln |t| - t)$$

(c)

$$\frac{d}{dx} \left(\ln(x + \sqrt{x^2 - 1}) \right)$$

(d)

$$\frac{d}{dx} \left(\ln(x + \sqrt{x^2 + 1}) \right)$$

(e)

$$\frac{d}{dx} \left(\ln(x + \sqrt{x^2 - 1}) \right)$$

(f)

$$\frac{d}{dt} \left(\ln \sqrt{\frac{1+t}{1-t}} \right)$$

Trails and Hanging Chains, Part 1

Trails and catenaries.

1 Angles of Inclination

Question 493 Find the measure of the acute (between 0 and $\pi/2$) angle the tangent line to the curve

$$y = \ln |\sec \theta|$$

at the point $(2\pi/7, \ln |\sec(2\pi/7)|)$ makes with the x -axis. Give an exact answer without using a calculator.

Desmos link: <https://www.desmos.com/calculator/yyyj3pjmk2>

152: Log Secant

Question 494 Find the measure of the acute angle the tangent lines to the curve

$$y = 4 \tan(x/20)$$

make with the x -axis at the points with y -coordinate $y = 3$. Give an exact answer, fully simplified, without using a calculator.

Desmos link: <https://www.desmos.com/calculator/df7hjgabzb>

152: Angle of Inclination

Explanation. Since we are given the y -coordinate of the point of tangency, we should express the derivative of our function not in terms of its input, but in terms of its output.

As a start, note that if $z = \tan \theta$, then

$$\begin{aligned} \frac{dz}{d\theta} &= \frac{d}{d\theta} (\tan \theta) \\ &= \sec^2 \theta \\ &= 1 + \tan^2 \theta \\ &= 1 + z^2. \end{aligned}$$

Learning outcomes:
Author(s):

For our function $y = 4 \tan(x/20)$, let $u = x/20$. Then

$$y = 4 \tan u$$

and by the chain rule

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= \frac{d}{du}(4 \tan u) \cdot \frac{d}{dx} \left(\frac{1}{20}x \right) \\ &= 4(1 + \tan^2 u) \left(\frac{1}{20} \right) \\ &= \frac{1 + (y/4)^2}{5}. \end{aligned}$$

So the tangent lines at the points on the curve $y = 4 \tan(x/20)$ with y -coordinate $y = 3$ have slope

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{y=3} &= \left(\frac{1 + (y/4)^2}{5} \right) \Big|_{y=3} \\ &= 5/16. \end{aligned}$$

These tangent lines therefore cut the x -axis at the acute angle $\phi = \arctan(5/16)$.

2 A Climber's Trail

Question 495 A loading ramp of length L meters runs from the ground to a truck. The ramp makes an angle of ϕ radians with the ground.

Find an expression for the function

$$h = f(s), \quad 0 \leq s \leq L,$$

that expresses the height of the ramp (in meters) in terms of the distance along the ramp, measured in meters from the bottom of the ramp.

Question 496 The function

$$h = f(s) = 100 + 250s - 10s^3, \quad 0 \leq s \leq 5$$

expresses the altitude (in meters) of a climber's trail in terms of the distance from the trailhead (in km).

- (a) Evaluate the derivative

$$\left. \frac{dh}{ds} \right|_{s=1}.$$

Include units.

- (b) Interpret the meaning of the above derivative using the language of small changes for a specific small change.
- (c) Find the exact angle the trail makes with the horizontal one kilometer from the trailhead.

Desmos link: <https://www.desmos.com/calculator/ljcugfqmee>

152: Hiking Trail 54

Explanation. (a) The derivative is

$$\begin{aligned} \frac{dh}{ds} &= \frac{d}{ds}(100 + 250s - 10s^3) \\ &= 250 - 30s^2 \end{aligned}$$

and

$$\left. \frac{dh}{ds} \right|_{s=1} = 220 \text{ meters/km.}$$

- (b) The derivative relates a small change

$$\Delta s = s - 1$$

in distance along the trail from kilometer marker $s = 1$ and the corresponding small change

$$\Delta h = h - f(1) = h - 340$$

in altitude (measured in meters).

More specifically, if $\Delta s \sim 0$, then

$$\begin{aligned} \Delta h &\sim \left. \frac{dh}{ds} \right|_{s=1} \cdot \Delta s \\ &= (220 \text{ meters/km})(\Delta s \text{ km}) \\ &= 220\Delta s \text{ meters.} \end{aligned}$$

So for example, if we walk 1 meter (ie. 0.001 km) farther along the trail from kilometer marker $s = 1$ km, then our altitude increases by approximately

$$\Delta h \sim 220(0.001) \text{ meters} = 0.2 \text{ meters.}$$

- (c) The description of the derivative's meaning tells us how to compute the trail's inclination angle at kilometer marker $s = 1$ km. Since we gain altitude at the rate of

$$220 \text{ meters/km} = 0.22 \text{ meters in elevation/meter walked}$$

one kilometer from the trailhead, the trail looks locally like a ramp inclined at the angle

$$\phi = \arcsin(0.22) \text{ rad}$$

above the horizontal there.

Remark: It's important to distinguish between the graph of the function $h = f(s)$ that expresses altitude in terms of distance along the trail and the trail profile. The latter is like a picture of the actual trail, assuming the trail stays confined to a vertical plane. For most trails, the two are indistinguishable. The worksheet below, for example, shows our original graph $h = f(s)$ along with the corresponding trail profile (in red).

Desmos link: <https://www.desmos.com/calculator/e5kuf1je8p>

152: Hiking Trail 54B

But for steeper trails, like one an ant might climb, there is a clear difference between the two.

Desmos link: <https://www.desmos.com/calculator/lktn53d7ju>

152: Hiking Trail 54C

3 A Hanging Chain

Question 497 Which of the following curves have the same shape as the curve

$$y = \sin \theta?$$

Select all that apply.

Select All Correct Answers:

- (a) $y = 2 \sin \theta$
- (b) $y = 2 \sin(2\theta)$
- (c) $y = 2 \sin(\theta/2)$ ✓

(d) $y = 0.2 \sin(0.2\theta)$

(e) $y = 0.2 \sin(5\theta)$ ✓

Note: Two curves have the same shape if you can scale a photograph of one to get an exact copy of the other.

Question 498 A chain of uniform density hanging under its own weight in a uniform gravitational field takes the shape of a the curve $y = \cosh x$ (we'll see why later in the course).

Desmos link: <https://www.desmos.com/calculator/iifkdeyq3y>

152: Hanging Chain 2

- (a) All that's needed to describe a particular chain is to scale the size of the curve $y = \cosh x$ by some factor. Test this by dragging the slider a in Line 6 to make the graph of the function

$$f(x) = -a + a \cosh(x/a)$$

match the highlighted (orange) chain below.

- (b) What are the units of a ? How do you know?
- (c) Using the value of a from part (a), find the angle the chain makes with the horizontal at the points two meters above the low point of the chain. Give an exact expression, fully simplified, without using a calculator.

See the chapter

Hyperbolic Trigonometry

of our class notes for information on the hyperbolic trig functions.

The Mercator Map

The Mercator map and its relation to Lambert's equal-area map.

1 Scaling Factors and Elastic Bands

The interpretation of the derivative as a rate of change is limited. A more universal description of the derivative, and one that generalizes to higher dimensions, is as a scaling factor. Multiplying the derivative by a sufficiently small change in the input to a function gives a good approximation to the change in the output. Of all possible scaling factors the derivative is the best in the sense that the error in the approximation approaches zero faster than the change in the output.

We can visualize the derivative as a scaling factor by regarding a function's domain as an elastic band and the function as acting on that band by stretching or compressing it.

For example, imagine a thin elastic band of length 4 meters running along the horizontal L -axis from $L = 0$ meters to $L = 4$ meters. Now hold the left end fixed at $L = 0$ and stretch the band by moving right end an additional four meters to the right. Then the function

$$H = g(L) = 2L, 0 \leq L \leq 4,$$

describes this stretching action. It takes as an input the distance (in meters) of a point on the band from the origin ($L = 0$) and returns as an output the distance between the origin and the corresponding point on the stretched band. The exploration below shows this stretching action.

Exploration 499 Drag the slider k in Line 2 below from $k = 1$ to $k = 2$ to illustrate the stretching action for the function g .

Desmos link: <https://www.desmos.com/calculator/qejivz36ui>

151: Rubber Band 1

The *global stretching factor* for a linear function like the one above,

$$H = g(L) = 2L, 0 \leq L \leq 4,$$

Learning outcomes:
Author(s):

is the slope of its graph. We can calculate this factor as an average stretching factor (ie. an average rate of change) between the points $L = a$ meters and $L = b$ meters from the origin. For the function $H = f(L) = 2L$, the stretching factor is

$$\begin{aligned}\frac{\Delta H}{\Delta L} &= \frac{f(b) - f(a)}{b - a} \\ &= \frac{2(b - a)}{b - a} \\ &= 2,\end{aligned}$$

This tells us that any two points on the stretched band are twice as far apart as they were on the unstretched band.

Question 500 (a) What are the units of the stretching factor?

Free Response:

(b) Find an expression for the inverse function

$$L = f^{-1}(H).$$

Include a domain.

(c) Interpret the inverse function as a deformation of a thin elastic band. What is the global stretching factor for this function?

Free Response:

Example 85. Here's an example

$$H = f(L) = 10 - \sqrt{100 - L^2}, \quad 0 \leq L \leq 10,$$

of a non-linear stretching function (where L and H are measured in meters as before). Like most functions in this class, it acts like a linear function near almost all points in its domain.

To stretch the elastic band in the demonstration below, drag the slider u in Line 2 from $u = 0$ to $u = 1$. Then zoom in close enough to the point $H = f(9.5)$ in the stretched band (highlighted in black) to make the stretching function look linear.

Desmos link: <https://www.desmos.com/calculator/xk8dvfcgwi>

152: Rubber Band Ladder 5B

- (a) Use the close-up view of the stretched band to approximate the local stretching factor at the input $L = 9.5$.
- (b) Parts of the elastic band get stretched, others compressed. Identify these.

2 Lambert's Equal-Area Map

You probably saw the idea of a stretching (scaling) factor long before taking calculus. Look at a map of Seattle and you'll see a scale, probably near the bottom. It might be 2 miles/inch, or something like that. But you will not see such a scale on a map of the earth for the simple reason that one does not exist. The scale varies from place to place and is usually a function of latitude. Even at a specific location, there is often not one but infinitely many scaling factors that depend on the direction in which you head.

The worksheet below shows a way to create a map of the earth by wrapping a cylinder around its equator. The map sends each point of the earth to its nearest point on the cylinder (ie. straight out, directly away from the earth's axis of rotation). Unwrapping the cylinder gives Lambert's equal-area map, from the earth to the points in a rectangle.

Exploration 501 Drag the slider in Line 6 from $r_1 = 0$ to $r_1 = 1$ to unwrap the cylinder. Drag sliders θ_1, ϕ_1 in Lines 2 and 4 to vary the longitude and latitude of the point on the sphere.

Desmos3D link: <https://www.desmos.com/3d/ehyvvtdeo>

152: Mercator 0

Suppose the earth has radius one. Lambert's map sends the circle of latitude ϕ (the angle between the plane of the equator and the segments from the sphere's center to points on the circle) to a circle on the cylinder with unit radius. So the map stretches the circle of latitude ϕ , with radius $\cos \phi$, by the factor

$$\lambda_p = \frac{1}{\cos \phi}.$$

This is the scaling factor along the circle in the east-west direction at latitude ϕ .

In the north-south direction the scaling factor is also a function of latitude. The key point is that the map sends a point on the circle of latitude ϕ to signed height

$$h = f(\phi) = \sin \phi$$

above the plane of the equator. So the scaling factor at latitude ϕ in the north-south direction is

$$\lambda_m = \frac{dh}{d\phi} = \cos \phi.$$

Drag the slider m in Line 2 of the worksheet below to visualize the scaling factor.

Desmos link: <https://www.desmos.com/calculator/kbryuogdaj>

151: Lambert Map Cosine Function

3 The Mercator Map

Watch this video.

Equal Earth Projection

4 Loxodromes

Rhumb Lines on the Sphere

For a curve on a sphere of radius a that cuts the meridians at a fixed angle α ,

$$d\theta = \frac{\tan \alpha d\phi}{\cos \phi},$$

where ϕ is the latitude and θ the longitude. Taking $\theta = 0$ at the equator where $\phi = 0$,

$$\begin{aligned}\theta &= \tan \alpha \int_0^{\phi^*} \sec \phi^* d\phi^* \\ &= \tan \alpha \operatorname{arctanh}(\sin \phi).\end{aligned}$$

So

$$\sin \phi = \tanh(k\theta),$$

where $k = \cot \alpha$.

Because $-\pi/2 < \phi < \pi/2$,

$$\cos \phi = \frac{1}{\cosh(k\theta)},$$

a parameterization of the rhumb line by longitude is

$$(x, y, z) = \left(\frac{a \cos \theta}{\cosh(k\theta)}, \frac{a \sin \theta}{\cosh(k\theta)}, a \tanh(k\theta) \right).$$

And a parameterization by latitude,

$$(x, y, z) = (a \cos \phi, a \cos \phi, a \sin \phi)$$

Desmos3D link: <https://www.desmos.com/3d/6cqnjsuew7>

The Mercator Map

152: Mercator 1

Desmos link: <https://www.desmos.com/calculator/rbcbxdrvcp>

Geogebra link: <https://www.geogebra.org/m/mvp9zvge>

152: Mercator

Riemann Sums with Geometric Partitions

Using Riemann sums with geometric partitions to evaluate definite integrals.

1 Integrating $1/x$

Before using Riemann sums and limits to find an expression for the integral

$$g(u) = \int_1^u \frac{1}{x} dx,$$

let's first compare the integrals

$$\int_1^2 \frac{1}{x} dx \tag{9}$$

and

$$\int_3^6 \frac{1}{x} dx. \tag{10}$$

We'll do this geometrically, without a computation.

The idea is to first describe a composition of transformations that takes the curve $y = 1/x$ to itself and the interval $[1, 2]$ to the interval $[3, 6]$. Then we'll look at how this composition affects the area.

Exploration 502 (a) *Drag the slider v in Line 2 of the worksheet below from $v = 1$ to $v = 3$ and describe what happens to the graph of $y = 1/x$.*

Free Response:

- (b) *What does the transformation do to the area of the shaded rectangle (or any rectangle)?*

It multiplies the area by a factor of 3.

- (c) *Express an equation of the new curve (dotted) in terms of v .*

An equation is

$$y = v/x.$$

- (d) Next drag the slider w in Line 4 of the worksheet from $w = 1$ to $w = 3$ and describe what happens to the graph of $y = v/x$.

Free Response:

- (e) What does the transformation do the area of the shaded rectangle (or any rectangle)?

It multiplies the area by a factor of $1/3$.

- (f) What is your conclusion about the relationship between the integrals (9) and (10) above? Explain your reasoning.

Free Response:

Desmos link: <https://www.desmos.com/calculator/al95rowj4h>

152: Log 1 Transformations

<https://www.desmos.com/calculator/al95rowj4h>

2 Computing

To evaluate

$$\int_1^2 \frac{1}{x} dx$$

with a limit of Riemann sums, we will *not* partition the interval $[1, 2]$ of integration into subintervals of equal length. Instead, we'll use the partition

$$x_i = 2^{i/n}$$

Desmos link: <https://www.desmos.com/calculator/rbcxdrvcp>

152: Log 1

Quiz Solutions Fall 2025

Quiz solutions.

1 Quiz1B

Question 503 The table below shows your gas mileage at different readings of the trip odometer reading (in miles) during a stretch of a 200-mile car trip.

Odometer Reading (miles)	Gas Mileage (miles/gallon)
23	23
35	25
40	30
50	32

Assume the gas mileage is an increasing function of the odometer reading and find the best possible (ie. the least) upper bound for the number of gallons of gas your car burned between odometer readings 23 miles and 50 miles. Explain your reasoning thoroughly. Your concluding sentence should start with “Between odometer readings 23 miles and 50 miles the car burned less than ...”

Explanation. We begin by considering the constant case. Here that means assuming your gas mileage is constant. Suppose for example our car gets 25 miles/gallon over a 50 mile trip. Then over these 50 miles our car burns

$$\frac{50 \text{ miles}}{25 \text{ miles/gal}} = 2 \text{ gallons}$$

of gas.

Now we would burn the most gas when our car gets the worst (ie. least) gas mileage. So to find the least possible upper bound we should assume the gas mileage is constant over each interval of distance and equal to its minimum over that interval. Since the gas mileage is assumed to increase throughout the trip, this means we should choose the left endpoint of each interval.

So between odometer readings 23 miles and 50 miles the car burned less than

$$\frac{(35 - 23) \text{ miles}}{23 \text{ miles/gal}} + \frac{(40 - 35) \text{ miles}}{25 \text{ miles/gal}} + \frac{(50 - 40) \text{ miles}}{30 \text{ miles/gal}} < 1.06 \text{ gallons}$$

Learning outcomes:

Author(s):

of gas.

2 Quiz 2A

Question 504 The table below shows the inclination angle of a mountain trail at several elevations along the trail as it ascends to Snow Lake. The inclination angle is the angle the trail makes with the horizontal.

Inclination Angle (radians)	Elevation (feet)
0.25	2300
0.2	2340
0.18	2400

Assume the inclination angle is a decreasing function of altitude and find the best possible (ie. the greatest) lower bound for the length of the trail between elevations 2300 feet and 2400 feet. Explain your reasoning thoroughly.

Explanation. We begin by considering the constant case. Here that means supposing the inclination angle is constant. Call the angle θ and consider a portion of the trail with length Δs miles and change in elevation Δh miles. Then (draw a picture) since

$$\sin \theta = \frac{\Delta h}{\Delta s},$$

we know that

$$\Delta s = \frac{\Delta h}{\sin \theta}.$$

Now the steeper the trail, the shorter the distance it takes to gain a given change in elevation. So to find a lower bound for the trail length, over each interval we should assume the trail has a constant inclination angle equal to the greatest angle in that interval. Since we've assumed the inclination angle to be a decreasing function over above section of trail, we should choose the left endpoints in our Riemann sum.

So between elevations 2300 feet and 2400 feet the length of the trail is at least

$$\frac{(2340 - 2300) \text{ ft}}{\sin(0.25)} + \frac{(2400 - 2340) \text{ ft}}{\sin(0.20)} > 463.68 \text{ ft.}$$

3 Quiz 2B

Question 505 The function

$$\theta = g(s) = \frac{s}{2} - \frac{s^3}{12}, \quad 0 \leq s \leq 3,$$

expresses the inclination angle (in radians) of the trail to Nada Lake in terms of the distance from the trailhead (measured along the trail in miles). The inclination angle is the angle the trail makes with the horizontal. It is positive (negative) when the trail slopes upward (downward) in the direction away from the trailhead.

The trail is at an elevation of 3200 feet 1.6 miles from the trailhead.

Use summation notation for an expression with n equal intervals of distance to find a lower bound for the trail's elevation (in feet) at Nada Lake (3 miles from the trailhead).

Use the graph of the function $\theta = g(s)$ shown below to help with your explanation.

Desmos link: <https://www.desmos.com/calculator/nzhemrajow>

152: Quiz 3A

Explanation. We begin by considering the constant case. Here that means supposing the inclination angle is constant. Call the angle θ and consider a portion of the trail with length Δs miles and change in elevation Δh miles. Then (draw a picture) since

$$\sin \theta = \frac{\Delta h}{\Delta s},$$

we know that

$$\Delta h = \Delta s \sin \theta.$$

Now the less steep the trail, the less the elevation gain over a given distance along the trail. So to find a lower bound for the elevation of Nada Lake, over each interval we should assume the trail has a constant inclination angle equal to the smallest angle in that interval. Since the inclination angle is a decreasing function over the trail from mile-marker 1.6 to Nada lake, we should choose the right endpoints in our Riemann sum.

With n -equal subintervals over the trail distance

$$\Delta s = (3 - 1.6) \text{ miles} = 1.4 \text{ miles},$$

each subinterval has length

$$\frac{\Delta s}{n} \text{ miles} = \frac{1.4}{n} \text{ miles}.$$

So the change in elevation from the 1.6-mile marker to Nada Lake is at least

$$\frac{1.4}{n} \sum_{i=1}^n \sin \left(g \left(1.6 + \frac{1.4}{n} i \right) \right)$$

miles.

Since the elevation of the trail is 3200 feet at the 1.6 mile-marker and there are 5280 feet in one mile, the elevation of Nada Lake is at least

$$3200 + 5280 \left(\frac{1.4}{n} \right) \sum_{i=1}^n \sin \left(g \left(1.6 + \frac{1.4}{n} i \right) \right)$$

feet.

4 Quiz 3A

Question 506 The function

$$r = g(t) = 12 - 2|t - 11|, 0 \leq t \leq 20,$$

expresses a balloon's rate of ascent (in meters/min) in terms of the number of minutes past noon.

- (a) Use geometry, not the Fundamental Theorem, to evaluate the integral

$$\int_{20}^3 g(t) dt.$$

Include a graph of the function to help with your explanation. Shade the region of integration and indicate the direction of integration.

- (b) Interpret the meaning of the integral, in its form above, in this particular scenario.

Explanation. (a) The key to graphing the function is to realize $g(t)$ has its maximum value when $2|t - 11|$ is as small as possible. This happens when $t = 11$.

Desmos link: <https://www.desmos.com/calculator/wioqqjwyt9>

152: Quiz 3A

Adding the signed area of the three triangles and integrating from right to left gives

$$\begin{aligned} \int_{20}^3 g(t) dt &= \frac{1}{2}(-3 \text{ min})(-6 \text{ meters/min}) + \frac{1}{2}(-12 \text{ min})(12 \text{ meters/min}) + \\ &= \frac{1}{2}(-2 \text{ min})(-4 \text{ meters/min}) \\ &= -59 \text{ ft.} \end{aligned}$$

- (b) The meaning of the integral is that going backward in time from 12:20pm to 12:03pm the balloon loses 59 feet of altitude. More simply stated, *the balloon is 59 feet lower at 12:03pm than it is at 12:20pm.*

5 Quiz 3B

Question 507 Evaluate the definite integral

$$\int_1^4 \frac{2 - \sqrt{t}}{t^2} dt.$$

Explanation. The key is to rewrite the integrand as

$$\frac{2 - \sqrt{t}}{t^2} = 2t^{-2} - t^{-3/2}.$$

That way we use the Fundamental Theorem and undo the power rule. Here's the computation.

$$\begin{aligned} \int_1^4 \frac{2 - \sqrt{t}}{t^2} dt &= \int_1^4 (2t^{-2} - t^{-3/2}) dt \\ &= \left(-2t^{-1} + 2t^{-1/2} \right) \Big|_1^4 \\ &= \left(-\frac{1}{2} + 2 \right) - (-1 + 2) \\ &= \frac{1}{2}. \end{aligned}$$

6 Quiz 4A

This was almost identical to Quiz 3B.

7 Quiz 4B

Question 508 Use the graph of the function

$$y = \int_2^x f(t) dt$$

shown below to answer the following questions. Explain your reasoning thoroughly in complete sentences.

Desmos link: <https://www.desmos.com/calculator/zoyrvkqbv8>

152: Quiz 4B

- (a) Solve the inequality $f(x) > 0$. Explain your reasoning.
- (b) For what value of $x \in [0, 5]$ is $f(x)$ a minimum? Explain your reasoning.
- (c) Find the minimum value of $f(x)$, $x \in [0, 5]$. Explain your reasoning and show your work.

Explanation. The key point to answering these questions is to recognize that

$$\frac{d}{dx} \left(\int_2^x f(t) dt \right) = f(x).$$

- (a) Solving the inequality $f(x) > 0$ means finding the values of x where the function graphed above is increasing. The solution set is therefore

$$\{0 < x < 1 \text{ or } 4 < x \leq 5\}.$$

- (b) The minimum value of $f(x)$ occurs where the function graphed above is decreasing at the fastest rate. This occurs at approximate $x = 3$.
- (c) The minimum value of $f(x)$ is the slope of the tangent line to the above graph at $x = 3$. Using the approximate coordinates of the points $(3, -1.6)$ and $(3.2, -2)$ on the graph, the minimum value is approximately

$$\frac{\Delta y}{\Delta x} \sim \frac{-0.4}{0.2} = -2.$$

We can make better sense of this problem by adding some context. So we'll suppose the function $f(t)$ expresses the rate of ascent (in ft/sec) of a balloon in terms of the number of seconds past noon. Then (changing the variable from x to t) the function

$$y = g_1(t) = \int_2^t f(x) dx,$$

expresses the balloon's change in height (measured in feet) from time $t = 2$ to time t seconds past noon. Adding the balloon's height at time $t = 2$, say h_0 feet, gives us the height (in feet)

$$h = g_2(t) = h_0 + \int_2^t f(x) dx$$

at time t seconds past noon.

Then since the functions g_1 and g_2 differ only by a constant, the derivatives

$$dh/dt = dy/dt$$

give the balloon's rate of ascent $f(t)$ function.

So the questions are asking us to determine

- (a) when the balloon is ascending
- (b) when the balloon is descending at the fastest rate
- (c) the fastest rate of descent, expressed as a negative rate.

8 Quiz 5A

Question 509 (a) Evaluate the definite integral

$$\int_1^2 \frac{5x}{1+9x^2} dx.$$

Use substitution as in the homework. Do not skip steps. No credit without using substitution.

Explanation. It helps to write the integral as

$$\int_1^2 \frac{5x}{1+9x^2} dx = \int_1^2 (1+9x^2)^{-1} 5x dx.$$

Then the integrand is a product of two functions,

$$(1+9x^2)^{-1}$$

and

$$5x.$$

The first, $(1+9x^2)^{-1}$, is a composition of two functions. And the second $5x$ is a scalar multiple of the derivative

$$\frac{d}{dx} (1+9x^2) = 18x$$

of the inside function of the composition.

This tells us that the substitution

$$u = 1+9x^2$$

will undo the chain rule.

So with $u = 1+9x^2$, we have

$$du = 18x dx$$

and

$$5x dx = \frac{5}{18} du.$$

When $x = 1$,

$$u = 1 + 9(1)^2 = 10$$

and when $x = 2$,

$$u = 1 + 9(2)^2 = 37.$$

So

$$\begin{aligned} \int_1^2 \frac{5x}{1+9x^2} dx &= \int_1^2 (1+9x^2)^{-1} (5x dx) \\ &= \int_{10}^{37} u^{-1} \left(\frac{5}{18} du \right) \\ &= \frac{5}{18} \int_{10}^{37} u^{-1} du \\ &= \frac{5}{18} \ln |u| \Big|_{u=10}^{u=37} \\ &= \frac{5}{18} (\ln |37| - \ln |10|) \\ &= \frac{5}{18} \ln (3.7). \end{aligned}$$

9 Quiz 5B

Question 510

Evaluate the definite integral

$$\int_0^1 \frac{8e^{-3t}}{\sqrt{9-4e^{-3t}}} dt.$$

Explanation. It helps to write the integral as

$$\int_0^1 \frac{8e^{-3t}}{\sqrt{9-4e^{-3t}}} dt = \int_0^1 8e^{-3t} (9-4e^{-3t})^{-1/2} dt.$$

Then the integrand is a product of two functions,

$$(9-4e^{-3t})^{-1/2}$$

and

$$8e^{-3t}.$$

The first, $(9-4e^{-3t})^{-1/2}$, is a composition of two functions. And the second, $8e^{-3t}$, is a scalar multiple of the derivative

$$\frac{d}{dt} (9-4e^{-3t}) = 12e^{-3t}$$

of the inside function of the composition.

This tells us that the substitution

$$u = 9 - 4e^{-3t}$$

will undo the chain rule.

So with $u = 9 - 4e^{-3t}$, we have

$$du = 12e^{-3t} dt$$

and

$$8e^{-3t} dt = \frac{2}{3} du.$$

When $t = 0$,

$$u = 9 - 4e^0 = 5$$

and when $t = 1$,

$$u = 9 - 4e^{-3}.$$

So

$$\begin{aligned} \int_0^1 \frac{8e^{-3t}}{\sqrt{9 - 4e^{-3t}}} dt &= \int_5^{9-4e^{-3}} (9 - 4e^{-3t})^{-1/2} (8e^{-3t} dt) \\ &= \int_5^{9-4e^{-3}} u^{-1/2} \left(\frac{2}{3} du \right) \\ &= \frac{2}{3} \int_5^{9-4e^{-3}} u^{-1/2} du \\ &= \frac{2}{3} (2)(u^{1/2}) \Big|_{u=5}^{u=9-4e^{-3}} \\ &= \frac{4}{3} \left(\sqrt{9 - 4e^{-3}} - \sqrt{5} \right). \end{aligned}$$

10 Quiz 6A

Question 511 A parabola with its vertex at the origin and symmetric about the y -axis passes through the point $P(-3, 6)$.

Find the area of the region bounded by the parabola, the tangent line to the parabola at P , and the x -axis.

Do this by slicing the region perpendicular to the y -axis.

Explanation. Since the parabola has its vertex at the origin and is symmetric about the y -axis, its equation is of the form

$$y = cx^2$$

for some constant $c \neq 0$.

Since the point $(-3, 6)$ lies on the parabola

$$6 = c(-3)^2$$

and

$$c = 2/3.$$

So the parabola has equation

$$y = \frac{2}{3}x^2.$$

To find an equation of the tangent line at $(-3, 6)$ we first compute the slope by evaluating the derivative dy/dx at $x = -3$. The slope is

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{x=-3} &= \left. \frac{d}{dx} \left(\frac{2}{3}x^2 \right) \right|_{x=-3} \\ &= \left. \frac{4}{3}x \right|_{x=-3} \\ &= -4. \end{aligned}$$

So an equation of the tangent line at $(-3, 6)$ is

$$y = 6 - 4(x + 3)$$

or

$$y = -6 - 4x.$$

Now slice the region perpendicular to the y -axis to get the differential rectangle AB . The endpoints share the same y -coordinate. Call it y .

The left endpoint A lies on the tangent line with equation

$$y = -6 - 4x$$

and has x -coordinate

$$x = -\frac{6 + y}{4}.$$

So A has coordinates

$$\left(-\frac{6 + y}{4}, y \right).$$

The right endpoint B lies on the parabola

$$y = \frac{2}{3}x^2.$$

So

$$x = \pm \sqrt{\frac{3}{2}}y.$$

But since the region is in the second quadrant, the x -coordinate of B is negative. So B has x -coordinate

$$x = -\sqrt{\frac{3}{2}}y$$

and coordinates

$$\left(-\sqrt{\frac{3}{2}}y, y\right).$$

So the differential rectangle AB has width dy , length

$$L = -\sqrt{\frac{3}{2}}y + \frac{6+y}{4}$$

and differential area

$$dA = L \, dy = \left(-\sqrt{\frac{3}{2}}y + \frac{6+y}{4}\right) dy.$$

The area of the region bounded is then the sum

$$\int_0^6 \left(-\sqrt{\frac{3}{2}}y + \frac{6+y}{4}\right) dy.$$

of the differential areas.

Using the fundamental theorem, the area is

$$\begin{aligned} \int_0^6 \left(-\sqrt{\frac{3}{2}}y + \frac{6+y}{4}\right) dy &= \left(-\frac{2}{3}\sqrt{\frac{3}{2}}y^{3/2} + \frac{(6+y)^2}{4}\right) \Big|_{y=-3}^{y=0} \\ &= 3/2. \end{aligned}$$

11 Quiz 6B

Question 512

A region is bounded by the coordinate axes, the line $x = -3$ and the curve

$$y = \frac{70}{4x - 15}.$$

- (a) Sketch a reasonably accurate graph of the curve and shade the region.
- (b) Write an integral that gives the area of the region.
- (c) Use substitution to evaluate the integral and compute the area of the region.

Explanation. (a) The curve

$$y = f(x) = \frac{70}{4x - 15}$$

is a linear transformation of the curve $y = 1/x$ with a vertical asymptote at $x = 15/4$ and no x -intercept. So to graph the function over the interval $x \in [-3, 0]$ we need only plot the points

$$(0, f(0)) = (1, -14/3)$$

and

$$(-3, f(-3)) = (-3, -70/27).$$

This tells us the function is negative over the interval $x \in [-3, 0]$. So a differential rectangle perpendicular to the x -axis has upper endpoint $A(x, 0)$ and lower endpoint $B(x, 70/(4x - 15))$. So with a width dx and length

$$L = 0 - \frac{70}{4x - 15} = \frac{-70}{4x - 15},$$

the rectangle has differential area

$$dA = L dx = \frac{-70}{4x - 15} dx.$$

- (b) The area of the region is the sum

$$A = \int_{-3}^0 \frac{-70}{4x - 15} dx$$

of the differential areas dA .

- (c) To evaluate the above integral, we make the substitution

$$u = 4x - 15.$$

Then

$$du = 4x dx$$

and

$$dx = \frac{1}{4} du.$$

When $x = -3$,

$$u = 4(-3) - 15 = -27$$

and when $x = 0$,

$$u = 4(0) - 15 = -15.$$

So the area of the region is

$$\begin{aligned} \int_{-3}^0 \frac{-70}{4x-15} dx &= - \int_{-27}^{-15} \frac{70}{u} \left(\frac{1}{4} du \right) \\ &= -\frac{35}{2} \ln |u| \Big|_{u=-27}^{u=-15} \\ &= \frac{35}{2} (\ln |-27| - \ln |-15|) \\ &= \frac{35}{2} \ln(27/15). \end{aligned}$$

12 Quiz8A

Question 513 The base of a solid is the parabolic region below with base $b = 2a$ and height h . Cross-sections perpendicular to the y -axis are rectangles half as tall (in the direction perpendicular to the base) as they are wide.

Desmos link: <https://www.desmos.com/calculator/mbxrsmkho8>

152: Parabola 23

Express the volume of the solid in terms of b and h . Explain your reasoning.

Explanation. Let the right endpoint B of the dashed segment above have coordinates (x, y) . Then the left endpoint A has coordinates $(-x, y)$.

The rectangular cross-section above segment $\overline{AB}$ has width $2x$, height x , and area

$$A = 2x^2.$$

So corresponding slice with differential width dy has differential volume

$$dV = A dy = 2x^2 dy.$$

To express dV in terms of y , we need an equation of the parabola. This equation has the form $y = cx^2$ and since the parabola passes through the point (a, h) ,

$$h = ca^2$$

and

$$\begin{aligned} c &= \frac{h}{a^2} \\ &= \frac{h}{(b/2)^2} \\ &= \frac{4h}{b^2}. \end{aligned}$$

So the parabola has equation

$$y = \frac{4h}{b^2}x^2.$$

Then

$$2x^2 = \frac{2b^2}{h}y$$

and

$$\begin{aligned} dV &= 2x^2 dy \\ &= \frac{2b^2}{h}y dy. \end{aligned}$$

The volume of the solid is the sum of these differential volumes as y runs from $y = 0$ to $y = h$. So the volume is

$$\begin{aligned} V &= \int_0^h \frac{2b^2}{h}y dy \\ &= \frac{b^2}{h}y^2 \Big|_{y=0}^{y=h} \\ &= b^2h. \end{aligned}$$

13 Quiz 8B

Question 514 Use integration to express the volume of the red cone in terms of the area A of its base and its height h .

Do this as follows:

- Establish a one-dimensional coordinate system as shown below.
- Use similar cones to express the area of the cross-section at coordinate w in terms of w , h , and A .
- Find an expression for the differential volume of the slice.

(d) *Integrate to find the volume of the cone.*

Desmos3D link: <https://www.desmos.com/3d/6pzlieihca>

152: Cone

Explanation. Let w be the distance from the vertex to the slicing plane and let B be the area of the cross-section. Then since the small and large cones have the same shape

$$\frac{B}{A} = \left(\frac{w}{h}\right)^2.$$

So

$$B = A \left(\frac{w}{h}\right)^2$$

and the slice with thickness differential dw has differential volume

$$dV = A \left(\frac{w}{h}\right)^2 dw.$$

Sum these volumes to get the volume of the cone,

$$\begin{aligned} V &= A \int_0^h \left(\frac{w}{h}\right)^2 dw \\ &= \frac{A}{h^2} \int_0^h w^2 dw \\ &= \frac{1}{3} Ah. \end{aligned}$$