Identifiability of B's row from endpoint counts: what can be proved, and what the E0 experiments actually show

BIO-SBI — note accompanying `report_sequential`, Part I

2026-11-07

Abstract

The sequential report (Part I: E0, X5a, X5c, E0 at 4096 clones) concludes that, with B divisions free, B's row is *structurally* non-identifiable from endpoint counts. We study the Galton–Watson (GW) model formally and with the exact likelihood (probability generating functions, no simulation). The results split the claim in two. (i) *What is true and provable*: B's row enters the mean dynamics only through its mean offspring vector, so first moments are blind to a 3-dimensional family that trades switching for division (Theorem 2); within that family the row is fixed by its second factorial moments, and B divisions always inflate clone-size variance (Theorem 3); and the variance of the Kesten–Stigum limit sees the second moments of *both* rows through a single scalar, so A's row can absorb B's (Theorem 4). (ii) *What is not true*: the model is not structurally non-identifiable. The exact Fisher information at the truth is positive definite at every endpoint $n = 2, \ldots, 12$ (Computation 6), so the parameters are locally identifiable. The non-identifiability is *practical*. At the real-like design the Cramér–Rao bound on B→A is 0.28, larger than the prior sd. Globally, the best model in which B divides with probability 0.5 is only 0.36 nats worse than the truth over all 266 organoids (Computation 9). At 4096 clones on three endpoints the bound is 0.023, and the exact likelihood rejects the NPE answer by 116–178 nats on the very datasets used in the report; the `moments` summaries still separate it (97% accuracy). That failure is in the amortised inference, not in the model. Finally, the information about B's row decays geometrically with the endpoint generation: one endpoint at generation 2–3 gives 10–15 times the information of the real-like design with the same number of organoids.

Contents

1 Setting 1

2 Exact results on moments 2

3 Exact likelihood: local identifiability and its strength 4

4 E0 at 4096 clones revisited 5

5 Profile: how flat is the likelihood away from $\theta^\star$? 6

6 Suggested changes to the report's wording 7

7 Scope, caveats and open questions 7

1 Setting

Types A (= LGR5$^+$, index 1) and B (= KRT20$^+$, index 2). Each living cell has one event per generation, drawn from its type's *row*: death, persist, switch, or division into AA, AB, BB. Write the rows as

$$\pi_A = (d_A, p_A, t_A, a_A, b_A, c_A), \qquad \pi_B = (d_B, p_B, q, a_B, b_B, c_B),$$

in the order death, persist, switch (A→B resp. B→A), div_AA, div_AB, div_BB. With $s = (x, y)$ marking A and B, the offspring probability generating functions (pgfs) are

$$f_A(x, y) = d_A + p_A x + t_A y + a_A x^2 + b_A xy + c_A y^2,$$
$$f_B(x, y) = d_B + p_B y + q\, x + a_B x^2 + b_B xy + c_B y^2.$$

$Z_n = (Z_{n,A}, Z_{n,B})$ is the clone at generation n from one A founder; its pgf is F_n^A with $F_n = f \circ F_{n-1}$, $F_0 = \mathrm{id}$. An endpoint n contributes N_n i.i.d. clones from the law of Z_n *conditioned on* $Z_n \neq 0$ (detected clones, perfect capture). A design is $\mathcal{D} = \{(n, N_n)\}$. The free parameter is $\theta = (\pi_A, \pi_B) \in \Delta_5 \times \Delta_5$ (dimension 10; `two_type`). Fixing $a_B = b_B = c_B = 0$ gives `reversible_ab` (dimension 7).

The truth used throughout E0/X5/E4 is

$$\theta^\star : \ \pi_A = (.03, .03, .05, .55, .05, .29), \quad \pi_B = (.10, .10, .80, 0, 0, 0),$$

which lies on the boundary of Δ_5 for B (no B divisions). Mean matrix $M_{kj} = \partial_j f_k(1, 1)$, Perron root ρ, right/left Perron vectors r, u ($Mr = \rho r$, $uM = \rho u$, $u \cdot r = 1$). At $\theta^\star$: $M = \left(\begin{smallmatrix} 1.18 & 0.68 \\ 0.80 & 0.10 \end{smallmatrix}\right)$, $\rho = 1.554$, $\lambda_2 = -0.274$, $r \propto (1, 0.550)$.

Three notions of identifiability. We separate them because the report's word "structural" conflates them.

Definition 1. (a) θ is *structurally (globally) identifiable* from design $\mathcal{D}$ if $\theta \mapsto (\mathcal{L}(Z_n \mid Z_n \neq 0))_{n \in \mathcal{D}}$ is injective. (b) It is *locally identifiable* at $\theta^\star$ if this holds in a neighbourhood; a sufficient condition is a positive definite Fisher information $I_{\mathcal{D}}(\theta^\star) = \sum_n N_n I_n(\theta^\star)$. (c) A parameter is *practically non-identifiable* at design $\mathcal{D}$ if it is locally identifiable but the attainable precision, e.g. the Cramér–Rao (CR) bound $\sqrt{g^\top I_{\mathcal{D}}^{-1} g}$ for a functional with gradient g, is no better than the prior spread.

A model that is structurally non-identifiable stays so for every N. A practically non-identifiable one becomes identifiable as N grows, at rate $N^{-1/2}$ (Bernstein–von Mises), and better designs move it.

2 Exact results on moments

Theorem 2 (Mean fibres)**.** *For every n, $\mathbb{E}Z_n = e_A M^n$, and B's row enters M only through its mean offspring vector*

$$\mu_B = (M_{BA}, M_{BB}) = (q + 2a_B + b_B,\ p_B + b_B + 2c_B).$$

For fixed μ_B the admissible B rows form a polytope of dimension 3 (when μ_B is interior), spanned locally by the moves

$e_1 : \ q \to q - 2\varepsilon,\ a_B \to a_B + \varepsilon,\ d_B \to d_B + \varepsilon$ *(switch → divide-into-AA and die),*
$e_2 : \ p_B \to p_B - 2\varepsilon,\ c_B \to c_B + \varepsilon,\ d_B \to d_B + \varepsilon$ *(persist → divide-into-BB and die),*
$e_3 : \ q \to q - \varepsilon,\ p_B \to p_B - \varepsilon,\ b_B \to b_B + \varepsilon$ *(switch and persist → divide-into-AB).*

Every expected count at every endpoint, and therefore every statistic that depends on the data only through mean counts, is constant on these fibres.

Proof. $\mathbb{E}[Z_n \mid Z_{n-1}] = Z_{n-1} M$ gives $\mathbb{E}Z_n = e_A M^n$. The two entries of μ_B are linear in π_B, and π_B lies in a 5-dimensional simplex. So the fibre is the intersection of the simplex with an affine subspace of codimension 2, which has dimension 3 at interior points. Each e_i keeps $\sum \pi_B = 1$ and both entries of μ_B; they are linearly independent. □

At the truth only the directions with $\varepsilon \geq 0$ are admissible (B divisions ≥ 0). The C2ST alternative of E0 (B$\to$AA $= a$, B$\to$A $= 0.8 - 2a$, B death $= 0.1 + a$) is exactly e_1.

Theorem 3 (Second moments fix the row; B divisions inflate variance)**.** *Let $G_k = \mathbb{E}[\xi_k \xi_k^\top] - \mathrm{diag}(\mu_k)$ be the second factorial moment matrix of type k's offspring ξ_k. Then*

$$G_k = \begin{pmatrix} 2a_k & b_k \\ b_k & 2c_k \end{pmatrix} \qquad (k = A, B),$$

and $\pi_k \mapsto (\mu_k, G_k)$ is an affine bijection from Δ_5 onto its image. Moreover, with $V_k = \mathrm{Cov}(\xi_k)$ and $m_j = e_A M^j$,

$$\mathrm{Cov}(Z_n) = \sum_{j=0}^{n-1} (M^\top)^{n-1-j} \Big(\sum_k m_j(k) V_k \Big) M^{n-1-j}. \tag{1}$$

Along a mean fibre (Theorem 2) only $V_B = G_B + \mathrm{diag}(\mu_B) - \mu_B \mu_B^\top$ changes, by ΔG_B. Starting from a row without B divisions (as $\theta^\star$), every admissible move has $\Delta G_B = \left(\begin{smallmatrix} 2a_B & b_B \\ b_B & 2c_B \end{smallmatrix}\right)$ with $a_B, b_B, c_B \geq 0$. Hence, for every $w \in \mathbb{R}^2_{>0}$ (e.g. $w = (1,1)$, total clone size) and $n \geq 2$,

$$\mathrm{Var}(w \cdot Z_n) \text{ strictly increases with the B-division mass } \delta = a_B + b_B + c_B.$$

Proof. The formula for G_k is a direct computation over the six outcomes: only the divisions have two daughters, and $\mathbb{E}[\xi_A(\xi_A - 1)] = 2a_k$, $\mathbb{E}[\xi_A \xi_B] = b_k$, $\mathbb{E}[\xi_B(\xi_B - 1)] = 2c_k$. Given (μ_k, G_k), the three division probabilities are read off G_k, then persist and switch off μ_k, and death is the remainder; so the map is an affine bijection. Formula (1) is the standard covariance recursion $\mathrm{Cov}(Z_n) = M^\top \mathrm{Cov}(Z_{n-1}) M + \sum_k \mathbb{E} Z_{n-1,k} V_k$ (Harris 1963; Athreya and Ney 1972, Ch. V), unrolled. Along a fibre M and μ_B are fixed, so the change is

$$\Delta \,\mathrm{Var}(w \cdot Z_n) = \sum_{j=1}^{n-1} m_j(B)\, v_j^\top \Delta G_B v_j, \qquad v_j = M^{n-1-j} w > 0,$$

and $v^\top \Delta G_B v = 2a_B v_A^2 + 2b_B v_A v_B + 2c_B v_B^2 > 0$ whenever $v > 0$ and $\delta > 0$. Since $t_A + b_A + c_A > 0$ we have $m_j(B) > 0$ for $j \geq 1$. □

Theorems 2–3 explain why the E0 C2ST found the switching/division difference only in clone-size *variability*. They also explain why `CohortMeans` cannot work. Its 6 features for 10 parameters are rank deficient by construction, and E0 showed that variance carries the information.

Theorem 4 (The variance of the growth limit sees one scalar)**.** *Let $\rho > 1$ and $W = \lim_n r \cdot Z_n / \rho^n$ (a.s. and in L^2, since the offspring have finite variance). Then*

$$\mathrm{Var}\, W = \sum_{k \in \{A,B\}} \beta_k \, r^\top V_k r, \qquad \beta_k = \sum_{j \geq 0} \rho^{-2(j+1)} m_j(k).$$

Hence, on the product of the two mean fibres (both rows moved with M fixed), $\mathrm{Var}\, W$ depends on the six second factorial moments (G_A, G_B) only through the scalar

$$\beta_A \, r^\top G_A r + \beta_B \, r^\top G_B r.$$

An increase of B's division variance can be cancelled exactly by fewer A divisions. At $\theta^\star$, $\beta = (1.000, 0.294)$.

Proof. $W_n = r \cdot Z_n / \rho^n$ is a martingale. Its increments $W_{j+1} - W_j = \rho^{-(j+1)} \sum_{\text{cells } c \text{ in gen } j} (r \cdot \xi_c - \rho r_{k(c)})$ are uncorrelated with conditional variance $\rho^{-2(j+1)} \sum_k Z_{j,k} \, r^\top V_k r$. Take expectations and sum (orthogonality of martingale increments). $W_0 = r_A$ is deterministic. Along fibres M, hence ρ, r, m_j and β, are fixed, and $r^\top V_k r = r^\top G_k r + \sum_i \mu_k(i) r_i^2 - (\mu_k \cdot r)^2$ changes only through $r^\top G_k r$. □

Check against the exact law (`scripts/b_row_theory/varW.py`): $\mathrm{Var}(r \cdot Z_n/\rho^n) = 0.3263$ ($n = 10$), 0.3286 ($n = 12$) against the formula $\mathrm{Var}\, W = 0.3302$.

Remark 5 (What Theorem 4 does and does not explain). For large clones $Z_n \approx \rho^n W u$. The dominant second-order feature of late endpoints, the spread of clone sizes, therefore constrains A's and B's division variances only jointly. This is the mechanism by which leaving B divisions free biases A's row (E0 at 4096: A persist 0.03 → 0.16, A→BB 0.29 → 0.21, §4). It is *not* a non-identifiability. The type composition, the early endpoints and the higher moments carry further constraints. A naive compensating move that matches M and $\mathrm{Var}\, W$ exactly (A: div_AA and div_BB −0.22, div_AB +0.44) is *easier* to detect than the uncompensated B move: KL 0.17–0.28 vs 0.011–0.016 nats per clone. The cheapest compensations are found numerically in §5. They raise A *death*, not A persist: B→AA instead of B→A makes extra A cells, and A death removes them. The NPE answer at 4096 (A persist 0.15) is a suboptimal version of the same compensation.

3 Exact likelihood: local identifiability and its strength

The law of Z_n is computed exactly by evaluating F_n^A on an $L \times L$ grid of roots of unity and applying a 2-D FFT. The polynomial F_n^A has degree $\le 2^n$ in each variable, so $L = 2^n + 1$ is alias-free. For $n = 10$, $L = 384$ already reproduces the alias-free result to 10^{-17}, because the mass beyond 384 cells is below 10^{-14}. Derivatives $\partial_\theta F_n$ are propagated by forward-mode differentiation through the iteration $F_n = f(F_{n-1})$ and checked against finite differences (relative error 10^{-11}). Means agree with $e_A M^n$ to machine precision. The Fisher information of the *conditioned* law $\tilde{p} = p/(1 - p_{00})$ is $I_n = \sum_{z \neq 0} \tilde{p}_z \nabla \log \tilde{p}_z \nabla \log \tilde{p}_z^\top$, in the 10 coordinates "entry minus death" of each row. Because $\theta^\star$ is on the boundary, B-division directions are one-sided. A positive definite I_n still gives $\mathrm{KL}(\theta^\star \| \theta) \ge c\,|\theta - \theta^\star|^2$ on the admissible cone near $\theta^\star$. Code: `scripts/b_row_theory/`.

Computation 6 (Local identifiability at $\theta^\star$)**.** For every single endpoint $n \in \{2, \dots, 10, 12\}$ the per-clone Fisher information $I_n(\theta^\star)$ of the 10-parameter model is positive definite (Table 1). Hence $\theta^\star$ is locally identifiable from *any single* endpoint distribution. The smallest eigenvalue is far above floating-point error: its ratio to the largest is $\ge 5 \times 10^{-7}$, against a relative accuracy of $\approx 10^{-12}$ for the computed probabilities.

n	$\lambda_{\min}(I_n)$	cond. I_n	CR var. B→A	CR var. δ	N for sd 0.05
2	0.362	1.3×10^2	1.8	1.7	730
3	0.331	1.9×10^2	2.3	1.8	920
4	0.205	5.1×10^2	3.6	3.6	1 440
5	0.051	3.2×10^3	13	14	5 300
6	0.037	6.6×10^3	18	19	7 200
7	0.015	2.3×10^4	41	39	16 000
8	0.0090	5.3×10^4	64	54	26 000
9	0.0044	1.4×10^5	124	91	50 000
10	0.0023	3.5×10^5	218	143	87 000
12	0.00067	1.8×10^6	686	413	274 000

Table 1: Exact per-clone Fisher information at $\theta^\star$, single endpoint, 10 free parameters. CR variance of B→A and of the B-division mass $\delta = a_B + b_B + c_B$ with all other parameters free (profiled), per clone: divide by N. Last column: N for a CR sd of 0.05 on B→A from that endpoint alone.

The weak direction. The eigenvector of the smallest eigenvalue of $I_{\mathcal{D}}$ is, at every design checked, dominated by B→A (≈ 0.8) against B→AA and B→AB (−0.2 to −0.5), with a smaller B persist component. This is the switching-versus-division direction of Theorem 2 (move e_1, partly e_3), now as a property of the full likelihood rather than of the means.

Proposition 7 (Practical non-identifiability at realistic designs)**.** *CR bounds for B→A and for δ (all other parameters free) are listed in Table 2. At the real-like and E0 designs the bound on B→A is 0.27–0.28. The uniform Dirichlet prior has sd 0.14 (Beta(1,5)) and the flat prior on B→A has sd 0.29. So no estimator that is unbiased near $\theta^\star$ can be more precise than the prior, and the posterior is necessarily prior-dominated. This is the regime of E0, X5a and X5c, and their conclusions hold there.*

design (endpoint: clones)	organoids	CR sd B→A	CR sd δ
real-like $\{6{:}165, 10{:}101\}$	266	0.28	0.28
E0 $\{4, 8, 12\} \times 32$	96	0.27	0.28
$\{4, 6, 10\} \times 512$	1 536	0.065	0.068
E0 at 4096: $\{4, 6, 10\} \times 4096$	12 288	0.023	0.024
$\{4\} \times 266$	266	0.12	0.12
$\{3\} \times 266$	266	0.093	0.083
$\{2, 3\} \times 133$	266	0.077	0.066
$\{3, 6, 10\} \times \{89, 89, 88\}$	266	0.12	0.12
$\{2, 4, 6, 10\} \times \approx 66$	266	0.093	0.091
$\{3, 4\} \times 1000$	2 000	0.036	0.033

Table 2: Cramér–Rao sd at $\theta^\star$ (exact Fisher information, GW, generations). Top: designs used in the report. Bottom: same 266 organoids redistributed to early endpoints.

Proposition 8 (The information about B's row is in early generations)**.** *Per clone, the information about B→A and δ decreases geometrically with the endpoint. From $n = 5$ to $n = 12$ the CR variance of B→A grows by a factor 1.4–2.3 per generation (geometric mean 1.76; Table 1), and the condition number of I_n by a factor 2–3.5. Between $n = 2$ and $n = 4$ the loss is milder (factor 2 in total).*

Interpretation. At small n the counts are small integers, and individual events are visible in the law. A clone $(2, 0)$ at $n = 2$ arises from A→B→AA but not from A→B→A. As n grows the law of Z_n approaches $\rho^n W u$ plus Gaussian composition noise. Its first moments are blind to the fibres of Theorem 2, and its main second moment ($\mathrm{Var}\, W$) mixes the two rows (Theorem 4). The geometric rate is a numerical observation over $n \leq 12$, not a proved asymptotic.
Consequence for design. With the same 266 organoids, an endpoint at generation 2–3 gives a CR sd of 0.08–0.09 on B→A, against 0.28 at $\{6, 10\}$: 10–15 times the Fisher information. Adding one early endpoint to $\{6, 10\}$ already helps (0.12 for $\{3, 6, 10\}$). Whether generation 2–3 is reachable in the wet lab (day 2–3 for a ≈20 h A cycle) and whether BH behaves alike are open (§7).

4 E0 at 4096 clones revisited

The report (§ "E0 revisited at 4096 clones") concludes "structural non-identifiability" because, at $\{4, 6, 10\} \times 4096$ with a 20k-simulation NPE, B→A stays at 0.17–0.34 (truth 0.8) and the B-division mass at its prior mean. Three exact checks contradict the "structural" reading for GW.

1. The bound. The CR sd is 0.023 (Table 2). NPE reported a posterior sd of 0.11 and a bias of 0.47, which is 20 CR sd.

2. The exact likelihood rejects the NPE answer on the report's own data. For each run we take the NPE posterior mean $\hat{\theta}$, renormalise each row, and evaluate the exact conditioned log-likelihood on the stored reference data (`reference.npz`, 3×4096 clones):

run (seed)	0	1	2	3	4
20k: $\ell(\theta^\star) - \ell(\hat\theta)$ [nats]	149.8	137.7	122.4	151.7	149.6
5k: $\ell(\theta^\star) - \ell(\hat\theta)$ [nats]	168.9	164.3	116.5	177.6	170.3
expected, $4096 \sum_n \mathrm{KL}_n(\theta^\star \| \hat\theta)$ (20k)	124	141	130	132	139

A likelihood ratio of e^{116}–e^{178} is decisive. The NPE posterior concentrates on parameters that the data exclude. The clean PPC does not contradict this: PPC compares *summaries* within 95% bands, a far coarser check than the likelihood.

3. The summaries carry the information; the network does not use it. We simulated 300 datasets at $\theta^\star$ and 300 at $\hat\theta$ (seed 0, 20k) with the `moments` summary. A regularised logistic regression separates them with 5-fold CV accuracy **0.968** (0.722 at 165 clones per endpoint). The most shifted features belong to the earliest endpoint (h4: sd of $\log(1+B)$, median of $\log(1+A)$) and to the A–B correlation. In Gaussian-equivalent terms, accuracy 0.968 corresponds to an expected log-LR of ≈ 7 nats. So the summary keeps about 5% of the exact 124 nats along this direction, and still more than enough to reject $\hat\theta$ ($\approx$ 900:1). The amortised NPE, trained on 20k draws spread over a 10-dimensional prior, does not resolve this.

Conclusion. In GW the B row is not structurally non-identifiable. It is *practically* non-identifiable at the designs of E0, X5a, X5c and the real-like design (Proposition 7). At 4096 clones per endpoint it is identifiable from the data, from the exact likelihood and to a useful extent from the summaries, and the failure reported there is an inference failure (amortised NPE with uninformative coverage of the relevant region). The report's recommendations survive: fix or measure B divisions, because realistic designs cannot decide them. Their justification and wording should change (§6).

5 Profile: how flat is the likelihood away from $\theta^\star$?

The Fisher information is local. For the global picture we compute the profile

$$K_{\mathcal{D}}(\delta) = \min \Big\{ \sum_n w_n \,\mathrm{KL}\left(\tilde p_n(\theta^\star) \,\|\, \tilde p_n(\theta)\right) :\ \theta \in \Delta_5 \times \Delta_5,\ a_B + b_B + c_B = \delta \Big\},$$

the per-clone cost of the best-fitting model in which B divides with probability δ. All other 9 coordinates are optimised (L-BFGS, three starts, exact law). Then $N\,K(\delta)$ is the expected log-likelihood ratio against the best "B divides" explanation with N organoids in total, split in the design's proportions ($w_n = N_n/N$).

B-division mass δ	0.02	0.05	0.1	0.2	0.3	0.5
real-like {6:165, 10:101} (weights 165/266, 101/266)						
$K(\delta)$ per clone	$9.8 \cdot 10^{-6}$	$5.4 \cdot 10^{-5}$	$1.9 \cdot 10^{-4}$	$5.8 \cdot 10^{-4}$	$9.3 \cdot 10^{-4}$	$1.36 \cdot 10^{-3}$
$266\,K(\delta)$ [nats]	0.003	0.014	0.05	0.15	0.25	0.36
best B→A / A death	0.78 / 0.04	0.76 / 0.05	0.71 / 0.07	0.62 / 0.10	0.55 / 0.12	0.39 / 0.16
E0 at 4096 {4, 6, 10}, equal weights						
$K(\delta)$ per clone	$2.9 \cdot 10^{-5}$	$1.6 \cdot 10^{-4}$	$5.5 \cdot 10^{-4}$	$1.6 \cdot 10^{-3}$	$2.8 \cdot 10^{-3}$	$4.2 \cdot 10^{-3}$
$12\,288\,K(\delta)$ [nats]	0.35	1.9	6.7	20	34	51
best B→A / A death	0.78 / 0.04	0.76 / 0.05	0.72 / 0.07	0.64 / 0.10	0.54 / 0.12	0.33 / 0.15

Table 3: Profile KL $K(\delta)$ (truth $\delta = 0$, B→A = 0.8, A death 0.03) and the B→A and A death of the best-fitting model at each δ. Expected log-LR of truth vs. that model = number of organoids $\times K(\delta)$. Full minimisers: `scripts/b_row_theory/profile.jsonl`.

Computation 9 (Global profile)**.** (i) $K(\delta) > 0$ for every δ on the grid. Near 0 it matches the Fisher prediction $\delta^2/(2\sigma_\delta^2)$, with σ_δ^2 the per-clone CR variance of the design (E0 at 4096: predicted $2.8{\cdot}10^{-5}$ at $\delta = 0.02$, computed $2.9{\cdot}10^{-5}$). Above $\delta \approx 0.2$, K grows sub-quadratically: explanations in which B divides a lot are relatively cheap. (ii) *Real-like design:* the best model with $\delta = 0.5$ (B→A = 0.39) has an expected log-LR of only 0.36 nats against the truth, a likelihood ratio of about 1.4. Every $\delta \leq 0.5$ is statistically indistinguishable from $\delta = 0$ with 266 organoids. This is the non-local form of Proposition 7. Three nats of expected evidence against $\delta = 0.5$ would need $\approx 2\,200$ organoids in the real-like split, or ≈ 720 split equally over $\{4, 6, 10\}$. (iii) *E0 at 4096:* the same alternative costs 51 nats, so it is excluded. Its parameters (B→A 0.33, A death 0.15, B→AA 0.34) are close to the NPE posterior mean (B→A 0.32, A death 0.09, B→AA 0.21). NPE found the bottom of the valley but not its depth. (iv) *The compensation* is mainly move e_1 (switch → divide-into-AA) in B's row, with the extra death *moved to A's row.* A death rises from 0.03 to 0.16 while B death stays at 0.06–0.10. This is the precise form of "the confounding leaks into A's row".

X5c ({6,10}×{32,128,512}) is consistent with this. Using the real-like profile as an approximation (its split differs slightly), 1 024 organoids give ≈ 1.4 nats of expected evidence against $\delta = 0.5$. So its failure is mainly the likelihood's, not only the network's.

6 Suggested changes to the report's wording

- Replace "structural non-identifiability" (E0 at 4096, Part I summary, "Decided: structural") with: *"practically non-identifiable at realistic designs (Cramér–Rao sd of B→A 0.27–0.28 > prior sd); locally identifiable in principle (positive definite exact Fisher information); at 4096 clones per endpoint the data and the summaries identify it, but amortised NPE does not."*
- "A clean PPC is no evidence that the B row is right" stays true. Add: the exact likelihood ratio against the NPE answer is e^{120}–e^{180} on the same data.
- "Biological constraints help more than data" holds at realistic sizes, with the caveat that more *early* data help much more than more late data (Proposition 8).
- E0's first attribution (prior tail) and its revision stay as they are. The revision "the confounding is the root cause" is correct as a description of practical non-identifiability.

7 Scope, caveats and open questions

- **Global identifiability** (injectivity) is not proved. The profile of §5 is numerical evidence over a grid of δ, from finitely many starts.
- **One truth.** All computations are at $\theta^\star$. Theorems 2–4 hold for every θ; the Fisher numbers do not. The same scripts compute them at any other truth (e.g. the X5a truths).
- **Boundary.** $\theta^\star$ has $\delta = 0$ on the boundary. The MLE of δ is asymptotically a censored normal, and the CR bound is the local scale, not an exact sd.
- **CR vs posterior.** The CR bound concerns unbiased estimators. Posteriors with informative priors can have smaller sd but are then biased, which is the E0/X5 picture. Asymptotically (data dominating) the posterior sd approaches the CR sd.
- **Bellman–Harris.** Not analysed. Continuous lifetimes add a free time scale. The BH ridge of Part III and the larger B→A error under BH at 4096 suggest weaker information there. An exact BH analysis would need the age-dependent pgf (integral equation), which is not done here.

- **Generations vs days.** The early-endpoint recommendation is in GW generations. Its translation to days, and to what can be counted and typed in 2–8-cell organoids, is a wet-lab question.
- **Inference fix at 4096.** The analysis suggests likelihood-based inference (exact GW likelihood here, or sequential/local NPE around the data), or summaries targeted at early endpoints. It does not show that a particular NPE configuration would succeed.

Reproducibility. `scripts/b_row_theory/`: `gwexact.py` (exact law, derivatives, Fisher information, KL), `fisher.py`, `fisher2.py` (Tables 1, 2), `llr.py` (§4, item 2; reads `results/e4/gw_xxl_two_type_*`), `summary_twopoint.py` (item 3), `varW.py` (Theorem 4 check), `profile.py` (§5). Run with `uv run python scripts/b_row_theory/<script>.py` from the repository root.