




Charge Equilibration via Neural Network Techniques (CENT)

Somayeh Faraji

Institute for Advanced Studies in Basic Sciences (IASBS), Zanjan

2nd workshop on Machine Learning in Physics,
University of Tehran, Tehran
October 5, 2018




Charge transfer at organic/metal interfaces

Introduction

Methods

CENT

Applications



Multi-orbital charge transfer at nickel tetraphenyl porphyrin molecules adsorbed on Cu(100)

DFT and the Hellmann Feynman Theorem

$$\hat{H}_{tot}\psi(\mathbf{r}, \mathbf{R}) = E\psi(\mathbf{r}, \mathbf{R})$$

Born-Openheimer approximation:

$$(\hat{H}_e(\{\mathbf{R}\}) - E_e(\{\mathbf{R}\}))\psi_e(\mathbf{r}; \mathbf{R}) = 0 \text{ where } \hat{H}_e = \hat{T}_e + \hat{V}_{ee} + \hat{V}_{e-I}$$

Hohenberg-Kohn theorems:

$$E[\rho] = T[\rho] + \int \rho(\mathbf{r})V_{ext}(\mathbf{r})d\mathbf{r} + E_{int}[\rho]$$

Kohn-Sham equations:

$$E_{KS} = - \sum_{i=1}^{occ} \frac{|\nabla_i^2|}{2} + \int V_{ext}(\mathbf{r})\rho(\mathbf{r})d\mathbf{r} + \frac{1}{2} \int \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' + E_{xc}[\rho(\mathbf{r})]$$

$$\left[-\frac{1}{2} \nabla^2 + V_{ext}(\mathbf{r}) + V_H(\mathbf{r}) + V_{xc}(\mathbf{r}) \right] \psi_i(\mathbf{r}) = \varepsilon_i \psi_i(\mathbf{r})$$

$$E_{gs}(\{\mathbf{R}\}) = \sum_{i=1}^{occ} \varepsilon_i - \int \rho(\mathbf{r})V_{xc}(\mathbf{r})d\mathbf{r} + \frac{1}{2} \int \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' + E_{xc}[\rho]$$

Hellmann Feynmann theorem

$$F = - \int \frac{\rho(\mathbf{r}')(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d\mathbf{r}'$$

A neural network scheme for ionic systems based on charge equilibration method

Introduction

Methods

CENT

Applications

$$U_{tot}(\{q_i\}) = \sum_{i=1}^N \left(E_i^0 + \chi_i q_i + \frac{1}{2} J_{ii} q_i^2 \right) + \frac{1}{2} \int \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}'$$

Simplest atomic charge distribution: Gaussian

$$\rho_i(\mathbf{r}) = \frac{q_i}{\alpha_i^3 \pi^{\frac{3}{2}}} \exp\left(-\frac{|\mathbf{r} - \mathbf{R}_i|^2}{\alpha_i^2}\right)$$

Resulting analytic energy expression:

$$U_{tot}(\{q_i\}, \{\mathbf{R}_i\}) = \sum_{i=1}^N \left(E_i^0 + \chi_i q_i + \frac{1}{2} \left(J_{ii} + \frac{2\gamma_{ii}}{\sqrt{\pi}} \right) q_i^2 \right) + \sum_{i>j} q_i q_j \frac{\text{erf}(\gamma_{ij} R_{ij})}{R_{ij}}$$

$$\gamma_{ij} = \frac{1}{\sqrt{\alpha_i^2 + \alpha_j^2}}$$

Charge equilibration process

Introduction

Methods

CENT

Applications

$$\frac{\partial U_{tot}}{\partial q_i} = 0, \forall i = 1, \dots, N \Rightarrow \sum_{j=1}^N A_{ij}q_j + \chi_i = 0.$$

where

$$A_{ij} = \begin{cases} J_{ii} + \frac{2\gamma_{ii}}{\sqrt{(\pi)}} & \text{for } i = j \\ \frac{erf(\gamma_{ij}r_{ij})}{r_{ij}} & \text{for } i \neq j \end{cases}$$

Adding the constraint total charge via the Lagrange multipliers leads

$$q_{tot} = \sum_i^N q_i \quad \tilde{A}\mathbf{Q} = -\boldsymbol{\chi}$$
$$\left(\begin{array}{ccc|c} A_{i,j} & & & 1 \\ & \vdots & & \\ & 1 & & 0 \end{array} \right) \left(\begin{array}{c} q_1 \\ \vdots \\ \frac{q_N}{\lambda} \end{array} \right) = \left(\begin{array}{c} -\chi_1 \\ \vdots \\ -\chi_N \\ q_{tot} \end{array} \right)$$


Schematic illustration of the charge equilibration via neural network technique (CENT)

Introduction

Methods

CENT

Applications



- Explicit short range environment dependence of the χ_i 's.
- Implicit long range environment dependence of the q_i 's.


Transferability: Testing CENT for bulk CaF₂

Introduction

Methods


CENT

Applications




- 2800 data points during the training process
 - *FHI-aims* all-electron full-potential code
 - PBE xc-functional


Transferability: Testing CENT for bulk CaF₂




(a) Fm-3m



(b) Pnma



(c) Pmc21



(d) P4/mmm

• Transferability

- Bulk properties (lattice constants, elastic properties etc.)
 - Heat capacities
 - Structural phase transition at high pressure
 - Phonon dispersion of fluorite structure
 - Vacancies in fluorite structure
 - Surfaces of fluorite structure
-
- DFT
 - PBE (XC functional used for reference data points)
 - LDA
 - Born-Mayer-Huggins (BMH) potential

The average errors of formation energies

CENT: 1 %

LDA: 11 %

BMH: 56 %


Structural phase transition at high pressure

Introduction

Methods

CENT

Applications




Phonon dispersion of fluorite structure

Introduction

Methods

CENT

Applications



- Phonopy code is used for the calculation of phonon dispersion
- Acoustic modes: CENT results agrees with those of PBE
- Optical modes: results of all methods are the same qualitatively and differ from each other quantitatively

New Crystal Phase of CaF_2

Introduction



Methods

CENT

Applications

- New Found Crystal Phase

- Space group: P-62m
- Lattice vectors from PBE (in Ang):
 - a:(3.55, 0, 0)
 - b:(0, 6.09, 0)
 - c:(0, 3.04, 5.28)




Energy landscape of ZnO

Introduction

Methods

CENT

Applications




Energy landscape of ZnO

Introduction

Methods

CENT

Applications





Phonon dispersion of new phases of ZnO

Introduction

Methods

CENT

Applications







Two-dimensional hexagonal nano sheet(HNS) of TiO_2

Introduction

Methods

CENT

Applications



Thank you...

Introduction

Methods

CENT

Applications