



Royal Institute of
Technology

DGM MLE AND POSTERIOR

ORDER OF PRODUCTS

$$p(x | \theta) = \prod_{v=1}^V p(x_v | x_{\text{pa}(v)}, \theta_v)$$

$$\begin{aligned} p(\mathcal{D} | \theta) &= \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \\ &= \prod_{v=1}^V \prod_{n=1}^N p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) = \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \theta_v) \end{aligned}$$

CATEGORICAL – NOTATION

★ For a $v \in [M]$,

values $k \in S_v$

combined values

$$c \in C_v = \prod_{s \in \text{pa}(v)} S_s$$


Cartesian product

★ Cat CPDs

where $P(x_v | x_{\text{pa}(v)} = c) = \text{Cat}(\theta_{vc})$

and $\theta_{vck} = P(X_v = k | X_{\text{pa}(v)} = c)$

NOTATION EXAMPLE



$$C_g = \prod_{s \in \text{pa}(g)} S_s = S_i \times S_d = \{\langle 0,0 \rangle, \langle 0,1 \rangle, \langle 1,0 \rangle, \langle 1,1 \rangle\}$$

THE DGM LIKELIHOOD FACTORIZES

★ Data

$$\mathcal{D} = \{x^1, \dots, x^N\} \quad x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$p(\mathcal{D} | \theta) = \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v)$$

★ Notation

$$p(X_v = k | X_{\text{pa}} = c) = \theta_{vck}$$

★ Counts

$$N_{vck} = \sum_{n=1}^N I(x_v^n = k, x_{\text{pa}(v)}^n = c)$$

$$N_{vc} = \sum_{n=1}^N I(x_{\text{pa}(v)}^n = c)$$

THE DGM LIKELIHOOD FACTORIZES

★ Complete data

$$\mathcal{D} = \{x^1, \dots, x^N\}$$

$$x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$p(\mathcal{D} | \theta) = \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v)$$

v's CPD



THE DGM LIKELIHOOD FACTORIZES

★ Complete data

$$\mathcal{D} = \{x^1, \dots, x^N\}$$

$$x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$\begin{aligned} p(\mathcal{D} | \theta) &= \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \\ &= \prod_{v=1}^V \prod_{n=1}^N p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \end{aligned}$$

THE DGM LIKELIHOOD FACTORIZES

★ Complete data

$$\mathcal{D} = \{x^1, \dots, x^N\}$$

$$x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$\begin{aligned} p(\mathcal{D} | \theta) &= \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \\ &= \prod_{v=1}^V \prod_{n=1}^N p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) = \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \theta_v) \end{aligned}$$

values of v values of v 's parents

Called: decomposable likelihood (factorizes into family-factors)

PARAMETER AND COUNTS

$$N_{vc} = \sum_{n=1}^N I(x_{\text{pa}(v)}^n = c)$$

$$N_{vck} = \sum_{n=1}^N I(x_v^n = k, x_{\text{pa}(v)}^n = c)$$

★ Complete data

$$\mathcal{D} = \{x^1, \dots, x^N\}$$

$$x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$\begin{aligned} p(\mathcal{D} | \theta) &= \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \\ &= \prod_{v=1}^V \prod_{n=1}^N p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) = \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \theta_v) \\ &= \prod_{v=1}^V \prod_{c \in C_v} \prod_{k \in S_v} \theta_{vck}^{N_{vck}} \end{aligned}$$

THE LIKELIHOOD FACTORIZES

$$N_{vc} = \sum_{n=1}^N I(x_{\text{pa}(v)}^n = c)$$

$$N_{vck} = \sum_{n=1}^N I(x_v^n = k, x_{\text{pa}(v)}^n = c)$$

★ Complete data

$$\mathcal{D} = \{x^1, \dots, x^N\}$$

$$x^n = x_1^n, \dots, x_V^n$$

★ Likelihood

$$\begin{aligned} p(\mathcal{D} | \theta) &= \prod_{n=1}^N p(x^n | \theta) = \prod_{n=1}^N \prod_{v=1}^V p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) \\ &= \prod_{v=1}^V \prod_{n=1}^N p(x_v^n | x_{\text{pa}(v)}^n, \theta_v) = \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \theta_v) \end{aligned}$$

$$= \prod_{v=1}^V \prod_{c \in C_v} \prod_{k \in S_v} \theta_{vck}^{N_{vck}}$$

So ML estimate

$$\theta_{vck} = N_{vck} / N_{vc}$$

BAYESIAN PARAMETER LEARNING

- ★ Decomposable prior

$$p(\boldsymbol{\theta}) = \prod_{v=1}^V p(\boldsymbol{\theta}_v) = \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Dir}(\boldsymbol{\theta}_{vc} | \boldsymbol{\alpha}_{vc})$$

- ★ Gives decomposable posterior

$$\begin{aligned} p(\boldsymbol{\theta} | \mathcal{D}) &\propto p(\mathcal{D} | \boldsymbol{\theta}) p(\boldsymbol{\theta}) \\ &= \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \boldsymbol{\theta}_v) p(\boldsymbol{\theta}_v) \end{aligned}$$

BAYESIAN PARAMETER LEARNING

- ★ Decomposable prior

$$p(\boldsymbol{\theta}) = \prod_{v=1}^V p(\boldsymbol{\theta}_v) = \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Dir}(\boldsymbol{\theta}_{vc} | \boldsymbol{\alpha}_{vc})$$

- ★ Gives decomposable posterior

$$\begin{aligned} p(\boldsymbol{\theta} | \mathcal{D}) &\propto p(\mathcal{D} | \boldsymbol{\theta}) p(\boldsymbol{\theta}) \\ &= \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \boldsymbol{\theta}_v) p(\boldsymbol{\theta}_v) \\ &= \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Cat}(N_{vc1}, \dots, N_{vc|S_v} | \boldsymbol{\theta}_{vc}) \text{Dir}(\boldsymbol{\theta}_{vc} | \boldsymbol{\alpha}_{vc}) \\ &\propto \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Dir}(\boldsymbol{\theta}_{vc} | N_{vc1} + \alpha_{vc1}, \dots, N_{vc|S_v} + \alpha_{vc|S_v}) \end{aligned}$$

BAYESIAN PARAMETER LEARNING

- ★ Decomposable prior

$$p(\boldsymbol{\theta}) = \prod_{v=1}^V p(\boldsymbol{\theta}_v) = \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Dir}(\boldsymbol{\theta}_{vc} | \boldsymbol{\alpha}_{vc})$$

- ★ Gives decomposable posterior

$$\begin{aligned} p(\boldsymbol{\theta} | \mathcal{D}) &\propto p(\mathcal{D} | \boldsymbol{\theta}) p(\boldsymbol{\theta}) \\ &= \prod_{v=1}^V p(\mathcal{D}_v | \mathcal{D}_{\text{pa}(v)}, \boldsymbol{\theta}_v) p(\boldsymbol{\theta}_v) \\ &= \prod_{\substack{v \in [V] \\ c \in S_{\text{pa}(v)}}} \text{Cat}(N_{vc1}, \dots, N_{vc|S_v} | \boldsymbol{\theta}_{vc}) \text{Dir}(\boldsymbol{\theta}_{vc} | \boldsymbol{\alpha}_{vc}) \end{aligned}$$