

# 2D boundary integral equations and the Nyström method

**Alex Barnett<sup>1</sup>** and **Fruzsina Agocs<sup>1</sup>**

Computational Tools 2024 BIE workshop. Day 1, 6/10/24

<sup>1</sup>Center for Computational Mathematics, Flatiron Institute, Simons Foundation

## Integral equations on 1D interval

Given: i) function  $\sigma(t)$  defined on interval  $[0, 2\pi)$ ,    periodic:  $\sigma(2\pi) = \sigma(0)$ , etc  
ii) “kernel” function  $k(t, s)$  defined on square  $[0, 2\pi)^2$ ,

Integral *operator*  $K$  acts on  $\sigma$  to give another function  $K\sigma$ :

$$(K\sigma)(t) := \int_0^{2\pi} k(t, s)\sigma(s)ds, \quad t \in [0, 2\pi)$$

continuous analog of  
matrix-vector prod.  $Ax$

## Integral equations on 1D interval

Given: i) function  $\sigma(t)$  defined on interval  $[0, 2\pi)$ ,    periodic:  $\sigma(2\pi) = \sigma(0)$ , etc  
 ii) “kernel” function  $k(t, s)$  defined on square  $[0, 2\pi)^2$ ,


Integral operator  $K$  acts on  $\sigma$  to give another function  $K\sigma$ :

$$(K\sigma)(t) := \int_0^{2\pi} k(t, s)\sigma(s)ds, \quad t \in [0, 2\pi)$$

continuous analog of  
matrix-vector prod.  $Ax$

Integral equation:  $(I + K)\sigma = f$ , ie

$$\sigma(t) + \int_0^{2\pi} k(t, s)\sigma(s)ds = f(t), \quad t \in [0, 2\pi)$$



analog of lin. sys.  $Ax = b$

Fredholm “second kind” (due to presence of  $I$ , otherwise called “first kind”)

# Integral equations on 1D interval

Given: i) function  $\sigma(t)$  defined on interval  $[0, 2\pi)$ ,    periodic:  $\sigma(2\pi) = \sigma(0)$ , etc  
 ii) “kernel” function  $k(t, s)$  defined on square  $[0, 2\pi)^2$ ,


Integral operator  $K$  acts on  $\sigma$  to give another function  $K\sigma$ :

$$(K\sigma)(t) := \int_0^{2\pi} k(t, s)\sigma(s)ds, \quad t \in [0, 2\pi)$$

continuous analog of matrix-vector prod.  $Ax$

Integral equation:  $(I + K)\sigma = f$ , ie

$$\sigma(t) + \int_0^{2\pi} k(t, s)\sigma(s)ds = f(t), \quad t \in [0, 2\pi)$$



analog of lin. sys.  $Ax = b$

Fredholm “second kind” (due to presence of  $I$ , otherwise called “first kind”)

If kernel continuous, or “weakly” singular (integrable),  $K$  is **compact**:

- eigenvalues  $(K\phi_k = \lambda_k\phi_k)$  discrete, with  $\lim_{k \rightarrow \infty} \lambda_k = 0$   
unless some  $\lambda_k = -1$ , 2nd kind IE has at most one soln:  $\text{Nul}(I + K) = \{0\}$
- $\text{Nul}(I + K) = \{0\} \Rightarrow$  existence of solution for any  $f$  Fredholm Alternative  
like square matrix (finite-dim), recall: uniqueness  $\Rightarrow$  consistent for any RHS

## Integral equations on 1D interval

- Given: i) function  $\sigma(t)$  defined on interval  $[0, 2\pi)$ ,    periodic:  $\sigma(2\pi) = \sigma(0)$ , etc  
ii) “kernel” function  $k(t, s)$  defined on square  $[0, 2\pi)^2$ ,


Integral operator  $K$  acts on  $\sigma$  to give another function  $K\sigma$ :

$$(K\sigma)(t) := \int_0^{2\pi} k(t, s)\sigma(s)ds, \quad t \in [0, 2\pi)$$

continuous analog of  
matrix-vector prod.  $Ax$

Integral equation:  $(I + K)\sigma = f$ , ie

$$\sigma(t) + \int_0^{2\pi} k(t, s)\sigma(s)ds = f(t), \quad t \in [0, 2\pi)$$



analog of lin. sys.  $Ax = b$

Fredholm “second kind” (due to presence of  $I$ , otherwise called “first kind”)

If kernel continuous, or “weakly” singular (integrable),  $K$  is **compact**:

- eigenvalues  $(K\phi_k = \lambda_k\phi_k)$  discrete, with  $\lim_{k \rightarrow \infty} \lambda_k = 0$   
unless some  $\lambda_k = -1$ , 2nd kind IE has at most one soln:  $\text{Nul}(I + K) = \{0\}$
- $\text{Nul}(I + K) = \{0\} \Rightarrow$  existence of solution for any  $f$     Fredholm Alternative  
like square matrix (finite-dim), recall: uniqueness  $\Rightarrow$  consistent for any RHS

Contrast 1st kind IE  $K\sigma = f$  is ill-posed problem (unstable)!

## Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For  $f$  smooth, superalgebraically convergent ("spectral"): error =  $\mathcal{O}(N^{-p})$  for any  $p$

## Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For  $f$  smooth, superalgebraically convergent ("spectral"): error =  $\mathcal{O}(N^{-p})$  for any  $p$

Apply quadr. to integral in 2nd kind IE:

call the resulting approx soln  $\tilde{\sigma}$

$$\tilde{\sigma}(t) + \sum_{j=1}^N k(t, t_j) w_j \tilde{\sigma}(t_j) = f(t), \quad t \in [0, 2\pi) \quad (*)$$

## Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For  $f$  smooth, superalgebraically convergent ("spectral"): error =  $\mathcal{O}(N^{-p})$  for any  $p$

Apply quadr. to integral in 2nd kind IE:

call the resulting approx soln  $\tilde{\sigma}$

$$\tilde{\sigma}(t) + \sum_{j=1}^N k(t, t_j) w_j \tilde{\sigma}(t_j) = f(t), \quad t \in [0, 2\pi) \quad (*)$$

Holds for all  $t$ ; in particular, holds at each  $t_i$ ,  $i = 1, \dots, N$ , giving:

$$\sigma_i + \sum_{j=1}^N k(t_i, t_j) w_j \sigma_j = f(t_i), \quad i = 1, \dots, N \quad \text{where } \sigma_j := \tilde{\sigma}(t_j)$$



## Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For  $f$  smooth, superalgebraically convergent ("spectral"): error =  $\mathcal{O}(N^{-p})$  for any  $p$

Apply quadr. to integral in 2nd kind IE:

call the resulting approx soln  $\tilde{\sigma}$

$$\tilde{\sigma}(t) + \sum_{j=1}^N k(t, t_j) w_j \tilde{\sigma}(t_j) = f(t), \quad t \in [0, 2\pi) \quad (*)$$

Holds for all  $t$ ; in particular, holds at each  $t_i$ ,  $i = 1, \dots, N$ , giving:

$$\sigma_i + \sum_{j=1}^N k(t_i, t_j) w_j \sigma_j = f(t_i), \quad i = 1, \dots, N \quad \text{where } \sigma_j := \tilde{\sigma}(t_j)$$

Write as:  $A\sigma = f$   $N \times N$  lin sys, entries  $a_{ij} = \delta_{ij} + k(t_i, t_j) w_j$ , and  $f_j := f(t_j)$

solve? dense direct  $\mathcal{O}(N^3)$ ; dense iter.  $\mathcal{O}(N^2)$ ; fast iter.  $\approx \mathcal{O}(N)$ ; fast direct  $\approx \mathcal{O}(N^{(d+1)/2})$

Why want 2nd kind? eigs( $A$ ) accumulate only at  $+1 \Rightarrow$  iterative converges fast

# Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For  $f$  smooth, superalgebraically convergent ("spectral"): error =  $\mathcal{O}(N^{-p})$  for any  $p$

Apply quadr. to integral in 2nd kind IE:

call the resulting approx soln  $\tilde{\sigma}$

$$\tilde{\sigma}(t) + \sum_{j=1}^N k(t, t_j) w_j \tilde{\sigma}(t_j) = f(t), \quad t \in [0, 2\pi) \quad (*)$$

Holds for all  $t$ ; in particular, holds at each  $t_i$ ,  $i = 1, \dots, N$ , giving:

$$\sigma_i + \sum_{j=1}^N k(t_i, t_j) w_j \sigma_j = f(t_i), \quad i = 1, \dots, N \quad \text{where } \sigma_j := \tilde{\sigma}(t_j)$$

Write as:  $A\sigma = f$   $N \times N$  lin sys, entries  $a_{ij} = \delta_{ij} + k(t_i, t_j) w_j$ , and  $f_j := f(t_j)$


solve? dense direct  $\mathcal{O}(N^3)$ ; dense iter.  $\mathcal{O}(N^2)$ ; fast iter.  $\approx \mathcal{O}(N)$ ; fast direct  $\approx \mathcal{O}(N^{(d+1)/2})$

Why want 2nd kind? eigs( $A$ ) accumulate only at  $+1 \Rightarrow$  iterative converges fast

Sometimes for BIE (eg, far-field eval), node values  $\{\sigma_j\}_{j=1}^N$  suffice.

If not, interpolate from them to any  $t \in [0, 2\pi)$ . Two approaches:

- either: rearrange (\*) to give  $\tilde{\sigma}(t) = \dots$ , called "Nyström interpolant" (rare)
- or (common): use local high-order Lagrange (or global spectral) interpolation:






# Demo Nyström on interval (1D)

day1/code/nyst1d\_demos.m

```
kfun = @(t,s) exp(3*cos(t-s));  
ffun = @(t) cos(5*t+1);  
N = 30;  
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);  
A = eye(N) + bsxfun(kfun,t',t)*diag(w);  
rhs = ffun(t');  
sigmaj = A\rhs;
```

*% smooth convolutional kernel, periodic domain [0,2pi)  
% smooth data (RHS) func  
% number of unknowns: should study convergence as N grows...  
% PTR nodes and weights, row vecs  
% identity plus fill k(t\_i,t\_j)w\_j for i,j=1..N  
% col vec  
% dense direct square solve (pivoted LU), gives col vec*



“self-convergence”:  
use  $N=40$  as “true”

$f$  and  $k$  smooth

⇒  $\sigma$  smooth

⇒ spectral conv?




**Thm.** (Anselone, Kress,...): error at node values (and Nyström interpolant) same order as that of quadrature rule applied to integrand  $k(t, \cdot)\sigma$ .

# Demo Nyström on interval (1D)

day1/code/nyst1d\_demos.m

```
kfun = @(t,s) exp(3*cos(t-s));
ffun = @(t) cos(5*t+1);
N = 30;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
A = eye(N) + bsxfun(kfun,t',t)*diag(w);
rhs = ffun(t');
sigma_j = A\rhs;
```

```
% smooth convolutional kernel, periodic domain [0,2pi)
% smooth data (RHS) func
% number of unknowns: should study convergence as N grows...
% PTR nodes and weights, row vecs
% identity plus fill k(t_i,t_j)w_j for i,j=1..N
% col vec
% dense direct square solve (pivoted LU), gives col vec
```





“self-convergence”:  
use  $N=40$  as “true”

$f$  and  $k$  smooth  
 $\Rightarrow \sigma$  smooth  
 $\Rightarrow$  spectral conv?

**Thm.** (Anselone, Kress,...): error at node values (and Nyström interpolant) same order as that of quadrature rule applied to integrand  $k(t, \cdot)\sigma$ .

- Then,  $f$  or  $k$  nonsmooth?  
worse (here *algebraic*) convergence using plain PTR rule:

Qu: why does order appear to improve at end?



## Laplace fundamental solution in $\mathbb{R}^2$

Eg PDE: Poisson eqn  $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$  Laplacian

notation:  $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$  is a point. This frees up  $\mathbf{y} \in \mathbb{R}^2$  as another point (not y-coord!)

not well-posed unless add BC! BIEs are good for *homogeneous* PDEs (driving  $g \equiv 0$ )

# Laplace fundamental solution in $\mathbb{R}^2$

Eg PDE: Poisson eqn  $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$  Laplacian

notation:  $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$  is a point. This frees up  $\mathbf{y} \in \mathbb{R}^2$  as another point (not y-coord!)

not well-posed unless add BC! BIEs are good for *homogeneous* PDEs (driving  $g \equiv 0$ )

A well-posed\* BVP:

$$\Delta u = 0 \text{ in } \Omega$$

PDE ( $u$  harmonic)

$$u = f \text{ on } \Gamma$$

Dirichlet BC

\*exists, unique,  
continuous w.r.t. data



# Laplace fundamental solution in $\mathbb{R}^2$

Eg PDE: Poisson eqn  $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$  Laplacian

notation:  $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$  is a point. This frees up  $\mathbf{y} \in \mathbb{R}^2$  as another point (not y-coord!)

not well-posed unless add BC! BIEs are good for *homogeneous* PDEs (driving  $g \equiv 0$ )

A well-posed\* BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE (} u \text{ harmonic)}$$

$$u = f \text{ on } \Gamma \quad \text{Dirichlet BC}$$


\*exists, unique,  
continuous w.r.t. data



Laplace fundamental soln:  $\Phi(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi} \log \frac{1}{r}$  where  $r := \|\mathbf{x} - \mathbf{y}\|$

obeys  $-\Delta_{\mathbf{x}} \Phi = -\Delta_{\mathbf{y}} \Phi = \delta_{\mathbf{x}}$   $\Phi$  harmonic except unit point-mass at  $\mathbf{0}$

notation:  $\mathbf{n}$  points outwards,  $\|\mathbf{n}\| = 1$ ,  $u_n := \mathbf{n} \cdot \nabla u$



Green's 2nd identity:  $\int_{\Gamma} v u_n - v_n u \, ds = \int_{\Omega} v \Delta u - (\Delta v) u \, d\mathbf{y}$

calculus

# Laplace fundamental solution in $\mathbb{R}^2$

Eg PDE: Poisson eqn  $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$  Laplacian

notation:  $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$  is a point. This frees up  $\mathbf{y} \in \mathbb{R}^2$  as another point (not y-coord!)

not well-posed unless add BC! BIEs are good for *homogeneous* PDEs (driving  $g \equiv 0$ )

A well-posed\* BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE (} u \text{ harmonic)}$$

$$u = f \text{ on } \Gamma \quad \text{Dirichlet BC}$$


\*exists, unique,  
continuous w.r.t. data



Laplace fundamental soln:  $\Phi(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi} \log \frac{1}{r}$  where  $r := \|\mathbf{x} - \mathbf{y}\|$

obeys  $-\Delta_{\mathbf{x}} \Phi = -\Delta_{\mathbf{y}} \Phi = \delta_{\mathbf{x}}$   $\Phi$  harmonic except unit point-mass at  $\mathbf{0}$

notation:  $\mathbf{n}$  points outwards,  $\|\mathbf{n}\| = 1$ ,  $u_n := \mathbf{n} \cdot \nabla u$



Green's 2nd identity:  $\int_{\Gamma} v u_n - v_n u \, ds = \int_{\Omega} v \Delta u - (\Delta v) u \, d\mathbf{y}$

calculus

warm-up: set  $u =$  BVP soln,  $v \equiv 1$ , G2I becomes  $\int_{\Gamma} u_n \, ds - 0 = 0 - 0$ : so  $u$  has zero flux



# Laplace fundamental solution in $\mathbb{R}^2$

Eg PDE: Poisson eqn  $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$  Laplacian

notation:  $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$  is a point. This frees up  $\mathbf{y} \in \mathbb{R}^2$  as another point (not y-coord!)

not well-posed unless add BC! BIEs are good for *homogeneous* PDEs (driving  $g \equiv 0$ )

A well-posed\* BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE (} u \text{ harmonic)}$$

$$u = f \text{ on } \Gamma \quad \text{Dirichlet BC}$$


\*exists, unique,  
continuous w.r.t. data



Laplace fundamental soln:  $\Phi(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi} \log \frac{1}{r}$  where  $r := \|\mathbf{x} - \mathbf{y}\|$

obeys  $-\Delta_{\mathbf{x}} \Phi = -\Delta_{\mathbf{y}} \Phi = \delta_{\mathbf{x}} = \delta_{\mathbf{y}}$   $\Phi$  harmonic except unit point-mass at 0

notation:  $\mathbf{n}$  points outwards,  $\|\mathbf{n}\| = 1$ ,  $u_n := \mathbf{n} \cdot \nabla u$



Green's 2nd identity:  $\int_{\Gamma} v u_n - v_n u \, ds = \int_{\Omega} v \Delta u - (\Delta v) u \, dy$

calculus

warm-up: set  $u = \text{BVP soln}$ ,  $v \equiv 1$ , G2I becomes  $\int_{\Gamma} u_n \, ds - 0 = 0 - 0$ : so  $u$  has zero flux


Now some fun: fix "target"  $\mathbf{x} \in \Omega$ , let  $v = \Phi(\mathbf{x}, \cdot)$ , G2I gives:

Green's representation formula:

$$\int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) u_n(\mathbf{y}) - \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial n_{\mathbf{y}}} u(\mathbf{y}) \, ds_{\mathbf{y}} = u(\mathbf{x}) \quad \text{for } \mathbf{x} \in \Omega$$

recovers soln from "Cauchy data"  $(u, u_n)|_{\Gamma}$

also versions for Helmholtz, Stokes, Maxwell,...





## Layer potentials and their jump relations

Representations of harmonic functions off a curve  $\Gamma$ : “density”  $\sigma$

Single-layer potential  $(\mathcal{S}\sigma)(\mathbf{x}) := \int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}}$  charge sheet

Double-layer potential  $(\mathcal{D}\sigma)(\mathbf{x}) := \int_{\Gamma} \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}_{\mathbf{y}}} \sigma(\mathbf{y}) ds_{\mathbf{y}}$  dipole sheet



# Layer potentials and their jump relations

Representations of harmonic functions off a curve  $\Gamma$ : “density”  $\sigma$

Single-layer potential  $(\mathcal{S}\sigma)(\mathbf{x}) := \int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$  charge sheet

Double-layer potential  $(\mathcal{D}\sigma)(\mathbf{x}) := \int_{\Gamma} \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}_{\mathbf{y}}} \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$  dipole sheet

interior (-) / exterior (+) limits:

$$u^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

$$u_n^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} \mathbf{n}_{\mathbf{x}} \cdot \nabla u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

Jump relations:



$(\mathcal{S}\sigma)^{\pm} = S\sigma$   $S$  (Roman font) means *restriction* of  $S$  to  $\Gamma$ : a bdry int. op.

$(\mathcal{D}\sigma)^{\pm} = (D \pm I/2)\sigma$  jump in potential equal to  $\sigma$ ;  $D$  restriction to  $\Gamma$  in P.V. sense

$(\mathcal{S}\sigma)_n^{\pm} = (D^T \mp I/2)\sigma$  jump in normal derivative

$(\mathcal{D}\sigma)_n^{\pm} = T\sigma$   $T$  hypersingular, kernel  $\partial^2 \Phi(\mathbf{x}, \mathbf{y}) / \partial \mathbf{n}_{\mathbf{x}} \partial \mathbf{n}_{\mathbf{y}} \sim 1/r^2$

- $D$  smooth kernel on smooth  $\Gamma$ , while  $S$  always log (weakly) singular





# Layer potentials and their jump relations

Representations of harmonic functions off a curve  $\Gamma$ : “density”  $\sigma$

Single-layer potential  $(\mathcal{S}\sigma)(\mathbf{x}) := \int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$  charge sheet

Double-layer potential  $(\mathcal{D}\sigma)(\mathbf{x}) := \int_{\Gamma} \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}_{\mathbf{y}}} \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$  dipole sheet



interior (-) / exterior (+) limits:

$$u^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

$$u_n^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} \mathbf{n}_{\mathbf{x}} \cdot \nabla u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

Jump relations:

$(\mathcal{S}\sigma)^{\pm} = S\sigma$   $S$  (Roman font) means *restriction* of  $S$  to  $\Gamma$ : a bdry int. op.

$(\mathcal{D}\sigma)^{\pm} = (D \pm I/2)\sigma$  jump in potential equal to  $\sigma$ ;  $D$  restriction to  $\Gamma$  in P.V. sense

$(\mathcal{S}\sigma)_n^{\pm} = (D^T \mp I/2)\sigma$  jump in normal derivative

$(\mathcal{D}\sigma)_n^{\pm} = T\sigma$   $T$  hypersingular, kernel  $\partial^2 \Phi(\mathbf{x}, \mathbf{y}) / \partial \mathbf{n}_{\mathbf{x}} \partial \mathbf{n}_{\mathbf{y}} \sim 1/r^2$

- $D$  smooth kernel on smooth  $\Gamma$ , while  $S$  always log (weakly) singular

Recap GRF in LP notation:  $u$  harmonic in  $\Omega \Rightarrow \mathcal{S}u_n^- - \mathcal{D}u^- = u$  in  $\Omega$

## Converting BVP to BIE and solving

Say wish to solve interior

Dirichlet Laplace BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form?

## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form? Fred. Alt.: BIE has soln  $\forall f$ ! BVP & BIE equivalent ☺

(had we picked  $u = \mathcal{S}\sigma$ , would get 1st kind, poorly conditioned but can have its uses)



## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form? Fred. Alt.: BIE has soln  $\forall f$ ! BVP & BIE equivalent ☺

(had we picked  $u = \mathcal{S}\sigma$ , would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on  $\Gamma$  using arc-length measure  $ds_\gamma$ . Usually not how  $\Gamma$  described...


## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form? Fred. Alt.: BIE has soln  $\forall f$ ! BVP & BIE equivalent ☺

(had we picked  $u = \mathcal{S}\sigma$ , would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on  $\Gamma$  using arc-length measure  $ds_y$ . Usually not how  $\Gamma$  described...

**Parameterize** the bdry  $\mathbf{y}(t)$   $\mathbf{y} : \mathbb{R} \rightarrow \mathbb{R}^2$ ,  $2\pi$ -periodic,  $\Gamma = \{\mathbf{y}(t) : t \in [0, 2\pi)\}$

change variable  $ds_y = \|\mathbf{y}'(t)\|dt$  abuse notation  $\sigma(t) = \sigma(\mathbf{y}(t))$


## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form? Fred. Alt.: BIE has soln  $\forall f$ ! BVP & BIE equivalent ☺

(had we picked  $u = \mathcal{S}\sigma$ , would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on  $\Gamma$  using arc-length measure  $ds_y$ . Usually not how  $\Gamma$  described...

**Parameterize** the bdry  $\mathbf{y}(t)$   $\mathbf{y} : \mathbb{R} \rightarrow \mathbb{R}^2$ ,  $2\pi$ -periodic,  $\Gamma = \{\mathbf{y}(t) : t \in [0, 2\pi)\}$

change variable  $ds_y = \|\mathbf{y}'(t)\|dt$  abuse notation  $\sigma(t) = \sigma(\mathbf{y}(t))$

Get 1D IE:  $\sigma(t) - 2 \int_0^{2\pi} \frac{\partial \Phi(\mathbf{y}(t), \mathbf{y}(s))}{\partial \mathbf{n}_{\mathbf{y}(s)}} \sigma(s) \|\mathbf{y}'(s)\| ds = -2f(t), \quad t \in [0, 2\pi)$

familiar form  $(I + K)\sigma = -2f$ , with kernel  $k(s, t) = \frac{-2}{2\pi} \frac{\mathbf{n}_{\mathbf{y}(s)} \cdot (\mathbf{y}(t) - \mathbf{y}(s))}{\|\mathbf{y}(t) - \mathbf{y}(s)\|^2} \|\mathbf{y}'(s)\|$

formula on diagonal:  $k(t, t) = \lim_{s \rightarrow t} k(t, s) = \kappa(t)/2\pi$ ,  $\kappa$  curvature of  $\Gamma$  (check!)


## Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**:  $u = \mathcal{D}\sigma$ , look up its **JR** for BC:  $u^- = (D - I/2)\sigma$

Insert the BC to get BIE:  $(I - 2D)\sigma = -2f$  scaled to 2nd kind form

This shows: let  $\sigma$  solve BIE, then  $u = \mathcal{D}\sigma$  solves BVP (i.e., no spurious solns)

But how know *all* solns  $u$  of this form? Fred. Alt.: BIE has soln  $\forall f$ ! BVP & BIE equivalent ☺

(had we picked  $u = \mathcal{S}\sigma$ , would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on  $\Gamma$  using arc-length measure  $ds_y$ . Usually not how  $\Gamma$  described...

**Parameterize** the bdry  $\mathbf{y}(t)$   $\mathbf{y} : \mathbb{R} \rightarrow \mathbb{R}^2$ ,  $2\pi$ -periodic,  $\Gamma = \{\mathbf{y}(t) : t \in [0, 2\pi)\}$

change variable  $ds_y = \|\mathbf{y}'(t)\|dt$  abuse notation  $\sigma(t) = \sigma(\mathbf{y}(t))$

Get 1D IE:  $\sigma(t) - 2 \int_0^{2\pi} \frac{\partial \Phi(\mathbf{y}(t), \mathbf{y}(s))}{\partial \mathbf{n}_{\mathbf{y}(s)}} \sigma(s) \|\mathbf{y}'(s)\| ds = -2f(t), \quad t \in [0, 2\pi)$

familiar form  $(I + K)\sigma = -2f$ , with kernel  $k(s, t) = \frac{-2}{2\pi} \frac{\mathbf{n}_{\mathbf{y}(s)} \cdot (\mathbf{y}(t) - \mathbf{y}(s))}{\|\mathbf{y}(t) - \mathbf{y}(s)\|^2} \|\mathbf{y}'(s)\|$

formula on diagonal:  $k(t, t) = \lim_{s \rightarrow t} k(t, s) = \kappa(t)/2\pi$ ,  $\kappa$  curvature of  $\Gamma$  (check!)

Now Nyström discretize with PTR, solve lin. sys. for  $\boldsymbol{\sigma} := \{\sigma_j\}_{j=1}^N$

Finally evaluate soln:  $u(\mathbf{x}) = (\mathcal{D}\sigma)(\mathbf{x}) \stackrel{\text{PTR}}{\approx} \sum_{j=1}^N \frac{\mathbf{n}_{\mathbf{y}(t_j)} \cdot (\mathbf{x} - \mathbf{y}(t_j))}{2\pi \|\mathbf{x} - \mathbf{y}(t_j)\|^2} \|\mathbf{y}'(t_j)\| w_j \sigma_j$

# Interior Laplace Dirichlet BVP solve demo

demo\_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```

*% shape params (note a=1,b=0 unit circle)*  
*% kite parameterization y(t)*  
*% y', analytic*  
*% y'', analytic*  
  
*% PTR nodes & weights*  
*% bdry nodes, 2-by-N*  
*% bdry normals*  
*% bdry curvatures*  
*% matrix of r=x-y (two vec cmpnts)*  
*% off-diag (-1/pi) n.r/r^2*  
*% overwrite diag elements*  
*% note Id gets no "speed weights"*  
*% test u(x) = x\_1(x\_2-0.3), not symmetric!*  
*% read off its Dirichlet data*  
  
*% solve. Leave u = D.sigma eval to reader*

# Interior Laplace Dirichlet BVP solve demo


demo\_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```

```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic

% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data

% solve. Leave u = D.sigma eval to reader
```



# Interior Laplace Dirichlet BVP solve demo


demo\_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```


```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic
```

```
% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data
```

```
% solve. Leave u = D.sigma eval to reader
```



$\log_{10} |u - u_{ex}|$ : naive PTR quadr. eval.



error: "5h" rule.  
Note: special  
quadratures  
can fix near  $\Gamma$   
(Helsing, QBX...)

# Interior Laplace Dirichlet BVP solve demo


demo\_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```


```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic

% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data

% solve. Leave u = D.sigma eval to reader
```



log<sub>10</sub> |u - u<sub>ex</sub>|: naive PTR quadr. eval.



error: "5h" rule.

Note: special quadratures can fix near  $\Gamma$  (Helsing, QBX...)

Debug:  $\sigma \equiv -1 \Rightarrow u \equiv 1$ , then test data from (generic!) soln  $u$ , and...

- ① check/plot  $n, \kappa$ . First test unit circle!
- ② check Nyström matrix smooth at diag (before add I)



## Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg  $u = \mathcal{D}\sigma$ ), get BIE from JRs

## Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg  $u = \mathcal{D}\sigma$ ), get BIE from JRs

Alternative is “direct”: take limit of GRF on  $\Gamma$ , rearrange to get BIE:

$$\text{GRF } u = Su^- - \mathcal{D}u_n^- \xrightarrow{\text{JR}_s} u_n^- = (D^T + I/2)u_n^- - Tu^- \xrightarrow{\text{BC}} (D^T - I/2)u_n^- = Tf$$

Needs hypersingular apply ☹. Then solve BIE for  $u_n^-$ , eval  $u$  via GRF (needs two LP evals)

## Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg  $u = \mathcal{D}\sigma$ ), get BIE from JRs

Alternative is “direct”: take limit of GRF on  $\Gamma$ , rearrange to get BIE:

$$\text{GRF } u = Su^- - \mathcal{D}u_n^- \xrightarrow{\text{JR}_s} u_n^- = (D^T + I/2)u_n^- - Tu^- \xrightarrow{\text{BC}} (D^T - I/2)u_n^- = Tf$$

Needs hypersingular apply  $\odot$ . Then solve BIE for  $u_n^-$ , eval  $u$  via GRF (needs two LP evals)

Notice BIO  $(D^T - I/2)$  **adjoint** of that for indirect  $(D - I/2)$

generally true. So, spectra the same, thus iterative convergence rates too

## Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg  $u = \mathcal{D}\sigma$ ), get BIE from JRs

Alternative is “direct”: take limit of GRF on  $\Gamma$ , rearrange to get BIE:

$$\text{GRF } u = \mathcal{S}u^- - \mathcal{D}u_n^- \xrightarrow{\text{JR}_s} u_n^- = (D^T + I/2)u_n^- - Tu^- \xrightarrow{\text{BC}} (D^T - I/2)u_n^- = Tf$$

Needs hypersingular apply  $\ominus$ . Then solve BIE for  $u_n^-$ , eval  $u$  via GRF (needs two LP evals)

Notice BIO  $(D^T - I/2)$  **adjoint** of that for indirect  $(D - I/2)$

generally true. So, spectra the same, thus iterative convergence rates too

Indirect BIE	Direct BIE
unknown density (unphysical)	unknown is physical
RHS is plain data	RHS needs BIO apply to data
eval the representation (may be simpler)	eval the GRF

- indirect: more flexibility, but need math to prove equivalence to BVP
- accuracy differences for domains with corners (Hoskins–Rachh...)

## Indirect 2nd-kind BIE for Neumann, exterior

recap: Laplace int. Dir.

$$\Delta u = 0 \text{ in } \Omega$$

$$u^- = f \text{ on } \Gamma$$

uniqueness, existence  $\forall f$

- $u = \mathcal{D}\sigma$       rep.  
 $(D - I/2)\sigma = f$       BIE: well-cond.

Laplace int. Neu.

$$\Delta u = 0 \text{ in } \Omega$$

$$u_n^- = g \text{ on } \Gamma$$

$$\text{require } \int_{\Gamma} g ds = 0$$

unique only up to a const.

- $u = \mathcal{S}\sigma$        $\swarrow$  kernel  $\equiv 1$ , kills nullspace  
 $(D^T + I/2 + 11^T)\sigma = g$       well-cond.

## Indirect 2nd-kind BIE for Neumann, exterior

recap: Laplace int. Dir.

$$\Delta u = 0 \text{ in } \Omega$$

$$u^- = f \text{ on } \Gamma$$

uniqueness, existence  $\forall f$

- $u = \mathcal{D}\sigma$  rep.  
 $(D - I/2)\sigma = f$  BIE: well-cond.

Laplace int. Neu.

$$\Delta u = 0 \text{ in } \Omega$$

$$u_n^- = g \text{ on } \Gamma$$

$$\text{require } \int_{\Gamma} g ds = 0$$

unique only up to a const.

- $u = \mathcal{S}\sigma$   $\checkmark$  kernel  $\equiv 1$ , kills nullspace  
 $(D^T + I/2 + 11^T)\sigma = g$  well-cond.

Laplace ext. Dir.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u^+ = f \text{ on } \Gamma$$

$$u_{\infty} := \lim_{\|x\| \rightarrow \infty} u(x) \text{ exists}$$

uniqueness, existence  $\forall f$

- $u = \mathcal{D}\sigma + \int_{\Gamma} \sigma ds$  modified rep.  
 $(D + I/2 + 11^T)\sigma = f$  well-cond.

Laplace ext. Neu.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u_n^+ = g \text{ on } \Gamma$$

$$\text{require } \int_{\Gamma} g ds = 0 \text{ and } u_{\infty} = 0$$

unique

- $u = \mathcal{S}\sigma$   
 $(D^T - I/2)\sigma = g$  well-cond.

# Indirect 2nd-kind BIE for Neumann, exterior

recap: Laplace int. Dir.

$$\Delta u = 0 \text{ in } \Omega$$

$$u^- = f \text{ on } \Gamma$$

uniqueness, existence  $\forall f$

•  $u = \mathcal{D}\sigma$  rep.

$(D - I/2)\sigma = f$  BIE: well-cond.

Laplace int. Neu.

$$\Delta u = 0 \text{ in } \Omega$$

$$u_n^- = g \text{ on } \Gamma$$

require  $\int_{\Gamma} g ds = 0$

unique only up to a const.

•  $u = \mathcal{S}\sigma$

$(D^T + I/2 + 11^T)\sigma = g$  kernel  $\equiv 1$ , kills nullspace  
well-cond.

Laplace ext. Dir.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u^+ = f \text{ on } \Gamma$$

$$u_{\infty} := \lim_{\|x\| \rightarrow \infty} u(x)$$

uniqueness, existence  $\forall f$

•  $u = \mathcal{D}\sigma + \int_{\Gamma} \sigma ds$  modified rep.

$(D + I/2 + 11^T)\sigma = f$  well-cond.

Laplace ext. Neu.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u_n^+ = g \text{ on } \Gamma$$

require  $\int_{\Gamma} g ds = 0$  and  $u_{\infty} = 0$

unique

•  $u = \mathcal{S}\sigma$

$(D^T - I/2)\sigma = g$  well-cond.

complementary BVPS

# Indirect 2nd-kind BIE for Neumann, exterior

recap: Laplace int. Dir.

$$\Delta u = 0 \text{ in } \Omega$$

$$u^- = f \text{ on } \Gamma$$

uniqueness, existence  $\forall f$

•  $u = \mathcal{D}\sigma$  rep.

$(D - I/2)\sigma = f$  BIE: well-conv.

Laplace int. Neu.

$$\Delta u = 0 \text{ in } \Omega$$

$$u_n^- = g \text{ on } \Gamma$$

require  $\int_{\Gamma} g ds = 0$

unique only up to a const.

•  $u = \mathcal{S}\sigma$

$(D^T + I/2 + 11^T)\sigma = g$  kernel  $\equiv 1$ , kills nullspace  
well-conv.

Laplace ext. Dir.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u^+ = f \text{ on } \Gamma$$

$$u_{\infty} := \lim_{\|x\| \rightarrow \infty} u(x) \text{ exists}$$

uniqueness, existence  $\forall f$

•  $u = \mathcal{D}\sigma + \int_{\Gamma} \sigma ds$  modified rep.

$(D + I/2 + 11^T)\sigma = f$  well-conv.

Laplace ext. Neu.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u_n^+ = 0 \text{ on } \Gamma$$

require  $\int_{\Gamma} g ds = 0$  and  $u_{\infty} = 0$

unique

•  $u = \mathcal{S}\sigma$

$(D^T - I/2)\sigma = g$  well-conv.

complementary BVPs



# Indirect 2nd-kind BIE for Neumann, exterior

recap: Laplace int. Dir.

$$\Delta u = 0 \text{ in } \Omega$$

$$u^- = f \text{ on } \Gamma$$

uniqueness, existence  $\forall f$

- $u = \mathcal{D}\sigma$  rep.

$$(D - I/2)\sigma = f \quad \text{BIE: well-conv.}$$

Laplace int. Neu.

$$\Delta u = 0 \text{ in } \Omega$$

$$u_n^- = g \text{ on } \Gamma$$

requires  $\int_{\Gamma} g ds = 0$

unique only up to a const.

- $u = \mathcal{S}\sigma$

$$(D^T + I/2 + 11^T)\sigma = g \quad \text{well-conv.} \quad \checkmark \text{kernel} \equiv 1, \text{ kills nullspace}$$

Laplace ext. Dir.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u^+ = f \text{ on } \Gamma$$

$$u_{\infty} := \lim_{\|x\| \rightarrow \infty} u(x) \text{ exists}$$

uniqueness, existence  $\forall f$

- $u = \mathcal{D}\sigma + \int_{\Gamma} \sigma ds$  modified rep.

$$(D + I/2 + 11^T)\sigma = f \quad \text{well-conv.}$$

Laplace ext. Neu.

$$\Delta u = 0 \text{ in } \mathbb{R}^2 \setminus \overline{\Omega}$$

$$u_n^+ = 0 \text{ on } \Gamma$$

requires  $\int_{\Gamma} g ds = 0$  and  $u_{\infty} = 0$

unique

- $u = \mathcal{S}\sigma$

$$(D^T - I/2)\sigma = g \quad \text{well-conv.}$$

complementary BVPs

③ Exterior: don't test with  $u = \log r!$  why not? BVPs enforce zero net charge

# Helmholtz — introduction and connection to Maxwell


$$(\Delta + \omega^2)u = 0$$

time-harmonic scalar waves

comes from scalar wave equation  $\Delta u - u_{tt} = 0$  when  $u(\mathbf{x}, t) = u(\mathbf{x})e^{-i\omega t}$   
 $\omega$  is the wavenumber spatial frequency, related to wavelength via  $\lambda = 2\pi/\omega$

Also used for Maxwell's equations in cylindrical symm (z-invariance):


1. Assume  $\mathbf{E}, \mathbf{H}(x, y, z) = \mathbf{E}, \mathbf{H}(x, y)$
2. Write Maxwell's eqs:  $\nabla \times \mathbf{E} = i\omega\mu\mathbf{H}$ ,  $\nabla \times \mathbf{H} = -i\omega\varepsilon\mathbf{E}$ ,
3. Notice only  $E_z, H_z$  are indep  $\rightarrow$  2 polarizations, TE or TM:  $E_z = 0, H_z = 0$  resp.
4. Choose TE and let  $u := H_z$ , then:  $\mathbf{H} = (0, 0, u)$ ,  
 $\mathbf{E} = \frac{1}{i\omega\varepsilon}(\partial_x u, -\partial_y u, 0)$ , and  $(\Delta + n^2\omega^2)u = 0$  with  $n^2 = \varepsilon\mu$



Dirichlet BC in TE formalism = PEC perfect electric conductor;  $\mathbf{E} \perp$  to surface

# Helmholtz — scattering formalism

Split total potential into incident (known) and scattered (unknown) parts,  $u^{\text{tot}} = u^{\text{inc}} + u$



BVP for  $u$ :

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^d \setminus \overline{\Omega} \quad \text{PDE}$$

$$u = -u_i \quad \text{on } \Gamma \quad \text{Dirichlet BC, } u_n = -(u_i)_n \text{ for Neumann}$$


$$\lim_{r \rightarrow \infty} \left( \frac{\partial u}{\partial r} - iku \right) = 0 \quad r := |\mathbf{x} - \mathbf{y}|, \text{ Sommerfeld radiation condition for uniqueness}$$

Fundamental solution  $\Phi(\mathbf{x}, \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$


$$\text{Asymptotics: } \lim_{r \rightarrow 0} \Phi(r) = \frac{1}{2\pi} \log \frac{1}{r} + \mathcal{O}(1)$$

$$\lim_{r \rightarrow \infty} \Phi(r) = \sqrt{\frac{2}{\pi r}} e^{i(r - \nu\pi/2 - \pi/4)} + \mathcal{O}(r^{-1})$$


Same singularity as Laplace  $\rightarrow$  same JRs!



Layer potentials



SLP



DLP

# Helmholtz — interior resonances and how to avoid them

Try the ext Dir BVP with  $u = \mathcal{D}\sigma$  ( $(\Delta + \omega)^2 u = 0$  in  $\mathbb{R}^2 \setminus \overline{\Omega}$ ,  $u = -u_i$  on  $\Gamma$ , SRC for  $u$ )

Observe that for some  $\omega$ , condition # of BIE blows up, not always solvable

Why? Suppose  $\phi \not\equiv 0$  s.t. 
$$\begin{cases} (\Delta + \omega^2)\phi = 0 & \text{in } \Omega \\ \phi_n = 0 & \text{on } \Gamma \end{cases}$$
  $\phi$  is interior Neumann eigenfunction with eigenvalue  $\omega^2$

Then by (interior) GRF (same as for Laplace),  $\mathcal{S}\phi|_{\Gamma} - \mathcal{D}\phi|_{\Gamma} = u$  in  $\Omega$ .

Take  $\mathbf{x} \rightarrow \Gamma^-$  and use JR:  $(-D - I/2)\phi|_{\Gamma} = \phi_{\Gamma}$ , i.e.  $(I + 2D)\phi|_{\Gamma} = 0$ .

Since  $\phi|_{\Gamma}$  was nontrivial (otherwise  $\phi = 0$  by GRF), nullity of  $I + 2D > 0$ , i.e. singular, by FA not solvable  $\forall f(u_i)$ .

We made use of the **complementary BVP** (int Neu), this is an “internal resonance”.

Fix:  $u = (\mathcal{D} - i\eta\mathcal{S})\sigma$  combined field integral eq (CFIE), same # unknowns, new kernel  
ext Dir BIE becomes  $(I + 2D - 2i\eta\mathcal{S})\sigma = -2u_i$  on  $\Gamma$

Proof: Let  $\tau$  solve  $(I/2 + D - i\eta\mathcal{S})\tau = 0$ , wish to show  $\tau = 0$ .

From  $\tau$  construct potential  $v := (\mathcal{D} - i\eta\mathcal{S})\tau$ , then  $v^+ = 0$  by construction.

Then  $v = 0$  in  $\mathbb{R}^2 \setminus \overline{\Omega}$  by uniqueness of the complementary BVP (ext Dir)

Then  $v_n^+$  on  $\Gamma$ , and by JRs and Green's 1st thm (exercise for the reader ☺),  $\tau = 0$ .

## Helmholtz — Dirichlet demo

Dirichlet demo, plots only: Solve BVP for  $u$  via PTR + Nyström, with new diag limit for  $k(t, t)$ , show  $1/N^3$  convergence if use naive PTR with correct diag limit (see M126 HW?)

④ debug BVP with known data from a radiative soln [sources inside  \$\Omega\$](#)   
(don't demo CFIE since requires  $S$  w/ log-singularity).

## Helmholtz transmission BVP

refractive indices in  $\Omega$  vs exterior

Matching? (more effort, needs  $\mathcal{S}\sigma + \mathcal{D}\tau \dots$ )

difference kernels at most log-singular.

# Helmholtz

Getting spectral-acc Nyström for log-singular kernels: beyond today.

eg kernel-split or product quadratures (Kress, Helsing, . . . )

close-eval: kernel-split, QBX, etc.

see libraries: chunkie, BIE2D, etc

## More debug ideas

TO DISCARD

Other tests:

- ⑤ Test SLP & DLP evaluators via GRF for any harmonic  $u$  in  $\Omega$



# Summary

Covered BIE basics for smooth curves with global quadrature:

- Well-posed Laplace & Helmholtz BVPs exterior need condition as  $\|\mathbf{x}\| \rightarrow \infty$
- Choosing representation to get 2nd kind BIE if poss., equivalent to BVP if poss.  
Can switch interior/exterior, Laplace/Helmholtz/etc, via simple code changes
- Nyström discretization high-order/spectral convergence, if poss.
- Build/debug codes via well-chosen sequence of test cases also for libraries

practise! write out theory yourself + try HW exer. in repo + run demos

# Summary

Covered BIE basics for smooth curves with global quadrature:

- Well-posed Laplace & Helmholtz BVPs exterior need condition as  $\|x\| \rightarrow \infty$
- Choosing representation to get 2nd kind BIE if poss., equivalent to BVP if poss.  
Can switch interior/exterior, Laplace/Helmholtz/etc, via simple code changes
- Nyström discretization high-order/spectral convergence, if poss.
- Build/debug codes via well-chosen sequence of test cases also for libraries

practise! write out theory yourself + try HW exer. in repo + run demos

Useful 2D tools we did not yet cover: in libraries, eg chunkie, BIE2D

- panel (composite) quadratures essential for adaptivity
- high-order quadratures for log-singular kernel SLP, Helmholtz, etc
- special quadratures for evaluation close to the curve  
some need interpolation of  $\sigma(t)$  off the nodes  $t_j$ , some not
- corners, open arcs, slits, multi-material junctions

## Resources

Many numerical analysis (mathematics heavy). Somewhat accessible:

- *Linear Integral Equations*, R. Kress, (1999, Springer). Ch. 6 & 12.
- *The Numerical Solution of Integral Equations of the Second Kind*, K. E. Atkinson, (1997, CUP).

Fewer on the practical/tutorial side, few with modern devels:

- “High-order accurate methods for Nyström discretization of integral equations on smooth curves in the plane”, S Hao, AH Barnett, PG Martinsson, P Young. *Adv. Comput. Math.* **40**, 245–272 (2014).

focuses on quadrature for logarithmic singularities, eg SLP, Helmholtz

- <https://users.flatironinstitute.org/~ahb/BIE/>
- <https://github.com/ahbarnett/BIEbook>

in progress...