

2D boundary integral equations and the Nyström method

Alex Barnett¹ and **Fruzsina Agocs¹**

Computational Tools 2024 BIE workshop. Day 1, 6/10/24

¹Center for Computational Mathematics, Flatiron Institute, Simons Foundation

Integral equations on 1D interval

- Given: i) function $\sigma(t)$ defined on interval $[0, 2\pi)$, periodic: $\sigma(2\pi) = \sigma(0)$, etc
 ii) “kernel” function $k(t, s)$ defined on square $[0, 2\pi)^2$,


Integral operator K acts on σ to give another function $K\sigma$:

$$(K\sigma)(t) := \int_0^{2\pi} k(t, s)\sigma(s)ds, \quad t \in [0, 2\pi)$$

continuous analog of
matrix-vector prod. Ax

Integral equation: $(I + K)\sigma = f$, ie

$$\sigma(t) + \int_0^{2\pi} k(t, s)\sigma(s)ds = f(t), \quad t \in [0, 2\pi)$$




analog of lin. sys. $Ax = b$

Fredholm “second kind” (due to presence of I , otherwise called “first kind”)

If kernel continuous, or “weakly” singular (integrable), K is *compact*:

- eigenvalues $(K\phi_k = \lambda_k\phi_k)$ discrete, with $\lim_{k \rightarrow \infty} \lambda_k = 0$
unless some $\lambda_k = -1$, 2nd kind IE has at most one soln: $\text{Nul}(I + K) = \{0\}$
- $\text{Nul}(I + K) = \{0\} \Rightarrow$ existence of solution for *any* f Fredholm Alternative
like square matrix (finite-dim), recall: uniqueness \Rightarrow consistent for any RHS

Contrast 1st kind IE $K\sigma = f$ is ill-posed problem (unstable)!  **FLATIRON**
INSTITUTE

See references for lots of beautiful theory, precise statements

Nyström discretization of 2nd kind IE on interval

Simplest quadrature for periodic funcs: periodic trapezoid rule (PTR)

$$\int_0^{2\pi} f(t) dt \approx \sum_{j=1}^N \frac{2\pi}{N} f\left(\frac{2\pi j}{N}\right) = \sum_{j=1}^N w_j f(t_j) \quad w_j = \text{weights}, \quad t_j = \text{nodes}$$

For f smooth, superalgebraically convergent ("spectral"): error = $\mathcal{O}(N^{-p})$ for any p

Apply quad to integral in 2nd kind IE:

call the resulting approx soln $\tilde{\sigma}$

$$\tilde{\sigma}(t) + \sum_{j=1}^N k(t, t_j) w_j \tilde{\sigma}(t_j) = f(t), \quad t \in [0, 2\pi) \quad (*)$$

Holds for all t ; in particular, holds at each t_i , $i = 1, \dots, N$, giving:

$$\sigma_i + \sum_{j=1}^N k(t_i, t_j) w_j \sigma_j = f(t_i), \quad i = 1, \dots, N \quad \text{where } \sigma_j := \tilde{\sigma}(t_j)$$

Write as: $A\sigma = f$ $N \times N$ lin sys, entries $a_{ij} = \delta_{ij} + k(t_i, t_j) w_j$, and $f_j := f(t_j)$


solve? dense direct $\mathcal{O}(N^3)$; dense iter. $\mathcal{O}(N^2)$; fast iter. $\approx \mathcal{O}(N)$; fast direct $\approx \mathcal{O}(N^{(d+1)/2})$

Why 2nd kind? eigs(A) accumulate only at $+1$, iterative fast conv.

Sometimes for BIE (eg, far-field eval), node values $\{\sigma_j\}_{j=1}^N$ suffice.

If not, interpolate from them to any $t \in [0, 2\pi)$. Two approaches:

- either: rearrange $(*)$ to give $\tilde{\sigma}(t) = \dots$, called "Nyström interpolant" (rare)
- or (common): use local high-order Lagrange (or global spectral) interpolation:






Demo Nyström on interval (1D)

day1/code/nyst1d_demos.m

```
kfun = @(t,s) exp(3*cos(t-s));  
ffun = @(t) cos(5*t+1);  
N = 30;  
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);  
A = eye(N) + bsxfun(kfun,t',t)*diag(w);  
rhs = ffun(t');  
sigmaj = A\rhs;
```

*% smooth convolutional kernel, periodic domain [0,2pi)
% smooth data (RHS) func
% number of unknowns: should study convergence as N grows...
% PTR nodes and weights, row vecs
% identity plus fill k(t_i,t_j)w_j for i,j=1..N
% col vec
% dense direct square solve (pivoted LU), gives col vec*



“self-convergence”:
use $N=40$ as “true”

f and k smooth

⇒ σ smooth

⇒ spectral conv?




Thm. (Anselone, Kress,...): error at node values (and Nyström interpolant) same order as that of quadrature rule applied to integrand $k(t, \cdot)\sigma$.

Demo Nyström on interval (1D)

day1/code/nyst1d_demos.m

```
kfun = @(t,s) exp(3*cos(t-s));
ffun = @(t) cos(5*t+1);
N = 30;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
A = eye(N) + bsxfun(kfun,t',t)*diag(w);
rhs = ffun(t');
sigma_j = A\rhs;
```

```
% smooth convolutional kernel, periodic domain [0,2pi)
% smooth data (RHS) func
% number of unknowns: should study convergence as N grows...
% PTR nodes and weights, row vecs
% identity plus fill k(t_i,t_j)w_j for i,j=1..N
% col vec
% dense direct square solve (pivoted LU), gives col vec
```





“self-convergence”:
use $N=40$ as “true”

f and k smooth
 $\Rightarrow \sigma$ smooth
 \Rightarrow spectral conv?

Thm. (Anselone, Kress,...): error at node values (and Nyström interpolant) same order as that of quadrature rule applied to integrand $k(t, \cdot)\sigma$.

- Then, f or k nonsmooth?
worse (here *algebraic*) convergence using plain PTR rule:

Qu: why does order appear to improve at end?



Fundamental solution in \mathbb{R}^2

Eg PDE: Poisson eqn $\Delta u = g$

$\Delta := (\partial/\partial x_1)^2 + (\partial/\partial x_2)^2$ Laplacian

Notation: $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$ is a point. This frees up $\mathbf{y} \in \mathbb{R}^2$ as another point (not y-coord!)

Not well-posed prob. unless add BC! BIEs are good for *homogeneous* PDEs (driving $g \equiv 0$)

Eg well-posed* BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE (} u \text{ harmonic)}$$


$$u = f \text{ on } \Gamma \quad \text{Dirichlet BC}$$

*exists, unique, continuous

w.r.t. data



Laplace fundamental soln: $\Phi(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi} \log \frac{1}{r}$ where $r := \|\mathbf{x} - \mathbf{y}\|$



obeys $-\Delta_{\mathbf{x}} \Phi = -\Delta_{\mathbf{y}} \Phi = \delta_{\mathbf{x}}$ Φ harmonic except unit point-mass at 0

Normal \mathbf{n} points outwards, $\|\mathbf{n}\| = 1$ normal deriv. notation $u_n := \mathbf{n} \cdot \nabla u$

Green's 2nd identity: $\int_{\Gamma} v u_n - v_n u \, ds = \int_{\Omega} v \Delta u - (\Delta v) u \, d\mathbf{y}$

calculus

warm-up: set $u = \text{BVP soln}$, $v \equiv 1$, G2I becomes $\int_{\Gamma} u_n \, ds - 0 = 0 - 0$: so u has zero flux


more fun: fix "target" $\mathbf{x} \in \Omega$, let $v = \Phi(\mathbf{x}, \cdot)$, G2I gives:

Green's representation formula:

$$\int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) u_n(\mathbf{y}) - \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial n_{\mathbf{y}}} u(\mathbf{y}) \, ds_{\mathbf{y}} = u(\mathbf{x}) \quad \text{for } \mathbf{x} \in \Omega$$

Gets soln from "Cauchy data" $(u, u_n)|_{\Gamma}$

also versions for Helmholtz, Stokes, Maxwell, ...



Layer potentials and their jump relations

Representations of harmonic functions off a curve Γ : “density” σ

Single-layer potential $(\mathcal{S}\sigma)(\mathbf{x}) := \int_{\Gamma} \Phi(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$ charge sheet

Double-layer potential $(\mathcal{D}\sigma)(\mathbf{x}) := \int_{\Gamma} \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial \mathbf{n}_{\mathbf{y}}} \sigma(\mathbf{y}) d\mathbf{s}_{\mathbf{y}}$ dipole sheet

interior (-) / exterior (+) limits:

$$u^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

$$u_n^{\pm}(\mathbf{x}) := \lim_{h \rightarrow 0^+} \mathbf{n}_{\mathbf{x}} \cdot \nabla u(\mathbf{x} \pm h \mathbf{n}_{\mathbf{x}})$$

Jump relations:

$(\mathcal{S}\sigma)^{\pm} = S\sigma$ S (Roman font) means *restriction* of S to Γ : a bdry int. op.


$(\mathcal{D}\sigma)^{\pm} = (D \pm I/2)\sigma$ jump in potential equal to σ ; D restriction to Γ in P.V. sense

$(\mathcal{S}\sigma)_n^{\pm} = (D^T \mp I/2)\sigma$ jump in normal derivative

$(\mathcal{D}\sigma)_n^{\pm} = T\sigma$ T hypersingular, kernel $\partial^2 \Phi(\mathbf{x}, \mathbf{y}) / \partial \mathbf{n}_{\mathbf{x}} \partial \mathbf{n}_{\mathbf{y}} \sim 1/r^2$

- D smooth kernel on smooth Γ , while S always log (weakly) singular

Recap GRF in LP notation: u harmonic in $\Omega \Rightarrow \mathcal{S}u_n^- - \mathcal{D}u^- = u$ in Ω



Converting BVP to BIE and solving

Say wish to solve interior

Dirichlet Laplace BVP:

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form?

Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form? Fred. Alt.: BIE has soln $\forall f$! BVP & BIE equivalent ☺

(had we picked $u = \mathcal{S}\sigma$, would get 1st kind, poorly conditioned but can have its uses)

Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form? Fred. Alt.: BIE has soln $\forall f$! BVP & BIE equivalent ☺

(had we picked $u = \mathcal{S}\sigma$, would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on Γ using arc-length measure ds . Usually not how Γ described...


Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form? Fred. Alt.: BIE has soln $\forall f$! BVP & BIE equivalent ☺

(had we picked $u = \mathcal{S}\sigma$, would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on Γ using arc-length measure ds . Usually not how Γ described...

Parameterize the bdry $y(t)$

$$y : \mathbb{R} \rightarrow \mathbb{R}^2, \text{ } 2\pi\text{-periodic}, \Gamma = \{y(t) : t \in [0, 2\pi)\}$$

change variable $ds_y = \|y'(t)\|dt$ abuse notation $\sigma(t) = \sigma(y(t))$


Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form? Fred. Alt.: BIE has soln $\forall f$! BVP & BIE equivalent ☺

(had we picked $u = \mathcal{S}\sigma$, would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on Γ using arc-length measure ds . Usually not how Γ described...

Parameterize the bdry $\mathbf{y}(t)$

$$\mathbf{y} : \mathbb{R} \rightarrow \mathbb{R}^2, 2\pi\text{-periodic}, \Gamma = \{\mathbf{y}(t) : t \in [0, 2\pi)\}$$

change variable $ds_{\mathbf{y}} = \|\mathbf{y}'(t)\|dt$ abuse notation $\sigma(t) = \sigma(\mathbf{y}(t))$

Get 1D IE: $\sigma(t) - 2 \int_0^{2\pi} \frac{\partial \Phi(\mathbf{y}(t), \mathbf{y}(s))}{\partial \mathbf{n}_{\mathbf{y}(s)}} \sigma(s) \|\mathbf{y}'(s)\| ds = -2f(t), \quad t \in [0, 2\pi)$

familiar form $(I + K)\sigma = -2f$, with kernel $k(s, t) = \frac{-2}{2\pi} \frac{\mathbf{n}_{\mathbf{y}(s)} \cdot (\mathbf{y}(t) - \mathbf{y}(s))}{\|\mathbf{y}(t) - \mathbf{y}(s)\|^2} \|\mathbf{y}'(s)\|$

formula on diagonal: $k(t, t) = \lim_{s \rightarrow t} k(t, s) = \kappa(t)/2\pi$, κ curvature of Γ (check!)


Converting BVP to BIE and solving

Say wish to solve interior

$$\Delta u = 0 \text{ in } \Omega \quad \text{PDE}$$

Dirichlet Laplace BVP:

$$u^- = f \text{ on } \Gamma \quad \text{BC}$$



Pick **representation**: $u = \mathcal{D}\sigma$, look up its **JR** for BC: $u^- = (D - I/2)\sigma$

Insert the BC to get BIE: $(I - 2D)\sigma = -2f$ scaled to 2nd kind form

This shows: let σ solve BIE, then $u = \mathcal{D}\sigma$ solves BVP (i.e., no spurious solns)

But how know *all* solns u of this form? Fred. Alt.: BIE has soln $\forall f$! BVP & BIE equivalent ☺

(had we picked $u = \mathcal{S}\sigma$, would get 1st kind, poorly conditioned but can have its uses)

Above BIE expressed on Γ using arc-length measure ds . Usually not how Γ described...

Parameterize the bdry $\mathbf{y}(t)$

$$\mathbf{y} : \mathbb{R} \rightarrow \mathbb{R}^2, 2\pi\text{-periodic}, \Gamma = \{\mathbf{y}(t) : t \in [0, 2\pi)\}$$

change variable $ds_{\mathbf{y}} = \|\mathbf{y}'(t)\|dt$ abuse notation $\sigma(t) = \sigma(\mathbf{y}(t))$

Get 1D IE: $\sigma(t) - 2 \int_0^{2\pi} \frac{\partial \Phi(\mathbf{y}(t), \mathbf{y}(s))}{\partial \mathbf{n}_{\mathbf{y}(s)}} \sigma(s) \|\mathbf{y}'(s)\| ds = -2f(t), \quad t \in [0, 2\pi)$

familiar form $(I + K)\sigma = -2f$, with kernel $k(s, t) = \frac{-2}{2\pi} \frac{\mathbf{n}_{\mathbf{y}(t)} \cdot (\mathbf{y}(t) - \mathbf{y}(s))}{\|\mathbf{y}(t) - \mathbf{y}(s)\|^2} \|\mathbf{y}'(s)\|$

formula on diagonal: $k(t, t) = \lim_{s \rightarrow t} k(t, s) = \kappa(t)/2\pi$, κ curvature of Γ (check!)

Now Nyström discretize with PTR, solve lin. sys. for $\boldsymbol{\sigma} := \{\sigma_j\}_{j=1}^N$

Finally evaluate soln: $u(\mathbf{x}) = (\mathcal{D}\sigma)(\mathbf{x}) \stackrel{\text{PTR}}{\approx} \sum_{j=1}^N \frac{\mathbf{n}_{\mathbf{y}(t_j)} \cdot (\mathbf{x} - \mathbf{y}(t_j))}{2\pi \|\mathbf{x} - \mathbf{y}(t_j)\|^2} \|\mathbf{y}'(t_j)\| w_j \sigma_j$

Interior Laplace Dirichlet BVP solve demo

demo_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```

% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic

% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data

% solve. Leave u = D.sigma eval to reader

Interior Laplace Dirichlet BVP solve demo


demo_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```

```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic

% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data

% solve. Leave u = D.sigma eval to reader
```



Interior Laplace Dirichlet BVP solve demo


demo_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```


```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic
```

```
% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data
```

```
% solve. Leave u = D.sigma eval to reader
```



$\log_{10} |u - u_{ex}|$: naive PTR quadr. eval.



“5h” rule:
special eval.
quadratures
can fix near Γ
(Helsing, QBX...)

Interior Laplace Dirichlet BVP solve demo


demo_lapintdir.m

```
a=0.7; b=1.0;
Y = @(t) [a*cos(t)+b*cos(2*t); sin(t)];
Yp = @(t) [-a*sin(t)-2*b*sin(2*t); cos(t)];
Ypp = @(t) [-a*cos(t)-4*b*cos(2*t); -sin(t)];
N = 100;
t = 2*pi/N*(1:N); w = 2*pi/N*ones(1,N);
y = Y(t);
n = [0 1;-1 0]*Yp(t); speed = sqrt(sum(n.^2,1)); n = n./speed;
kappa = -sum(Ypp(t) .* n,1)./speed.^2;
r1 = y(1,:)'-y(1,:); r2 = y(2,:)'-y(2,:);
A = (-1/pi)*(n(1,:).*r1 + n(2,:).*r2) ./ (r1.^2+r2.^2);
A(diagind(A)) = kappa/(2*pi);
A = eye(N) + A*diag(speed.*w);
uex = @(x) ([1 0]*x) .* ([0 1]*x-0.3);
f = @(t) uex(Y(t));
rhs = -2*f(t)';
sigma = A\rhs;
```


```
% shape params (note a=1,b=0 unit circle)
% kite parameterization y(t)
% y', analytic
% y'', analytic
```

```
% PTR nodes & weights
% bdry nodes, 2-by-N
% bdry normals
% bdry curvatures
% matrix of r=x-y (two vec cmpnts)
% off-diag (-1/pi) n.r/r^2
% overwrite diag elements
% note Id gets no "speed weights"
% test u(x) = x_1(x_2-0.3), not symmetric!
% read off its Dirichlet data
```

```
% solve. Leave u = D.sigma eval to reader
```



$\log_{10} |u - u_{ex}|$: naive PTR quadr. eval.



"5h" rule:
special eval.
quadratures
can fix near Γ
(Helsing, QBX...)

Debug: $\sigma \equiv -1 \Rightarrow u \equiv 1$, then test data from (generic!) soln u , and...

- ① check/plot n, κ . First test unit circle!
- ② check Nyström matrix smooth at diag (before add I)

Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg $u = \mathcal{D}\sigma$), get BIE from JRs

Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg $u = \mathcal{D}\sigma$), get BIE from JRs

Alternative is “direct”: take limit of GRF on Γ , rearrange to get BIE:

$$\text{GRF } u = Su^- - \mathcal{D}u_n^- \xrightarrow{\text{JR}_s} u_n^- = (D^T + I/2)u_n^- - Tu^- \xrightarrow{\text{BC}} (D^T - I/2)u_n^- = Tf$$

Needs hypersingular apply \ominus . Then solve BIE for u_n^- , eval u via GRF (needs two LP evals)

Notice BIO $(D^T - I/2)$ **adjoint** of that for indirect $(D - I/2)$ general fact
spectra the same, hence iterative convergence rates

Indirect vs direct formulations

using Laplace interior Dirichlet BVP

So far “indirect” BIE: pick representation (eg $u = \mathcal{D}\sigma$), get BIE from JRs

Alternative is “direct”: take limit of GRF on Γ , rearrange to get BIE:

$$\text{GRF } u = Su^- - \mathcal{D}u_n^- \xrightarrow{\text{JRs}} u_n^- = (D^T + I/2)u_n^- - Tu^- \xrightarrow{\text{BC}} (D^T - I/2)u_n^- = Tf$$

Needs hypersingular apply \oplus . Then solve BIE for u_n^- , eval u via GRF (needs two LP evals)

Notice BIO $(D^T - I/2)$ **adjoint** of that for indirect $(D - I/2)$ general fact spectra the same, hence iterative convergence rates

indirect BIE	direct BIE
unknown density (unphysical)	unknown is physical
RHS is plain data	RHS needs BIO apply to data
eval the representation (may be simpler)	eval the GRF

- indirect: more flexibility but need math to prove equivalence to BVP
- accuracy differences for domains with corners (Hoskins–Rachh...)

Generalizations: Exterior, Neumann

some laplace aspects, confusions w/ BVP, briefly
notation for exterior domain

subtlety of decay in 2D

For int Neu or ext Dir, need kernel $k(t, s) + 1$

- ③ Neumann BC case: contour lines orthog to Γ

Helmholtz

$(\Delta + \kappa^2)u = 0$ arises from scalar wave equation $u_{tt} - \Delta u = 0$

κ “wavenumber”; wavelength $\lambda = 2\pi/\kappa$

Also used for 2D Maxwell (z-invar); TE vs TM

Dirichlet BC = PEC in TE (check)

Scattering formalism:

$$u^{\text{tot}} = u^{\text{inc}} + u.$$

BVP for u (3 lines: PDE, BC, SRC)

Solve BVP for u via PTR + Nyström

- 4 debug BVP with known data from a radiative soln sources inside Ω

Helmholtz transmission BVP

refractive indices in Ω vs exterior

DECIDE IF TRANSMISSION

Matching? (more effort, needs $\mathcal{S}\sigma + \mathcal{D}\tau \dots$)

Helmholtz

fund sol,

Dirichlet demo, plots only

(don't demo CFIE since requires S w/ log-singularity). Demo DLP only,
see $1/N^3$ conv if use naive PTR with correct diag limit (see M126 HW?)

Getting spectral-acc Nyström for log-singular kernels: beyond today.

eg kernel-split or product quadratures (Kress, Helsing, ...)

close-eval: kernel-split, QBX, etc.

see libraries: chunkie, BIE2D, etc

More debug ideas

TO DISCARD

Other tests:

- ⑤ Test SLP & DLP evaluators via GRF for any harmonic u in Ω

Recap

*** TO EDIT WITH FRUZZINA:

BIE involves several steps: write out yourself + try HW exer. in repo

- we covered basics for one smooth curve in 2D
code yourself, if only to understand what libraries do better!
- Interior/exterior, Laplace/Helmholtz/etc, somewhat simple changes
spectral conv once you have good log-singular kernel quadrature
- Fancier quadratures needed for singular kernels and/or close eval
- Transmission BVPs may need twice the unknowns
- Nyström is not only way to discretize: Galerkin, collocation
but: simplest and no less accurate

Resources

Many numerical analysis (mathematics heavy). Somewhat accessible:

- *Linear Integral Equations*, R. Kress, (1999, Springer). Ch. 6 & 12.
- *The Numerical Solution of Integral Equations of the Second Kind*, K. E. Atkinson, (1997, CUP).

Fewer on the practical/tutorial side, few with modern devels:

- “High-order accurate methods for Nyström discretization of integral equations on smooth curves in the plane”, S Hao, AH Barnett, PG Martinsson, P Young. *Adv. Comput. Math.* **40**, 245–272 (2014).

focuses on quadrature for logarithmic singularities, eg SLP, Helmholtz

- <https://users.flatironinstitute.org/~ahb/BIE/>
- <https://github.com/ahbarnett/BIEbook>

in progress...