

NUMERICAL LINEAR ALGEBRA

Exercises from previous exams on *linear equation solving*, the *least square problem* and the *singular value decomposition*.

1. Let A^{-1} be the inverse matrix of A .

- (1) Show that $A^{-1} = [c_1 \ \cdots \ c_n]$ where i -th column c_i verifies

$$(0.1) \quad Ac_i = e_i$$

with e_i the i -th vector in the standard basis of \mathbb{R}^n .

- (2) Given a PLU factorization of A , show that solving each equation in (0.1) costs $4n^2/3$ flops.
 (3) Show that the time complexity of obtaining A^{-1} by computing a PLU factorization of A and then solving the equations (0.1) is $2n^3 + O(n^2)$.

2. Let $A \in \mathbb{R}^{m \times n}$ be a matrix of rank n and $b \in \mathbb{R}^m$.

- (1) Explain how to use the QR factorization of A to solve the least square problem (LSP) that asks to find the vector $x_{\min} \in \mathbb{R}^n$ minimizing the quantity $\|Ax - b\|_2$ for $x \in \mathbb{R}^n$, and give the expression for the residual error $\|Ax_{\min} - b\|_2$.
 (2) Find the affine function $\ell(x) = \alpha x + \beta$ whose graph fits better the points $(-1, 1)$, $(0, 0)$ and $(1, 1)$, in the sense that the Euclidean norm of the vector

$$(\ell(-1) - 1, \ell(0) - 0, \ell(1) - 1) \in \mathbb{R}^3$$

is minimal among all possible choices of $\alpha, \beta \in \mathbb{R}$.

3. Consider the matrix

$$A = \begin{pmatrix} 4 & 1 \\ -2 & -1 \end{pmatrix}$$

- (1) Compute its QR factorization using Householder reflexions.
 (2) Compute the same factorization, but this time using Givens rotations instead of reflexions.

5. Consider the *singular value decomposition* (SVD)

$$A = \begin{pmatrix} 1 & 1 & 0.41 \\ -1 & 0 & 0.41 \\ 0 & 1 & -0.41 \end{pmatrix} = \begin{pmatrix} -0.82 & 0 & -0.58 \\ 0.41 & -0.71 & -0.58 \\ -0.41 & -0.71 & 0.58 \end{pmatrix} \cdot \begin{pmatrix} 1.73 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0.71 \end{pmatrix} \cdot \begin{pmatrix} -0.71 & -0.71 & 0 \\ 0.71 & -0.71 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

- (1) Compute the condition number of the matrix A with respect to the 2-norm.
 (2) Compute the best rank 1 and rank 2 approximations of A with respect to the same norm, and determine the distance to A of these approximations.

4. Let $A \in \mathbb{R}^{2 \times 2}$ such that the eigenvalues of $A \cdot A^T$ are 9 and $\frac{1}{4}$ with respective eigenvectors

$$\begin{pmatrix} \sqrt{2}/2 \\ -\sqrt{2}/2 \end{pmatrix} \approx \begin{pmatrix} 0.71 \\ -0.71 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \sqrt{2}/2 \\ \sqrt{2}/2 \end{pmatrix} \approx \begin{pmatrix} 0.71 \\ 0.71 \end{pmatrix},$$

and the eigenvalues of $A^T \cdot A$ are 9 and $\frac{1}{4}$ with respective eigenvectors

$$\begin{pmatrix} 1/2 \\ -\sqrt{3}/2 \end{pmatrix} \approx \begin{pmatrix} 0.50 \\ -0.87 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \sqrt{3}/2 \\ 1/2 \end{pmatrix} \approx \begin{pmatrix} 0.87 \\ 0.50 \end{pmatrix}.$$

- (1) Compute a singular value decomposition (SVD) of A .
 (2) Using this SVD, compute condition number of A with respect to the 2-norm.
 (3) Determine the image of the unit disk of \mathbb{R}^2 under the linear map defined by A .