

Chapter 5

Convex sets and functions

Optimality conditions

Definition of convex sets in \mathbb{R}^n

Definition. We say that $\mathcal{C} \subset \mathbb{R}^n$ is a **convex set** if for any two points $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{C}$ we have

$$\alpha \mathbf{u}_1 + (1 - \alpha) \mathbf{u}_2 \in \mathcal{C}, \quad \alpha \in [0, 1]$$

Exercise. Any (open and closed) hyper-cube and any (open and closed) hyper-ball in \mathbb{R}^n are convex sets.

$$Q := \{\mathbf{x} \in \mathbb{R}^n \mid 0 < |x_j| < 1, j = 1, \dots, n\}$$


$$B := \{\mathbf{x} \in \mathbb{R}^n \mid \|\mathbf{x}\| < 1\}$$

Hyperplanes

Definition. Let $\mathbf{a} \in \mathbb{R}^n \setminus \{\vec{0}\}$, and let $b \in \mathbb{R}$. The set

$$\mathbb{H}_a := \mathbb{H} = \left\{ \mathbf{x} \mid \mathbf{a}^T \mathbf{x} = b \right\}$$

is called a **hyperplane** of \mathbb{R}^n . Alternatively, \mathbb{H} is the set of all all the vectors $\mathbf{x} \in \mathbb{R}^n$ such that its scalar product with $\mathbf{a} \in \mathbb{R}^n \setminus \{\vec{0}\}$ is constant.



Hyperplanes

Exercise. If \mathbf{x}_0 i \mathbf{x}_1 are two points in \mathbb{H}_a , then

$$\mathbf{a}^T(\mathbf{x}_1 - \mathbf{x}_0) = 0$$

Definition. The vector \mathbf{a} is called the **normal vector** of \mathbb{H}_a .

Exercise. The set \mathbb{H}_a is a convex set. Moreover the set \mathbb{H}_a defines two convex open half-spaces and two closed half-spaces given by

$$\begin{aligned}\mathbb{H}_a^+ &= \{\mathbf{x} \mid \mathbf{a}^T \mathbf{x} > b\}, \quad \mathbb{H}_a^- = \{\mathbf{x} \mid \mathbf{a}^T \mathbf{x} < b\}, \quad \text{and} \\ \overline{\mathbb{H}}_a^+ &= \{\mathbf{x} \mid \mathbf{a}^T \mathbf{x} \geq b\}, \quad \overline{\mathbb{H}}_a^- = \{\mathbf{x} \mid \mathbf{a}^T \mathbf{x} \leq b\}\end{aligned}$$

The convex hull

Lemma. The intersection of an arbitrary family of convex sets is also a convex set.

Definition. Let $G \subset \mathbb{R}^n$ be an arbitrary set. The intersection of all convex sets containing G is called the **convex hull of A** , and it will be denoted by $\mathcal{C}(G)$.

Corollary. For any given $G \subset \mathbb{R}^n$, the set $\mathcal{C}(G)$ is a convex set.

Exercise. Compute $\mathcal{C}(G)$ for

$$G = \{\cup_{n=1}^3 (x_n, y_n)\} \subset \mathbb{R}^2.$$

Separating hyperplanes

Let G_1 and G_2 be nonempty subsets of \mathbb{R}^n .

Definition. We say that \mathbb{H}_a separates G_1 from/and G_2 if

$$G_1 \subset \left\{ \mathbf{x} \mid \mathbf{a}^T \mathbf{x} \geq b \right\} \quad \text{and} \quad G_2 \subset \left\{ \mathbf{x} \mid \mathbf{a}^T \mathbf{x} \leq b \right\}$$

The set \mathbb{H}_a strictly separates G_1 and G_2 if the inequalities are strict.

Theorem (separation theorem). Let $G_j \in \mathbb{R}^n$, $j = 1, 2$ be two disjoint nonempty convex sets. Then there exists a hyperplane that separates them. Moreover, if we assume that C_2 is compact then there exists a hyperplane that strictly separates them.

Farkas Lemma

Theorem (Farkas' Lemma). Let A be an $m \times n$ real matrix and let $\mathbf{b} \in \mathbb{R}^n$. The inequality $\mathbf{b}^T \mathbf{y} \geq 0$ holds for all vectors $\mathbf{y} \in \mathbb{R}^n$ satisfying $A\mathbf{y} \geq 0$ if and only if there exists a vector $\boldsymbol{\rho} \in \mathbb{R}^m$ with $\boldsymbol{\rho} \geq 0$, such that $A^T \boldsymbol{\rho} = \mathbf{b}$

Proof. The statement is equivalent to

$$\left. \begin{array}{l} A\mathbf{y} \geq 0 \\ \mathbf{b}^T \mathbf{y} < 0 \end{array} \right\} \text{ has a solution if and only if } \left. \begin{array}{l} A^T \boldsymbol{\rho} = \mathbf{b} \\ \boldsymbol{\rho} \geq 0 \end{array} \right\} \text{ has no solution}$$

\Leftrightarrow) Then, the nonempty convex sets

$$C_1 = \left\{ \mathbf{x} \in \mathbb{R}^n \mid \mathbf{x} = A^T \boldsymbol{\rho}, \boldsymbol{\rho} \geq 0 \right\} \quad \text{and} \quad C_2 = \{ \mathbf{b} \}$$

are disjoint. Note that C_2 is compact. According to the **Strict Separation Theorem**, there exist $\mathbf{c} \in \mathbb{R}^n$, $\mathbf{c} \neq 0$ and $\alpha \in \mathbb{R}$ such that the hyperplane $H = \{ \mathbf{x} \in \mathbb{R}^n \mid \mathbf{c}^T \mathbf{x} = \alpha \}$ separates them. This is

$$\left\{ \begin{array}{l} \mathbf{c}^T \mathbf{b} < \alpha \\ \forall \mathbf{x} \in C_1, \mathbf{c}^T \mathbf{x} > \alpha \end{array} \right\} \Leftrightarrow \forall \boldsymbol{\rho} \geq 0, \mathbf{c}^T A^T \boldsymbol{\rho} > \alpha$$

Farkas Lemma

Proof (continue). This is

$$\left\{ \begin{array}{l} \mathbf{c}^T \mathbf{b} < \alpha \\ \forall \mathbf{x} \in C_1, \mathbf{c}^T \mathbf{x} > \alpha \end{array} \right\} \Leftrightarrow \forall \boldsymbol{\rho} \geq 0, \mathbf{c}^T A^T \boldsymbol{\rho} > \alpha$$

- (a) Claim: $\mathbf{c}^T \mathbf{b} = \mathbf{b}^T \mathbf{c} < 0$. To see this claim, take $\boldsymbol{\rho} = 0$ above. Then $\alpha < 0$.
- (b) Claim: $\mathbf{c}^T A^T \geq 0$. To this this claim notice that if for a certain k we have that $(\mathbf{c}^T A^T)_k < 0$, then, choosing $\boldsymbol{\rho} = (0, \dots, 0, \rho_k, 0, \dots, 0)$ with $\rho_k \rightarrow +\infty$, we have that $\mathbf{c}^T A^T \boldsymbol{\rho} \rightarrow -\infty$, in contradiction with $\mathbf{c}^T A^T \boldsymbol{\rho} > \alpha$.

Accordingly the vector \mathbf{c} is a solution of

$$\left. \begin{array}{l} A\mathbf{y} \geq 0 \\ \mathbf{b}^T \mathbf{y} < 0 \end{array} \right\},$$

as desired.

Farkas Lemma

Proof (continue).

\Rightarrow) We should prove that

$$\left. \begin{array}{l} Ay \geq 0 \\ b^T y < 0 \end{array} \right\} \text{ has a solution implies } \left. \begin{array}{l} A^T \rho = b \\ \rho \geq 0 \end{array} \right\} \text{ has no solution}$$

(We prove the negative version.) Assume there are ρ and y such that:

$A^T \rho = b$, $\rho \geq 0$ (and $Ay \geq 0$). Then $b^T y = \rho^T Ay \geq 0$. So


$$\left. \begin{array}{l} Ay \geq 0 \\ b^T y < 0 \end{array} \right\} \text{ has no solution.}$$

Convex functions: The epigraph of f .

Definition. Let $D \subset \mathbb{R}^n$ and let $f : D \rightarrow \mathbb{R}$ be a function defined on D with values in the extended reals $\overline{\mathbb{R}}$; this is, $f(\mathbf{x})$, $\mathbf{x} \in D$, is either a real number or it is $\pm\infty$. The subset of \mathbb{R}^{n+1} defined as

$$P(f) = \{(\mathbf{x}, \alpha) \in D \times \mathbb{R} \mid f(\mathbf{x}) \leq \alpha\} \subset \mathbb{R}^{n+1}$$

is called the **epigraph** of f . We say f is a **convex function** if $P(f)$ is a convex set.



Convex functions: The epigraph of f .

Consider a convex function f defined in a subset $D \subset \mathbb{R}^n$. Let

$$f_1(\mathbf{x}) = \begin{cases} f(\mathbf{x}) & \text{if } \mathbf{x} \in D \\ +\infty & \text{if } \mathbf{x} \notin D \end{cases}$$

The **epigraph** of $f|_D$ is identical to the one of $f_1|_{\mathbb{R}^n}$. Hence we can always extend a convex function f (over D), to be a convex function defined throughout all \mathbb{R}^n .

Remark. Let $a \in \mathbb{R}$, $\mathbf{b} \in \mathbb{R}^n$. Then

$$f_1(\mathbf{x}) = \begin{cases} a & \text{if } \mathbf{x} = \mathbf{b} \\ +\infty & \text{if } \mathbf{x} \neq \mathbf{b} \end{cases}$$

is a convex (not continuous) function defined over all \mathbb{R}^n .

Convex functions: The effective domain of f .

Definition. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. The **effective domain** of f is the set

$$\text{ED}(f) = \{\mathbf{x} \in \mathbb{R}^n \mid f(\mathbf{x}) < +\infty\}$$

Exercises.

- (a) Show that $\text{ED}(f)$ is the projection of $P(f)$ over \mathbb{R}^n (the first component).
- (b) If f is a convex function, then $\text{ED}(f)$ is a convex set.
- (c) Show that the converse (of statement (b)) is not necessarily true.

Definition. We say that f is a **proper convex function** if f is convex, $f(\mathbf{x}) > -\infty$ for every \mathbf{x} , and $\text{ED}(f) \neq \emptyset$.

An equivalent definition for convexity

Theorem. Let $q_1, \dots, q_s \in \mathbb{R}$ with $q_j \geq 0$, $j = 1, \dots, s$ and $\sum_{j=1}^s q_j = 1$.

Then, f is a (proper) convex function on \mathbb{R}^n if and only if for all $\mathbf{x}_1, \dots, \mathbf{x}_s \in \mathbb{R}^n$ we have

$$f(q_1 \mathbf{x}_1 + \dots + q_s \mathbf{x}_s) \leq q_1 f(\mathbf{x}_1) + \dots + q_s f(\mathbf{x}_s) \quad (1)$$

Proof (\Rightarrow).

- (a) If $f(\mathbf{x}_j) = +\infty$ for some $j = 1, \dots, s$, then (1) trivially holds.
- (b) Assume now that $f(\mathbf{x}_j) < +\infty$ for all $j = 1, \dots, s$. Since f is convex, then $P(f)$ is a convex set. That is,

$$(\mathbf{x}_1, \alpha_1) \in P(f), \dots, (\mathbf{x}_s, \alpha_s) \in P(f) \Rightarrow (q_1 \mathbf{x}_1 + \dots + q_s \mathbf{x}_s, q_1 \alpha_1 + \dots + q_s \alpha_s) \in P(f).$$

This is to say that

$$f(q_1 \mathbf{x}_1 + \dots + q_s \mathbf{x}_s) \leq q_1 \alpha_1 + \dots + q_s \alpha_s$$

- (c) Since $(\mathbf{x}_i, \alpha_i) \in P(f) \Rightarrow f(\mathbf{x}_i) \leq \alpha_i$, we can take $\alpha_i = f(\mathbf{x}_i)$, for $i = 1, \dots, n$, and (1) follows.

Linear combinations of convex functions

Lemma. Let f and g be convex functions. Let $\lambda \in \mathbb{R}_+$. Then the functions λf and $f + g$ are also convex functions (provided that the operation $+\infty + (-\infty)$ is avoided).

In particular, every linear combination $\lambda_1 f_1 + \cdots + \lambda_k f_k$ of convex functions with $\lambda_j \geq 0$ for all $j = 1, \dots, k$ is also a convex function.

Exercise. Prove the above statements.

Composition and convex functions

Definition. Let $\Psi : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ be a function defined on \mathbb{R} with values in the extended reals. We say that Ψ is **non-decreasing** if for every $x_1 < x_2$ we have $\Psi(x_1) \leq \Psi(x_2)$.

Theorem. Let f be a real convex function defined on \mathbb{R}^n , and let Ψ be a non-decreasing proper convex function defined on \mathbb{R} . Then $\Psi \circ f$ is convex on \mathbb{R}^n .

Proof. Since f is convex and Ψ is non-decreasing we have $(0 \leq q_1 \leq 1)$

$$f(q_1 x_1 + (1 - q_1)x_2) \leq q_1 f(x_1) + (1 - q_1)f(x_2), \text{ and} \\ \Psi(f(q_1 x_1 + (1 - q_1)x_2)) \leq \Psi(q_1 f(x_1) + (1 - q_1)f(x_2))$$

Finally by the convexity of Ψ we have

$$\Psi(f(q_1 x_1 + (1 - q_1)x_2)) \leq \Psi(q_1 f(x_1) + (1 - q_1)f(x_2)) \leq q_1 \Psi(f(x_1)) + (1 - q_1)\Psi(f(x_2)).$$

The maximum of convex functions

Theorem. Let f_j , $j = 1, \dots, m$ be a finite collection of convex functions on \mathbb{R}^n . Then the function

$$F(x) := \max_j f_j(x)$$

is a **convex function** (i.e., $P(F)$ is a convex set).

Proof. The sets $P(f_j)$, $j = 1, \dots, m$ (epigraphs) are convex sets and so their intersection is convex as well. By definition

$$\begin{aligned} \bigcap_j P(f_j) &= \{(x, \alpha) \in \mathbb{R}^n \times \mathbb{R} \mid \max_j f_j(x) \leq \alpha, \text{ for all } j = 1, \dots, m\} = \\ &= \{(x, \alpha) \in \mathbb{R}^n \times \mathbb{R} \mid \max_j f_j(x) = F(x) \leq \alpha\} = P(F). \end{aligned}$$

Two important results

Theorem. A real valued function f defined on \mathbb{R}^n is convex if and only if for every $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^n$, the function $\phi : [0, 1] \rightarrow \mathbb{R}$ defined by

$$\phi(\lambda) = f(\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2)$$

is convex.

Theorem. A real-valued convex function on \mathbb{R}^n is continuous everywhere.

Convex differentiable functions

Definition. Let $D \in \mathbb{R}^n$ an open set and let $\mathbf{x}_0 \in D$. Let $f : D \rightarrow \mathbb{R}$. Let $\mathbf{v} \in \mathbb{R}^n$ a unitary vector. We define the **\mathbf{v} -directional derivative of f at the point \mathbf{x}_0** by

$$Df(\mathbf{x}_0, \mathbf{v}) := \lim_{t \rightarrow 0} \frac{f(\mathbf{x}_0 + t\mathbf{v}) - f(\mathbf{x}_0)}{t}.$$



When we consider the above limit with $t \rightarrow 0^+$ and $t \rightarrow 0^-$ we denote them by $D^+f(\mathbf{x}_0, \mathbf{v})$ and $D^-f(\mathbf{x}_0, \mathbf{v})$ and we called them **right-sided (left-sided) \mathbf{v} -directional derivative of f at the point \mathbf{x}_0** , respectively.

Remark. According to previous notation and results we have

$$Df(\mathbf{x}_0; \mathbf{v}) = \mathbf{v}^T \nabla f(\mathbf{x}_0) = \nabla f(\mathbf{x}_0)^T \mathbf{v}$$

Convex differentiable functions

Definition. A function f is said to be **positively homogeneous of degree $k \geq 1$** if for every $\mathbf{x} \in \mathbb{R}^n$ and every $t \in \mathbb{R}^+$ we have

$$f(t\mathbf{x}) = t^k f(\mathbf{x})$$

Theorem. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex (finite) function. Then

- (a) For any unitary $\mathbf{v} \in \mathbb{R}^n$ there exist the right-sided and left-sided derivatives of f at every \mathbf{x} .
- (b) D^+f and D^-f are positively homogeneous convex functions of \mathbf{v} of degree one.
- (c) The following inequality holds:

$$D^+f(\mathbf{x}; \mathbf{y}) \geq D^-f(\mathbf{x}; \mathbf{y})$$

Convex differentiable functions: Subgradients

Definition. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function. A **subgradient of f at a point $\mathbf{x}_0 \in \mathbb{R}^n$** , is a vector $\boldsymbol{\xi} \in \mathbb{R}^n$ such that



$$f(\mathbf{y}) \geq f(\mathbf{x}_0) + \boldsymbol{\xi}^T (\mathbf{y} - \mathbf{x}_0) \quad (2)$$

for every $\mathbf{y} \in \mathbb{R}^n$.

Remark. A subgradient of a convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at a $\mathbf{x}_0 \in \mathbb{R}^n$ may be a unique vector or several vectors.

Notation and definition. We denote by $\partial f(\mathbf{x})$ the set of all subgradients of a convex function f at a given point \mathbf{x} . In some books $\partial f(\mathbf{x})$ is called **subdifferential**.

Convex differentiable functions: Subgradients



Theorem. Let f be a convex function. A vector $\xi \in \partial f(\mathbf{x})$ if and only if

$$D^+ f(\mathbf{x}; \mathbf{v}) \geq \xi^T \mathbf{v} \quad \forall \mathbf{v} \in \mathbb{R}^n \quad (3)$$

Convex differentiable functions: Subgradients

Proof. If $\xi \in \partial f(x)$, then it satisfies $f(y) \geq f(x) + \xi^T(y - x)$ for all $y \in \mathbb{R}^n$. If we write $y = x + tz$, with $t > 0$, then the previous inequality writes as

$$f(x + tz) \geq f(x) + t\xi^T z \quad \text{or} \quad \frac{f(x + tz) - f(x)}{t} \geq \xi^T z$$

for every $z \in \mathbb{R}^n$ and $t > 0$. We deduce from above that $D^+f(x; z) \geq \xi^T z$ since $D^+f(x; z)$ is the limit of the incremental quotients ($t > 0$).

The other implication follows similarly.