

A formalized extension of the substitution lemma in Coq

Maria J. D. Lima

Departamento de Ciência da Computação
Universidade de Brasília, Brasília, Brazil
majuhdl@gmail.com

Flávio L. C. de Moura

Departamento de Ciência da Computação
Universidade de Brasília, Brasília, Brazil
flaviomoura@unb.br

The substitution lemma is a renowned theorem within the realm of λ -calculus theory and concerns the interactional behaviour of the metasubstitution operation. In this work, we augment the λ -calculus's grammar with an uninterpreted explicit substitution operation, which allows the use of our framework for different calculi with explicit substitutions. Our primary contribution lies in verifying that, despite these modifications, the substitution lemma continues to remain valid. This confirmation was achieved using the Coq proof assistant. Our formalization methodology employs a nominal approach, which provides a remarkably direct implementation of the α -equivalence concept. Despite this simplicity, the strategy involved in variable renaming within the proofs presents a substantial challenge, ensuring a comprehensive exploration of the implications of our extension to the grammar of the λ -calculus.

1 Introduction

In this work, we present a formalization of the substitution lemma[5] in a general framework that extends the λ -calculus with an explicit substitution operator. The formalization is done in the Coq proof assistant[25] and the source code is available at:

https://github.com/flaviodemoura/lx_confl/tree/m_subst_lemma

The substitution lemma is an important result concerning the composition of the substitution operation, and is usually presented as follows in the context of the λ -calculus:

Let t, u and v be λ -terms. If $x \notin FV(v)$ (i.e. x does not occur in the set of free variables of the term v) then $\{y := v\}\{x := u\}t =_{\alpha} \{x := \{y := v\}u\}\{y := v\}t$.

This is a well known result already formalized in the context of the λ -calculus [7]. Nevertheless, in the context of λ -calculi with explicit substitutions its formalization is not straightforward due to the interaction between the metasubstitution and the explicit substitution operator. Our formalization is done in a nominal setting that uses the MetaLib¹ package of Coq, but no particular explicit substitution calculi is taken into account because the expected behaviour between the metasubstitution operation with the explicit substitution constructor is the same regardless the calculus. The contributions of this work are twofold:

1. The formalization is modular in the sense that no particular calculi with explicit substitutions is taken into account. Therefore, we believe that this formalization could be seen as a generic framework for proving properties of these calculi that uses the substitution lemma in the nominal setting[18, 22, 24];

¹<https://github.com/plclub/metilib>

2. A solution to a circularity problem in the proofs is given. It adds an axiom to the formalization that replaces the set equality by the syntactic equality. In this way, we are allowed to replace/rewrite sets of (free) variables by another sets of (free) variables in arbitrary contexts.

This document is built directly from a Coq script using the CoqDoc² tool. In the following section, we present the general framework and the basics of the nominal approach. In Section 3, we present our definition of metasubstitution and some of its properties. In Section 4, we present the main theorem, *i.e.* the substitution lemma, and we conclude in Section 5.

2 A syntactic extension of the λ -calculus

In this section, we present the framework of the formalization, which is based on a nominal approach[15] where variables use names. In the nominal setting, variables are represented by atoms that are structure-less entities with a decidable equality:

Parameter `eq_dec` : forall `x y` : atom, {`x = y`} + {`x <> y`}.

therefore different names mean different atoms and different variables. The nominal approach is close to the usual paper and pencil notation used in λ -calculus lectures, whose grammar of terms is given by:

$$t ::= x \mid \lambda_x.t \mid t \ t \quad (1)$$

and its main rule, named β -reduction, is given by:

$$(\lambda_x.t) \ u \rightarrow_\beta \{x := u\}t \quad (2)$$

where $\{x := u\}t$ represents the term obtained from t after replacing all its free occurrences of the variable x by u in a way that renaming of bound variable is done in order to avoid variable capture. In other words, $\{x := u\}t$ is a metanotation for a capture free substitution. For instance, the λ -term $(\lambda_x \lambda_y. x \ y) \ y$ has both bound and free occurrences of the variable y . In order to β -reduce it one has to replace (or substitute) the free variable y for all free occurrences of the variable x in the term $(\lambda_y. x \ y)$. But a straight substitution will capture the free variable y , *i.e.* this means that the free occurrence of y before the β -reduction will become bound after the β -reduction step. A renaming of bound variables is done to avoid such capture, so in this example, one can take an α -equivalent³ term, say $(\lambda_z. x \ z)$, and perform the β -step correctly as $(\lambda_x \lambda_y. x \ y) \ y \rightarrow_\beta \lambda_z. y \ z$. The renaming of variables in the nominal setting is done via a name-swapping, which is formally defined as follows:

$$((x \ y))z := \begin{cases} y, & \text{if } z = x; \\ x, & \text{if } z = y; \\ z, & \text{otherwise.} \end{cases}$$

This notion can be extended to λ -terms in a straightforward way:

$$(x \ y)t := \begin{cases} ((x \ y))z, & \text{if } t = z; \\ \lambda_{((x \ y))z}. (x \ y)t_1, & \text{if } t = \lambda_z. t_1; \\ (x \ y)t_1 \ (x \ y)t_2, & \text{if } t = t_1 \ t_2 \end{cases} \quad (3)$$

In the previous example, one could apply a swap to avoid the variable capture in a way that, a swap is applied to the body of the abstraction before applying the metasubstitution to it: $(\lambda_x \lambda_y. x \ y) \ y \rightarrow_\beta \{x :=$

²<https://coq.inria.fr/refman/using/tools/coqdoc.html>

³A formal definition of this notion will be given later in this section.

$y\}((y\ z)(\lambda_y.x\ y)) = \{x := y\}(\lambda_z.x\ z) = \lambda_z.y\ z$. Could we have used a variable substitution instead of a swapping in the previous example? Absolutely. We could have done the reduction as $(\lambda_x.\lambda_y.x\ y)\ y \rightarrow_\beta \{x := y\}(\{y := z\}(\lambda_y.x\ y)) = \{x := y\}(\lambda_z.x\ z) = \lambda_z.y\ z$, but as we will shortly see, variable substitution is not stable under α -equivalence, while the swapping is stable under α -equivalence, thereby rendering it a more fitting choice when operating modulo α -equivalence.

In what follows, we will adopt a mixed-notation approach, intertwining metanotation with the equivalent Coq notation. This strategy aids in elucidating the proof steps of the upcoming lemmas, enabling a clearer and more detailed comprehension of each stage in the argumentation. The corresponding Coq code for the swapping of variables, named *swap_var*, is defined as follows:

Definition *vswap* (*x:atom*) (*y:atom*) (*z:atom*) := if (z == x) then y else if (z == y) then x else z.

therefore, the swap $((x\ y)z)$ is written in Coq as *vswap* *x* *y* *z*. A short example to acquaint ourselves with the Coq notation, let us show how we will write the proofs:

Lemma *swap_var_id*: $\forall\ x\ y, \text{vswap}\ x\ x\ y = y$.

Proof. The proof is done by case analysis, and it is straightforward in both cases, when $x = y$ and $x \neq y$. \square

2.1 An explicit substitution operator

The extension of the swap operation to terms require an additional comment because we will not work with the grammar (1), but rather, we will extend it with an explicit substitution operator:

$$t ::= x \mid \lambda_x.t \mid t\ t \mid [x := u]t \quad (4)$$

where $[x := u]t$ represents a term with an operator that will be evaluated with specific rules of a calculus. The intended meaning of the explicit substitution is that it will simulate the metasubstitution. This formalization aims to be a generic framework applicable to any calculi with explicit substitutions in named notation for variables. Therefore, we will not specify rules about how one can simulate the metasubstitution, but it is important to be aware that this is not a trivial task as one can easily lose important properties of the original λ -calculus[20, 16].

Calculi with explicit substitutions are formalisms that deconstruct the metasubstitution operation into more granular steps, thereby functioning as an intermediary between the λ -calculus and its practical implementations. In other words, these calculi shed light on the execution models of higher-order languages. In fact, the development of a calculus with explicit substitutions faithful to the λ -calculus, in the sense of the preservation of some desired properties were the main motivation for such a long list of calculi with explicit substitutions invented in the last decades[1, 8, 6, 11, 21, 17, 9, 12, 19].

The following inductive definition corresponds to the grammar (4), where the explicit substitution constructor, named *n_sub*, has a special notation. Instead of writing *n_sub* *t* *x* *u*, we will write $[x := u]$ *t* similarly to (4). Therefore, *n_sexp* is used to denote the set of nominal expressions equipped with an explicit substitution operator, which, for simplicity, we will refer to as just “terms”.

Inductive *n_sexp* : Set :=

| *n_var* (*x:atom*)
 | *n_abs* (*x:atom*) (*t:n_sexp*)
 | *n_app* (*t1:n_sexp*) (*t2:n_sexp*)
 | *n_sub* (*t1:n_sexp*) (*x:atom*) (*t2:n_sexp*).

Notation “[*x := u*] *t*” := (*n_sub* *t* *x* *u*) (at level 60).

The *size* and the set *fv_nom* of the free variables of a term are defined as usual:

```

Fixpoint size (t : n_sexp) : nat :=
  match t with
  | n_var x ⇒ 1
  | n_abs x t ⇒ 1 + size t
  | n_app t1 t2 ⇒ 1 + size t1 + size t2
  | n_sub t1 x t2 ⇒ 1 + size t1 + size t2
  end.

Fixpoint fv_nom (t : n_sexp) : atoms :=
  match t with
  | n_var x ⇒ { {x} }
  | n_abs x t1 ⇒ remove x (fv_nom t1)
  | n_app t1 t2 ⇒ fv_nom t1 'union' fv_nom t2
  | n_sub t1 x t2 ⇒ (remove x (fv_nom t1)) 'union' fv_nom t2
  end.

```

The action of a permutation on a term, written $(x\ y)t$, is inductively defined as in (3) with the additional case for the explicit substitution operator:

$$(x\ y)t := \begin{cases} ((x\ y))v, & \text{if } t \text{ is the variable } v; \\ \lambda_{((x\ y))z}.(x\ y)t_1, & \text{if } t = \lambda_z.t_1; \\ (x\ y)t_1\ (x\ y)t_2, & \text{if } t = t_1\ t_2; \\ [((x\ y))z := (x\ y)t_2](x\ y)t_1, & \text{if } t = [z := t_2]t_1. \end{cases}$$

The corresponding Coq definition is given by the following recursive function:

```

Fixpoint swap (x:atom) (y:atom) (t:n_sexp) : n_sexp :=
  match t with
  | n_var z ⇒ n_var (vswap x y z)
  | n_abs z t1 ⇒ n_abs (vswap x y z) (swap x y t1)
  | n_app t1 t2 ⇒ n_app (swap x y t1) (swap x y t2)
  | n_sub t1 z t2 ⇒ n_sub (swap x y t1) (vswap x y z) (swap x y t2)
  end.

```

The *swap* function has many interesting properties, but we will focus on the ones that are more relevant to the proofs related to the substitution lemma. Nevertheless, all lemmas can be found in the source code of the formalization⁴. The next lemma shows that the *swap* function preserves the size of terms. It is proved by induction on the structure of the term *t*:

Lemma *swap_size_eq* : $\forall\ x\ y\ t, \text{size}(\text{swap}\ x\ y\ t) = \text{size}\ t$.

The *swap* function is involutive, which is also proved done by structural induction on the term *t*:

Lemma *swap_involutive* : $\forall\ t\ x\ y, \text{swap}\ x\ y\ (\text{swap}\ x\ y\ t) = t$.

The shuffle property given by the following lemma is also proved by structural induction on the structure of *t*: **Lemma *shuffle_swap*** : $\forall\ w\ y\ z\ t,$

$$w \neq z \rightarrow y \neq z \rightarrow$$

⁴https://github.com/flaviodemoura/lx_confl/tree/m_subst_lemma

$$(\text{swap } w \ y \ (\text{swap } y \ z \ t)) = (\text{swap } w \ z \ (\text{swap } w \ y \ t)).$$

Equivariance is another important property of the *swap* function. It states that a swap can uniformly be propagated over the structure of a term:

Lemma *vswap_equivariance* : $\forall v \ x \ y \ z \ w, \text{vswap } x \ y \ (\text{vswap } z \ w \ v) = \text{vswap } (\text{vswap } x \ y \ z) \ (\text{vswap } x \ y \ w) \ (\text{vswap } x \ y \ v).$

Lemma *swap_equivariance* : $\forall t \ x \ y \ z \ w, \text{swap } x \ y \ (\text{swap } z \ w \ t) = \text{swap } (\text{vswap } x \ y \ z) \ (\text{vswap } x \ y \ w) \ (\text{swap } x \ y \ t).$

If a variable, say *z*, is not in the set of free variables of a term *t* and one swaps *z* with another variable, say *y*, then *y* is not in the set of free variables of the term *t*. This is the content of the following lemma that can easily be proved using induction on the structure of the term *t*: **Lemma** *fv_nom_swap* : $\forall z \ y \ t, z \text{ 'notin' } \text{fv_nom } t \rightarrow y \text{ 'notin' } \text{fv_nom } (\text{swap } y \ z \ t).$

The standard proof strategy for the non trivial properties is induction on the structure of the terms. Nevertheless, the builtin induction principle automatically generated for the inductive definition *n_sexp* is not strong enough due to swappings. In fact, in general, the induction hypothesis in the abstraction case, for instance, refer to the body of the abstraction, while the goal involves a swap acting on the body of the abstraction. In order to circumvent this problem, we use an induction principle based on the size of terms:

Lemma *n_sexp_induction*:

$$\begin{aligned} & \forall P : n_sexp \rightarrow \text{Prop}, \\ & (\forall x, P \ (n_var \ x)) \rightarrow \\ & (\forall t1 \ z, (\forall t2 \ x \ y, \text{size } t2 = \text{size } t1 \rightarrow P \ (\text{swap } x \ y \ t2)) \rightarrow P \ (n_abs \ z \ t1)) \rightarrow \\ & (\forall t1 \ t2, P \ t1 \rightarrow P \ t2 \rightarrow P \ (n_app \ t1 \ t2)) \rightarrow \\ & (\forall t1 \ t3 \ z, P \ t3 \rightarrow (\forall t2 \ x \ y, \text{size } t2 = \text{size } t1 \rightarrow P \ (\text{swap } x \ y \ t2)) \rightarrow P \ (n_sub \ t1 \ z \ t3)) \rightarrow \\ & (\forall t, P \ t). \end{aligned}$$

We will use this induction principle to prove that if a certain variable, say *x'*, is not in the set of free variables of a term *t* then the variable obtained after applying any swap to *x'* also is not in the set of free variables of the term obtained from *t* after applying the same swap to *t*:

Lemma *notin_fv_nom_equivariance* : $\forall t \ x' \ x \ y, x' \text{ 'notin' } \text{fv_nom } t \rightarrow \text{vswap } x \ y \ x' \text{ 'notin' } \text{fv_nom } (\text{swap } x \ y \ t).$

Proof. The proof is by induction on the size of the term *t*. If *t* is a variable, say *z*, then we have that $x' \neq z$ by hypothesis, and we conclude by lemma *swap_neq*. If $t = n_abs \ z \ t1$ then we have that $x' \notin \text{fv}(t1) \setminus \{z\}$ by hypothesis. This means that either $x' = z$ or x' is not in $\text{fv}(t1)$, i.e. *fv_nom t1* in the Coq language. If $x' = z$ then we have to prove that a certain element is not in a set where it was removed, and we are done by lemma *notin_remove_3*⁵. Otherwise, x' is not in $\text{fv}(t1)$, and we conclude using the induction hypothesis. The application case is straightforward from the induction hypothesis. The case of the explicit substitution, i.e. when $t = [z := t2]t1$ then we have to prove that $\text{vswap } x \ y \ x' \text{ 'notin' } \text{fv_nom } (\text{swap } x \ y \ ([z := t2]t1))$. We then propagate the swap over the explicit substitution operator and, by the definition of *fv_nom*, we have prove that both $\text{vswap } x \ y \ x' \text{ 'notin' } \text{remove } (\text{vswap } x \ y \ z) \ (\text{fv_nom } (\text{swap } x \ y \ t1))$ and $\text{vswap } x \ y \ x' \text{ 'notin' } \text{fv_nom } (\text{swap } x \ y \ t2)$. In the former case, the hypothesis $x' \text{ 'notin' } \text{remove } z \ (\text{fv_nom } t1)$ generates two cases, either $x' = z$ or x' is not in $\text{fv}(t1)$, and we conclude with the same strategy of the abstraction case. The later case is straightforward by the induction hypothesis. \square

⁵This is a lemma from Metalib library and it states that forall (x y : atom) (s : atoms), x = y -> y 'notin' remove x s.

2.2 α -equivalence

As usual in the standard presentations of the λ -calculus, we work with terms modulo α -equivalence. This means that λ -terms are identified up to the name of bound variables. For instance, all the terms $\lambda_x.x$, $\lambda_y.y$ and $\lambda_z.z$ are seen as the same term which corresponds to the identity function. Formally, the notion of α -equivalence is defined by the following inference rules:

$$\begin{array}{c}
\frac{}{x =_{\alpha} x} \text{ (aeq_var)} \qquad \frac{t_1 =_{\alpha} t_2}{\lambda_x.t_1 =_{\alpha} \lambda_x.t_2} \text{ (aeq_abs_same)} \\
\\
\frac{x \neq y \quad x \notin \text{fv}(t_2) \quad t_1 =_{\alpha} (y\ x)t_2}{\lambda_x.t_1 =_{\alpha} \lambda_y.t_2} \text{ (aeq_abs_diff)} \\
\\
\frac{t_1 =_{\alpha} t'_1 \quad t_2 =_{\alpha} t'_2}{t_1\ t_2 =_{\alpha} t'_1\ t'_2} \text{ (aeq_app)} \qquad \frac{t_1 =_{\alpha} t'_1 \quad t_2 =_{\alpha} t'_2}{[x := t_2]t_1 =_{\alpha} [x := t'_2]t'_1} \text{ (aeq_sub_same)} \\
\\
\frac{t_2 =_{\alpha} t'_2 \quad x \neq y \quad x \notin \text{fv}(t'_1) \quad t_1 =_{\alpha} (y\ x)t'_1}{[x := t_2]t_1 =_{\alpha} [y := t'_2]t'_1} \text{ (aeq_sub_diff)}
\end{array}$$

Each of these rules correspond to a constructor in the *aeq* inductive definition below:

Inductive *aeq* : *n_sexp* \rightarrow *n_sexp* \rightarrow **Prop** :=

| *aeq_var* : $\forall x, \text{aeq } (n_var\ x) (n_var\ x)$
| *aeq_abs_same* : $\forall x\ t1\ t2, \text{aeq } t1\ t2 \rightarrow \text{aeq } (n_abs\ x\ t1) (n_abs\ x\ t2)$
| *aeq_abs_diff* : $\forall x\ y\ t1\ t2, x \neq y \rightarrow x \text{ 'notin' } \text{fv_nom } t2 \rightarrow \text{aeq } t1\ (\text{swap } y\ x\ t2) \rightarrow \text{aeq } (n_abs\ x\ t1) (n_abs\ y\ t2)$
| *aeq_app* : $\forall t1\ t2\ t1'\ t2', \text{aeq } t1\ t1' \rightarrow \text{aeq } t2\ t2' \rightarrow \text{aeq } (n_app\ t1\ t2) (n_app\ t1'\ t2')$
| *aeq_sub_same* : $\forall t1\ t2\ t1'\ t2'\ x, \text{aeq } t1\ t1' \rightarrow \text{aeq } t2\ t2' \rightarrow \text{aeq } ([x := t2]\ t1) ([x := t2']\ t1')$
| *aeq_sub_diff* : $\forall t1\ t2\ t1'\ t2'\ x\ y, \text{aeq } t2\ t2' \rightarrow x \neq y \rightarrow x \text{ 'notin' } \text{fv_nom } t1' \rightarrow \text{aeq } t1\ (\text{swap } y\ x\ t1') \rightarrow \text{aeq } ([x := t2]\ t1) ([y := t2']\ t1')$.

Notation "t =a u" := (*aeq* *t* *u*) (at level 60).

In what follows, we use an infix notation for α -equivalence in the Coq code. Therefore, we write *t* =a *u* instead of *aeq* *t* *u*. The above notion defines an equivalence relation over the set *n_sexp* of nominal expressions with explicit substitutions, i.e. the *aeq* relation is reflexive, symmetric and transitive. In addition, α -equivalent terms have the same size, and the same set of free variables:

Lemma *aeq_size*: $\forall t1\ t2, t1 =a t2 \rightarrow \text{size } t1 = \text{size } t2$.

Lemma *aeq_fv_nom*: $\forall t1\ t2, t1 =a t2 \rightarrow \text{fv_nom } t1 = \text{fv_nom } t2$.

The key point of the nominal approach is that the swap operation is stable under α -equivalence in the sense that, $t_1 =_{\alpha} t_2$ if, and only if $(x\ y)t_1 =_{\alpha} (x\ y)t_2, \forall t_1, t_2, x, y$. Note that this is not true for renaming substitutions: in fact, $\lambda_x.z =_{\alpha} \lambda_y.z$, but $\{z := x\}(\lambda_x.z) = \lambda_x.x \neq_{\alpha} \{z := x\}\lambda_y.x(\lambda_y.z)$, assuming that $x \neq y$. This stability result is formalized as follows:

Corollary *aeq_swap*: $\forall t_1 t_2 x y, t_1 =_a t_2 \leftrightarrow (\text{swap } x y t_1) =_a (\text{swap } x y t_2).$

When both variables in a swap does not occur free in a term, it eventually rename bound variables only, *i.e.* the action of this swap results in a term that is α -equivalent to the original term. This is the content of the followin lemma:

Lemma *swap_reduction*: $\forall t x y, x \text{ 'notin' } \text{fv_nom } t \rightarrow y \text{ 'notin' } \text{fv_nom } t \rightarrow (\text{swap } x y t) =_a t.$

There are several other interesting auxiliary properties that need to be proved before achieving the substitution lemma. In what follows, we refer only to the tricky or challenging ones, but the interested reader can have a detailed look in the source files⁶. Note that, swaps are introduced in proofs by the rules *aeq_abs_diff* and *aeq_sub_diff*. As we will see, the proof steps involving these rules are trick because a naïve strategy can easily get blocked in a branch without proof. We conclude this section, with a lemma that gives the conditions for two swaps with a common variable to be merged:

Lemma *aeq_swap_swap*: $\forall t x y z, z \text{ 'notin' } \text{fv_nom } t \rightarrow x \text{ 'notin' } \text{fv_nom } t \rightarrow (\text{swap } z x (\text{swap } x y t)) =_a (\text{swap } z y t).$

Proof. Initially, observe the similarity of the LHS of the α -equation with the lemma *shuffle_swap*. In order to use it, we need to have that both $z \neq y$ and $x \neq y$. If $z = y$ then the RHS reduces to t because the swap is trivial, and the LHS also reduces to t since swap is involutive. When $z \neq y$ then we proceed by comparing x and y . If $x = y$ then both sides of the α -equation reduces to $\text{swap } z y t$, and we are done. Finally, when $x \neq y$, we can apply the lemma *shuffle_swap* and use lemma *aeq_swap* to reduce the current goal to $\text{swap } z x t =_a t$, and we conclude by lemma *swap_reduction* since both z and x are not in the set of free variables of the term t . \square

3 The metasubstitution operation of the λ -calculus

The main operation of the λ -calculus is the β -reduction that express how to evaluate a function, say $(\lambda_x.t)$, applied to an argument u : $(\lambda_x.t) u \rightarrow_\beta \{x := u\}t$, where $\{x := u\}t$ is called a β -contractum and represents the result of the evaluation of the function $(\lambda_x.t)$ with argument u . In other words, $\{x := u\}t$ is the result of substituting u for the free occurrences of the variable x in t . Moreover, it is a capture free substitution in the sense that no free variable becomes bound after a β -reduction. This operation is in the meta level because it is outside the grammar of the λ -calculus, and that's why it is called metasubstitution. As a metaoperation, its definition usually comes with a degree of informality. For instance, Barendregt[5] defines it as follows:

$$\{x := u\}t = \begin{cases} u, & \text{if } t = x; \\ y, & \text{if } t = y \text{ and } x \neq y; \\ \{x := u\}t_1 \{x := u\}t_2, & \text{if } t = \{x := u\}(t_1 t_2); \\ \lambda_y.(\{x := u\}t_1), & \text{if } t = \lambda_y.t_1. \end{cases}$$

where it is assumed the so called “Barendregt’s variable convention”:

If t_1, t_2, \dots, t_n occur in a certain mathematical context (e.g. definition, proof), then in these terms all bound variables are chosen to be different from the free variables.

⁶https://github.com/flaviodemoura/lx_confl/tree/m_subst_lemma

This means that we are assuming that both $x \neq y$ and $y \notin \text{fv}(u)$ in the case $t = \lambda_y.t_1$. This approach is very convenient in informal proofs because it avoids having to rename bound variables. In order to formalize the capture free substitution, *i.e.* the metasubstitution, there exists different possible approaches. In our case, we perform a renaming of bound variables whenever the metasubstitution is propagated inside a binder. In our case, there are two binders: the abstraction and the explicit substitution.

Let t and u be terms, and x a variable. The result of substituting u for the free occurrences of x in t , written $\{x := u\}t$ is defined as follows:

$$\{x := u\}t = \begin{cases} u, & \text{if } t = x; \\ y, & \text{if } t = y \text{ and } x \neq y; \\ \{x := u\}t_1 \{x := u\}t_2, & \text{if } t = \{x := u\}(t_1 t_2); \\ \lambda_x.t_1, & \text{if } t = \lambda_x.t_1; \\ \lambda_z.(\{x := u\}((y z)t_1)), & \text{if } t = \lambda_y.t_1, x \neq y \text{ and } z \notin \text{fv}(t) \cup \text{fv}(u) \cup \{x\}; \\ [x := \{x := u\}t_2]t_1, & \text{if } t = [x := t_2]t_1; \\ [z := \{x := u\}t_2]\{x := u\}((y z)t_1), & \text{if } t = [y := t_2]t_1, x \neq y \text{ and } z \notin \text{fv}(t) \cup \text{fv}(u) \cup \{x\}. \end{cases} \quad (5)$$

and the corresponding Coq code is as follows:

```
Function subst_rec_fun (t:n_sexp) (u :n_sexp) (x:atom) {measure size t} : n_sexp :=
  match t with
  | n_var y => if (x == y) then u else t
  | n_abs y t1 => if (x == y) then t else let (z,-) :=
    atom_fresh (fv_nom u 'union' fv_nom t 'union' {{x}}) in n_abs z (subst_rec_fun (swap y z t1) u x)
  | n_app t1 t2 => n_app (subst_rec_fun t1 u x) (subst_rec_fun t2 u x)
  | n_sub t1 y t2 => if (x == y) then n_sub t1 y (subst_rec_fun t2 u x) else let (z,-) :=
    atom_fresh (fv_nom u 'union' fv_nom t 'union' {{x}}) in
    n_sub (subst_rec_fun (swap y z t1) u x) z (subst_rec_fun t2 u x)
end.
```

Note that this function is not structurally recursive due to the swaps in the recursive calls. A structurally recursive version of the function `subst_rec_fun` can be found in the file `nominal.v` of the *Metalib* library⁷, but it uses the size of the term in which the substitution will be performed as an extra argument that decreases with each recursive call. We write $[x:=u]t$ instead of `subst_rec_fun t u x` in the Coq code to represent $\{x := u\}t$.

The following lemma states that if $x \notin \text{fv}(t)$ then $\{x := u\}t =_\alpha t$. In informal proofs the conclusion of this lemma is usually stated as a syntactic equality, *i.e.* $\{x := u\}t = t$ instead of the α -equivalence, but the function `subst_rec_fun` renames bound variables whenever the metasubstitution is propagated inside an abstraction or an explicit substitution, even in the case that the metasubstitution has no effect in a subterm. That's why the syntactic equality does not hold here.

Lemma `m_subst_notin`: $\forall t u x, x \text{ 'notin' } \text{fv_nom } t \rightarrow \{x := u\}t =_\alpha t$.

Proof. The proof is done by induction on the size of the term t using the `n_sexp_induction` principle. The interesting cases are the abstraction and the explicit substitution. We focus in the abstraction case, *i.e.* when $t = \lambda_y.t_1$ and $x \neq y$. In this case, we have to prove that $\{x := u\}(\lambda_y.t_1) =_\alpha \lambda_y.t_1$. The induction hypothesis express the fact that every term with the same size as the body of the abstraction t_1 satisfies the property to be proven:

⁷<https://github.com/plclub/metalib>

$\forall t' \ x \ y, |t'| = |t_1| \rightarrow \forall u \ x', x' \notin fv((x \ y)t') \rightarrow \{x' := u\}((x \ y)t') =_\alpha (x \ y)t'$. Therefore, according to the function *subst_rec_fun*, the variable y will be renamed to a new name, say z , such that $z \notin fv(\lambda_y.t_1) \cup fv(u) \cup \{x\}$, and we have to prove that $\{x := u\}\lambda_z.((z \ y)t_1) =_\alpha \lambda_y.t_1$. Since $z \notin fv(\lambda_y.t_1) = fv(t_1) \setminus \{y\}$, there are two cases, either $z = y$ or $z \in fv(t_1)$:

1. $z = y$: In this case, we have to prove that $\{x := u\}\lambda_z.((z \ z)t_1) =_\alpha \lambda_z.t_1$. By the rule *aeq_abs_same* we get $\{x := u\}((z \ z)t_1) =_\alpha t_1$, but in order to apply the induction hypothesis the body of the metasubstitution and the term in the right hand side need to be the same and both need to be a swap. For this reason, we use the transitivity of α -equivalence with $(z \ z)t_1$ as intermediate term. The first subcase is proved by the induction hypothesis, and the second one is proved by the reflexivity of α -equivalence.
2. $z \neq y$: In this case, $x \notin fv(t)$ and we can apply the rule *aeq_abs_diff*. The new goal is $\{x := u\}((z \ y)t_1) =_\alpha (z \ y)t_1$ which holds by the induction hypothesis, since $|(z \ y)t_1| = |t_1|$ and $x \notin fv((z \ y)t_1)$ because $x \neq z$, $x \neq y$ and $x \notin fv(t)$.

The explicit substitution case is also interesting, but it follows a similar strategy used in the abstraction case for t_1 . For t_2 the result follows from the induction hypothesis. \square

We will now prove some stability results for the metasubstitution w.r.t. α -equivalence. More precisely, we will prove that if $t =_\alpha t'$ and $u =_\alpha u'$ then $\{x := u\}t =_\alpha \{x := u'\}t'$, where x is any variable and t, t', u and u' are any *n_sexp* terms. This proof is split in two steps: firstly, we prove that if $u =_\alpha u'$ then $\{x := u\}t =_\alpha \{x := u'\}t, \forall x, t, u, u'$; secondly, we prove that if $t =_\alpha t'$ then $\{x := u\}t =_\alpha \{x := u\}t', \forall x, t, t', u$. These two steps are then combined through the transitivity of the α -equivalence relation. Nevertheless, this task were not straightforward. Let's follow the steps of our first trial.

Lemma *aeq_m_subst_in_trial*: $\forall t \ u \ u' \ x, u =_\alpha u' \rightarrow ([x := u] \ t) =_\alpha ([x := u'] \ t)$.

Proof. The proof is done by induction on the size of the term t . The interesting case is when t is an abstraction, i.e. $t = \lambda_y.t_1$. We need to prove that $\{x := u\}(\lambda_y.t_1) =_\alpha \{x := u'\}(\lambda_y.t_1)$. If $x = y$ then the result is trivial. Suppose $x \neq y$. The metasubstitution will be propagated inside the abstraction on each side of the α -equation, after generating a new name for each side. The new goal is then $\lambda_{x_0}.\{x := u\}((y \ x_0)t_1) =_\alpha \lambda_{x_1}.\{x := u'\}((y \ x_1)t_1)$, where $x_0 \notin fv(\lambda_y.t_1) \cup fv(u) \cup \{x\}$ and $x_1 \notin fv(\lambda_y.t_1) \cup fv(u') \cup \{x\}$. The variables x_0 and x_1 are either the same or different. In the former case the result is trivial because $u =_\alpha u'$. In the latter case, $x_0 \neq x_1$ and we need to prove that $\{x := u\}((y \ x_0)t_1) =_\alpha (x_0 \ x_1)(\{x := u'\}((y \ x_1)t_1))$. Therefore, we need to propagate the swap over the metasubstitution before been able to apply the induction hypothesis. The propagation of the swap over the metasubstitution is stated by the following lemma:

Lemma 1 *Let t, u be terms, and x, y, z variables. Then $(y \ z)(\{x := u\}t) =_\alpha \{((y \ z))x := (y \ z)u\}((y \ z)t)$.*

whose corresponding Coq version is given by:

Lemma *swap_m_subst*: $\forall t \ u \ x \ y \ z, \text{swap } y \ z ([x := u]t) =_\alpha ([(\text{vswap } y \ z \ x) := (\text{swap } y \ z \ u)] (\text{swap } y \ z \ t))$.

Proof. The proof is by induction on the size of the term t . The interesting case is the abstraction, where we need to prove that $(y \ z)(\{x := u\}(\lambda_w.t_1)) =_\alpha \{((y \ z))x := (y \ z)u\}((y \ z)\lambda_w.t_1)$. On the left hand side, we can propagate the metasubstitution over the abstraction in the case that $x \neq w$ (the other is straightforward) and the new goal after the propagation of the swap over the abstraction is $\lambda_{(y \ z)w'}.(y \ z)(\{x := u\}(w \ w')t_1) =_\alpha \{((y \ z))x := (y \ z)u\}(\lambda_{(y \ z)w}.(y \ z)t_1)$, where $w' \notin fv(\lambda_w.t_1) \cup fv(u) \cup \{x\}$. Now we propagate the metasubstitution over the abstraction in the right hand side term. Since $x \neq w$, we get $((y \ z))x \neq ((y \ z))w$ and a renaming is necessary. After the renaming to a new name, say w'' , such that $w'' \notin fv(\lambda_{(y \ z)w}.(y \ z)t_1) \cup fv((y \ z)u) \cup \{((y \ z))x\}$, we get the following goal

$\lambda_{((y\ z))w'}.(y\ z)(\{x := u\}(w\ w')t_1) =_{\alpha} \lambda_{w''}.\{((y\ z))x := (y\ z)u\}((w''\ ((y\ z))w)((y\ z)t_1))$. We consider two cases: either $w'' = ((y\ z))w'$ or $w'' \neq ((y\ z))w'$. In the former case, we can apply the rule *aeq_abs_same* and we get $(y\ z)(\{x := u\}((w\ w')t_1)) =_{\alpha} \{((y\ z))x := (y\ z)u\}((w''\ ((y\ z))w)((y\ z)t_1))$ that can be proved by the induction hypothesis. When $w'' \neq ((y\ z))w'$, the application of the rule *aeq_abs_diff* generates the goal $(w''\ ((y\ z))w')(y\ z)(\{x := u\}(w\ w')t_1) =_{\alpha} \{((y\ z))x := (y\ z)u\}((w''\ ((y\ z))w)((y\ z)t_1))$. We can use the induction hypothesis to propagate the swap inside the metasubstitution, and then we get an α -equality with metasubstitution as main operation on both sides, and whose correspondent components are α -equivalent. In a more abstract way, we have to prove an α -equality of the form $\{x := u\}t =_{\alpha} \{x := u'\}t'$, where $t =_{\alpha} t'$ and $u =_{\alpha} u'$. The problem is that we cannot rewrite α -equalities inside metasubstitution unless we prove some special lemmas stating the compatibilities between them using the *Equations* library or something similar. Alternatively, if we decide to analyse the metasubstitution componentwise, *i.e.* as stated in a lemma similar to *aeq_m_subst_in_trial*, we get a circular proof problem because both *aeq_m_subst_in_trial* and *swap_m_subst* depend on each other to be proved. We will present a solution that do not use any additional library, but it adds the following axiom to the formalization:

Axiom *Eq_implies_equality*: $\forall s\ s': \text{atoms}, s [=] s' \rightarrow s = s'$.

This axiom transform a set equality into a syntactic equality. This will allow us to rewrite sets of atoms in a more flexible way. To show how it works, we will start proving the lemma *aeq_m_subst_in* without the need of the lemma *swap_m_subst*:

Lemma *aeq_m_subst_in*: $\forall t\ u\ u'\ x, u =_{\alpha} u' \rightarrow ([x := u]\ t) =_a ([x := u']\ t)$.

Proof. The proof is by induction on the size of the term t . The interesting case is the abstraction. We have by hypothesis that $u =_{\alpha} u'$ therefore both u and u' have the same set of free variables by lemma *aeq_fv_nom*. With the axiom *Eq_implies_equality*, we can replace the set $fv(u)$ by $fv(u')$, or vice-versa, in such a way that instead of generating two new names for the propagation of the metasubstitutions inside the abstractions, we need just one new name and there is no more the case where the binders of the abstractions were different names. The case of the explicit substitution is similar, and with this strategy we avoid the rules *aeq_abs_diff* and *aeq_sub_diff* that introduce swappings. \square

The next lemma, named *aeq_m_subst_out* will benefit the strategy used in the previous proof, but it is not straightforward.

Lemma *aeq_m_subst_out*: $\forall t\ t'\ u\ x, t =_{\alpha} t' \rightarrow ([x := u]\ t) =_a ([x := u]\ t')$.

Proof. The proof is by induction on the size of the term t . The interesting case is the abstraction. There are two cases based on the definition of the α -equivalence relation: either the binders have the same name or they are different. In the former case, we have to prove $([x := u]\ n_abs\ y\ t1) =_a ([x := u]\ n_abs\ y\ t2)$ assuming that $t1 =_{\alpha} t2$. In both sides of the α -equation, the metasubstitution need to be propagated over the abstraction, and according to our definition of metasubstitution, one name will be generated for each propagation. The new name to be generate for the term $[x := u]\ (n_abs\ y\ t1)$ (lhs) is such that it is not in the set $fv(\lambda_y.t_1) \cup fv(u) \cup \{x\}$, while the new name to be generated for the term $[x := u]\ (n_abs\ y\ t2)$ (rhs) is such that it is not in the set $fv(\lambda_y.t_2) \cup fv(u) \cup \{x\}$. Since $t1 =_{\alpha} t2$, by lemma *aeq_fv_nom* the set of free variables of $t1$ is equal to the set of free variables of $t2$, and therefore, we can generate just one new name for both propagations of the metasubstitution. If this new name is $x0$ then the new goal to be proved is $n_abs\ x0\ (subst_rec_fun\ (swap\ y\ x0\ t1)\ u\ x) =_a n_abs\ x0\ (subst_rec_fun\ (swap\ y\ x0\ t2)\ u\ x)$, which can be proved by the induction hypothesis. If $y = x$ then $x \neq y0$ and the metasubstitution $[x := u]$ has no effect on the LHS, but it can be propagated on the RHS, *i.e.* over the abstraction $(n_abs\ y0\ t2)$ but it also has no effect in $t2$ because x does not occur free in $t2$. If $y \neq x$ then the metasubstitution can

be propagated over the abstraction of the LHS, and similarly we compare x with $y0$ to see what happens in the RHS. When $y0 = x$ then the metasubstitution has no effect on the abstraction of the RHS. On the LHS the metasubstitution is propagated since $y \neq x$ but, as in the previous case, it has no effect in $t1$ because $y0$ does not occur free in $t1$. Now we have to prove that $[x := u](n_abs\ y\ t1) = a\ [x := u](n_abs\ y0\ t2)$, when $y \neq x$, $y0 \neq x$ and $n_abs\ y\ t1 = a\ n_abs\ y0\ t2$. Since the set of free variables of $n_abs\ y\ t1$ is equal to the set of free variables of $n_abs\ y0\ t2$, we can as in the previous case generate only one new name, say $x0$ that fulfill the condition to propagate the metasubstitution on both sides of the α -equation. More precisely, $x0 \notin fv(\lambda_y.t1) \cup fv(u) \cup \{x\}$, and the goal to be proved is $n_abs\ x0\ ([x := u](swap\ y\ x0\ t1)) = a\ n_abs\ x0\ ([x := u](swap\ y0\ x0\ t2))$. As just one new name was generated, there is no case where the binders of the abstractions are different. Remember that abstractions with different binders were the cause of the circularity problem in the proofs because the application of the rule *aeq-abs-diff* introduces a new swap that will be outside the metasubstitution in this case, whose solution would require a lemma of the form of *swap-m-subst*. Therefore, after an application of the rule *aeq-abs-same*, we conclude with the induction hypothesis. The explicit substitution operation is also interesting, but the proof strategy is similar to the one used in the abstraction case. \square

As a corollary, one can join the lemmas *aeq-m-subst-in* and *aeq-m-subst-out* as follows:

Corollary *aeq-m-subst-eq*: $\forall t\ t'\ u\ u'\ x, t = a\ t' \rightarrow u = a\ u' \rightarrow ([x := u]\ t) = a\ ([x := u']\ t')$.

Now, we show how to propagate a swap inside metasubstitutions using the decomposition of the metasubstitution provided by the corollary *aeq-m-subst-eq*.

Lemma *swap_subst_rec_fun*: $\forall x\ y\ z\ t\ u, swap\ x\ y\ (subst_rec_fun\ t\ u\ z) = a\ subst_rec_fun\ (swap\ x\ y\ t)\ (swap\ x\ y\ u)\ (vswap\ x\ y\ z)$.

Proof. Firstly, we compare x and y , since the case $x = y$ is trivial. The proof proceeds by induction on the size of the term t , assuming that $x \neq y$. The tricky cases are the abstraction and explicit substitution. In the abstraction case, i.e. when $t = \lambda_{y'}.t1$ then we must prove that $swap\ x\ y\ ([z := u](n_abs\ y'\ t1)) = a\ [(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ x\ y\ (n_abs\ y'\ t1))$, and the induction hypothesis states that a swap can be propagated inside a metasubstitution whose body is a term with the same size as $t1$. Firstly, we compare the variables y' and z to check whether we should propagate the metasubstitution inside the abstraction of the LHS. When $y' = z$ the metasubstitution is erased according to the definition (5) on both sides of the α -equation and we are done. When $y' \neq z$ then the metasubstitutions on both sides of the α -equation need to be propagated inside the corresponding abstractions. In order to do so, a new name need to be created. Note that in this case, it is not possible to create a unique name for both sides because the name of the LHS cannot belong to the set $fv(\lambda_{y'}.t1) \cup fv(u) \cup \{z\}$, while the name of the RHS cannot belong to the set $fv((x\ y)\lambda_{y'}.t1) \cup fv((x\ y)u) \cup \{(x\ y)z\}$. Let $x0$ be a new name that is not in the set $fv(\lambda_{y'}.t1) \cup fv(u) \cup \{z\}$, and $x1$ a new name that is not in the set $fv((x\ y)\lambda_{y'}.t1) \cup fv((x\ y)u) \cup \{(x\ y)z\}$. After renaming and propagating the metasubstitutions inside the abstractions, the current goal is $n_abs\ (vswap\ x\ y\ x0)\ (swap\ x\ y\ ([z := u](swap\ y'\ x0\ t1))) = a\ n_abs\ x1\ ([[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ (vswap\ x\ y\ y')\ x1\ (swap\ x\ y\ t1))])$. We proceed by comparing $x1$ with $(vswap\ x\ y\ x0)$. If $x1 = (vswap\ x\ y\ x0)$ then we use the induction hypothesis to propagate the swap inside the metasubstitution in the LHS and the current goal is $[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ x\ y\ (swap\ y'\ x0\ t1)) = a\ [(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ (vswap\ x\ y\ y')\ (vswap\ x\ y\ x0)\ (swap\ x\ y\ t1))$ that is proved by the swap equivariance lemma *swap-equivariance*. If $x1 \neq (vswap\ x\ y\ x0)$ then by the rule *aeq-abs-diff* we have to prove that the variable $vswap\ x\ y\ x0$ is not in the set of free variables of the term $[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ (vswap\ x\ y\ y')\ x1\ (swap\ x\ y\ t1))$ and that $swap\ x\ y\ ([z := u](swap\ y'\ x0\ t1)) = a\ swap\ x1\ (vswap\ x\ y\ x0)\ ([[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap$

$(vswap\ x\ y'\ y)\ x1\ (swap\ x\ y'\ t1)))$. The former condition is routine. The later condition is proved using the induction hypothesis twice to propagate the swaps inside the metasubstitutions on each side of the α -equality. This swap has no effect on the variable of the metasubstitution, therefore we can apply lemma *aeq-m_subst_eq* and each generated case is proved by routine manipulation of swaps. The case of the explicit substitution follows a similar strategy of the abstraction. The initial goal is to prove that $swap\ x\ y\ ([z := u](n_sub\ t1\ y'\ t2)) =_a [(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ x\ y\ (n_sub\ t1\ y'\ t2))$ and we start comparing the variables y' and z . When $y' = z$, the metasubstitution has no effect on the body of the metasubstitution but it can still be propagated to the term $t2$. Therefore, this case is proved using the induction hypothesis over $t2$. When $y' \neq z$, then the metasubstitutions are propagated on both sides of the α -equation. Analogously to the abstraction case, one new name for each propagation is necessary. Let $x0$ be a new name not in the set $fv([y' := t2]t1) \cup fv(u) \cup \{z\}$, and $x1$, a new name not in the set $fv([(x\ y)]y' := (x\ y)t2)(x\ y)t1) \cup fv((x\ y)u) \cup \{(x\ y)z\}$. After the propagation step, we have the goal $n_sub\ (swap\ x\ y\ ([z := u](swap\ y'\ x0\ t1)))\ (vswap\ x\ y\ x0)\ (swap\ x\ y\ ([z := u]t2)) =_a n_sub\ ([[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ (vswap\ x\ y\ y')\ x1\ (swap\ x\ y\ t1)))\ x1\ ([[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ x\ y\ t2))$. We proceed by comparing $x1$ and $(swap\ x\ y\ x0)$. If $x1 = vswap\ x\ y\ x0$ then after an application of the rule *aeq-sub_same*, each component of the explicit substitution is proved by the induction hypothesis. If $x1 \neq vswap\ x\ y\ x0$ then we apply the rule *aeq-sub_diff* to decompose the explicit substitution in its components. The second component is straightforward by the induction hypothesis. The first component follows the strategy used in the abstraction case. The current goal, obtained after the application of the rule *aeq-sub_diff* is $swap\ x\ y\ ([z := u](swap\ y'\ x0\ t1)) =_a swap\ x1\ (vswap\ x\ y\ x0)\ ([[(vswap\ x\ y\ z) := (swap\ x\ y\ u)](swap\ (vswap\ x\ y\ y')\ x1\ (swap\ x\ y\ t1)))$. The induction hypothesis is used twice to propagate the swap on both the LHS and RHS of the α -equality. This swap has no effect on the variable of the metasubstitution, therefore we can apply lemma *aeq-m_subst_eq* and each generated case is proved by routine manipulation of swaps. \square

The following lemmas state, respectively, what happens when the variable in the meta-substitution is equal or different from the one in the abstraction. When it is equal, the meta-substitution is irrelevant. When they are different, we take a new variable that does not occur freely in the substituted term in the meta-substitution nor in the abstraction and is not the variable in the meta-substitution, and the abstraction of this new variable using the meta-substitution of the swap of the former variable in the meta-substitution is alpha-equivalent to the original meta-substitution of the abstraction. The proofs were straightforward from the definition of the meta-substitution, each case being respectively each one in the definition.

Lemma *m_subst_abs_eq* : $\forall\ u\ x\ t, [x := u](n_abs\ x\ t) = n_abs\ x\ t$.

Lemma *m_subst_abs_neq* : $\forall\ t\ u\ x\ y\ z, x \neq y \rightarrow z \text{ 'notin' } fv_nom\ u \text{ 'union' } fv_nom\ (n_abs\ y\ t) \text{ 'union' } \{x\} \rightarrow [x := u](n_abs\ y\ t) =_a n_abs\ z\ ([x := u](swap\ y\ z\ t))$.

The following lemmas are the equivalent version of the two previous lemmas for the case of the explicit substitution. They state, respectively, what happens when the variable in the meta-substitution is equal or different from the one in the explicit substitution. When it is equal, the meta-substitution is irrelevant on $t1$, but it is applied to $e2$. When they are different, we take a new variable that does not occur freely in the substituted term in the meta-substitution nor in the substitution and is not the variable in the meta-substitution, and the explicit substitution of this new variable using the meta-substitution of the swap of the former variable in the meta-substitution in $e11$ and the application of the original meta-substitution in $e12$ is alpha-equivalent to the original meta-substitution of the explicit substitution. The proofs are straightforward from the definition of the meta-substitution, each case being respectively

each one in the definition.

Lemma $m_subst_sub_eq$: $\forall u x t1 t2, [x := u](n_sub t1 x t2) = n_sub t1 x ([x := u] t2)$.

Lemma $m_subst_sub_neq$: $\forall t1 t2 u x y z, x \neq y \rightarrow z \text{ 'notin' } fv_nom u \text{ 'union' } fv_nom (n_sub t1 y t2) \text{ 'union' } \{\{x\}\} \rightarrow [x := u](n_sub t1 y t2) = a n_sub ([x := u](swap y z t1)) z ([x := u] t2)$.

4 The substitution lemma for the metasubstitution

In the pure λ -calculus, the substitution lemma is probably the first non trivial property. In our framework, we have defined two different substitution operation, namely, the metasubstitution denoted by $[x:=u]t$ and the explicit substitution that has n_sub as a constructor. In what follows, we present the main steps of our proof of the substitution lemma for the metasubstitution operation:

Lemma m_subst_lemma : $\forall e1 e2 x e3 y, x \neq y \rightarrow x \text{ 'notin' } (fv_nom e3) \rightarrow ([y := e3]([x := e2]e1)) = a ([x := ([y := e3]e2)]([y := e3]e1))$.

Proof. The proof is by induction on the size of the term $e1$. The interesting cases are the abstraction and the explicit substitution. In the abstraction case, the initial goal is $([y := e3]([x := e2] n_abs z e1)) = a ([x := [y := e3] e2]([y := e3] n_abs z e1))$, assuming that $x \neq y$ and $x \text{ 'notin' } fv_nom e3$. We start comparing z with x aiming to apply the definition of the metasubstitution on the LHS of the goal. When $z = x$, the subterm $[x := e2](n_abs x e1)$ reduces to $(n_abs x e1)$ and then the LHS reduces to $([y := e3] n_abs x e1)$. The RHS reduces to the same term because x does not occur free neither in $(n_abs x e1)$ nor in $e3$. When $y = z$ then the subterm $([y := e3] n_abs z e1)$ reduces to $(n_abs z e1)$. On the LHS, we propagate the internal metasubstitution over the abstraction taking a fresh name for the binder. The external metasubstitution cannot be propagated inside the abstraction because they have the same binder. The resulting terms are α -equivalent, and although the strategy is similar to the one used in the lemmas $aeq_m_subst_in$, $aeq_m_subst_out$ and $swap_subst_rec_fun$ the proof requires much more steps. In the explicit substitution case, we used the same approach used in the abstraction for the left side and the same as the application for the right side of the substitution. It consisted of comparing the variable in the meta substitution and the one in the substitution. We used the auxiliary lemmas on the equality and inequality of the meta-substitution applied to explicit substitutions and it was necessary to create new variables in each use of the inequality. This is due to the attempt of removing the explicit substitution from inside the meta-substitution. When tis removal was made, the proof consisted in proving a similar case for the abstraction in the left side of the explicit substitution and the one similar to the application was used for the right part of it. \square

5 Conclusion and Future work

In this work, we presented a formalization of the substitution lemma in a framework that extends the λ -calculus with an explicit substitution operator. Calculi with explicit substitutions are important frameworks to study properties of the λ -calculus and have been extensively studied in the last decades[1, 2, 3, 4, 10, 14].

The formalization is modular in the sense that the explicit substitution operator is generic and could be instantiated with any calculi with explicit substitutions in a nominal setting. The main contribution of this work, besides the formalization itself, is the solution to a circular proof problem. Several auxiliary (minor) results were not included in this document, but they are numerous and can be found in the

source file of the formalization that is available in a GitHub repository from the following url https://github.com/flaviodemoura/lx_confl/tree/m_subst_lemma.

As future work, we plan to integrate this formalization with another one related to the Z property [13] to prove properties of calculi with explicit substitutions such as confluence [22, 23, 18]. In

References

- [1] M. Abadi, L. Cardelli, P.-L. Curien & J.-J. Lévy (1991): *Explicit Substitutions*. *Journal of Functional Programming* 1(4), pp. 375–416.
- [2] B. Accattoli (2012): *An Abstract Factorization Theorem for Explicit Substitutions*. In: *RTA*, pp. 6–21.
- [3] M. Ayala-Rincón, F.C. de Moura & F. Kamareddine (2002): *Comparing Calculi of Explicit Substitutions with Eta-reduction*. In R.J.G.B. de Queiroz, editor: *Proceedings Ninth Workshop on Logic, Language, Information and Computation (WoLLIC 2002)*, ENTCS 67, Elsevier Science Publishers, pp. 77–96.
- [4] M. Ayala-Rincón, F. L. C. de Moura & F. Kamareddine (2005): *Comparing and Implementing Calculi of Explicit Substitutions with Eta-Reduction*. *Annals of Pure and Applied Logic* 134, pp. 5–41.
- [5] H. P. Barendregt (1984): *The Lambda Calculus : Its Syntax and Semantics (Revised Edition)*. North Holland.
- [6] Z.-el-A. Benaissa, D. Briaud, P. Lescanne & J. Rouyer-Degli (1996): *λ_{epsilon} , a Calculus of Explicit Substitutions Which Preserves Strong Normalization*. *JFP* 6(5), pp. 699–722.
- [7] Stefan Berghofer & Christian Urban (2007): *A Head-to-Head Comparison of de Bruijn Indices and Names*. *Electronic Notes in Theoretical Computer Science* 174(5), pp. 53–67, doi:[10.1016/j.entcs.2007.01.018](https://doi.org/10.1016/j.entcs.2007.01.018).
- [8] R. Bloo & K. Rose (1995): *Preservation of Strong Normalisation in Named Lambda Calculi with Explicit Substitution and Garbage Collection*. In: *CSN-95: COMPUTER SCIENCE IN THE NETHERLANDS*, pp. 62–72.
- [9] Roel Bloo & Herman Geuvers (1999): *Explicit Substitution: On the Edge of Strong Normalization*. *Theoretical Computer Science* 211(1-2), pp. 375–395, doi:[10.1016/s0304-3975\(97\)00183-7](https://doi.org/10.1016/s0304-3975(97)00183-7).
- [10] E. Bonelli (2001): *Perpetuality in a Named Lambda Calculus With Explicit Substitutions*. *Mathematical Structures in Computer Science* 11(1), pp. 47–90, doi:[10.1017/s0960129500003248](https://doi.org/10.1017/s0960129500003248).
- [11] Pierre-Louis Curien, Thérèse Hardin & Jean-Jacques Lévy (1996): *Confluence Properties of Weak and Strong Calculi of Explicit Substitutions*. *Journal of the ACM* 43(2), pp. 362–397, doi:[10.1145/226643.226675](https://doi.org/10.1145/226643.226675).
- [12] R. David & B. Guillaume (2001): *A Lambda-Calculus with Explicit Weakening and Explicit Substitution*. *Mathematical Structures in Computer Science* 11(1), pp. 169–206.
- [13] Flávio L. C. de Moura & Leandro O. Rezende (2021): *A Formalization of the (Compositional) z Property*. In: *Fifth Workshop on Formal Mathematics for Mathematicians*.
- [14] Ken-etsu Fujita & Koji Nakazawa (2016): *Church-Rosser Theorem and Compositional Z-Property*.
- [15] M. Gabbay & A. Pitts (1999): *A New Approach to Abstract Syntax Involving Binders*. In: *14th Symposium on Logic in Computer Science (LICS'99)*, IEEE, Washington - Brussels - Tokyo, pp. 214–224.
- [16] B. Guillaume (2000): *The λ_{Se} -Calculus Does Not Preserve Strong Normalization*. *J. of Func. Programming* 10(4), pp. 321–325.
- [17] Fairouz Kamareddine & Alejandro Ríos (1997): *Extending a Lambda-Calculus with Explicit Substitution Which Preserves Strong Normalisation Into a Confluent Calculus on Open Terms*. *J. Funct. Program.* 7(4), pp. 395–420.
- [18] D. Kesner (2008): *Perpetuality for Full and Safe Composition (in a Constructive Setting)*. In: *Automata, Languages and Programming, 35th International Colloquium, ICALP 2008, Reykjavik, Iceland, July 7-11, 2008, Proceedings, Part II - Track B: Logic, Semantics, and Theory of Programming & Track C: Security and Cryptography Foundations*, pp. 311–322, doi:[10.1007/978-3-540-70583-3_26](https://doi.org/10.1007/978-3-540-70583-3_26).

- [19] D. Kesner (2009): *A Theory of Explicit Substitutions with Safe and Full Composition*. *Logical Methods in Computer Science* 5(3:1), pp. 1–29.
- [20] P.-A. Mellès (1995): *Typed λ -Calculi with Explicit Substitutions May Not Terminate* in *Proceedings of TLCA'95*. *LNCS* 902.
- [21] C. A. Muñoz (1996): *Confluence and Preservation of Strong Normalisation in an Explicit Substitutions Calculus*. In: *Proceedings, 11th Annual IEEE Symposium on Logic in Computer Science*, New Brunswick, New Jersey, USA, July 27-30, 1996, pp. 440–447, doi:[10.1109/LICS.1996.561460](https://doi.org/10.1109/LICS.1996.561460).
- [22] Koji Nakazawa & Ken-etsu Fujita (2016): *Compositional Z: Confluence Proofs for Permutative Conversion*. *Studia Logica* 104(6), pp. 1205–1224, doi:[10.1007/s11225-016-9673-0](https://doi.org/10.1007/s11225-016-9673-0).
- [23] Koji Nakazawa & Ken-etsu Fujita (2017): *Z for Call-by-Value*. In: *6th International Workshop on Confluence (IWC 2017)*, pp. 57–61.
- [24] Koji Nakazawa, Ken-etsu Fujita & Yuta Imagawa (2023): *Z Property for the Shuffling Calculus*. *Mathematical Structures in Computer Science*, pp. 1–13, doi:[10.1017/S0960129522000408](https://doi.org/10.1017/S0960129522000408).
- [25] The Coq Development Team (2021): *The Coq Proof Assistant*. Zenodo, doi:[10.5281/ZENODO.5704840](https://doi.org/10.5281/ZENODO.5704840).