GEOGLAM Period	Approx. Maize Stage	Waterlogging Duration	Expected Yield Loss	Confidence
Planting → Vegetative	V2–V3	3–4 days	10–25%	High
Planting → Vegetative	V2–V3	5–6 days	25–35%	High
Planting → Vegetative	V2–V3	≥7 days	35–60%	Moderate
Vegetative → Harvest	V6 / Jointing	4–6 days	20–30%	High
Vegetative → Harvest	Tasseling / Silking	≥7 days	up to ~58%	Moderate
Late Season	10VT / Early Grain Fill	≥7 days	<20%	Moderate

Ren (2014) found V3 was the most sensitive stage and observed losses around 15–35% under 3–6 days of waterlogging.

Zhang (2023) found major physiological stress begins around 4–6 days.

Based on the literature, severe maize losses are approximately 35–60% depending on growth stage. The highest documented losses in the papers collected so far are around 58% during tasseling under prolonged waterlogging, while early vegetative stages typically show 25–35%

SWI > P95 for ≥ 7 days → Severe yield-loss risk (~35–60%)

Consecutive days with SWI > P95	Interpretation	Literature-based expected impact
0–3 days	Minor / short-term waterlogging	Limited to moderate impacts; damage starts but is generally less severe than longer events. [renr580pgp...weebly.com]
4–6 days	High waterlogging stress	Major physiological damage threshold; yield losses commonly around **20–35%** depending on stage and variety. [frontiersin.org], [renr580pgp...weebly.com]

≥7 days	Severe waterlogging	Prolonged stress associated with substantial yield loss; approximately **30–60%** depending on growth stage. The highest documented value in your papers is about **58% loss at tasseling** after 7 days of waterlogging. [geo.libretexts.org], [renr580pgp...weebly.com]

