

DTE 2025

Combining Finite Element Methods and Neural Networks to Solve Elliptic Problems on 2D Geometries

Hélène Barucq¹, Michel Duprez², Florian Faucher¹, Emmanuel Franck³,
Frédérique Lecourtier², Vanessa Lleras⁴, Victor Michel-Dansac³ and
Nicolas Victorion¹

¹Project-Team Makutu, Inria, TotalEnergies, Pau, France


²Project-Team MIMESIS, Inria, Strasbourg, France

³Project-Team MACARON, Inria, Strasbourg, France

⁴IMAG, University of Montpellier, Montpellier, France

February 20, 2025

Scientific context



Context : Create real-time digital twins of an organ (e.g. liver).

Objective : Develop an hybrid finite element / neural network method.
accurate quick + parameterized

Parametric elliptic convection/diffusion PDE : For one or several $\mu \in \mathcal{M}$, find $u : \Omega \rightarrow \mathbb{R}$ such that


$$\mathcal{L}(u; \mathbf{x}, \mu) = f(\mathbf{x}, \mu), \quad (\mathcal{P})$$

where \mathcal{L} is the parametric differential operator defined by

$$\mathcal{L}(\cdot; \mathbf{x}, \mu) : u \mapsto R(\mathbf{x}, \mu)u + C(\mu) \cdot \nabla u - \frac{1}{\text{Pe}} \nabla \cdot (D(\mathbf{x}, \mu) \nabla u),$$

and some Dirichlet, Neumann or Robin BC (which can also depend on μ).

Pipeline of the Enriched FEM



Remark : The PINN prediction enriched Finite element approximation spaces.

Physics-Informed Neural Networks

Standard PINNs¹ (Weak BC) : Find the optimal weights θ^* , such that

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \left(\omega_r J_r(\theta) + \omega_b J_b(\theta) \right), \quad (\mathcal{P}_\theta)$$

with

residual loss

$$J_r(\theta) = \int_{\mathcal{M}} \int_{\Omega} \left| \mathcal{L}(u_\theta(\mathbf{x}, \boldsymbol{\mu}); \mathbf{x}, \boldsymbol{\mu}) - f(\mathbf{x}, \boldsymbol{\mu}) \right|^2 d\mathbf{x} d\boldsymbol{\mu},$$

boundary loss

$$J_b(\theta) = \int_{\mathcal{M}} \int_{\partial\Omega} \left| u_\theta(\mathbf{x}, \boldsymbol{\mu}) - g(\mathbf{x}, \boldsymbol{\mu}) \right|^2 d\mathbf{x} d\boldsymbol{\mu},$$

where u_θ is a neural network, $g = 0$ is the Dirichlet BC.

In (\mathcal{P}_θ) , ω_r and ω_b are some weights.

Monte-Carlo method : Discretize the cost functions by random process.

¹[Raissi et al., 2019]

Physics-Informed Neural Networks

Improved PINNs¹ (Strong BC) : Find the optimal weights θ^* such that

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \left(\omega_r J_r(\theta) + \cancel{\omega_b J_b(\theta)} \right),$$

with $\omega_r = 1$ and

residual loss


$$J_r(\theta) = \int_{\mathcal{M}} \int_{\Omega} |\mathcal{L}(u_{\theta}(\mathbf{x}, \mu); \mathbf{x}, \mu) - f(\mathbf{x}, \mu)|^2 d\mathbf{x} d\mu,$$

where u_{θ} is a neural network defined by

$$u_{\theta}(\mathbf{x}, \mu) = \varphi(\mathbf{x}) w_{\theta}(\mathbf{x}, \mu) + g(\mathbf{x}, \mu),$$

with φ a level-set function, w_{θ} a NN and $g = 0$ the Dirichlet BC.

Thus, the Dirichlet BC is imposed exactly in the PINN : $u_{\theta} = g$ on $\partial\Omega$.



¹[Lagaris et al., 1998; Franck et al., 2024]

Finite Element Method¹

Variational Problem :

$$\text{Find } u_h \in V_h^0 \text{ such that, } \forall v_h \in V_h^0, a(u_h, v_h) = l(v_h), \quad (\mathcal{P}_h)$$

with h the characteristic mesh size, a and l the bilinear and linear forms given by

$$a(u_h, v_h) = \frac{1}{\text{Pe}} \int_{\Omega} D \nabla u_h \cdot \nabla v_h + \int_{\Omega} R u_h v_h + \int_{\Omega} v_h C \cdot \nabla u_h, \quad l(v_h) = \int_{\Omega} f v_h,$$

and V_h^0 the finite element space defined by

$$V_h^0 = \{v_h \in C^0(\Omega), \forall K \in \mathcal{T}_h, v_h|_K \in \mathbb{P}_k, v_h|_{\partial\Omega} = 0\},$$


where \mathbb{P}_k is the space of polynomials of degree at most k .

Linear system : Let $(\phi_1, \dots, \phi_{N_h})$ a basis of V_h^0 .

$$\text{Find } U \in \mathbb{R}^{N_h} \text{ such that} \quad AU = b$$

with

$$A = (a(\phi_i, \phi_j))_{1 \leq i, j \leq N_h} \quad \text{and} \quad b = (l(\phi_j))_{1 \leq j \leq N_h}.$$



$$\mathcal{T}_h = \{K_1, \dots, K_{N_e}\}$$

(N_e : number of elements)

¹[Ern and Guermond, 2004]

How improve PINN prediction with FEM ?

Additive approach

Variational Problem : Let $u_\theta \in H^{k+1}(\Omega) \cap H_0^1(\Omega)$.

Find $p_h^+ \in V_h^0$ such that, $\forall v_h \in V_h^0, a(p_h^+, v_h) = l(v_h) - a(u_\theta, v_h)$, (\mathcal{P}_h^+)


with the **enriched trial space** V_h^+ defined by

$$V_h^+ = \{u_h^+ = u_\theta + p_h^+, \quad p_h^+ \in V_h^0\}.$$

General Dirichlet BC : If $u = g$ on $\partial\Omega$, then

$$p_h^+ = g - u_\theta \quad \text{on } \partial\Omega,$$

with u_θ the PINN prior.



Convergence analysis

Theorem 1: Convergence analysis of the standard FEM [Ern and Guermond, 2004]

We denote $u_h \in V_h$ the solution of (\mathcal{P}_h) with V_h the standard trial space. Then,

$$|u - u_h|_{H^1} \leq C_{H^1} h^k |u|_{H^{k+1}},$$

$$\|u - u_h\|_{L^2} \leq C_{L^2} h^{k+1} |u|_{H^{k+1}}.$$

Theorem 2: Convergence analysis of the enriched FEM [Lecourtier et al., 2025]

We denote $u_h^+ \in V_h^+$ the solution of (\mathcal{P}_h^+) with V_h^+ the enriched trial space. Then,

$$|u - u_h^+|_{H^1} \leq \frac{|u - u_\theta|_{H^{k+1}}}{|u|_{H^{k+1}}} (C_{H^1} h^k |u|_{H^{k+1}}),$$

$$\|u - u_h^+\|_{L^2} \leq \frac{|u - u_\theta|_{H^{k+1}}}{|u|_{H^{k+1}}} (C_{L^2} h^{k+1} |u|_{H^{k+1}}).$$

Numerical results

2D Poisson problem on Square - Dirichlet BC

2D Anisotropic Elliptic problem on a Square - Dirichlet BC

2D Poisson problem on Annulus - Mixed BC

Numerical results

2D Poisson problem on Square - Dirichlet BC

2D Anisotropic Elliptic problem on a Square - Dirichlet BC

2D Poisson problem on Annulus - Mixed BC

Problem considered

Problem statement: We consider the Poisson problem in 2D with homogeneous Dirichlet boundary conditions:

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \times \mathcal{M}, \\ u = 0, & \text{on } \partial\Omega \times \mathcal{M}, \end{cases}$$

with $\Omega = [-0.5\pi, 0.5\pi]^2$ and $\mathcal{M} = [-0.5, 0.5]^2$ ($p = 2$ parameters).

We define the right-hand side f such that the solution is given by

$$u(\mathbf{x}, \boldsymbol{\mu}) = \exp\left(-\frac{(x - \mu_1)^2 + (y - \mu_2)^2}{2}\right) \sin(2x) \sin(2y),$$

with $\mathbf{x} = (x, y) \in \Omega$ and some parameters $\boldsymbol{\mu} = (\mu_1, \mu_2) \in \mathcal{M}$.

PINN training: MLP of 5 layers, trained with an LBFGs optimizer (5000 epochs). Imposing the Dirichlet BC exactly in the PINN with the levelset φ defined by


$$\varphi(\mathbf{x}) = (x + 0.5\pi)(x - 0.5\pi)(y + 0.5\pi)(y - 0.5\pi).$$

Training time : less than 1 hour on a laptop GPU.

Numerical results

Error estimates : 1 given parameter.


$$\mu^{(1)} = (0.05, 0.22)$$



Numerical results

Error estimates : 1 given parameter.

$$\mu^{(1)} = (0.05, 0.22)$$



Gains achieved : $n_p = 50$ parameters.

$$\mathcal{S} = \left\{ \mu^{(1)}, \dots, \mu^{(n_p)} \right\}$$

**Gains in L^2 rel error
of our method w.r.t. FEM**

k	min	max	mean
1	134.32	377.36	269.39
2	67.02	164.65	134.85
3	39.52	72.65	61.55

$N = 20$


$$\text{Gain} : \|u - u_h\|_{L^2} / \|u - u_h^+\|_{L^2}$$

Cartesian mesh : N^2 nodes.

Numerical results

Error estimates : 1 given parameter.

$$\mu^{(1)} = (0.05, 0.22)$$



Numerical costs of the two approaches :

N required to reach the same error e .

k	e	N	
		FEM	Add
1	$1 \cdot 10^{-3}$	119	8
	$1 \cdot 10^{-4}$	379	24
2	$1 \cdot 10^{-4}$	42	8
	$1 \cdot 10^{-5}$	89	17
3	$1 \cdot 10^{-5}$	28	10
	$1 \cdot 10^{-6}$	48	18

Numerical results

2D Poisson problem on Square - Dirichlet BC

2D Anisotropic Elliptic problem on a Square - Dirichlet BC

2D Poisson problem on Annulus - Mixed BC

Problem considered

Problem statement: We consider the Poisson problem in 2D with mixed BC:

$$\begin{cases} -\operatorname{div}(D\nabla u) = f, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

with $\Omega = [0, 1]^2$ and $\mathcal{M} = [0.4, 0.6] \times [0.4, 0.6] \times [0.01, 1] \times [0.1, 0.8]$ ($p = 4$).

We define the right-hand side f by

$$f(\mathbf{x}, \boldsymbol{\mu}) = \exp\left(-\frac{(x - \mu_1)^2 + (y - \mu_2)^2}{0.025\sigma^2}\right).$$

with $\mathbf{x} = (x, y) \in \Omega$ and some parameters $\boldsymbol{\mu} = (\mu_1, \mu_2, \epsilon, \sigma) \in \mathcal{M}$.

The diffusion matrix D (symmetric and positive definite) is given by

$$D(\mathbf{x}, \boldsymbol{\mu}) = \begin{pmatrix} \epsilon x^2 + y^2 & (\epsilon - 1)xy \\ (\epsilon - 1)xy & x^2 + \epsilon y^2 \end{pmatrix}.$$


PINN training: MLP with Fourier Features¹ of 5 layers, trained with an Adam optimizer (15000 epochs). Imposing the Dirichlet BC exactly in the PINN with a levelset function.

¹[Tancik et al., 2020]

Numerical results

Error estimates : 1 given parameter.


$$\mu^{(1)} = (0.51, 0.54, 0.52, 0.55)$$



Numerical results

Error estimates : 1 given parameter.

$$\mu^{(1)} = (0.51, 0.54, 0.52, 0.55)$$



Gains achieved : $n_p = 50$ parameters.

$$\mathcal{S} = \left\{ \mu^{(1)}, \dots, \mu^{(n_p)} \right\}$$

**Gains in L^2 rel error
of our method w.r.t. FEM**

k	min	max	mean
1	7.12	82.57	35.67
2	3.54	35.88	18.32
3	1.33	26.51	8.32

$N = 20$

$$\text{Gain} : \|u - u_h\|_{L^2} / \|u - u_h^+\|_{L^2}$$

Cartesian mesh : N^2 nodes.

Numerical results

2D Poisson problem on Square - Dirichlet BC

2D Anisotropic Elliptic problem on a Square - Dirichlet BC

2D Poisson problem on Annulus - Mixed BC

Problem considered

Problem statement: We consider the Poisson problem in 2D with mixed BC:

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \times \mathcal{M}, \\ u = g, & \text{on } \Gamma_E \times \mathcal{M}, \\ \frac{\partial u}{\partial n} + u = g_R, & \text{on } \Gamma_I \times \mathcal{M}, \end{cases}$$

with $\Omega = \{(x, y) \in \mathbb{R}^2, 0.25 \leq x^2 + y^2 \leq 1\}$ and $\mathcal{M} = [2.4, 2.6]$ ($p = 1$).

We define the right-hand side f such that the solution is given by

$$u(\mathbf{x}; \boldsymbol{\mu}) = 1 - \frac{\ln(\mu_1 \sqrt{x^2 + y^2})}{\ln(4)},$$

with $\mathbf{x} = (x, y) \in \Omega$ and some parameters $\boldsymbol{\mu} = \mu_1 \in \mathcal{M}$. The BC are given by

$$g(\mathbf{x}; \boldsymbol{\mu}) = 1 - \frac{\ln(\mu_1)}{\ln(4)} \quad \text{and} \quad g_R(\mathbf{x}; \boldsymbol{\mu}) = 2 + \frac{4 - \ln(\mu_1)}{\ln(4)}.$$


PINN training: MLP of 5 layers, trained with an LBFGs optimizer (4000 epochs). Imposing the mixed BC exactly in the PINN¹.

¹[Sukumar and Srivastava, 2022]

Numerical results

Error estimates : 1 given parameter.


$$\mu^{(1)} = \mu_1 = 2.51$$



Numerical results

Error estimates : 1 given parameter.

$$\mu^{(1)} = \mu_1 = 2.51$$



Gains achieved : $n_p = 50$ parameters.

$$\mathcal{S} = \left\{ \mu^{(1)}, \dots, \mu^{(n_p)} \right\}$$

**Gains in L^2 rel error
of our method w.r.t. FEM**

k	min	max	mean
1	15.12	137.72	55.5
2	31	77.46	58.41
3	18.72	21.49	20.6

$$h = 1.33 \cdot 10^{-1}$$

$$\text{Gain : } \|u - u_h\|_{L^2} / \|u - u_h^+\|_{L^2}$$

Conclusion

Conclusion and Perspectives

- PINNs are good candidates for the enriched approach.
- Numerical validation of the theoretical results.
- The enriched approach provides the same results as the standard FEM method, but with coarser meshes. \Rightarrow Reduction of the computational cost.

Perspectives :

- Validate the additive approach on more complex geometry.
- Consider non-linear problems.
- Use the PINN prediction to build an optimal mesh, via a posteriori error estimates.

Add QR code with the paper + Image of the bean testcase

References

- A. Ern and J.-L. Guermond. *Theory and Practice of Finite Elements*. Springer New York, 2004. doi: [10.1007/978-1-4757-4355-5](https://doi.org/10.1007/978-1-4757-4355-5).
- E. Franck, V. Michel-Dansac, and L. Navoret. Approximately well-balanced Discontinuous Galerkin methods using bases enriched with Physics-Informed Neural Networks. *J. Comput. Phys.*, 512:113144, 2024. ISSN 0021-9991. doi: [10.1016/j.jcp.2024.113144](https://doi.org/10.1016/j.jcp.2024.113144).
- I. E. Lagaris, A. Likas, and D. I. Fotiadis. Artificial neural networks for solving ordinary and partial differential equations. *IEEE Trans. Neural Netw.*, 9(5):987–1000, 1998. ISSN 1045-9227. doi: [10.1109/72.712178](https://doi.org/10.1109/72.712178).
- F. Lecourtier, H. Barucq, M. Duprez, F. Faucher, E. Franck, V. Lleras, V. Michel-Dansac, and N. Victorion. Enriching continuous lagrange finite element approximation spaces using neural networks, 2025.
- M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *J. Comput. Phys.*, 378:686–707, 2019. doi: [10.1016/j.jcp.2018.10.045](https://doi.org/10.1016/j.jcp.2018.10.045).
- N. Sukumar and A. Srivastava. Exact imposition of boundary conditions with distance functions in physics-informed deep neural networks. *Comput. Method. Appl. M.*, 389:114333, 2022. ISSN 0045-7825. doi: [10.1016/j.cma.2021.114333](https://doi.org/10.1016/j.cma.2021.114333).
- M. Tancik, P. Srinivasan, and al. Fourier Features Let Networks Learn High Frequency Functions in Low Dimensional Domains. In *Advances in Neural Information Processing Systems*, volume 33, pages 7537–7547. Curran Associates, Inc., 2020.