

Enriching continuous Lagrange finite element approximation spaces using neural networks

Michel Duprez¹, Emmanuel Franck², **Frédérique Lecourtier¹** and Vanessa Lleras³

¹Project-Team MIMESIS, Inria, Strasbourg, France

²Project-Team MACARON, Inria, Strasbourg, France

³IMAG, University of Montpellier, Montpellier, France


Joint work with:

H. Barucq, F. Faucher, N. Victorion and V. Michel-Dansac.


PhD student seminar

IMAG

INSTITUT MONTPELLIERAIN
ALEXANDER GROTHENDIECK



Scientific context



Context : Create real-time digital twins of an organ (e.g. liver).

Objective : Develop an hybrid finite element / neural network method.
accurate quick + parameterized

Parametric elliptic convection/diffusion PDE : For one or several $\mu \in \mathcal{M}$, find $u : \Omega \rightarrow \mathbb{R}$ such that

$$\mathcal{L}(u; \mathbf{x}, \mu) = f(\mathbf{x}, \mu), \quad (\mathcal{P})$$

where $\mathcal{L}$ is the parametric differential operator defined by

$$\mathcal{L}(\cdot; \mathbf{x}, \mu) : u \mapsto R(\mathbf{x}, \mu)u + C(\mu) \cdot \nabla u - \frac{1}{\text{Pe}} \nabla \cdot (D(\mathbf{x}, \mu) \nabla u),$$


and some Dirichlet, Neumann or Robin BC (which can also depend on μ).

Pipeline of the Enriched FEM

OFFLINE : PINN training

Inputs

$\mathbf{x} \in \Omega$: space coordinates
 $\boldsymbol{\mu} \in \mathcal{M}$: parameter



Output

$u_\theta(\mathbf{x}, \boldsymbol{\mu})$: prediction of
 the PDE solution $u(\mathbf{x}, \boldsymbol{\mu})$

ONLINE (fixed $\boldsymbol{\mu}$) : PINN evaluation + Enriched FEM resolution

PINN evaluation

$u_\theta(\cdot, \boldsymbol{\mu})$: prediction for
 the given parameter $\boldsymbol{\mu}$



FEM solver
 enriched by u_θ



Output

enriched FEM solution
 (depending on the mesh size h)

Complete ONLINE process : quick + accurate

LECOURTIER Frédérique

Classical approaches

Physics-Informed Neural Networks

Standard PINNs¹ (Weak BC) : Find the optimal weights θ^* , such that

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \left(\omega_r J_r(\theta) + \omega_b J_b(\theta) \right), \quad (\mathcal{P}_\theta)$$

with

residual loss

$$J_r(\theta) = \int_{\mathcal{M}} \int_{\Omega} \left| \mathcal{L}(u_\theta(\mathbf{x}, \boldsymbol{\mu}); \mathbf{x}, \boldsymbol{\mu}) - f(\mathbf{x}, \boldsymbol{\mu}) \right|^2 d\mathbf{x} d\boldsymbol{\mu},$$

boundary loss

$$J_b(\theta) = \int_{\mathcal{M}} \int_{\partial\Omega} \left| u_\theta(\mathbf{x}, \boldsymbol{\mu}) - g(\mathbf{x}, \boldsymbol{\mu}) \right|^2 d\mathbf{x} d\boldsymbol{\mu},$$

where u_θ is a neural network, $g = 0$ is the Dirichlet BC.

In $(\mathcal{P}_\theta)$, ω_r and ω_b are some weights.

Monte-Carlo method : Discretize the cost functions by random process.

¹[Raissi et al., 2019]

Physics-Informed Neural Networks

Improved PINNs¹ (Strong BC) : Find the optimal weights θ^* such that

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \left(\omega_r J_r(\theta) + \cancel{\omega_b J_b(\theta)} \right),$$

with $\omega_r = 1$ and

residual loss

$$J_r(\theta) = \int_{\mathcal{M}} \int_{\Omega} |\mathcal{L}(u_{\theta}(\mathbf{x}, \boldsymbol{\mu}); \mathbf{x}, \boldsymbol{\mu}) - f(\mathbf{x}, \boldsymbol{\mu})|^2 d\mathbf{x} d\boldsymbol{\mu},$$


$$\partial\Omega = \{\varphi = 0\}$$

where u_{θ} is a neural network defined by

$$u_{\theta}(\mathbf{x}, \boldsymbol{\mu}) = \varphi(\mathbf{x}) w_{\theta}(\mathbf{x}, \boldsymbol{\mu}) + g(\mathbf{x}, \boldsymbol{\mu}),$$

with φ a level-set function, w_{θ} a NN and $g = 0$ the Dirichlet BC.

Thus, the Dirichlet BC is imposed exactly in the PINN : $u_{\theta} = g$ on $\partial\Omega$.



¹[Lagaris et al., 1998; Franck et al., 2024]

Finite Element Method¹

Variational Problem :

$$\text{Find } u_h \in V_h^0 \text{ such that, } \forall v_h \in V_h^0, a(u_h, v_h) = l(v_h), \quad (\mathcal{P}_h)$$

with h the characteristic mesh size, a and l the bilinear and linear forms given by

$$a(u_h, v_h) = \frac{1}{\text{Pe}} \int_{\Omega} D \nabla u_h \cdot \nabla v_h + \int_{\Omega} R u_h v_h + \int_{\Omega} v_h C \cdot \nabla u_h, \quad l(v_h) = \int_{\Omega} f v_h,$$

and V_h^0 the finite element space defined by

$$V_h^0 = \{v_h \in C^0(\Omega), \forall K \in \mathcal{T}_h, v_h|_K \in \mathbb{P}_k, v_h|_{\partial\Omega} = 0\},$$


where $\mathbb{P}_k$ is the space of polynomials of degree at most k .

Linear system : Let $(\phi_1, \dots, \phi_{N_h})$ a basis of V_h^0 .

$$\text{Find } U \in \mathbb{R}^{N_h} \text{ such that} \quad AU = b$$

with

$$A = (a(\phi_i, \phi_j))_{1 \leq i, j \leq N_h} \quad \text{and} \quad b = (l(\phi_j))_{1 \leq j \leq N_h}.$$



$$\mathcal{T}_h = \{K_1, \dots, K_{N_e}\}$$

(N_e : number of elements)

¹[Ern and Guermond, 2004]

Enriched finite element method using PINN

Additive approach

Variational Problem : Let $u_\theta \in H^{k+1}(\Omega) \cap H_0^1(\Omega)$ be a PINN prior.

$$\text{Find } p_h^+ \in V_h^0 \text{ such that, } \forall v_h \in V_h^0, a(p_h^+, v_h) = l(v_h) - a(u_\theta, v_h), \quad (\mathcal{P}_h^+)$$

with the **enriched trial space** V_h^+ defined by


$$V_h^+ = \{u_h^+ = u_\theta + p_h^+, \quad p_h^+ \in V_h^0\}.$$

General Dirichlet BC : If $u = g$ on $\partial\Omega$, then

$$p_h^+ = g - u_\theta \quad \text{on } \partial\Omega,$$

with u_θ the PINN prior.

We hope that the modified problem will give the same results as the standard one on coarser meshes.



Convergence analysis

Theorem 1: Convergence analysis of the standard FEM [Ern and Guermond, 2004]

We denote $u_h \in V_h$ the solution of $(\mathcal{P}_h)$ with V_h the standard trial space. Then,

$$|u - u_h|_{H^1} \leq C_{H^1} h^k |u|_{H^{k+1}},$$

$$\|u - u_h\|_{L^2} \leq C_{L^2} h^{k+1} |u|_{H^{k+1}}.$$

Theorem 2: Convergence analysis of the enriched FEM [Lecourtier et al., 2025]

We denote $u_h^+ \in V_h^+$ the solution of $(\mathcal{P}_h^+)$ with V_h^+ the enriched trial space. Then,

$$|u - u_h^+|_{H^1} \leq \frac{|u - u_\theta|_{H^{k+1}}}{|u|_{H^{k+1}}} (C_{H^1} h^k |u|_{H^{k+1}}),$$

$$\|u - u_h^+\|_{L^2} \leq \frac{|u - u_\theta|_{H^{k+1}}}{|u|_{H^{k+1}}} (C_{L^2} h^{k+1} |u|_{H^{k+1}}).$$

Gains of the additive approach.

Numerical results

2D Poisson problem on Square - Dirichlet BC

3D Poisson problem on Cube - Dirichlet BC

2D Anisotropic Elliptic problem on Square - Dirichlet BC

Numerical results

2D Poisson problem on Square - Dirichlet BC

3D Poisson problem on Cube - Dirichlet BC

2D Anisotropic Elliptic problem on Square - Dirichlet BC

Problem considered

Problem statement: Consider the Poisson problem in 2D with Dirichlet BC:

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \times \mathcal{M}, \\ u = 0, & \text{on } \partial\Omega \times \mathcal{M}, \end{cases}$$

with $\Omega = [-0.5\pi, 0.5\pi]^2$ and $\mathcal{M} = [-0.5, 0.5]^2$ ($p = 2$ parameters).

Analytical solution :

$$u(\mathbf{x}, \boldsymbol{\mu}) = \exp\left(-\frac{(x - \mu_1)^2 + (y - \mu_2)^2}{2}\right) \sin(2x) \sin(2y).$$

PINN training:

MLP 5 layers (sine - 40, 60, 60, 60, 40); LBFGs optimizer (5000 epochs - 6000 col. pts).


Imposing the Dirichlet BC exactly in the PINN with the levelset φ defined by

$$\varphi(\mathbf{x}) = (x + 0.5\pi)(x - 0.5\pi)(y + 0.5\pi)(y - 0.5\pi).$$

Online phase - 1 set of parameters

$$\mu^{(1)} = (0.05, 0.22)$$

Error estimates in L^2 norm:




Online phase - 1 set of parameters

$$\mu^{(1)} = (0.05, 0.22)$$

Error estimates in L^2 norm:

N_{dofs} required to reach the same error e :



k	e	N_{dofs}	
		FEM	Add
1	$1 \cdot 10^{-3}$	14 161	64
	$1 \cdot 10^{-4}$	143 641	576
2	$1 \cdot 10^{-4}$	6 889	225
	$1 \cdot 10^{-5}$	31 329	1 089
3	$1 \cdot 10^{-5}$	6 724	784
	$1 \cdot 10^{-6}$	20 164	2 704

Parametric framework

Considering the set of $n_p = 50$ parameter instances

$$\mathcal{S} = \left\{ \boldsymbol{\mu}^{(1)}, \dots, \boldsymbol{\mu}^{(n_p)} \right\}.$$

Gains achieved: $\|u - u_h\|_{L^2} / \|u - u_h^+\|_{L^2}$

Gains in L^2 rel error of our method w.r.t. FEM			
k	min	max	mean
1	134.32	377.36	269.39
2	67.02	164.65	134.85
3	39.52	72.65	61.55

Cartesian mesh : N^2 nodes with $N = 20$.

Numerical results

2D Poisson problem on Square - Dirichlet BC

3D Poisson problem on Cube - Dirichlet BC

2D Anisotropic Elliptic problem on Square - Dirichlet BC

Problem considered

Problem statement: Consider the Poisson problem in 3D with Dirichlet BC:

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \times \mathcal{M}, \\ u = 0, & \text{on } \partial\Omega \times \mathcal{M}, \end{cases}$$

with $\Omega = [-0.5\pi, 0.5\pi]^3$ and $\mathcal{M} = [-0.5, 0.5]^3$ ($p = 3$ parameters).

Analytical solution :

$$u(\mathbf{x}, \boldsymbol{\mu}) = \exp\left(-\frac{(x - \mu_1)^2 + (y - \mu_2)^2 + (z - \mu_3)^2}{2}\right) \sin(2x) \sin(2y) \sin(2z).$$

PINN training:

MLP 5 layers (tanh - 40, 60, 60, 60, 40); LBFGs optimizer (5000 epochs - 40000 col. pts).

Imposing the Dirichlet BC exactly in the PINN with the levelset φ defined by


$$\varphi(\mathbf{x}) = (x + 0.5\pi)(x - 0.5\pi)(y + 0.5\pi)(y - 0.5\pi)(z + 0.5\pi)(z - 0.5\pi).$$

Training time : 12 minutes on NVIDIA RTX 2000 (8GB VRAM).

Online phase - 1 set of parameters I

$$\mu^{(1)} = (0.05, 0.22, 0.1)$$


Error estimates in L^2 norm:




Online phase - 1 set of parameters I

$$\mu^{(1)} = (0.05, 0.22, 0.1)$$

Error estimates in L^2 norm:



L^2 error w.r.t. execution time*:




* mesh + assembly + solve
 + prior derivatives evaluation
 + training time


Online phase - 1 set of parameters I

$$\mu^{(1)} = (0.05, 0.22, 0.1)$$

Error estimates in L^2 norm:



L^2 error w.r.t. execution time*:



* mesh + assembly + solve
+ prior derivatives evaluation
+ training time

Online phase - 1 set of parameters II

$$\mu^{(1)} = (0.05, 0.22, 0.1)$$

N_{dofs} and computation time required to reach the same error e :

e	N		N_{dofs}		Computation time*	
	FEM	Add	FEM	Add	FEM	Add
$1 \cdot 10^{-3}$	152	12	$3.51 \cdot 10^6$	$1.73 \cdot 10^3$	$(7.65 \cdot 10^1)$	$5.2 \cdot 10^{-2}$
$1 \cdot 10^{-4}$	484	39	$1.13 \cdot 10^8$	$5.93 \cdot 10^4$	$(2.8 \cdot 10^3)$	$9.26 \cdot 10^{-1}$
$1 \cdot 10^{-5}$	1539	122	$3.65 \cdot 10^9$	$1.82 \cdot 10^6$	$(1.02 \cdot 10^5)$	$(5.26 \cdot 10^1)$


— $\div 1471$ —→

Cartesian mesh : N^3 nodes; $\mathbb{P}_1$ elements.

* The results in brackets are extrapolations.

Parametric framework

How many parameters n_p are needed to balance training?



Online: assembly + solve + prior derivatives evaluation

Offline: mesh + training time

Numerical results

2D Poisson problem on Square - Dirichlet BC

3D Poisson problem on Cube - Dirichlet BC

2D Anisotropic Elliptic problem on Square - Dirichlet BC

Problem considered

Problem statement: Considering an Anisotropic Elliptic problem with Dirichlet BC:

$$\begin{cases} -\operatorname{div}(D\nabla u) = f, & \text{in } \Omega \times \mathcal{M}, \\ u = 0, & \text{on } \partial\Omega \times \mathcal{M}, \end{cases}$$

with $\Omega = [0, 1]^2$ and $\mathcal{M} = [0.4, 0.6]^2 \times [0.01, 1] \times [0.1, 0.8]$ ($p = 4$ parameters).

Right-hand side and diffusion matrix: $\mu = (\mu_1, \mu_2, \sigma, \epsilon) \in \mathcal{M}$

$$f(\mathbf{x}, \mu) = \exp\left(-\frac{(x - \mu_1)^2 + (y - \mu_2)^2}{0.025\sigma^2}\right).$$

$$D(\mathbf{x}, \mu) = \begin{pmatrix} \epsilon x^2 + y^2 & (\epsilon - 1)xy \\ (\epsilon - 1)xy & x^2 + \epsilon y^2 \end{pmatrix}.$$

PINN training: MLP with Fourier Features¹ of 5 layers (tanh - 40, 60, 60, 60, 40).

LBFGs optimizer (15000 epochs - 8000 col. pts).


Imposing the Dirichlet BC exactly in the PINN with the a level-set function.

¹[Tancik et al., 2020]

Online phase

Error estimates : 1 set of parameters.


$$\mu^{(1)} = (0.51, 0.54, 0.52, 0.55)$$



Online phase

Error estimates : 1 set of parameters.

$$\mu^{(1)} = (0.51, 0.54, 0.52, 0.55)$$



Gains achieved : $\|u - u_h\|_{L^2} / \|u - u_h^+\|_{L^2}$

Considering $n_p = 50$ parameter instances


$$\mathcal{S} = \{\mu^{(1)}, \dots, \mu^{(n_p)}\}.$$

**Gains in L^2 rel error
of our method w.r.t. FEM**

k	min	max	mean
1	7.12	82.57	35.67
2	3.54	35.88	18.32
3	1.33	26.51	8.32

Cartesian mesh : N^2 nodes with $N = 20$.

Numerical solutions and errors



$$\mu^{(2)} = (0.46, 0.52, 0.05, 0.12)$$

Conclusion

TODO

Preprint (linear)



References

- A. Ern and J.-L. Guermond. Theory and Practice of Finite Elements. Springer New York, 2004.
- E. Franck, V. Michel-Dansac, and L. Navoret. Approximately well-balanced Discontinuous Galerkin methods using bases enriched with Physics-Informed Neural Networks. J. Comput. Phys., 512:113144, 2024. ISSN 0021-9991.
- I. E. Lagaris, A. Likas, and D. I. Fotiadis. Artificial neural networks for solving ordinary and partial differential equations. IEEE Trans. Neural Netw., 9(5):987–1000, 1998. ISSN 1045-9227.
- F. Lecourtier, H. Barucq, M. Duprez, F. Faucher, E. Franck, V. Lleras, V. Michel-Dansac, and N. Victorion. Enriching continuous lagrange finite element approximation spaces using neural networks, 2025.
- M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. J. Comput. Phys., 378:686–707, 2019.
- M. Tancik, P. Srinivasan, and al. Fourier Features Let Networks Learn High Frequency Functions in Low Dimensional Domains. In Advances in Neural Information Processing Systems, volume 33, pages 7537–7547. Curran Associates, Inc., 2020.

Preprint (linear)





Appendix 1 : Sec1



A1 – Appendix 1

TODO